


Strand Invasion of Mixed-Sequence, Double-Helical B-DNA by γ -Peptide Nucleic Acids Containing G-Clamp Nucleobases under Physiological Conditions

Srinivas Rapireddy, Raman Bahal, and Danith H. Ly*

Department of Chemistry and Center for Nucleic Acids Science and Technology (CNAST), Carnegie Mellon University, 4400 Fifth Avenue, Pittsburgh, Pennsylvania 15213, United States

ABSTRACT: Peptide nucleic acids (PNAs) make up the only class of nucleic acid mimics developed to date that has been shown to be capable of invading double-helical B-form DNA. Recently, we showed that sequence limitation associated with PNA recognition can be relaxed by utilizing conformationally preorganized γ -peptide nucleic acids (γ PNAs). However, like all the previous studies, with the exception of triplex binding, DNA strand invasion was performed at relatively low salt concentrations. When physiological ionic strengths were used, little to no binding was observed. On the basis of this finding, it was not clear whether the lack of binding is due to the lack of base pair opening or the lack of binding free energy, either of which would result in no productive binding. In this work, we show that it is the latter. Under simulated physiological conditions, the DNA double helix is sufficiently dynamic to permit strand invasion by the designer oligonucleotide molecules provided that the required binding free energy can be met. This finding has important implications for the design oligonucleotides for recognition of B-DNA via direct Watson–Crick base pairing.



Double-helical DNA has traditionally been considered as a relatively static molecular entity, specifically designed for the storage and safeguard of genetic information.¹ Early theoretical estimations² and nuclear magnetic resonance (NMR) imino proton exchange studies^{3,4} placed the probability of internal base pair opening right around 10^{-5} , the open base pair lifetime around 10^{-7} s, and the base pair activation energy in the range of 43–65 kJ/mol at ambient temperature. These initial estimates cast a grim outlook on the prospect of being able to access the Watson–Crick (W–C) hydrogen bond donors and acceptors of nucleobases for recognition. As a consequence, efforts to establish sequence-specific interactions with double-stranded DNA (dsDNA) in the past have generally been focused on chemical groups that reside in the minor⁵ and major^{6,7} grooves, in large part, because of the ease of accessibility and the precedence for their recognition in nature. However, over the past two decades, peptide nucleic acids (PNAs),⁸ a particular class of nucleic acid mimics comprising a pseudopeptide backbone (Chart 1a, top), have been shown to be capable of invading dsDNA.^{8–10} This finding was significant because, contrary to the traditional belief, it demonstrates that W–C base pairing interactions can be established with intact dsDNA at physiological temperature. Though they are promising as antigene reagents, because of the generality and specificity of recognition, the originally designed achiral PNAs can recognize only homopurine¹¹ and homopyrimidine targets.¹² Mixed-sequence PNAs can invade topologically constrained supercoiled plasmid DNA^{13,14} and structurally perturbed regions of genomic DNA;¹⁵ however, they are unable to invade linear, double-helical B-form DNA (B-DNA).

Recently, two approaches, tail clamp^{16,17} and double-duplex invasion,¹⁸ have been developed, allowing mixed-sequence PNAs to invade B-DNA. However, they are not without limitations. The first approach still requires a homopyrimidine tract for anchoring the triplex binding, while the second approach, although more relaxed in sequence selection, is complicated by the need to use two separate strands of pseudocomplementary PNA to invade dsDNA. More recently, we showed that sequence restriction can be relaxed by utilizing conformationally preorganized γ PNAs as molecular reagents (Chart 1a, bottom).¹⁹ However, like all the other studies, with the exception of triplex binding by homopyrimidine PNAs,²⁰ strand invasion of linear double-helical B-DNA was performed in buffers containing relatively low salt concentrations.^{11,18,19,21,22} When physiological ionic strengths were used, little to no binding was observed.²³ On the basis of this result, it was not clear whether the lack of binding is due to the lack of base pair opening (accessibility) or the lack of binding free energy, because the kinetics of base pair breathing and the thermodynamic stability of the DNA duplex are tightly coupled. Knowing the answer to this question is important because it will tell us whether W–C recognition is a viable option for targeting mixed-sequence B-DNA under physiological conditions. If the former is true, there is little that can be done from a design standpoint. However, if the latter is true, this issue could be resolved through molecular design. Herein, we show that it is the latter. Under simulated

Received: March 9, 2011

Revised: April 7, 2011

Published: April 11, 2011




Figure 1. Result of a gel shift assay following incubation of a 171 bp DNA fragment at 0.2 μ M containing (a) a perfectly matched (PM, lanes 1–5) and single-base mismatched (MM, lanes 6 and 7) binding site with the indicated γ PNA oligomers at 1.0 μ M and (b) a PM binding site with the indicated concentrations of the PNA1-3X oligomer in a simulated physiological buffer at 37 °C for 16 h, followed by separation on nondenaturing PAGE and staining with SYBR-Gold.

separated on 10% urea–denaturing polyacrylamide gels. The cleavage patterns were visualized by autoradiography.

DEPC Chemical Probing Assay. The 5′-tcactataggcgcaattgga-3′/5′-gtactcaagctgtgtccgcctagtgcaca-3′ PCR primer set was used to PCR-amplify the DNA targets containing PM and MM binding sites from the corresponding plasmid vectors. The PCR products were digested with *Hind*III and 3′-labeled with P-32 under conditions identical to those described above. The invasion complex was prepared the same way as that for DNase-I footprinting. DEPC treatment was initiated via addition of a 10% volume of DEPC to the invasion samples. The mixtures were incubated at room temperature for 5 min, followed by precipitation and piperidine treatment. The samples were then separated on 10% urea–denaturing polyacrylamide gels, and the cleavage patterns were visualized by autoradiography.

RESULTS AND DISCUSSION

To address the question of accessibility versus thermodynamic stability, we synthesized a series of decameric γ PNA oligomers containing the same nucleobase sequence content but with a different number of cytosine (C) to G-clamp (X) nucleobase substitutions (Chart 1b,c). We then assessed their DNA strand invasion capability using a combination of gel shift, enzymatic footprinting, and chemical probing assays. The C \rightarrow X nucleobase substitutions were made in an attempt to systematically improve the binding free energy of γ PNAs without resorting to changing the nucleobase sequence or chain length. This particular cytosine analogue was chosen because prior studies showed that C \rightarrow X substitution can significantly enhance the thermodynamic stability of the PNA–DNA duplex.²⁷ Given the already strong binding affinity of γ PNAs,²² making such nucleobase substitutions should significantly further enhance their thermodynamic stability. Implementing such a design would allow us to survey the invasion capability of γ PNAs over a range of binding




Figure 2. Effects of biological buffers on DNA strand invasion by PNA1-3X. A gel shift assay was performed by incubating 0.2 μ M DNA containing a PM binding site with 1.0 μ M PNA1-3X in the indicated buffers at 37 °C for 16 h. Simulated: 10 mM NaP_i, 2 mM MgCl₂, and 150 mM KCl. NEB1 (New England Biolabs buffer 1, 1 \times): 10 mM Bis Tris propane-HCl, 10 mM MgCl₂, and 1 mM dithiothreitol. NEB2: 10 mM Tris-HCl, 50 mM NaCl, 10 mM MgCl₂, and 1 mM dithiothreitol. NEB3: 10 mM Tris-HCl, 100 mM NaCl, 10 mM MgCl₂, and 1 mM dithiothreitol. NEB4: 20 mM Tris-OAc, 50 mM KOAc, 10 mM MgCl₂, and 1 mM dithiothreitol. Replication: 10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl₂, and 0.1% Triton X-100. Transcription: 40 mM Tris-HCl, 6 mM MgCl₂, 10 mM dithiothreitol, and 2 mM spermidine.

free energies. Our working hypothesis is that if DNA strand invasion under simulated physiological conditions is predominated by thermodynamics, we would expect a gradual increase in the invasion efficiency with binding free energy, or in this case the number of C \rightarrow X nucleobase substitutions. On the other hand, if base pair accessibility is the determining factor in DNA strand invasion, we do not expect the correlation described above but instead expect a steep dependence on temperature, which has a direct effect on the rate of base pair opening.

A gel shift assay was performed by incubating a 171 bp, linear doubled-stranded B-DNA containing an internal, perfectly matched (PM) binding site (Chart 1d) with various γ PNA oligomers in a buffer containing physiological salt concentrations²⁸ [10 mM sodium phosphate, 2 mM MgCl₂, and 150 mM KCl (pH 7.4)] at 37 °C. The mixtures were separated on nondenaturing polyacrylamide gels and stained with SYBR-Gold for visualization. Inspection of Figure 1a reveals no binding for PNA1 or PNA1-1X, but a small amount for PNA1-2X, as evidenced by the formation of a shifted band following a 16 h incubation. This result is consistent with an earlier finding.²³ With a further increase in the number of C \rightarrow X nucleobase substitutions from two to three, we noticed a dramatic increase in the intensity of the shifted band, from \sim 3% with PNA1-2X (lane 4) to \sim 80% with PNA1-3X (lane 5), indicating that more of the bound complex was formed. Formation of this complex was sequence-specific, because no trace of the shifted band was observed with DNA containing an inverted, single-base mismatched (MM) binding site (compare lane 7 to lane 6). Nucleobase inversion is expected to have a minimal impact on the thermodynamic stability (or base pair breathing rate) of the DNA double helix. Thus, the difference in binding efficiency is predominantly due to sequence mismatch discrimination. We attribute the high level of sequence specificity to the competitive binding of the native cDNA strand. Our attempt to drive the reaction to completion by increasing the concentration of PNA1-3X was unsuccessful because beyond a 5:1 PNA1-3X:DNA ratio, the DNA bands gradually disappeared from the gel (in Figure 1b, compare lanes 6 and 7 to lane 5). This is probably due to aggregation and nonspecific binding of




Figure 3. (a) DNase-I footprinting and (b) DEPC chemical probing assay following incubation with 10000 cpm of the 3'-labeled DNA target containing a PM (lanes 1–4) and MM (lanes 5 and 6) binding site with the indicated concentrations of PNA1-3X in a simulated physiological buffer at 37 °C for 16 h. In panel a, the 3'-end of the target strand was labeled with P-32, and in panel b, the homologous strand was labeled with P-32.

PNA1-3X to DNA as the result of the charge-neutral polyamide backbone and hydrophobic character of the methyl group at the γ -backbone. In addition to the simulated physiological buffer, we have examined a number of other biologically relevant buffers, including those employed in restriction digestion, transcription, and replication to assess the generality of PNA1-3X binding. In all of these buffers, some of which contained concentrations of $MgCl_2$ as high as 10 mM, binding efficiencies of >60% were achieved within a 16 h incubation period (Figure 2). Longer incubation times did not produce additional binding, indicating that equilibrium had been reached. Taken together, these results show that PNA1-3X can invade dsDNA under physiologically relevant conditions. Strand invasion is unique to γ PNA, because no evidence of binding for the unmodified PNA containing the same nucleobase sequence as PNA1-3X has been found.

To confirm that binding of PNA1-3X to DNA occurred at the expected site and through a strand invasion mechanism, we performed DNase footprinting and DEPC (diethyl pyrocarbonate) chemical probing assays. DNase is an endonuclease enzyme that indiscriminately cleaves dsDNA. Selective binding of PNA1-3X to dsDNA is expected to block DNase digestion, resulting in a footprinting pattern following electrophoretic separation and detection of the DNA targets. DEPC, on the other hand, is a chemical reagent known to react selectively with adenine and to a lesser extent with the guanine nucleobase of single-stranded or perturbed regions of double-stranded DNA.²⁹ Strand invasion of DNA by PNA1-3X is expected to result in local displacement of the homologous DNA strand. The looped-out strand can be revealed in the form of strand cleavage following DEPC and subsequent piperidine treatments.




Figure 4. Effects of temperature and incubation time on DNA strand invasion by PNA1-3X. The fraction bound was determined by gel shift assays following incubation of 0.2 μ M DNA containing a PM binding site with 1.0 μ M PNA1-3X in a simulated physiological buffer at the indicated temperatures and time points. $t_{1/2}$ is defined as the time it took to reach 50% binding. The inset shows a plot of $-\ln(1 - C)$ vs incubation time; the data show that strand invasion of B-DNA by PNA1-3X at 37 $^{\circ}$ C obeys pseudo-first-order kinetics: $C = 1 - \exp(-k_{ps}t)$, where C is the fraction bound at time t and k_{ps} is the pseudo-first-order rate constant ($k_{ps,37\text{ }^{\circ}\text{C}} = 0.0075\text{ min}^{-1}$).

Consistent with these predictions, we observed a distinct footprinting pattern on the target strand, localized at the expected binding site following incubation of DNA with PNA1-3X (Figure 3a). The extent of the footprint became more pronounced as the PNA1-3X concentrations increased, and it was observed only with DNA containing a PM binding site (lanes 1–4). Likewise, we observed cleavage of the cDNA strand directly across from the binding site (Figure 3b). Strand cleavage occurred only with DNA containing a PM binding site (lanes 1–4). These results are consistent with PNA1-3X binding selectively to its target site through a strand invasion mechanism.

Next, we assessed the effects of temperature and incubation time on the efficiency of DNA strand invasion. Our result shows that the efficiency of strand invasion is strongly dependent on temperature in the range of 22–37 $^{\circ}$ C (Figure 4). This is not surprising because temperature has a direct effect on the rate of base pair opening. Strand invasion was extremely inefficient at ambient temperature, in which case <10% invasion took place after incubation for 24 h. The rate of strand invasion dramatically increased at 37 $^{\circ}$ C, reaching equilibrium after incubation for \sim 6 h. Further increases in temperature resulted in further increases in the rate of strand invasion, but the difference in the rates at 37 and 50 $^{\circ}$ C is not as dramatic as compared to the difference in the rates at 22 and 37 $^{\circ}$ C ($t_{1/2,22\text{ }^{\circ}\text{C}} \sim 170\text{ h}$; $t_{1/2,37\text{ }^{\circ}\text{C}} \sim 2\text{ h}$; $t_{1/2,45\text{ }^{\circ}\text{C}} \sim 30\text{ min}$). There is an only 4-fold difference in the $t_{1/2}$ in the higher-temperature regimes as compared to 135-fold in the lower-temperature regimes. Regardless of whether the samples were incubated at 37 or 50 $^{\circ}$ C, the invasion profiles reached a plateau corresponding to \sim 90% binding efficiency. This result indicates that at 37 $^{\circ}$ C, the DNA double helix is sufficiently dynamic to permit strand invasion by γ PNAs. However, whether successful strand invasion takes place

depends on whether γ PNA has the required binding free energy to displace the native cDNA strand.

CONCLUSION

In summary, we have shown that PNA1-3X can invade double-helical B-DNA under physiologically relevant conditions. Improvements in thermodynamic stability significantly enhance the efficiency of DNA strand invasion. We have already demonstrated how additional binding free energy may be achieved, through backbone preorganization and nucleobase substitution in this study, covalent attachment of a DNA intercalating agent, such as acridine, as demonstrated in prior studies,^{20,22} and extension of the length of oligonucleotides.¹⁹ Besides cytosine, other nucleobases, including thymine, could also be modified to improve their hydrogen bonding and base stacking capabilities. Important work in this area by Nielsen and co-workers^{30,31} has already begun in earnest. Replacement of natural nucleobases with synthetic analogues capable of forming additional hydrogen bonding interactions and larger and geometrically aligned hydrophobic cores would further expand the recognition repertoire of PNA and γ PNA. Issues with aggregation and nonspecific binding, as observed at high PNA and γ PNA concentrations, could be alleviated by installing a hydrophilic group at the γ -backbone.²⁴ We have already embarked on this front, the result of which will be reported in due course.

γ PNAs are attractive, as compared to other classes of antigene reagents developed to date,^{5,32–36} because they are relatively easy to synthesize and they hybridize to their targets (via strand invasion) in a highly sequence-specific and predictable manner in accordance with the Watson–Crick base pairing rules. Molecules that can recognize and bind double-stranded DNA in a sequence-specific manner are of considerable interest in biology, biotechnology, and medicine. They could be used as molecular tools to probe sequence information, manipulate a genome's structure and function, and regulate gene expression at the transcriptional level, which may be more effective than the antisense approach because it shuts down gene expression at the initial step, as well as potential therapeutic and diagnostic reagents for the treatment and detection of genetic diseases.

AUTHOR INFORMATION

Corresponding Author

*E-mail: dly@andrew.cmu.edu. Telephone: (412) 268-4010. Fax: (412) 268-5579.

Funding Sources

This work was supported by the National Institutes of Health (GM076251) and the National Science Foundation (CHE-1012467).

ACKNOWLEDGMENT

We thank Prof. B. A. Armitage, M. Ly, and J. Deems for reading the manuscript and for providing insightful comments and suggestions.

ABBREVIATIONS

PNA, peptide nucleic acid; γ PNA, γ -backbone-modified peptide nucleic acid; A, adenine; C, cytosine; G, guanine; T, thymine; X,

G-clamp; TFA, trifluoroacetic acid; TFMSA, trifluoromethanesulfonic acid; DEPC, diethyl pyrocarbonate.

REFERENCES

- (1) Frank-Kamenetskii, M. D. (1987) How the double helix breathes. *Nature* 328, 17–18.
- (2) Frank-Kamenetskii, M. D. (1985) Structure and Motion: Membranes, Nucleic Acids and Proteins. In *Proceedings of the International Symposium on the Structure, Dynamics, and Membranes of Nucleic Acids and Proteins* (Clementi, E., Corongiu, G., Sarma, M. H., and Sarma, R. H., Eds.) pp 417–432, Adenine Press, Rome.
- (3) Gueron, M., Kochoyan, M., and Leroy, J.-L. (1987) A single mode of DNA base-pair opening drives imino proton exchange. *Nature* 328, 89–92.
- (4) Leroy, J. L., Kochoyan, M., Huynh-Dinh, T., and Gueron, M. (1988) Characterization of base-pair opening in deoxynucleotide duplexes using catalyzed exchange of the imino proton. *J. Mol. Biol.* 200, 223–238.
- (5) Dervan, P. B. (2001) Molecular recognition of DNA by small molecules. *Bioorg. Med. Chem.* 9, 2215–2235.
- (6) Thuong, N. T., and Helene, C. (1993) Sequence-specific recognition and modification of double-helical DNA by oligonucleotides. *Angew. Chem., Int. Ed.* 32, 666–690.
- (7) Greisman, H. A., and Pabo, C. O. (1997) A general strategy for selecting high-affinity zinc finger proteins for diverse DNA target sites. *Science* 275, 657–661.
- (8) Nielsen, P. E., Egholm, M., Berg, R. H., and Buchardt, O. (1991) Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science* 254, 1497–1500.
- (9) Nielsen, P. E. (1999) Peptide nucleic acid. A molecule with two identities. *Acc. Chem. Res.* 32, 624–630.
- (10) Ishihara, T., and Corey, D. R. (1999) Rules for strand invasion by chemically modified oligonucleotides. *J. Am. Chem. Soc.* 121, 2012–2020.
- (11) Nielsen, P. E., and Christensen, L. (1996) Strand displacement binding of a duplex-forming homopurine PNA to a homopyrimidine duplex DNA target. *J. Am. Chem. Soc.* 118, 2287–2288.
- (12) Egholm, M., Nielsen, P. E., Buchardt, O., and Berg, R. H. (1992) Recognition of guanine and adenine in DNA by cytosine and thymine containing peptide nucleic acids (PNA). *J. Am. Chem. Soc.* 114, 9677–9678.
- (13) Bentin, T., and Nielsen, P. E. (1996) Enhanced peptide nucleic acid binding to supercoiled DNA: Possible implications for DNA “breathing” dynamics. *Biochemistry* 35, 8863–8869.
- (14) Zhang, X., Ishihara, T., and Corey, D. R. (2000) Strand invasion by mixed base PNAs and a PNA-peptide chimera. *Nucleic Acids Res.* 28, 3332–3338.
- (15) Janowski, B. A., Kaihatsu, K., Huffman, K. E., Schwartz, J. C., Ram, R., Hardy, D., Mendelson, C. R., and Corey, D. R. (2005) Inhibiting transcription of chromosomal DNA with antigene peptide nucleic acids. *Nat. Chem. Biol.* 1, 210–215.
- (16) Bentin, T., Larsen, H. J., and Nielsen, P. E. (2003) Combined triplex/duplex invasion of double-stranded DNA by “tail-clamp” peptide nucleic acid. *Biochemistry* 42, 13987–13995.
- (17) Kaihatsu, K., Shah, R. H., Zhao, X., and Corey, D. R. (2003) Extending recognition by peptide nucleic acids (PNAs): Binding to duplex DNA and inhibition of transcription by tail-clamp PNA-peptide conjugate. *Biochemistry* 42, 13996–14003.
- (18) Lohse, J., Dahl, O., and Nielsen, P. E. (1999) Double duplex invasion by peptide nucleic acid: A general principle for sequence-specific targeting of double-stranded DNA. *Proc. Natl. Acad. Sci. U.S.A.* 96, 11804–11808.
- (19) He, G., Rapireddy, S., Bahal, R., Sahu, B., and Ly, D. H. (2009) Strand invasion of extended, mixed sequence B-DNA by γ PNAs. *J. Am. Chem. Soc.* 131, 12088–12090.
- (20) Bentin, T., and Nielsen, P. E. (2003) Superior duplex DNA strand invasion by acridine conjugated peptide nucleic acids. *J. Am. Chem. Soc.* 125, 6378–6379.
- (21) Smolina, I. V., Demidov, V. V., Soldatenkov, V. A., Chasovskikh, S. G., and Frank-Kamenetskii, M. D. (2005) End invasion of peptide nucleic acids (PNAs) with mixed-sequence composition into linear DNA duplexes. *Nucleic Acids Res.* 33, e146.
- (22) Rapireddy, S., He, G., Roy, S., Armitage, B. A., and Ly, D. H. (2007) Strand invasion of mixed-sequence B-DNA by acridine-linked, γ -peptide nucleic acid (γ -PNA). *J. Am. Chem. Soc.* 129, 15596–15600.
- (23) Chenna, V., Rapireddy, S., Sahu, B., Ausin, C., Pedrosa, E., and Ly, D. H. (2008) A simple cytosine-to-G-clamp nucleobase substitution enables chiral γ -PNAs to invade mixed-sequence double helical B-form DNA. *ChemBioChem* 9, 2388–2391.
- (24) Dragulescu-Andrasi, A., Rapireddy, S., Frezza, B. M., Gayathri, C., Gil, R. R., and Ly, D. H. (2006) A simple γ -backbone modification preorganizes peptide nucleic acid into a helical structure. *J. Am. Chem. Soc.* 128, 10258–10267.
- (25) Christensen, L., Fitzpatrick, R., Gildea, B., Petersen, K. H., Hansen, H. F., Koch, T., Egholm, M., Buchardt, O., Nielsen, P. E., Coull, J., and Berg, R. H. (1995) Solid-phase synthesis of peptide nucleic acids. *J. Pept. Sci.* 1, 175–183.
- (26) Brummelkamp, T. R., Bernards, R., and Agami, R. (2002) A system for stable expression of short interfering RNAs in mammalian cells. *Science* 296, 550–553.
- (27) Ortega, J.-A., Blas, J. R., Orozco, M., Grandas, A., Pedrosa, E., and Robles, J. (2007) Binding affinities of oligonucleotides and PNAs containing phenoxazine and G-clamp cytosine analogues are unusually sequence dependent. *Org. Lett.* 9, 4503–4506.
- (28) Santoro, S. W., and Joyce, G. F. (1997) A general purpose RNA-cleaving DNA enzyme. *Proc. Natl. Acad. Sci. U.S.A.* 94, 4262–4266.
- (29) Furlong, J. C., and Lilley, D. M. J. (1986) Highly selective chemical modification of cruciform loops by diethyl pyrocarbonate. *Nucleic Acids Res.* 14, 3995–4007.
- (30) Eldrup, A. B., Nielsen, P. E., Haaima, G., Rasmussen, H., Kastrop, J. S., Christensen, C., and Nielsen, P. E. (2001) 1,8-Naphthyridin-2(1H)-ones: Novel bicyclic and tricyclic analogues of thymine in peptide nucleic acids (PNAs). *Eur. J. Org. Chem.*, 1781–1790.
- (31) Eldrup, A. B., Christensen, C., Haaima, G., and Nielsen, P. E. (2002) Substituted 1,8-naphthyridin-2(1H)-ones are superior to thymine in the recognition of adenine in duplex as well as triplex structure. *J. Am. Chem. Soc.* 124, 3254–3262.
- (32) Nielsen, P. E. (1991) Sequence-selective DNA recognition by synthetic ligands. *Bioconjugate Chem.* 2, 1–12.
- (33) Sun, J. S., and Helene, C. (1993) Oligonucleotide-directed triple-helix formation. *Curr. Opin. Struct. Biol.* 3, 345–356.
- (34) Beerli, R. R., and Barbas, C. F. I. (2002) Engineering polydactyl zinc-finger transcription factors. *Nat. Biotechnol.* 20, 135–141.
- (35) Jantz, D., Amann, B. T., Gatto, G. J., and Berg, J. M. (2004) The design of functional DNA-binding proteins based on zinc finger domains. *Chem. Rev.* 104, 789–800.
- (36) Jamieson, A. C., Miller, J. C., and Pabo, C. O. (2003) Drug discovery with engineered zinc-finger proteins. *Nat. Rev. Drug Discovery* 2, 361–368.