

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8537017>

Single Chain Variable Fragments against β -Amyloid ($A\beta$) Can Inhibit $A\beta$ Aggregation and Prevent $A\beta$ -Induced Neurotoxicity †

ARTICLE in BIOCHEMISTRY · JULY 2004

Impact Factor: 3.02 · DOI: 10.1021/bi049933o · Source: PubMed

CITATIONS

87

READS

46

9 AUTHORS, INCLUDING:



Bin Yuan

Arizona State University

6 PUBLICATIONS 280 CITATIONS

SEE PROFILE



Yuri L. Lyubchenko

University of Nebraska Medical Center

192 PUBLICATIONS 5,601 CITATIONS

SEE PROFILE



Michael R Sierks

Arizona State University

49 PUBLICATIONS 1,581 CITATIONS

SEE PROFILE

Proteolytic Antibody Light Chains Alter β -Amyloid Aggregation and Prevent Cytotoxicity[†]

Ruitian Liu,[‡] Chad McAllister,[§] Yuri Lyubchenko,[§] and Michael R. Sierks^{*,‡}

Department of Chemical and Materials Engineering and Department of Microbiology, Arizona State University, Tempe, Arizona 85287

Received April 16, 2004; Revised Manuscript Received June 1, 2004

ABSTRACT: β -Amyloid ($A\beta$), a peptide generated by proteolytic cleavage of the amyloid precursor protein (APP), is a major constituent of the neuritic plaques associated with Alzheimer's disease (AD). Up-regulation of α -secretase, which can hydrolyze $A\beta$ between Lys¹⁶ and Leu¹⁷, has been proposed as a potential therapeutic strategy in the treatment of AD. Previously, we identified two light-chain antibody fragments that had proteolytic activity against $A\beta$, one with α -secretase-like activity and one with carboxypeptidase-like activity. Here we show that cleavage of $A\beta$ 40 by hk14, the light-chain antibody having carboxypeptidase-like activity, alters aggregation of $A\beta$ and neutralizes any cytotoxic effects of the peptide. Cleavage of $A\beta$ 40 with c23.5, the light chain having α -secretase-like cleavage, substantially increases the aggregation rate of $A\beta$; however, it does not show any corresponding increase in cytotoxicity. The increase in aggregation resulting from hydrolysis by c23.5 can be attributed to the decreased solubility of the hydrolyzed products relative to the parent $A\beta$ 40, primarily the $A\beta$ 17–40 fragment. These results demonstrate that antibody fragment mediated proteolytic degradation of $A\beta$ peptide can be a potential therapeutic route to control $A\beta$ aggregation and toxicity in vivo. Our results also suggest that increasing α -secretase activity as a therapeutic route must be approached with some caution because this can lead to a substantial increase in aggregation.

The critical pathological features of Alzheimer's disease (AD)¹ are extracellular amyloid plaques and intracellular neurofibrillary tangles (1). The primary proteinaceous component of the amyloid plaques is β -amyloid ($A\beta$) (2). $A\beta$ is generated from the amyloid precursor protein (APP) by proteolytic cleavage at both the β - and γ -secretase sites, resulting in the 40/42 amino acid products (1). $A\beta$ will readily aggregate under appropriate conditions (3), and the different aggregate morphologies that can form, including soluble oligomers, filaments, and insoluble fibrils, have all been reported to play varying roles in the neurotoxic properties of the peptides (1, 4–7, 8–13). $A\beta$ is a normal product of cell metabolism and is found in the plasma and cerebrospinal fluid (CSF) of healthy humans (14). In patients with AD however, numerous different mechanisms have been proposed to explain why $A\beta$ forms toxic aggregates including increased production or decreased clearance of $A\beta$ (15), increased oxidative stress (16, 17), free-radical formation (18), disrupted calcium homeostasis (19), cytoskeletal anomalies (20), and metal ions (21). Therapeutic strategies aimed at reducing $A\beta$ aggregation include decreasing $A\beta$ production

rates by inhibiting activity of either β - or γ -secretases, enzymes that produce the $A\beta$ peptide (14), inhibiting aggregation of $A\beta$ with β -sheet blockers or competitive inhibitors (22, 23), increasing clearance of $A\beta$ from the cerebral cortex (24–27) or from blood (28, 29), and addition of chelators to bind metal ions such as Cu²⁺ and Zn²⁺ that promote aggregation (21, 30). Additional therapeutic approaches include using antiinflammatory agents to decrease the inflammatory response in the brain (31, 32) and using statins to reduce cholesterol levels (33, 34).

Three different proteases are involved in the processing of $A\beta$ from APP: β -secretase cleavage forms the amino terminal portion of $A\beta$, γ -secretase cleavage forms the carboxyl terminal portion, and α -secretase cleaves between what would be residues Lys16 and Leu17 of $A\beta$, precluding formation of full-length $A\beta$. Considerable effort has been placed to develop therapeutics that inhibit processing of APP by β - and γ -secretases or to enhance processing by α -secretase (35). Inhibiting either β - or γ -secretase activity has been shown to effectively decrease $A\beta$ levels in vivo (36–38). Cleavage of APP by α -secretase prevents formation of $A\beta$, releasing instead a large ectodomain of APP (sAPP α) from the cell surface that is reported to have neuroprotective and memory-enhancing properties (39–41). There is apparently a delicate balance in the brain between nonamyloidogenic cleavage of APP by α -secretase and amyloidogenic cleavage of APP by β -secretase, which when viewed in terms of the progression of AD and neurodegeneration represents a classic battle between good and bad processing of APP (35). Facilitating α -secretase processing of APP by supplementing

[†] This work was supported in part by grants from the NIH (Grant AG17984), American Health Assistance Foundation (Grant A2001-0043), and the Alzheimer's Association (Grant IIRG-01-2753).

^{*} Corresponding author. Mailing address: Chemical and Materials Engineering, Box 876006, Tempe, AZ 85287-6006. Fax: 480-965-0037. Phone: 480-965-2828. E-mail: sierks@asu.edu.

[‡] Department of Chemical and Materials Engineering.

[§] Department of Microbiology.

¹ Abbreviations: AD, Alzheimer's disease; $A\beta$, β -amyloid; scFv, single-chain variable fragment; ThT, Thioflavin T; AFM, atomic force microscopy; MTT, 3-(2,5-diphenyltetrazolium bromide).

or increasing α -secretase activity therefore represents a potentially promising route to mitigate amyloid formation (42, 43).

Degradation of A β by various proteolytic enzymes has been studied as a potential therapeutic route for treating AD. Neprilysin (endopeptidase 24.11) was shown to degrade both monomeric and oligomeric forms of A β 40 (44–46), while insulin-degrading enzyme (IDE) (47–49) and endothelin-converting enzyme (ECE) have also been reported to degrade A β (50, 51). We have previously shown that two different antibody light chains can hydrolyze A β : c23.5, which has α -secretase-like activity, and hk14, which has carboxypeptidase-like activity cleaving sequentially from the carboxyl terminus of A β (52). Here we extend these results to demonstrate that these proteolytic light-chain antibody fragments can alter A β aggregation and cytotoxicity. Proteolytic cleavage of A β by the hk14 antibody fragment inhibits A β aggregation and eliminates toxicity toward SH-SY5Y cells, while cleavage by c23.5 increases the rate and extent of A β aggregation but without any corresponding increase in toxicity.

EXPERIMENTAL PROCEDURES

Antibody Samples. Lyophilized samples of the c23.5 and hk14 light-chain antibodies purified from bacterial periplasmic extracts were generously provided by Dr. Sudhir Paul (University of Texas-Health Science Center, Houston, Texas). These light chains were previously reported to cleave vasoactive intestinal polypeptide (53). Each of the purified protein preparations contained a major ~29 kD band corresponding to intact light chain and minor degradation products stainable with anti-human light-chain antibody (Bence Jones proteins) (53). The antibody fragments were lyophilized after purification and dialysis with water, stored at 4 °C, and reconstituted in 1 \times PBS (0.137 M NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄, 1.4 mM KH₂PO₄, pH 7.4) to a concentration of 20 μ M. The purity of the c23.5 and hk14 samples were previously analyzed by SDS–PAGE, and catalytic activity toward A β 40 was verified by mass spectrometry (MS) (52).

A β Fragments and Preparation of Aggregates. A β 1–16 was purchased from Bachem (Torrance, CA), and A β 40 and A β 17–40 were purchased from Biosource (Camarillo, CA). For aggregation experiments, A β 40 was dissolved in 100% 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) to a concentration of 2 mg/mL, sonicated in a water bath for 10 min, aliquoted in microcentrifuge tubes, dried under vacuum, and stored at –20 °C. Immediately prior to use, the HFIP-treated A β 40 was dissolved in dimethyl sulfoxide (DMSO) to 20 mg/mL and diluted to 8 μ M in PBS, pH 7.4, and incubated at 37 °C without shaking in a 0.5 mL Eppendorf PCR tube. For initial studies with antibody fragment mixtures, A β 40 was mixed to a final concentration of 8 μ M, and aliquots of either c23.5 or hk14 were added at 50:1, 200:1, and 1000:1 molar dilutions. For studies using a mixture of peptide samples, A β 17–40 with A β 40, A β peptides were added in equimolar ratios to final total protein concentration of 8 μ M. For the three-peptide mixture, A β 1–16, A β 17–40, and A β 40, 4 μ M solutions of each peptide were mixed together.

Thioflavin T (ThT) Fluorescence Assay. Fluorescence emission of ThT is shifted when it binds to β -sheet aggregate structures such as amyloid fibrils (54). The extent of A β

aggregation in the samples prepared as described above was followed by periodically removing 30 μ L aliquots from the samples and adding them to 2 mL of 5 μ M ThT solution (50 mM phosphate buffer, pH 6.5). Fluorescence intensity was monitored at an excitation wavelength of 450 nm and an emission wavelength of 482 nm using a Shimadzu PF-3501PC spectrofluorophotometer (Shimadzu, Japan) using 1 cm light-path quartz cuvettes with both excitation and emission bandwidths of 5 nm. Each reading represented the average of three values determined by a time scan after subtracting out the fluorescence contribution from free ThT. The ThT readings obtained from samples of A β incubated with either hk14 or c23.5 represent the reading obtained on the sample minus the reading obtained on a control sample containing only the scFv. All ThT fluorescence experiments were performed in triplicate. The standard errors and *P* values were analyzed with Excel and MINITAB software, respectively.

Atomic Force Microscope (AFM) Imaging. APS-modified mica was used as an AFM substrate (55, 56). Five microliters of sample was placed on APS-mica for 2 min, rinsed with deionized water, and dried with argon as described earlier (55, 56). Images were acquired in air using a MultiMode SPM NanoScope III system (Veeco/Digital Instruments, Santa Barbara, CA) operating in tapping mode using silicon probes (Olympus). An analysis of the raw AFM data of the acquired images was performed using Femtoscanner software. This was used as a reflection of the relative size distributions of aggregate populations in the acquired images.

MTT Assay. Human neuroblastoma cells (SH-SY5Y) were maintained in medium containing 40% minimal essential medium (MEM), 40% Ham's modification of F-12, 18% fetal bovine serum (FBS), 1% L-glutamine (3.6 mM), and 1% penicillin/streptomycin antibiotics and grown in a 5% CO₂ atmosphere at 37 °C. Cells were harvested from flasks and plated in 96-well polystyrene plates (Corning Inc., Corning, NY) with approximately 10 000 cells per 100 μ L of medium per well. Plates were incubated at 37 °C for 24 h to allow cells to attach. Eight micromolar A β 40 with or without aliquots of antibody light chains c23.5 or hk14 were preincubated for 6, 12, and 18 days before addition to cultures. Samples were diluted with fresh medium. The final concentration of A β 40 in each culture well with or without added c23.5 or hk14 was 400 nM. Similarly, the final protein concentration in each culture well in the various peptide mixture and control samples was also 400 nM, except for the tripeptide mixture where it was 600 nM. The same volume of medium was added to all sample and control cultures. Plates were then incubated for an additional 72 h at 37 °C. Cell viability was determined using an MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) toxicity assay with addition of 10 μ L of 5 mg/mL MTT to each well (58). After incubation for 3 h at 37 °C, plates were centrifuged, and medium was aspirated from each well. One hundred microliters of MTT dissolvent (0.1 N HCl in 2-propanol) was added to each well. Plates were agitated at room temperature for 15 min to dissolve crystals. The absorbance at 560 nm was measured by a Victor Wallac multiwell assay plate reader (PerkinElmer, Gaithersburg, MD). Averages from six replicate wells were used for each sample and control, and each experiment was repeated three times. Cell viability was calculated by dividing the absor-




FIGURE 1: Effects on aggregation of A β 1–40 by hk14 or c23.5. The kinetics of A β 1–40 fibril formation was monitored by ThT fluorescence in the absence and presence of different concentrations of hk14 (A) or c23.5 (B). The samples were incubated at 37 °C, and 30 μ L of sample was removed periodically and added to 2 mL of 5 μ M ThT at 6, 12, and 18 days. Fluorescence intensity was measured at excitation wavelength 450 nm and emission wavelength 482 nm. The experiments were performed in triplicate. The standard errors and *P* values were analyzed with Excel and Minitab software, respectively, by one-way ANOVA (*, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001 compared to A β 40 alone).

bance of wells containing samples (corrected for background) by the absorbance of wells containing medium alone (corrected for background).

RESULTS

A β 1–40 Aggregation. A β 1–40 will aggregate into fibrillar structures in a time-dependent manner depending on concentration and pH (59, 60). A β 1–40 when incubated alone shows the expected increase in ThT fluorescence from 0 to 6, 12, and 18 days; however when the peptide is incubated with either of the two proteolytic light-chain antibody fragments, c23.5 or hk14, the aggregation process was dramatically altered (Figure 1). Co-incubation of A β 40 with hk14 inhibits aggregation of the peptide in a dose-dependent manner (Figure 1A). At the lowest hk14 dilution (50:1) aggregation of A β 40 was almost totally inhibited, even after 18 days. At the higher dilution of 200:1, aggregation of A β 40 was decreased by around 25% at each time point, while at the highest dilution tested (1000:1), no inhibition of aggregation was observed.

Essentially opposite results were obtained when A β 40 was co-incubated with the c23.5 light-chain fragment. Here aggregation of A β 40 increased in a dose-dependent manner, where at the lowest dilution of 50:1, there was a nearly 200% increase in aggregation at 6, 12, and 18 days, a substantially

smaller increase was observed at a dilution of 200:1, and no effect was observed at the highest dilution rate (Figure 1B).

Imaging of Aggregate Morphologies. Co-incubation of A β 40 with a 50:1 dilution of hk14 or c23.5 alters aggregation as evidenced by ThT staining and AFM imaging. AFM images of the control A β 40 sample taken after 18 days show formation of small oligomers and extensive formation of thin filaments (Figure 2A). Images of the sample co-incubated with hk14 indicate a change in the aggregation morphology with the formation of numerous very thin, almost fragmented filaments along with the small oligomers (Figure 2B). AFM images of samples of A β 40 co-incubated with c23.5 show extensive formation of filaments and small aggregates similar to the control sample (Figure 2C). These results are consistent with our previous study showing that A β 1–40 when co-incubated with A β 17–40 formed more but structurally similar filaments compared to A β 1–40 alone (57). Quantification of the height distributions of the aggregates in the AFM images indicates very similar distributions of particle heights in the control A β alone sample and the sample co-incubated with c23.5. However when A β is co-incubated with hk14, there is a significant increase in the percentage of larger aggregate sizes (Table 1).

Cytotoxicity of A β 40 Aggregate Samples. An MTT assay was used to measure the toxicity of the preincubated A β 40 samples with and without added c23.5 or hk14 light chains after preincubation at 37 °C for 6, 12, and 18 days. When incubated alone, A β 40 exhibited increasing cytotoxicity to SH-SY5Y cells as the preincubation time increases. Addition of a 50/1 dilution of the control protein, BSA, did not alter A β 40 cytotoxicity. However addition of hk14 produces a dose-dependent decrease in toxicity, showing nearly complete protection against A β 40 toxicity at a dilution of 50:1, partial protection at 200:1 dilution, and no protection at 1000:1 dilution (Figure 3A). However when A β 40 was co-incubated with c23.5, there was a slight increase in toxicity compared to A β 40 alone at 6 days for the higher c23.5 concentrations, but essentially no change in toxicity at 12 and 18 days at any concentration compared to A β 40 alone (Figure 3B). Even though c23.5 results in a substantial increase in A β 40 aggregation, it does not result in a corresponding increase in A β 40 cytotoxicity.

Identification of c23.5 Degradation Products That Promote Aggregation. We have previously shown that the light-chain antibody fragment c23.5 can cleave A β 40 at the α -secretase site between residues 16 and 17 and to a lesser extent can also cleave after the lysine 28 residue (52). Since incubation of A β 40 with c23.5 resulted in an increase in aggregation, we tested how A β 1–16 and A β 17–40, the two major hydrolytic fragments resulting from α -secretase cleavage of A β 40, would affect aggregation and cytotoxicity of the parent peptide. We incubated 8 μ M samples of A β 1–16, A β 17–40, and A β 1–40 alone for 18 days along with mixtures containing 4 μ M each of: (a) A β 17–40 and A β 1–40 and (b) A β 1–16, A β 17–40, and A β 1–40. In addition, we incubated a control sample containing 8 μ M A β 1–40 and 0.16 μ M BSA and a test sample containing 8 μ M of A β 1–40 and 0.16 μ M of c23.5. Aliquots of each of these samples were periodically removed and tested for fibrillar content as determined by ThT (Figure 4A) and cytotoxicity as determined by MTT assay (Figure 4B). A β 1–16 alone did not aggregate, nor was it cytotoxic, while




FIGURE 2: AFM imaging of A β 40 aggregation alone and with hk14 or c23.5. AFM analysis of 18-day samples of A β 40 aggregates when incubated alone (A), with hk14 (B), and with c23.5 (C). Scale bars = 1 μ m. Z-scale = 250 nm. Images were acquired in air using a MultiMode SPM NanoScope III system operating in tapping mode using silicon probes.

A β 17–40 and A β 40 both aggregated as expected and showed substantial toxicity. While there were some differences in aggregation between A β 17–40 and A β 40 after 6 and 12 days as measured by ThT, there was no difference in toxicity of the samples at any of the time points tested. We did not observe an increase in aggregation of A β 17–40

compared to A β 40 as previously reported (61) except for a small increase after 6 days. When either the two-peptide mixture, A β 17–40 and A β 1–40, or the three-peptide mixture of A β 1–16, A β 17–40, and A β 1–40 were co-incubated, there was a substantial increase in aggregation after 12 and 18 days, but no corresponding increase in

Table 1: Height Distribution Analysis (%) of Aggregates Obtained by AFM Images of Samples of A β 40 Alone and Co-incubated with hk14 or c23.5 for 18 days at 37 °C

sample	molecule height (nm)					
	0–1	1–2	2–3	3–4	4–5	5–6
A β 40 alone	95.6	3.4	0.4	0.2	0.1	0.1
A β 40 with hk14	89.7	6.6	1.7	0.7	0.3	0.3
A β 40 with c23.5	97.3	1.7	0.5	0.2	0.1	0.1




FIGURE 3: Comparison of toxicity of A β 40 and its mixture with hk14 or c23.5. Eight micromolar A β 40 with or without different concentrations of hk14 (A) or c23.5 (B) was incubated at 37 °C for 6, 12, 18 days. Ten microliters of sample was added to wells of 96-well plates. The final concentration of A β 40 in all wells was 400 nM. Data shown are expressed as percentage of control values (one control for each time point) from three independent experiments with each experimental value being the average of six trials. Standard errors of the mean are shown for each sample as bars above the mean (*, $P < 0.05$; **, $P < 0.01$ compared to A β 40 alone).

cytotoxicity; the three-peptide mixture actually showed a decrease in toxicity. The three-peptide mixture comes close to replicating the aggregation results observed when A β 40 is incubated with c23.5 and almost exactly duplicates the cytotoxicity studies. Addition of a nonspecific protein, BSA, had no effect on aggregation or toxicity compared to A β 40 alone.

DISCUSSION

The hydrophobic carboxyl terminal region of A β is a principal force driving the protein to aggregate (62). The longer carboxyl terminal form of the peptide (42 vs 40) favors aggregation (63, 64), as do deletions in the more hydrophilic amino terminal region (61, 65). The amino terminal deleted peptide, A β 17–40/42 (also referred to as P3) is very insoluble and will form cytotoxic aggregates that may




FIGURE 4: Aggregation and cytotoxicity of degradation products of A β 40 by c23.5. The kinetics of fibril formation was monitored by ThT fluorescence for the samples containing different A β peptides (A). The samples were incubated at 37 °C, and 30 μ L of samples was removed to 2 mL of 5 μ M ThT after 6, 12, and 18 days. Fluorescence intensity was measured at excitation wavelength 450 nm and emission wavelength 482 nm. The experiments were performed in triplicate (*, $P < 0.05$; **, $P < 0.01$ compared to A β 40 alone). The cytotoxicity of the different A β peptides samples when incubated for 6, 12, and 18 days was measured using an MTT assay (B). Ten microliters of sample was added to wells of 96-well plates. The final concentration of A β 40 and total protein of other peptide mixtures in all wells was 400 nM, except for the tripeptide mixture where it was 600 nM. Data shown are expressed as percentage of control values from three independent experiments with each experimental value being the average of six trials.

contribute to the neuronal apoptotic characteristic of AD (66, 67). Here we provide further evidence that A β 17–40 not only aggregates readily but also promotes aggregation when mixed with A β 40, possibly by providing nucleation sites as previously suggested (61). When A β 40 peptide is incubated with the light-chain fragment c23.5, which we have previously shown to possess α -secretase-like activity toward A β 40 (52), aggregation of A β 40 is stimulated since the less soluble 17–40 fragment is generated. Similar α -secretase-like cleavage of A β by insulin-degrading enzyme (IDE) was previously shown to also enhance oligomerization of A β at physiological concentrations in vitro due to the formation of more hydrophobic truncated N-terminal fragments (49).

β -secretase, or BACE (β -site APP-cleaving enzyme), is an aspartyl protease that cleaves membrane-bound APP releasing a soluble protein (β -APPs) and generating the N-terminal fragment of A β on a 99-residue membrane-bound C-terminal fragment (C99) (68, 69). APP can alternatively be cleaved by α -secretase, which cleaves inside the A β region (Lys¹⁶-Leu¹⁷) to produce α -APPs and a slightly shorter membrane-bound C-terminal fragment (C83) (39, 70). Both C99 and C83 are substrates for γ -secretase (43), which can

cleave the intermembrane region of APP to produce the 4-kD A β peptide from C99 and the 3-kD peptide (A β 17–40/42) called P3 from C83. Previous studies have shown that stimulating α -secretase-like cleavage of APP by promoting protein kinase C activity leads to a significant decrease in A β formation (71–74). Since α -secretase cleavage of APP prevents formation of A β but increases production of the P3 fragment, increasing α -secretase activity will decrease soluble A β levels through two different mechanisms: (1) directly decreasing A β production and (2) increasing aggregation of existing A β 40/42 by increasing P3 levels. Our results show that addition of P3 can significantly increase A β aggregation *in vitro*; however, there is no corresponding increase in cytotoxicity. These results indicate that if promoting α -secretase cleavage is to be used as a potential therapeutic approach for treating AD, it may also be beneficial to simultaneously decrease γ -secretase activity. There is still considerable confusion as to whether α -secretase levels play a role in the progression of AD as one study indicated that 80% of samples from AD patients showed decreased α -secretase levels (75), while a second report indicated that mRNA levels of a suggested α -secretase candidate ADAM 10 (disintegrin and metalloproteinase 10) was 2-fold higher in AD patients (76). Promotion of α -secretase activity toward APP by addition of protein kinase C (PKC) reduced formation of A β (77).

The p3 fragment, A β 17–40/42, may play a role in the progression of AD as it has been identified in diffuse plaques of AD patients (78), is neurotoxic (61), and A β 17–42 and A β 1–42 both induce cell death via a similar Fas-like/caspase-8 activation pathway (67, 79). Here we add to these results by showing that A β 17–40 when added directly to A β 1–40 or when formed by α -secretase like proteolytic hydrolysis of A β 1–40, substantially promotes aggregation of A β 1–40, but does not increase cytotoxicity toward SH-SY5Y cells, even despite the formation of extensive fibrillar structures. These results suggest that A β 17–40, while it aggregates more quickly than the parent peptide, favors formation of less toxic aggregate morphologies consistent with previous reports showing that enhanced aggregation of A β 40 did not correlate with increased toxicity to cells (80). Many different studies have indicated that the toxicity of A β may be due to intermediate structures in the aggregation pathway toward formation of insoluble fibrils, rather than the insoluble fibrils themselves (10, 81, 82). Apparently A β 17–40 when mixed with A β 40 favors formation of a less toxic aggregate morphology.

Our results also show that co-incubation of A β 40 with hk14 alters aggregation, forming a higher percentage of larger oligomers and extensive very thin filaments. When incubated alone, hk14 also formed oligomers and filaments as indicated by ThT staining and AFM images; however, it showed no toxicity to SH-SY5Y (data not shown). Therefore the low ThT values observed with the hk14/A β 40 sample is attributable to subtraction of the high fluorescence background of hk14 alone. The very thin filaments observed in Figure 2B may also be partly due to aggregation of hk14; however, the same thin filaments are observed even when A β 40 is incubated with 2-fold lower concentrations of hk14 (data not shown). The reduced toxicity of A β 40 when co-incubated with hk14 may be attributable to proteolytic degradation from the carboxyl terminus or to an altered aggregate morphology

caused by co-aggregation with hk14 or a combination of both mechanisms.

The amyloid cascade hypothesis (15), while still unproven, is a central tenet for AD therapeutic strategies aimed at reducing soluble or aggregated A β levels. Such strategies include active and passive immunization approaches to clear soluble and fibrillar A β , inhibition of A β production by inhibiting β - and γ -secretases, and increasing proteolytic degradation of A β . Active and passive immunization protocols can reduce amyloid deposits and decrease soluble A β levels (29, 83–85) and have been shown to reverse memory deficits (86–88). Unfortunately, active immunization strategies led to clinical signs and symptoms of meningoencephalitis in a small group of AD patients (89, 90). Postmortem examination of immunized animals with vascular amyloid also showed perimicrovascular hemorrhages and inflammation (91–93). Since AD is in part an inflammatory disease (94, 95) and immunization strategies may perturb the delicate balance in the brain between healthy maintenance and unhealthy inflammation (96), a proteolytic therapeutic approach has great promise. A β is normally secreted from cells, and in healthy brains, it is readily hydrolyzed before forming aggregates by various enzymes such as neutral endopeptidase (NEP) (46), angiotension-converting enzyme (ACE) (97), neprilysin (endopeptidase 24.11) (98), endothelin-converting enzyme (ECE) (50), and insulin-degrading enzyme (IDE) (48, 99). Since these enzymes have other functions, stimulating their activity to decrease A β levels may cause other unwanted side effects. Similarly, the proteolytic antibody light chains reported here, particularly hk14, which can alter A β aggregation and inhibit toxicity, in their current state may have very broad specificity and would not be suitable for therapeutic applications. However, recombinant proteolytic antibody light chains are readily adaptable to protein engineering techniques that can greatly increase target specificity. For example pairing of a parent proteolytic light chain with a target specific heavy chain increased the specificity of the resulting construct by 6.6-fold (100). Affinity maturation techniques have been effectively used to increase the affinity of parent antibody fragments over a 1000-fold, even into the femtomolar range (101) and diabody constructs containing a proteolytic antibody fragment along with a high-affinity antigen-specific antibody fragment can also be constructed. Since none of these constructs would contain the Fc fragment of the antibody, the dangers of aggravating an inflammatory response can be alleviated and the smaller size may facilitate transport across the blood–brain barrier. Therefore our current results indicating that proteolytic antibody fragments can potently alter aggregation and decrease cytotoxicity of A β hold great promise for the use of recombinant proteolytic antibody constructs as part of a potential noninflammatory therapeutic treatment and prevention of AD.

ACKNOWLEDGMENT

We would like to thank Dr. Sudhir Paul (University of Texas-Houston Medical School, Houston, Texas) for generously providing samples of the c23.5 and hk14 light-chain antibodies.

REFERENCES

- Selkoe, D. J. (1999) Translating cell biology into therapeutic advances in alzheimer's disease, *Nature* 399, A23–A31.
- Kirschner, D. A., Abraham, C., and Selkoe, D. J. (1986) X-ray diffraction from intraneuronal paired helical filaments and extra-neuronal amyloid fibers in alzheimer disease indicates cross-beta conformation, *Proc. Natl. Acad. Sci. U.S.A.* 83, 503–507.
- Zagorski, M. G., Yang, J., Shao, H., Ma, K., Zeng, H., and Hong, A. (1999) Methodological and chemical factors affecting amyloid beta peptide amyloidogenicity, *Methods Enzymol.* 309, 189–204.
- Klein, W. L. (2002) Abeta toxicity in alzheimer's disease: Globular oligomers (addls) as new vaccine and drug targets, *Neurochem. Int.* 41, 345–352.
- Hardy, J., and Selkoe, D. J. (2002) The amyloid hypothesis of alzheimer's disease: Progress and problems on the road to therapeutics, *Science* 297, 353–356.
- Koo, E. H., Lansbury, P. T., Jr., and Kelly, J. W. (1999) Amyloid diseases: Abnormal protein aggregation in neurodegeneration, *Proc. Natl. Acad. Sci. U.S.A.* 96, 9989–9990.
- Nichols, M. R., Moss, M. A., Reed, D. K., Lin, W. L., Mukhopadhyay, R., Hoh, J. H., and Rosenberry, T. L. (2002) Growth of beta-amyloid(1–40) protofibrils by monomer elongation and lateral association. Characterization of distinct products by light scattering and atomic force microscopy, *Biochemistry* 41, 6115–6127.
- Soto, C. (2003) Unfolding the role of protein misfolding in neurodegenerative diseases, *Nat. Rev. Neurosci.* 4, 49–60.
- Lashuel, H. A., Hartley, D. M., Balakhaneh, D., Aggarwal, A., Teichberg, S., and Callaway, D. J. (2002) New class of inhibitors of amyloid-beta fibril formation. Implications for the mechanism of pathogenesis in alzheimer's disease, *J. Biol. Chem.* 277, 42881–42890.
- Kayed, R., Head, E., Thompson, J. L., McIntire, T. M., Milton, S. C., Cotman, C. W., and Glabe, C. G. (2003) Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis, *Science* 300, 486–489.
- Rosenblum, W. I. (2002) Structure and location of amyloid beta peptide chains and arrays in alzheimer's disease: New findings require reevaluation of the amyloid hypothesis and of tests of the hypothesis, *Neurobiol. Aging* 23, 225–230.
- Ward, R. V., Jennings, K. H., Jepras, R., Neville, W., Owen, D. E., Hawkins, J., Christie, G., Davis, J. B., George, A., Karran, E. H., and Howlett, D. R. (2000) Fractionation and characterization of oligomeric, protofibrillar and fibrillar forms of beta-amyloid peptide, *Biochem. J.* 348 (Part 1), 137–144.
- Kirkitadze, M. D., Condrón, M. M., and Teplow, D. B. (2001) Identification and characterization of key kinetic intermediates in amyloid beta-protein fibrillogenesis, *J. Mol. Biol.* 312, 1103–1119.
- Haass, C., and De Strooper, B. (1999) The presenilins in alzheimer's disease—proteolysis holds the key, *Science* 286, 916–919.
- Selkoe, D. J. (2000) Toward a comprehensive theory for alzheimer's disease. Hypothesis: Alzheimer's disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein, *Ann. N. Y. Acad. Sci.* 924, 17–25.
- Guan, Z. Z., Yu, W. F., Shan, K. R., Nordman, T., Olsson, J., and Nordberg, A. (2003) Loss of nicotinic receptors induced by beta-amyloid peptides in pc12 cells: Possible mechanism involving lipid peroxidation, *J. Neurosci. Res.* 71, 397–406.
- Varadarajan, S., Kanski, J., Aksenova, M., Lauderback, C., and Butterfield, D. A. (2001) Different mechanisms of oxidative stress and neurotoxicity for alzheimer's a beta(1–42) and a beta(25–35), *J. Am. Chem. Soc.* 123, 5625–5631.
- Bartzokis, G. (2004) Age-related myelin breakdown: A developmental model of cognitive decline and alzheimer's disease, *Neurobiol. Aging* 25, 5–18; author reply 49–62.
- Abramov, A. Y., Canevari, L., and Duchon, M. R. (2003) Changes in intracellular calcium and glutathione in astrocytes as the primary mechanism of amyloid neurotoxicity, *J. Neurosci.* 23, 5088–5095.
- Yuasa, S., Nakajima, M., Aizawa, H., Sahara, N., Koizumi, K., Sakai, T., Usami, M., Kobayashi, S., Kuroyanagi, H., Mori, H., Koseki, H., and Shirasawa, T. (2002) Impaired cell cycle control of neuronal precursor cells in the neocortical primordium of presenilin-1-deficient mice, *J. Neurosci. Res.* 70, 501–513.
- Bush, A. I. (2003) The metallobiology of alzheimer's disease, *Trends Neurosci.* 26, 207–214.
- Mason, J. M., Kokkoni, N., Stott, K., and Doig, A. J. (2003) Design strategies for anti-amyloid agents, *Curr. Opin. Struct. Biol.* 13, 526–532.
- Moore, S. A., Huckerby, T. N., Gibson, G. L., Fullwood, N. J., Turnbull, S., Tabner, B. J., El-Agnaf, O. M., and Allsop, D. (2004) Both the d-(+) and l-(–) enantiomers of nicotine inhibit abeta aggregation and cytotoxicity, *Biochemistry* 43, 819–826.
- Soto, C., Kacsak, R. J., Saborio, G. P., Aucouturier, P., Wisniewski, T., Prelli, F., Kacsak, R., Mendez, E., Harris, D. A., Ironside, J., Tagliavini, F., Carp, R. I., and Frangione, B. (2000) Reversion of prion protein conformational changes by synthetic beta-sheet breaker peptides, *Lancet* 355, 192–197.
- Forloni, G., Colombo, L., Girola, L., Tagliavini, F., and Salmona, M. (2001) Anti-amyloidogenic activity of tetracyclines: Studies in vitro, *FEBS Lett.* 487, 404–407.
- LeVine, H. (2002) The challenge of inhibiting abeta polymerization, *Curr. Med. Chem.* 9, 1121–1133.
- Sacchettini, J. C., and Kelly, J. W. (2002) Therapeutic strategies for human amyloid diseases, *Nat. Rev. Drug Discovery* 1, 267–275.
- DeMattos, R. B., Bales, K. R., Cummins, D. J., Paul, S. M., and Holtzman, D. M. (2002) Brain to plasma amyloid-beta efflux: A measure of brain amyloid burden in a mouse model of alzheimer's disease, *Science* 295, 2264–2267.
- DeMattos, R. B., Bales, K. R., Cummins, D. J., Dodart, J. C., Paul, S. M., and Holtzman, D. M. (2001) Peripheral anti-a beta antibody alters cns and plasma a beta clearance and decreases brain a beta burden in a mouse model of alzheimer's disease, *Proc. Natl. Acad. Sci. U.S.A.* 98, 8850–8855.
- Cherny, R. A., Atwood, C. S., Xilinas, M. E., Gray, D. N., Jones, W. D., McLean, C. A., Barnham, K. J., Volitakis, I., Fraser, F. W., Kim, Y., Huang, X., Goldstein, L. E., Moir, R. D., Lim, J. T., Beyreuther, K., Zheng, H., Tanzi, R. E., Masters, C. L., and Bush, A. I. (2001) Treatment with a copper–zinc chelator markedly and rapidly inhibits beta-amyloid accumulation in alzheimer's disease transgenic mice, *Neuron* 30, 665–676.
- Stuve, O., Youssef, S., Steinman, L., and Zamvil, S. S. (2003) Statins as potential therapeutic agents in neuroinflammatory disorders, *Curr. Opin. Neurol.* 16, 393–401.
- Aisen, P. S. (2002) The potential of antiinflammatory drugs for the treatment of alzheimer's disease, *Lancet Neurol.* 1, 279–284.
- Jick, H., Zornberg, G. L., Jick, S. S., Seshadri, S., and Drachman, D. A. (2000) Statins and the risk of dementia, *Lancet* 356, 1627–1631.
- Sparks, D. L., and Schreurs, B. G. (2003) Trace amounts of copper in water induce {beta}-amyloid plaques and learning deficits in a rabbit model of alzheimer's disease, *Proc. Natl. Acad. Sci. U.S.A.* 100, 11065–11069.
- Neve, R. L. (2003) A new wrestler in the battle between alpha- and beta-secretases for cleavage of app, *Trends Neurosci.* 26, 461–463.
- Eriksen, J. L., Sagi, S. A., Smith, T. E., Weggen, S., Das, P., McLendon, D. C., Ozols, V. V., Jessing, K. W., Zavitz, K. H., Koo, E. H., and Golde, T. E. (2003) Nsais and enantiomers of flurbiprofen target gamma-secretase and lower abeta 42 in vivo, *J. Clin. Invest.* 112, 440–449.
- Takahashi, Y., Hayashi, I., Tominari, Y., Rikimaru, K., Morohashi, Y., Kan, T., Natsugari, H., Fukuyama, T., Tomita, T., and Iwatsubo, T. (2003) Sulindac sulfide is a noncompetitive gamma-secretase inhibitor that preferentially reduces abeta 42 generation, *J. Biol. Chem.* 278, 18664–18670.
- Basi, G., Frigon, N., Barbour, R., Doan, T., Gordon, G., McConlogue, L., Sinha, S., and Zeller, M. (2003) Antagonistic effects of beta-site amyloid precursor protein-cleaving enzymes 1 and 2 on beta-amyloid peptide production in cells, *J. Biol. Chem.* 278, 31512–31520.
- Meziane, H., Dodart, J. C., Mathis, C., Little, S., Clemens, J., Paul, S. M., and Ungerer, A. (1998) Memory-enhancing effects of secreted forms of the beta-amyloid precursor protein in normal and amnesic mice, *Proc. Natl. Acad. Sci. U.S.A.* 95, 12683–12688.
- Mattson, M. P., Guo, Z. H., and Geiger, J. D. (1999) Secreted form of amyloid precursor protein enhances basal glucose and glutamate transport and protects against oxidative impairment of glucose and glutamate transport in synaptosomes by a cyclic gmp-mediated mechanism, *J. Neurochem.* 73, 532–537.
- Allinson, T. M., Parkin, E. T., Turner, A. J., and Hooper, N. M. (2003) Adams family members as amyloid precursor protein alpha-secretases, *J. Neurosci. Res.* 74, 342–352.

42. Dewachter, I., and Van Leuven, F. (2002) Secretases as targets for the treatment of alzheimer's disease: The prospects, *Lancet Neurol.* 1, 409–416.
43. Esler, W. P., and Wolfe, M. S. (2001) A portrait of alzheimer secretases—new features and familiar faces, *Science* 293, 1449–1454.
44. Kanemitsu, H., Tomiyama, T., and Mori, H. (2003) Human neprilysin is capable of degrading amyloid beta peptide not only in the monomeric form but also the pathological oligomeric form, *Neurosci. Lett.* 350, 113–116.
45. Iwata, N., Tsubuki, S., Takaki, Y., Shirotani, K., Lu, B., Gerard, N. P., Gerard, C., Hama, E., Lee, H. J., and Saido, T. C. (2001) Metabolic regulation of brain abeta by neprilysin, *Science* 292, 1550–1552.
46. Iwata, N., Tsubuki, S., Takaki, Y., Watanabe, K., Sekiguchi, M., Hosoki, E., Kawashima-Morishima, M., Lee, H. J., Hama, E., Sekine-Aizawa, Y., and Saido, T. C. (2000) Identification of the major abeta1–42-degrading catabolic pathway in brain parenchyma: Suppression leads to biochemical and pathological deposition, *Nat. Med.* 6, 143–150.
47. Chesneau, V., Vekrellis, K., Rosner, M. R., and Selkoe, D. J. (2000) Purified recombinant insulin-degrading enzyme degrades amyloid beta-protein but does not promote its oligomerization, *Biochem. J.* 351 (Part 2), 509–516.
48. Morelli, L., Llovera, R., Gonzalez, S. A., Affranchino, J. L., Prelli, F., Frangione, B., Ghiso, J., and Castano, E. M. (2003) Differential degradation of amyloid beta genetic variants associated with hereditary dementia or stroke by insulin-degrading enzyme, *J. Biol. Chem.* 278, 23221–23226.
49. Vekrellis, K., Ye, Z., Qiu, W. Q., Walsh, D., Hartley, D., Chesneau, V., Rosner, M. R., and Selkoe, D. J. (2000) Neurons regulate extracellular levels of amyloid beta-protein via proteolysis by insulin-degrading enzyme, *J. Neurosci.* 20, 1657–1665.
50. Eckman, E. A., Reed, D. K., and Eckman, C. B. (2001) Degradation of the alzheimer's amyloid beta peptide by endothelin-converting enzyme, *J. Biol. Chem.* 276, 24540–24548.
51. Eckman, E. A., Watson, M., Marlow, L., Sambamurti, K., and Eckman, C. B. (2003) Alzheimer's disease beta-amyloid peptide is increased in mice deficient in endothelin-converting enzyme, *J. Biol. Chem.* 278, 2081–2084.
52. Rangan, S. K., Liu, R., Brune, D., Planque, S., Paul, S., and Sierks, M. R. (2003) Degradation of beta-amyloid by proteolytic antibody light chains, *Biochemistry* 42, 14328–14334.
53. Paul, S., Tramontano, A., Gololobov, G., Zhou, Y. X., Taguchi, H., Karle, S., Nishiyama, Y., Planque, S., and George, S. (2001) Phosphonate ester probes for proteolytic antibodies, *J. Biol. Chem.* 276, 28314–28320.
54. LeVine, H., 3rd (1993) Thioflavine t interaction with synthetic alzheimer's disease beta-amyloid peptides: Detection of amyloid aggregation in solution, *Protein Sci.* 2, 404–410.
55. Shlyakhtenko, L. S., Miloskeska, L., Potaman, V. N., Sinden, R. R., and Lyubchenko, Y. L. (2003) Intersegmental interactions in supercoiled DNA: Atomic force microscope study, *Ultramicroscopy* 97, 263–270.
56. Shlyakhtenko, L. S., Potaman, V. N., Sinden, R. R., Gall, A. A., and Lyubchenko, Y. L. (2000) Structure and dynamics of three-way DNA junctions: Atomic force microscopy studies, *Nucleic Acids Res.* 28, 3472–3477.
57. Liu, R., McAllister, C., Lyubchenko, Y., and Sierks, M. R. (2004) Residues 17–20 and 30–35 of beta-amyloid play critical roles in aggregation, *J. Neurosci. Res.* 75, 162–171.
58. Shearman, M. S., Hawtin, S. R., and Taylor, V. J. (1995) The intracellular component of cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (mtt) reduction is specifically inhibited by beta-amyloid peptides, *J. Neurochem.* 65, 218–227.
59. Stine, W. B., Jr., Dahlgren, K. N., Krafft, G. A., and LaDu, M. J. (2003) In vitro characterization of conditions for amyloid-beta peptide oligomerization and fibrillogenesis, *J. Biol. Chem.* 278, 11612–11622.
60. Kim, Y. S., Randolph, T. W., Stevens, F. J., and Carpenter, J. F. (2002) Kinetics and energetics of assembly, nucleation, and growth of aggregates and fibrils for an amyloidogenic protein. Insights into transition states from pressure, temperature, and cosolute studies, *J. Biol. Chem.* 277, 27240–27246.
61. Pike, C. J., Overman, M. J., and Cotman, C. W. (1995) Amino-terminal deletions enhance aggregation of beta-amyloid peptides in vitro, *J. Biol. Chem.* 270, 23895–23898.
62. Barrow, C. J., and Zagorski, M. G. (1991) Solution structures of beta peptide and its constituent fragments: Relation to amyloid deposition, *Science* 253, 179–182.
63. Jarrett, J. T., Berger, E. P., and Lansbury, P. T., Jr. (1993) The c-terminus of the beta protein is critical in amyloidogenesis, *Ann. N. Y. Acad. Sci.* 695, 144–148.
64. Barrow, C. J., Yasuda, A., Kenny, P. T., and Zagorski, M. G. (1992) Solution conformations and aggregational properties of synthetic amyloid beta-peptides of alzheimer's disease. Analysis of circular dichroism spectra, *J. Mol. Biol.* 225, 1075–1093.
65. He, W., and Barrow, C. J. (1999) The a beta 3-pyroglyutamyl and 11-pyroglyutamyl peptides found in senile plaque have greater beta-sheet forming and aggregation propensities in vitro than full-length a beta, *Biochemistry* 38, 10871–10877.
66. Szczepanik, A. M., Rampe, D., and Ringheim, G. E. (2001) Amyloid-beta peptide fragments p3 and p4 induce pro-inflammatory cytokine and chemokine production in vitro and in vivo, *J. Neurochem.* 77, 304–317.
67. Wei, W., Norton, D. D., Wang, X., and Kusiak, J. W. (2002) Abeta 17–42 in alzheimer's disease activates jnk and caspase-8 leading to neuronal apoptosis, *Brain* 125, 2036–2043.
68. Sinha, S., and Lieberburg, I. (1999) Cellular mechanisms of beta-amyloid production and secretion, *Proc. Natl. Acad. Sci. U.S.A.* 96, 11049–11053.
69. Vassar, R., Bennett, B. D., Babu-Khan, S., Kahn, S., Mendiaz, E. A., Denis, P., Teplow, D. B., Ross, S., Amarante, P., Loeloff, R., Luo, Y., Fisher, S., Fuller, J., Edenson, S., Lile, J., Jarosinski, M. A., Biere, A. L., Curran, E., Burgess, T., Louis, J. C., Collins, F., Treanor, J., Rogers, G., and Citron, M. (1999) Beta-secretase cleavage of alzheimer's amyloid precursor protein by the transmembrane aspartic protease bace, *Science* 286, 735–741.
70. Furukawa, K., Sopher, B. L., Rydel, R. E., Begley, J. G., Pham, D. G., Martin, G. M., Fox, M., and Mattson, M. P. (1996) Increased activity-regulating and neuroprotective efficacy of alpha-secretase-derived secreted amyloid precursor protein conferred by a c-terminal heparin-binding domain, *J. Neurochem.* 67, 1882–96.
71. Caporaso, G. L., Gandy, S. E., Buxbaum, J. D., Ramabhadran, T. V., and Greengard, P. (1992) Protein phosphorylation regulates secretion of alzheimer beta/a4 amyloid precursor protein, *Proc. Natl. Acad. Sci. U.S.A.* 89, 3055–3059.
72. Buxbaum, J. D., Koo, E. H., and Greengard, P. (1993) Protein phosphorylation inhibits production of alzheimer amyloid beta/a4 peptide, *Proc. Natl. Acad. Sci. U.S.A.* 90, 9195–9198.
73. Jacobsen, J. S., Spruyt, M. A., Brown, A. M., Sahasrabudhe, S. R., Blume, A. J., Vitek, M. P., Muenkel, H. A., and Sonnenberg-Reines, J. (1994) The release of alzheimer's disease beta amyloid peptide is reduced by phorbol treatment, *J. Biol. Chem.* 269, 8376–8382.
74. Hung, A. Y., Haass, C., Nitsch, R. M., Qiu, W. Q., Citron, M., Wurtman, R. J., Growdon, J. H., and Selkoe, D. J. (1993) Activation of protein kinase c inhibits cellular production of the amyloid beta-protein, *J. Biol. Chem.* 268, 22959–22962.
75. Tyler, S. J., Dawbarn, D., Wilcock, G. K., and Allen, S. J. (2002) Alpha- and beta-secretase: Profound changes in alzheimer's disease, *Biochem. Biophys. Res. Commun.* 299, 373–376.
76. Gatta, L. B., Albertini, A., Ravid, R., and Finazzi, D. (2002) Levels of beta-secretase bace and alpha-secretase adam10 mRNAs in Alzheimer hippocampus, *Neuroreport* 13, 2031–2033.
77. Zhu, G., Wang, D., Lin, Y. H., McMahon, T., Koo, E. H., and Messing, R. O. (2001) Protein kinase c epsilon suppresses abeta production and promotes activation of alpha-secretase, *Biochem. Biophys. Res. Commun.* 285, 997–1006.
78. Higgins, L. S., Murphy, G. M., Jr., Forno, L. S., Catalano, R., and Cordell, B. (1996) P3 beta-amyloid peptide has a unique and potentially pathogenic immunohistochemical profile in alzheimer's disease brain, *Am. J. Pathol.* 149, 585–596.
79. Wei, W., Wang, X., and Kusiak, J. W. (2002) Signaling events in amyloid beta-peptide-induced neuronal death and insulin-like growth factor i protection, *J. Biol. Chem.* 277, 17649–17656.
80. Lowe, T. L., Strzelec, A., Kiessling, L. L., and Murphy, R. M. (2001) Structure–function relationships for inhibitors of beta-amyloid toxicity containing the recognition sequence klvff, *Biochemistry* 40, 7882–7889.
81. Caghey, B., and Lansbury, P. T. (2003) Protofibrils, pores, fibrils, and neurodegeneration: Separating the responsible protein aggregates from the innocent bystanders, *Annu. Rev. Neurosci.* 26, 267–298.

82. Walsh, D. M., Klyubin, I., Fadeeva, J. V., Cullen, W. K., Anwyl, R., Wolfe, M. S., Rowan, M. J., and Selkoe, D. J. (2002) Naturally secreted oligomers of amyloid beta protein potentially inhibit hippocampal long-term potentiation in vivo, *Nature* 416, 535–539.
83. Bard, F., Cannon, C., Barbour, R., Burke, R. L., Games, D., Grajeda, H., Guido, T., Hu, K., Huang, J., Johnson-Wood, K., Khan, K., Kholodenko, D., Lee, M., Lieberburg, I., Motter, R., Nguyen, M., Soriano, F., Vasquez, N., Weiss, K., Welch, B., Seubert, P., Schenk, D., and Yednock, T. (2000) Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of alzheimer disease, *Nat. Med.* 6, 916–919.
84. Wilcock, D. M., DiCarlo, G., Henderson, D., Jackson, J., Clarke, K., Ugen, K. E., Gordon, M. N., and Morgan, D. (2003) Intracranially administered anti- β antibodies reduce β -amyloid deposition by mechanisms both independent of and associated with microglial activation, *J. Neurosci.* 23, 3745–3751.
85. Bard, F., Barbour, R., Cannon, C., Carretto, R., Fox, M., Games, D., Guido, T., Hoenow, K., Hu, K., Johnson-Wood, K., Khan, K., Kholodenko, D., Lee, C., Lee, M., Motter, R., Nguyen, M., Reed, A., Schenk, D., Tang, P., Vasquez, N., Seubert, P., and Yednock, T. (2003) Epitope and isotype specificities of antibodies to β -amyloid peptide for protection against alzheimer's disease-like neuropathology, *Proc. Natl. Acad. Sci. U.S.A.* 100, 2023–2028.
86. Dodart, J. C., Bales, K. R., Gannon, K. S., Greene, S. J., DeMattos, R. B., Mathis, C., DeLong, C. A., Wu, S., Wu, X., Holtzman, D. M., and Paul, S. M. (2002) Immunization reverses memory deficits without reducing brain β burden in alzheimer's disease model, *Nat. Neurosci.* 5, 452–457.
87. Kotilinek, L. A., Bacskai, B., Westerman, M., Kawarabayashi, T., Younkin, L., Hyman, B. T., Younkin, S., and Ashe, K. H. (2002) Reversible memory loss in a mouse transgenic model of alzheimer's disease, *J. Neurosci.* 22, 6331–6335.
88. Hock, C., Konietzko, U., Streffer, J. R., Tracy, J., Signorell, A., Muller-Tillmanns, B., Lemke, U., Henke, K., Moritz, E., Garcia, E., Wollmer, M. A., Umbricht, D., de Quervain, D. J., Hofmann, M., Maddalena, A., Papassotiropoulos, A., and Nitsch, R. M. (2003) Antibodies against β -amyloid slow cognitive decline in alzheimer's disease, *Neuron* 38, 547–554.
89. Schenk, D. (2002) Amyloid- β immunotherapy for alzheimer's disease: The end of the beginning, *Nat. Rev. Neurosci.* 3, 824–828.
90. Senior, K. (2002) Dosing in phase ii trial of alzheimer's vaccine suspended, *Lancet Neurol.* 1, 3.
91. Furlan, R., Brambilla, E., Sanvito, F., Roccatagliata, L., Olivieri, S., Bergami, A., Pluchino, S., Uccelli, A., Comi, G., and Martino, G. (2003) Vaccination with amyloid- β peptide induces autoimmune encephalomyelitis in c57/bl6 mice, *Brain* 126, 285–291.
92. Nicoll, J. A., Wilkinson, D., Holmes, C., Steart, P., Markham, H., and Weller, R. O. (2003) Neuropathology of human alzheimer disease after immunization with amyloid- β peptide: A case report, *Nat. Med.* 9, 448–452.
93. Pfeifer, M., Boncristiano, S., Bondolfi, L., Stalder, A., Deller, T., Staufenbiel, M., Mathews, P. M., and Jucker, M. (2002) Cerebral hemorrhage after passive anti- β immunotherapy, *Science* 298, 1379.
94. Lue, L. F., Walker, D. G., and Rogers, J. (2001) Modeling microglial activation in alzheimer's disease with human postmortem microglial cultures, *Neurobiol. Aging* 22, 945–956.
95. Zhang, Q., Powers, E. T., Nieva, J., Huff, M. E., Dendle, M. A., Bieschke, J., Glabe, C. G., Eschenmoser, A., Wentworth, P., Jr., Lerner, R. A., and Kelly, J. W. (2004) Metabolite-initiated protein misfolding may trigger alzheimer's disease, *Proc. Natl. Acad. Sci. U.S.A.*
96. Shen, Y., Lue, L., Yang, L., Roher, A., Kuo, Y., Strohmeier, R., Goux, W. J., Lee, V., Johnson, G. V., Webster, S. D., Cooper, N. R., Bradt, B., and Rogers, J. (2001) Complement activation by neurofibrillary tangles in alzheimer's disease, *Neurosci Lett* 305, 165–168.
97. Hu, J., Igarashi, A., Kamata, M., and Nakagawa, H. (2001) Angiotensin-converting enzyme degrades alzheimer amyloid β -peptide (a β); retards a β aggregation, deposition, fibril formation; and inhibits cytotoxicity, *J. Biol. Chem.* 276, 47863–47868.
98. Iwata, N., Mizukami, H., Shirotani, K., Takaki, Y., Muramatsu, S., Lu, B., Gerard, N. P., Gerard, C., Ozawa, K., and Saido, T. C. (2004) Presynaptic localization of neprilysin contributes to efficient clearance of amyloid- β peptide in mouse brain, *J. Neurosci.* 24, 991–998.
99. Leissring, M. A., Farris, W., Chang, A. Y., Walsh, D. M., Wu, X., Sun, X., Frosch, M. P., and Selkoe, D. J. (2003) Enhanced proteolysis of β -amyloid in app transgenic mice prevents plaque formation, secondary pathology, and premature death, *Neuron* 40, 1087–1093.
100. Sun, M., Gao, Q. S., Kimarskiy, L., Rees, A., and Paul, S. (1997) Cleavage specificity of a proteolytic antibody light chain and effects of the heavy chain variable domain, *J. Mol. Biol.* 271, 374–385.
101. Boder, E. T., Midelfort, K. S., and Wittrup, K. D. (2000) Directed evolution of antibody fragments with monovalent femtomolar antigen-binding affinity, *Proc. Natl. Acad. Sci. U.S.A.* 97, 10701–10705.

BI0492354