

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7703958>

# PhosphoThr Peptide Binding Globally Rigidifies Much of the FHA Domain from Arabidopsis Receptor Kinase-Associated Protein Phosphatase †,‡

ARTICLE *in* BIOCHEMISTRY · SEPTEMBER 2005

Impact Factor: 3.02 · DOI: 10.1021/bi050414a · Source: PubMed

---

CITATIONS

19

---

READS

13

## 6 AUTHORS, INCLUDING:



Fabio Gallazzi

University of Missouri

51 PUBLICATIONS 1,825 CITATIONS

[SEE PROFILE](#)



Steven R Van Doren

University of Missouri

60 PUBLICATIONS 1,390 CITATIONS

[SEE PROFILE](#)



# NIH Public Access

## Author Manuscript

*Biochemistry*. Author manuscript; available in PMC 2010 January 29.

Published in final edited form as:  
*Biochemistry*. 2005 August 2; 44(30): 10119. doi:10.1021/bi050414a.

## PhosphoThr Peptide Binding Globally Rigidifies much of the FHA Domain from Arabidopsis Receptor Kinase-Associated Protein Phosphatase

**Zhaofeng Ding<sup>†</sup>,**

Department of Biochemistry, 117 Schweitzer Hall, University of Missouri, Columbia, Missouri, 65211

**Gui-in Lee<sup>†,‡</sup>,**

Department of Biochemistry, 117 Schweitzer Hall, University of Missouri, Columbia, Missouri, 65211

**Xiangyang Liang,**

Department of Biochemistry, 117 Schweitzer Hall, University of Missouri, Columbia, Missouri, 65211

**Fabio Gallazzi,**

Molecular Biology Program, 125 Chemistry, 601 S. College Ave., University of Missouri, Columbia, Missouri, 65211 USA

**A. Arunima, and**

Department of Biochemistry, 117 Schweitzer Hall, University of Missouri, Columbia, Missouri, 65211

**Steven R. Van Doren\***

Department of Biochemistry, 117 Schweitzer Hall, University of Missouri, Columbia, Missouri, 65211

### Abstract

A net increase in the backbone rigidity of the kinase-interacting FHA domain (KI-FHA) from the Arabidopsis receptor kinase-associated protein phosphatase (KAPP) accompanies the binding of a phosphoThr peptide from its CLV1 receptor-like kinase partner, according to  $^{15}\text{N}$  NMR relaxation at 11.7 and 14.1 T. All of the loops of free KI-FHA display evidence of nsec-scale motions. Many of these same residues have residual dipolar couplings that deviate from structural predictions.

Binding of the CLV1 pT868 peptide seems to reduce nsec-scale fluctuations of all loops, including half of the residues of recognition loops. Residues important for affinity are found to be rigid, i.e. conserved residues and residues of the subsite for the key pT+3 peptide position. –This behavior parallels SH2 and PTB domain recognition of pTyr peptides. PhosphoThr peptide binding increases KI-FHA backbone rigidity ( $S^2$ ) of three recognition loops, a loop nearby, seven strands from the  $\beta$ -sandwich, and a distal loop. Compensating the trend of increased rigidity, binding enhances fast mobility at a few sites in four loops on the periphery of the recognition surface and in two loops on the far side of the  $\beta$ -sandwich. Line broadening evidence of  $\mu\text{sec}$  to msec-scale fluctuations occurs across the six-stranded  $\beta$ -sheet and nearby edges of the  $\beta$ -sandwich; this forms a network connected by packing of interior side chains and H-bonding. A patch of the slowly fluctuating residues coincides with the site of segment-swapped dimerization in crystals of the FHA domain of human Chfr.

Phosphopeptide binding introduces  $\mu\text{sec}$  to msec-scale fluctuations to more residues of the long 8/9

\*To whom correspondence should be addressed, E-mail: vandorens@missouri.edu, Phone: 1 (573) 882-5113, FAX: 1 (573) 884-4812.

<sup>†</sup>These authors contributed equally to this work.

<sup>‡</sup>Current address: Department of Chemistry, 225 Prospect St., Yale University, New Haven, CT 06520

The NMR relaxation data and their model-free interpretation are available for free and bound KI-FHA under BMRB accession codes 5841 and 6474 at [www.bmrb.wisc.edu](http://www.bmrb.wisc.edu).

recognition loop of KI-FHA. The rigidity of this FHA domain appears to couple as a whole to pThr peptide binding.

## Keywords

FHA domain; phosphoThr-binding module; protein-protein interactions; NMR relaxation; relaxation dispersion; dynamics; residual dipolar couplings

FHA domains bind phosphothreonine- and phosphoserine-containing partners in diverse eukaryotic signaling pathways that include DNA damage repair, cell proliferation, pre-mRNA splicing, forkhead transcription factors, Ring-finger proteins, and kinesins (1). Several high-resolution structures of FHA domains have appeared (2–8), but no detailed studies of their dynamics. Flexibility often appears to correlate with binding events, including affinity of protein modules for peptides (9–11). Residue-specific effects on the affinity of SH2 domains for phosphotyrosine peptides are not fully explained by the structures but do correlate with rigidity of interfacial side chains (12). Peptide-binding dependent changes in backbone motions of the c-Src SH3 domain help explain its unfavorable entropy of peptide binding and show that the binding stabilizes the SH3 domain both near and far from the interface (13). Investigation of the dynamic character of an FHA domain should add insights into behaviors affecting binding and stability.

The FHA domain investigated herein is from *Arabidopsis* kinase-associated protein phosphatase (KAPP), a key downstream regulator of receptor-like protein kinase (RLK) signaling pathways in plants (14). KAPP has a C-terminal protein phosphatase 2C (PP2C) domain that dephosphorylates RLK partners to attenuate the activation of the receptor complex (15). KAPP was shown to interact with these RLKs from *Arabidopsis* that include HAESA (formerly RLK5) (14), CLAVATA1 (16), RLK4 (17), TMK1 (17), WAK1 (18), FLS2 (19), BAK1 (20), and SERK1 (21). KAPP uses its kinase-interacting FHA domain (KI-FHA) to bind epitopes of RLKs activated by phosphorylation of serine or threonine residues. KI-FHA minimally comprises 119 amino acids, i.e. residues 180 to 298 of KAPP (22). The solution structure of KI-FHA is a sandwich of a five-stranded mixed β sheet with six-stranded anti-parallel β-sheet (7), like most other structures of FHA domains (2,3,5) and SMAD MH2 domains(23), despite sequence identities of less than 20% (7). The six-stranded β-sheet is an array of three β-hairpins: the 1/2, 10/11, and 7/8 β-hairpins (see below). The 5/6 β-hairpin lies in the middle of the five-stranded β-sheet. Sequence identity among diverse FHA domains is restricted to Gly211, Arg212, Ser226, His229 and Asn250 in KI-FHA numbering (22) of the 3/4, 4/5, and 6/7 loops on the edge of the five-stranded β-sheet that forms a surface for binding phosphorylated peptides and proteins (3,4,7). The 8/9 and 10/11 loops also appear to contribute to the phosphoprotein (RLK) recognition surface of KI-FHA from KAPP (7). A second surface on an independent face of FHA domains was proposed to be a potential self-recognition surface (7) and coincides with the β-hairpin that rearranges to make the segment-swapped dimer reported in the FHA domain of Chfr (6). Crystallographic B factors of FHA domains are low, especially in the β-strands (5), suggesting rigidity. Phosphopeptide binding does not appear to alter the structure of FHA domains significantly (4,6,24–26). This suggests that FHA domains may use a rigid, pre-formed surface for recognizing phosphoprotein partners.

Many protein-protein interaction sites do, however, have some flexibility revealed by NMR relaxation, often fluctuating over the μsec to msec range (12,13,27–33). NMR relaxation rates result from the density of motions that reorient the bond probed, described by spectral density functions that are Fourier transforms of time correlation functions of the bond reorientation. The model-free formalism estimates primarily the amplitude and secondarily the time scale of intramolecular motions faster than the rotation diffusion of globular proteins (several nsec),

by fitting spectral density functions (34,35). The amplitude of fast internal bond reorientations can be estimated from the generalized order parameter  $S^2$  that ranges from a value of 1.0 for full restriction to 0.0 for full lack of restriction (34,36). The model-free approach also can yield a line broadening term that results from fluctuations occurring on a time scale of  $\mu$ sec to msec. Alternatively,  $^{15}\text{N}$  relaxation data are occasionally interpreted by the reduced spectral density approach. Examples of its application in molecular recognition are refs (33,37). Without explicit modeling of protein tumbling, this approach suggests frequency distribution of amide bond reorientations at  $^1\text{H}$  and  $^{15}\text{N}$  Larmor precession frequencies, as well as much lower frequencies (38–41). Another means of finding structural averaging from psec to tens of msec is to evaluate agreement of residual dipolar couplings (RDCs), measured from weakly aligned NMR samples, with the static structural model (42,43). Comparison of the RDC fits with relaxation data can help assess structural averaging (44).

This first investigation of binding-linked mobility changes of an FHA domain compares the backbone dynamics of representative KI-FHA from Arabidopsis KAPP, free and bound to a pThr peptide we identified from its CLAVATA1 (CLV1) receptor-like kinase partner. CLV1 is required in Arabidopsis for balancing proliferation and differentiation of shoot and floral meristems. KAPP negatively regulates CLV1 in plant development (45).  $^{15}\text{N}$  relaxation data from two magnetic fields are interpreted using both the model-free formalism (34–36) and a reduced spectral density function. Structural fitting of residual dipolar couplings provides an independent assay sensitive to additional time scales. All the loops manifest fluctuations in nsec in the free state, but fewer do in the bound state. In the bound state, many residues have increased rigidity or  $S^2$  whether near or far from the phosphopeptide binding site. Around seven residues have clearly decreased rigidity or  $S^2$  as well. Fluctuations on the  $\mu$ sec to msec scale map to a larger swath and a smaller patch of the six-stranded  $\beta$ -sheet.

## Materials and Methods

### Preparation of KI-FHA

GST-tagged KI-FHA, containing Arabidopsis KAPP residues 180–313, was expressed in *E. coli*, enriched with  $^{15}\text{N}$ , and purified as described (46). After proteolytic removal of the N-terminal GST tag, a linker of Gly-Pro-Leu-Gly-Ser remained at the N-terminus of KI-FHA; the NMR-observable amides of the Leu-Gly-Ser of this linker are assigned KAPP residue numbers of 177–179. Purified KI-FHA was exchanged into 20 mM sodium phosphate (pH 6.3), 120 mM NaCl and 7%  $\text{D}_2\text{O}$  for NMR. NMR samples of KI-FHA were 0.33 to 0.6 mM, the solubility limit.

### Synthesis, purification, and NMR sample uses of CLV1 pT868 peptide

The pT868 peptide, comprising residues 863–875 of the receptor kinase CLAVATA1 of Arabidopsis, was synthesized with an Advanced ChemTech 396 multiple peptide synthesizer (Louisville, KY) using standard Fmoc chemistry and solid phase synthesis. Each Fmoc amino acid was coupled at least twice, except for the building blocks for the phosphothreonine and the subsequent amino acid that were coupled three times. Cleavage and side chain deprotection were achieved by treating the resin with 87.5% trifluoroacetic acid / 2.5% thioanisole / 2.5% phenol / 2.5% water / 2.5% ethanedithiol / 2.5% triisopropylsilane. All reagents were HPLC or peptide synthesis grade and obtained from Fluka (Milwaukee, WI), Novabiochem (San Diego, CA), or VWR Scientific (St. Louis, MO). The crude product obtained was characterized by HPLC (Beckmann Coulter, Fullerton, CA) and LC-ESI-MS (Thermo Finnigan, San Jose, CA). The product was purified to greater than 80% by semi-preparative HPLC using an in-house-optimized multi-step gradient.  $^{15}\text{N}$ - $^1\text{H}$  TROSY at 22°C monitored the effects of the CLV1 pT868 peptide on the KI-FHA amide spectrum when titrated up to a 4:1 ratio of peptide:protein. A sample of 0.795 mM pT868 peptide and 0.53 mM KI-FHA, providing ~94%

saturation at the  $K_D$  of 20  $\mu\text{M}$ , was used to measure a full set of standard relaxation data at each of two magnetic field strengths. A sample of 1.16 mM pT868 peptide and 0.33 mM KI-FHA, providing ~98% saturation at the  $K_D$  of 20  $\mu\text{M}$ , was used to measure standard in-phase  $R_2$  and TROSY-enhanced, relaxation-compensated  $R_2$  at 600 MHz.

### **<sup>15</sup>N NMR relaxation measurements**

Spectra monitoring  $^{15}\text{N}$   $R_1$  and  $R_2$  relaxation rates and  $^{15}\text{N}\{^1\text{H}\}$  steady-state NOE were acquired at 25 °C at both 600 and 500 MHz. Relaxation dispersion experiments were collected at 22 °C at 600 MHz. A Varian Inova 600 MHz spectrometer (Univ. of Missouri) was used with either a high-sensitivity 5 mm  $^1\text{H}\{^{13}\text{C}/^{15}\text{N}\}$  cryogenic probe with shielded z gradient coil or a 5 mm HCN triple resonance probe fitted with an xyz gradient coil. The Bruker DRX-500 spectrometer at the Univ. of Missouri was used with an 8 mm triple resonance probe with an actively shielded z gradient coil (Nalorac, Martinez, CA). The Bruker DMX-500 at NMRFAM was used for  $^{15}\text{N}\{^1\text{H}\}$  steady-state NOE and  $R_1$  at 500 MHz as it is fitted with a very sensitive 5 mm Cryoprobe™ with shielded z gradient coil. The temperature settings of Missouri's 500 and 600 MHz instruments were calibrated using ethylene glycol from room temperature upward and using methanol below room temperature. The actual temperatures are within 1 °C of the nominal temperatures and within 1 °C between the two instruments. The 600 MHz  $R_1$  and  $R_2$  pulse sequences used sensitivity-enhanced gradient-based coherence selection (47), implemented in Biopack. Cross-correlation was suppressed from  $R_1$  and  $R_2$  spectra (48). For good water suppression with cryogenic probes and economy of pulses for the 8 mm probe, home-implemented 3-9-19 WATERGATE (49) versions of sequences of ref (50) were used for  $^{15}\text{N}\{^1\text{H}\}$  steady-state NOE, 500 MHz  $R_1$  and  $R_2$ . The relaxation-compensated CPMG pulse sequence of ref (51) was applied to the free state of KI-FHA and the TROSY-enhanced version (52) to the bound state, using the Inova 600 for each.  $\tau_{\text{cp}}$  delays were in both cases set for 2 ms and 7.5 ms  $\pi$  pulse spacing, to identify sites of exchange broadening on the msec scale.

Using relaxation-compensated CPMG sequences with  $\tau_{\text{cp}} = 1$  ms ( $\pi$  pulse spacing of 2 ms), total relaxation delays were 16, 32, 48, 64, 80, 96 ms using the sequence of ref (51) or 8, 32, 64, 96, 128, and 160 ms using the sequence of ref (52). At  $\tau_{\text{cp}} = 3.75$  ms ( $\pi$  pulse spacing of 7.5 ms) total delays were 30, 60, 90, and 120 ms using the sequence of ref (51) or 30, 60, 90, 120, 150, and 180 ms using the sequence of ref (52). Relaxation delay periods at 500 MHz were 20\*, 60, 120\*, 240, 400\*, 600 and 860\* ms for  $R_1$  and 16\*, 32, 48\*, 64, 80\*, 112, 144\* ms for  $R_2$ . Asterisks (\*) indicate points duplicated for estimating uncertainties in peak heights. 500 MHz  $R_1$  data of free KI-FHA were found to reproduce well between Missouri's DRX-500 with conventional probe and NMRFAM's DMX-500 with Cryoprobe™. Delay times at 600 MHz were 10, 20\*, 50, 100\*, 200, 350\*, 500, and 700\* ms for  $R_1$  and 10\*, 30, 50\*, 70, 90\*, 110, and 130\* ms for  $R_2$ . For KI-FHA in the free state, the 600 MHz  $R_1$  and  $R_2$  results from the conventional probe were repeated with the cryogenic probe and averaged. NMR spectra were processed using Sybyl 6.8 or NMRPipe 2.3 (53). Uncertainties in  $R_1$  and  $R_2$  relaxation rates were generated by relaxation fitting simulations with SPARKY 3 (54) using 40 iterations with addition of Gaussian-distributed random noise. To obtain  $^{15}\text{N}\{^1\text{H}\}$  NOEs, spectra were measured with and without 3.5 seconds of proton saturation, in an interleaved manner to maintain identical sample conditions. The non-saturated spectra reference spectra had relaxation delays of 9.2 seconds at 500 MHz and 12 seconds at 600 MHz to allow water magnetization to recover completely and avoid saturation transfer. Identical pairs of NOE spectra were collected in triplicate, except for the free state at 500 MHz collected in duplicate, to evaluate standard deviations for each residue. Standard rules of statistical error propagation were applied to arithmetic combinations of relaxation rates.

## Reduced spectral density analysis of $^{15}\text{N}$ relaxation

The spectral density function,  $J(\omega)$ , for an N-H bond vector describes the frequency spectrum of its reorientation.  $J(\omega)$  decreases monotonically with frequency  $\omega$  and has nearly zero slope at high frequency, i.e.  $J(\omega_{\text{H}}+\omega_{\text{N}})=J(\omega_{\text{H}})=J(\omega_{\text{H}}-\omega_{\text{N}})$ . (Recall that  $\omega_{\text{N}}$  is negative.) Consequently, three spectral density functions for each amide are sufficient and well-approximated from measurements of just  $^{15}\text{N}$   $R_1$  and  $R_2$  and  $\{^1\text{H}\}-^{15}\text{N}$  NOE enhancement. The three spectral density expressions resulting are (38–40):

$$J_{\text{eff}}(0)=[6/3d^2+4c^2](-R_1/2+R_2-3\sigma/5) \quad (1)$$

$$J(\omega_{\text{N}})=[4/3d^2+4c^2](R_1-7\sigma/5) \quad (2)$$

$$J(0.87\omega_{\text{H}})=4\sigma/(5d^2) \quad (3)$$

$$\sigma=(\text{NOE}-1)(\gamma_{\text{N}}/\gamma_{\text{H}})R_1 \quad (4)$$

where  $d=\mu_0h\gamma_{\text{N}}\gamma_{\text{H}}\langle r_{\text{NH}}^{-3} \rangle/(8\pi^2)$ ;  $c=\omega_{\text{N}}(\sigma_{\parallel}-\sigma_{\perp})/\sqrt{3}$ ;  $\sigma_{\parallel}-\sigma_{\perp}$  is the chemical shift anisotropy estimated to be -170 ppm;  $\gamma_{\text{H}}$  and  $\gamma_{\text{N}}$  are the gyromagnetic ratios of  $^1\text{H}$  and  $^{15}\text{N}$ ; and  $\omega_{\text{H}}$  and  $\omega_{\text{N}}$  are the Larmor frequencies of  $^1\text{H}$  and  $^{15}\text{N}$ .  $J(\omega)$  at  $\omega=0$ ,  $\omega_{\text{N}}$ , and  $0.87\omega_{\text{H}}$  were estimated from both 500 and 600 MHz relaxation data. The reduced spectral mapping approach needs no models of internal motion or of anisotropy in Brownian tumbling of the protein.

## Model-free analysis of $^{15}\text{N}$ relaxation

Model-free calculations were performed using the ModelFree 4.1 suite of programs provided by Prof. Arthur G. Palmer (55). Estimates of  $\tau_m$  and anisotropy of the rotational diffusion tensor were obtained from  $R_2/R_1$  ratios using the program TENSOR2 (56) after filter coarse- and fine-filtering out residues undergoing motion on the ps to ns or  $\mu\text{s}$  to ms timescales (57); see Results. The  $^{15}\text{N}$  CSA was set to the approximation of -170 ppm (9,58). To anticipate systematic effects upon  $S^2$  of the unknown variation in  $^{15}\text{N}$  CSA (59) of KI-FHA, parallel model-free calculations were fixed to a range of other CSA values. At -150 ppm,  $S^2$  averages  $0.015 \pm 0.019$  above  $S^2$  at -170 ppm. At -190 ppm,  $S^2$  averages  $0.029 \pm 0.026$  below  $S^2$  at -170 ppm. The uncertainties for  $R_1$  and  $R_2$  were set to 5% of their values as suggested (60). Uncertainties in the dynamics parameters were obtained using 400 steps Monte Carlo simulations carried out using Modelfree 4.1. Under the assumption of a wobbling-in-a-cone motional model, the amplitude of the maximal cone semi-angle of N-H bond excursions can be estimated from order parameter  $S$  using the relationship (34):

$$S=\frac{\cos(\theta)+\cos^2(\theta)}{2} \quad (5)$$

## Selection of spectral density expressions

Relaxation rates were fitted to one of five forms of spectral density functions (Table 1), where  $\tau_m$  and  $D_{\parallel}/D_{\perp}$  were fixed to estimates from  $R_2/R_1$  ratios (Table 2). To accommodate data from two magnetic fields leading to higher model-free  $\chi^2$  errors and to avoid under-fitting and bias

(61), long-established Akaike's Information Criterion (AIC) (62) was used as the principal method of statistical model selection as described (61). Akaike's information criterion is computed as  $\chi^2 + 2k$ , where  $k$  is the number of parameters evaluated from among  $S_s^2$ ,  $\tau_e$ ,  $R_{ex}$  and  $S_f^2$  (61,62). AIC was used to test need to let  $\tau_e$  range up to 2 nsec in expressions 2 and 4 of Table 1. By monitoring  $R_1^{500MHz} / R_1^{600MHz}$  after a correction for psec internal motion, the PINATA algorithm suggested nsec-scale motions in additional residues with ratio < 0.96 (63). The  $\tau_e$  term of these residues was allowed to range up to 1 nsec. The suitability of use of 1 to 2 nsec  $\tau_e$  was confirmed by elevated  $J(\omega_N=61\text{ MHz})$  (eq. 2) since the baseline at each spectral density frequency marks the rigid structural core (11). To justify use of the exchange broadening term  $R_{ex}$  (expressions 3 and 4 in Table 1), F-testing of model-free  $\chi^2$ , using data from the two fields, was required to show a P value < 0.25 in addition to satisfying AIC. Alternatively, relaxation dispersion data with increased line broadening at slower CPMG pulsing rate, i.e.  $R_2(7.5\text{ ms}) - R_2(2\text{ ms})$  beyond uncertainty above the baseline, was considered clear evidence of exchange on the msec scale and sufficient justification for use of  $R_{ex}$ .

### Residual dipolar coupling analysis

RDCs were measured using liquid crystals that align in the magnetic field, introducing a small net average alignment (< 0.001) of KI-FHA.  ${}^1D_{NH}$  values were measured using 0.5 mM KI-FHA in the presence of Pf1 phage (64) at 9.6 mg/ml and using 0.4 mM KI-FHA in 5% (w/v) C12E6 / hexanol liquid crystalline medium (65). RDCs were obtained by comparing coupled  ${}^{15}N$  IPAP-HSQC spectra of KI-FHA in the absence of orienting media against spectra of KI-FHA weakly oriented using the phage or PEG/hexanol media.  ${}^1D_{NH}$  values of successive sets of five neighboring residues were fitted by singular value decomposition (SVD) to the four lowest energy structural models of KI-FHA (PDB code 1MZK) to obtain alignment tensor parameters as described (44). The application here is novel in using a group of NMR structures rather than an X-ray structure. In simulating the range of uncertainty of the alignment parameter  $S_{ZZ}$ , estimated uncertainty in atomic coordinates of 0.3 Å and of 1 Hz in measurements of  ${}^1D_{NH}$  values were included in  $512 * 1024$  fits introducing random noise of this size. Repetition for the group of NMR structures in effect incorporated further structural uncertainty.

## RESULTS

### ${}^{15}N$ relaxation in free and phosphoThr peptide-bound states

${}^{15}N$  NMR relaxation parameters of KI-FHA, free and bound to an excess of a 13-mer pThr peptide from CLAVATA1 (CLV1), were determined at 298 K, at both 500 MHz and 600 MHz to enhance insight into the amplitude and timescale of backbone dynamics. The peptide, dubbed CLV1 pT868, comprises residues 863–875 of the Arabidopsis CLAVATA1 receptor-like kinase (RLK): PEYAY(pT)LKVDEKS. Multiple sequence alignment of kinase domains of RLKs indicates that the threonine that we phosphorylated in this CLV1 pT868 peptide corresponds to Ser1060 of the BRI1 RLK, one of two neighboring residues phosphorylated at the C-terminal end of the activation loop of kinases (66). This suggests pT868 of CLV1 might also be phosphorylated *in vivo*, making it the most likely of our KI-FHA-binding phosphopeptide candidates to be physiologically relevant. At 298K, CLV1 pT868 binds KI-FHA with apparent  $K_D$  of 21  $\mu\text{M}$  by NMR at pH 6.3 and 16  $\mu\text{M}$  by titration calorimetry at pH 7.5; details will be described elsewhere. Of 114 observable backbone amide peaks, 99 in the free state and 102 in the bound state were resolved well enough for quantitative analysis of their relaxation and dynamics. Single exponentials fit well the  $R_1$  and  $R_2$  relaxation rates. The exception is in 500 MHz  $R_1$  spectra where four to five residues have anomalously fast relaxation (marked with gray squares in Figure 1b, Figure 2b) that approaches or exceeds the theoretical limit of  $3.14\text{ s}^{-1}$  when fitted with a single exponential; their biphasic relaxation is possibly an artifact of the exchange broadening that these particular backbone peaks undergo. In the free state of KI-FHA, the peaks of the following were too overlapped for reliable fitting: I187 /

L210 / S223, D217 / I293, N235 / L283, V198 / S200 / E224 / S248, and M246 / L264. In the CLV1 pT868-bound state of KI-FHA, the spectral overlap remains in the first two peak clusters listed. With the 0.33 to 0.6 mM KI-FHA concentrations employed, particularly high S/N was assured by detection with a cryogenic probe at 600 MHz and a cryogenic probe for some of the 500 MHz data.

High backbone  $^{15}\text{N}\{\text{H}\}$  NOEs indicate that KI-FHA is generally rigid on the psec timescale in free and CLV1 pT868-bound states (Figure 1a, Figure 2a and Table 2). Negative NOE values and low  $R_2$  values (Figure 1a,c) at the termini indicate their very high mobility and that the folded domain extends from KAPP residue Ser180 through Ser295.  $^{15}\text{N}\{\text{H}\}$  NOEs, especially at 600 MHz, lower than 0.65 suggest such residues to be enriched in fluctuations on the psec to nsec scale. In free KI-FHA, such residues with higher amplitude fast motions are found in the 2/3 loop, 3/4 loop, 4/5 loop, and the long 8/9 loop (Figure 1a). Elevated  $R_1$  values were shown to correlate with nsec internal fluctuations (67). In the free state, there are residues in all the loops and termini that have higher-than-average  $R_1$  values, suggesting they may undergo nsec fluctuations. In the pT868-bound state of KI-FHA, all the loops show less of this trend of elevation of  $R_1$  (Figure 2b).

Sites of high  $R_2$  can suggest fluctuations on the  $\mu\text{sec}$  to msec timescale. In the free state, residues with at 600 MHz  $R_2$  exceeding  $13.8 \text{ s}^{-1}$ , i.e.  $> 1.5 \sigma$  above the coarse-filtered  $\bar{R}_2 = 12.6 \text{ s}^{-1}$  (Table 2), came under consideration for conformational exchange broadening processes on the  $\mu\text{sec}$  to msec scale. These residues lie in the 1/2  $\beta$ -hairpin, the 3/4 loop,  $\beta$ 4, the 4/5 loop, the 7/8 loop,  $\beta$ 8, and 10/11  $\beta$ -hairpin (Figure 1c). (The term  $\beta$ -hairpin refers to both anti-parallel strands and the intervening loop). Relaxation-compensated CPMG  $R_2$  measurements comparing averaged  $R_2$  between faster and slower pulsing of CPMG  $\pi$  pulses can exaggerate the line broadening of msec-scale chemical exchange processes (51). Elevated  $\Delta R_2 = R_2(7.5 \text{ ms}) - R_2(2 \text{ ms})$  in Figure 1d clearly suggest msec exchange broadening to occur in the 1/2  $\beta$ -hairpin, the 4/5 loop,  $\beta$ 8, the 9/10 loop, 10/11  $\beta$ -hairpin, and C-terminus. Most of these residues evidently with msec-scale motions are found in a swath across the six-stranded  $\beta$ -sheet, i.e. its  $\beta$ 1,  $\beta$ 2,  $\beta$ 10, and  $\beta$ 11 strands. Similar inspection of elevated 600 MHz  $R_2$  values with CLV1 pT868-bound present at 3.5-fold excess suggest the same segments of bound KI-FHA to undergo exchange broadening, plus the 6/7 recognition loop (Figure 2c). For the bound state, elevated  $\Delta R_2 = R_2(7.5 \text{ ms}) - R_2(2 \text{ ms})$  in Figure 2d is very similar to that of the free state (Figure 1d), including the patterns of msec exchange broadening in the 1/2  $\beta$ -hairpin, 10/11  $\beta$ -hairpin, and C-terminus.

### **$J(\omega_N)$ reduced spectral density evidence of nsec motions in KI-FHA**

Reduced spectral density fitting of the relaxation data (39,40) gives temporal insight into the backbone mobility of free and bound KI-FHA. Elevated  $J(\omega_N)$  values, where  $\omega_N = 60.8 \text{ MHz}$ , in free KI-FHA (Figure 3a) suggest that all of its loops and its termini have sites enriched in nsec-scale fluctuations.  $J(\omega_N)$  values of pT868-bound KI-FHA are lower overall with fewer elevated excursions (Figure 3a). Much of the general trend to lower  $J(\omega_N)$  values in the bound state may be a consequence of its slower tumbling and higher basal  $J_{eff}(0)$  (not shown). In the pThr peptide-bound state, however, locally more attenuated  $J(\omega_N)$  values suggest that all of the loops have residues with nsec fluctuations apparently quenched (Figure 3a). The following  $\beta$ -strands have residues with apparently fewer nsec fluctuations: especially  $\beta$ 1, but also one to two residues in each of  $\beta$ -strands 6 through 11 (Figure 3a).

### **Fluctuations in the loops of free KI-FHA are corroborated by deviations of $^{15}\text{N}-\text{H}$ residual dipolar couplings**

Residual dipolar couplings (RDCs) can respond to motional averaging over a wide, continuous range of timescales from psec to tens of msec (42,43), an advantage over spin relaxation studies.

Deviations of measured RDCs from RDCs predicted from the alignment of the protein structure can result from static or dynamic discrepancy from the protein structural coordinates (42,43). All  $^1\text{D}_{\text{NH}}$  RDCs were fitted to the whole of each of the ten lowest energy structures from the KI-FHA ensemble (PDB code 1mzk) using PALES (68). Deviations of individual  $^1\text{D}_{\text{NH}}$  RDCs from those predicted by fits using singular value decomposition (SVD) are represented in Figure 3b as a  $\chi^2$ -like quantity that is normalized by the  $D_a$  description of the range of observed RDCs. Several loops have  $^1\text{D}_{\text{NH}}$  RDCs that deviate from RDCs predicted by the structure, where L is for loop: L1/2, L2/3, L4/5, L5/6, L6/7, L7/8, L8/9, and L10/11 loops.  $\beta$ 8 and the C-terminus also have residues with deviating  $^1\text{D}_{\text{NH}}$  RDCs (Figure 3b). A majority of the sites of  $^1\text{D}_{\text{NH}}$  deviation have higher-than-average RMSD in the ensemble of structures. Most cases of  $(^1\text{D}_{\text{NH}, \text{meas}} - ^1\text{D}_{\text{NH}, \text{calc}})/D_a$  deviations in the free state (Figure 3b) have elevated  $J(\omega_N)$  evidence of nsec-scale motions (Figure 3ba). The  $(^1\text{D}_{\text{NH}, \text{meas}} - ^1\text{D}_{\text{NH}, \text{calc}})/D_a$  parameter appears to respond independently to many but not all possible sites of apparent nsec scale motion.

We also used the method of ref (44) to fit successive quintets of  $^1\text{D}_{\text{NH}}$  RDCs to the atomic coordinates of their respective fragments of the KI-FHA structural model. Deviations of molecular segments' alignment tensors from the overall protein alignment may result from dynamic or static deviations (43). The following segments of the backbone deviated beyond uncertainty from prediction from the coordinates using weak protein alignment in two different media: Glu185 – Gln195 that span the 1/2 hairpin loop, Ala197 to Lys209 in the 2/3 loop to  $\beta$ 3, Ala230 – Trp234 in  $\beta$ 5, Thr237 – Trp241 in the 5/6 loop to  $\beta$ 6, and Asp263 to Trp269 in the 8/9 loop (see Figure S1 and description in Supporting Information). All of these deviant segments, except for  $\beta$ -strand 5, are enriched in motions on the scales of either  $\mu\text{sec}$  – msec or nsec (Figure 1d, Figure 3a, Figure 4).

### Hydrodynamics of KI-FHA, free and bound to CLV1 pT868 peptide

Rotational correlation time  $\tau_m$  (overall tumbling) and rotational diffusion anisotropy can be estimated from  $R_2/R_1$  ratios (and atomic coordinates) using rigid residues where the  $R_2/R_1$  ratio depends only on  $\tau_m$  in the limit of small amplitude fast internal motions ( $\tau_e \ll \tau_m$ ) (69). An attractive two-stage procedure for identifying such residues without discarding  $R_2/R_1$  ratios made high or low by anisotropy of diffusion is that of ref (57). The results of the coarse filtering stage using those authors' NormaDyn software with the 500 and 600 MHz data collected for free and bound states are presented in Table 2. Typically about 80 candidate residues having acceptably high  $^{15}\text{N}\{^1\text{H}\}$  NOE and acceptably moderate  $R_2$  remained after coarse filtering. The purpose of the subsequent fine filtering stage is to remove additional residues poorly fit by simple model-free expressions 1 and 2 (Table 1) due to motions on the nsec or  $\mu\text{sec}$  to msec scales (57). We enhanced the fine filtering stage by removing (1) residues with msec scale exchange broadening according to relaxation dispersion (Figure 1d, Figure 2d), (2) residues likely to have nsec motion due to  $J(\omega_N) \geq \overline{J(\omega_N)} + 1\sigma$  where  $\omega_N = 60.8$  MHz (Figure 3a), or (3) residues requiring the exchange broadening term  $R_{\text{ex}}$  in preliminary model-free simulations according to statistical model selection. Around 40 residues survived this enhanced fine filtering (Table 2).

The trustworthy, fine-filtered data from 600 MHz with cryogenic probe were fitted to the NMR structure of free KI-FHA (PDB accession code 1MZK) to assess correlation time and anisotropy of rotational diffusion using the TENSOR2 program of ref (56). The ratio of the principal moments of inertia of free KI-FHA is 1.00:0.91:0.55. The free state is estimated to have  $\tau_m$  of  $7.57 \pm 0.08$  nsec with  $D_{||}/D_{\perp} = 0.81 \pm 0.07$  (oblate) and the pT868-bound state to have  $\tau_m$  of  $10.70 \pm 0.07$  with  $D_{||}/D_{\perp} = 1.21 \pm 0.06$  (prolate). (Note that a prolate to oblate shift was reported for peptide binding to an SH3 domain (13).) For the free and bound states, both oblate and prolate axially symmetric fits are acceptable with high statistical confidence and

have  $\chi^2$  similar to within 10%. Such two-minimum behavior suggests the possibility of fully anisotropic diffusion (70). Indeed, a fully anisotropic diffusion model is statistically significantly better for the bound state. The limitation to axial symmetry of Modelfree 4.1 used here and widely could add a very modest systematic bias, but little difference in  $S^2$  values results from switching between oblate and prolate models for KI-FHA (not shown).

These  $\tau_m$  estimates for KI-FHA at 25 °C are comparable to the  $\tau_m$  values of monomers of similar size and temperature (33,71,72). The unlabeled free state of KI-FHA is 124 residues and 13.3 kD by ESI-MS while the complex is 12% larger at 137 residues and 14.9 kD. Saturation with the 1.6 kDa, 13-mer pT868 peptide increases  $R_2/R_1$  more than expected (Table 2). The resulting increase in apparent Stokes radius with the pT868 peptide bound is around 9%. Why does the increase in  $\tau_m$  upon saturation with the 1.6 kDa pT868 peptide appear to be ~1.5 nsec more than expected? To investigate whether peptide binding promotes self-association, a complex of 0.53 mM KI-FHA with 1.5-fold excess of pT868 peptide was diluted 2.3-fold, resulting in perhaps a 4% average drop in  $R_2$  values (Table 2). This decrease parallels the decrease in pT868 peptide saturation from 94% at 0.53 mM KI-FHA to 88% at 0.23 mM KI-FHA, i.e. the enhanced weighting of the free state contribution to apparent  $\tau_m$  when diluted. If self-association were substantial, a larger drop in  $R_2$  and estimated  $\tau_m$  ought to have accompanied 2.3-fold dilution. A more likely alternative explanation could be that the greater extent of nsec fluctuations of the loops in the free state may decrease its effective Stokes radius relative to the bound state. (Note that a case of greater slowing of rotational diffusion upon peptide-binding to a monomer has been reported, where coupling between altered internal motion and diffusional shape seems larger (73).)

### Spectral density function selection for model-free simulations

Model-free simulations of the relaxation data proceeded with hydrodynamics parameters  $\tau_m$  and  $D_{\parallel}/D_{\perp}$  fixed at the values listed in Table 2. Use of relaxation data from two magnetic fields in the model-free approach more reliably defines the  $S^2$ ,  $R_{ex}$  (excess line broadening), and  $\tau_e$  (internal correlation time) parameters. Data at two magnetic fields also provides enough measurements to use three-parameter fits where needed. Five measurements were used for both the free and bound states of KI-FHA, namely 600 MHz  $R_1$ ,  $R_2$ , and NOE and 500 MHz  $R_1$  and NOE. Omission of 500 MHz  $R_1$  gave very similar results. The spectral density functions (Table 1) most appropriate for use in model-free simulations were chosen primarily using AIC-based statistics (61) augmented by other experimental identification of msec motion from relaxation dispersion parameter  $\Delta R_2(\tau_{cp})$  (ref 51) (Figure 1d, Figure 2d) and of nsec motion from the reduced spectral density function  $J(\omega_N=60.8 \text{ MHz})$  (ref 41) (Figure 3a), as described in Materials and Methods. The spectral functions selected from among the choices of Table 1 are listed in Table S1 of Supporting Information, by residue number for free and bound states of KI-FHA.

### General model-free outcomes for free and bound states

Excluding the termini,  $S^2$  values appear to average 0.902 and 0.946 for the free and pT868-bound states of KI-FHA, respectively. For the non-terminal residues compared in both states,  $\Delta S^2_{\text{bound-free}}$  averages 0.042 with a median  $\Delta S^2_{\text{bound-free}}$  of 0.036 and average uncertainty of 0.037. These suggest that phosphoThr binding results in enhanced overall rigidity on psec to nsec scale. Model-free  $\tau_e$  values (Figure 4c) are joined by elevation of both  $J(\omega_N)$  (ref 41) (Figure 3a) and  $R_1$  (ref 67) (Figure 1b) in suggesting nsec motions in all of the loops and termini of free KI-FHA except for the 1/2 loop. Fewer of these residues in the CLV1 pT868-bound state of KI-FHA show such evidence of nsec motions (Figure 2b, Figure 3a, Figure 5c, Figure 7), suggesting that peptide binding quenches nsec motions globally. Where the loops are enriched with nsec scale fluctuations, especially in the free state, the  $S^2$  values can be regarded as overestimates, as established by methods that account for coupling of internal nsec motion

with rotational diffusion (74, 75). The relaxation data and model-free results for KI-FHA, free and bound to CLV1 pT868, were deposited under BMRB accession codes 5841 and 6474, respectively, at [www.bmrb.wisc.edu](http://www.bmrb.wisc.edu).

### **pThr peptide binding surface and its slower dynamics**

Titration of  $^{15}\text{N}$ -enriched KI-FHA with the CLV pT868 peptide shifted the amide peak positions of loops on one edge of the  $\beta$ -sandwich of KI-FHA. Significant amide chemical shift changes occur at Leu210 - Val213 of the 3/4 loop; Lys221 - Ser223, Val225, and Gly227 - His229 of the 4/5 loop; Met246 - Asn250 and Thr252 of the 6/7 loop; Asp263, Gly265, Arg267, and Trp269 of the 8/9 loop; and Gly284 - Thr287 of the 10/11 loop (Figure 6c,d). Nsec-scale motions characterize the *free* state of several residues quite perturbed by the pT868 peptide: Arg212, Val213, Ser223, Val225, Lys228, Leu249, Asn250, Arg267, Thr285, and Thr286 (Figure 3a, Figure 4c). Much slower conformational fluctuations over  $\mu\text{sec}$  to msec are shared by the free and bound states of these pT868-perturbed residues: Val213, Ser216, Lys228, Arg267, and Thr286 (Figure 4b, Figure 5b). Exchange broadening  $> 1 \text{ s}^{-1}$  suggests that fluctuations in the  $\mu\text{sec}$  to msec regime also occur at Ser223 and Glu224 in the free state (Figure 4b) and at His229 and Gly251 in the bound state (Figure 5b).

### **Binding effects on flexibility on fast timescale in recognition interface**

Largely conserved recognition loop residues shown to be essential for binding phosphorylated receptor kinase domain (22) are clearly rigid. Their apparent  $S^2$  values in free / bound states are: 0.99 / 0.999 for Gly211, 0.92 / 0.92 for His229, and 0.96 / 0.99 for Asn250 (Figure 4a, Figure 5a). Binding appears to quench nsec fluctuations of Asn250 (Figure 3a, Figure 4c, and Figure 5c). Near the five most conserved residues in the recognition loops are non-conserved residues with greater mobility and significant changes of  $S^2$  upon binding of the pT868 peptide from CLV1. Fluctuations in the free state of greater-than-average amplitude are suggested by apparent  $S^2 \leq 0.8$  of residues of the 3/4, 4/5, and 8/9 recognition loops: Ser216, Asp222, Glu224, Lys228, Ser260, and Ser266 (Figure 4a). Lys228 and Glu224 of the 4/5 loop, Ser266 of the 8/9 loop, as well as Ser216 and Val213 of the 3/4 loop undergo striking increases in rigidity on the psec-nsec scale upon pT868 peptide binding (Figure 6). Leu220, Asp263, and Leu283 of these recognition loops are rigidified to a smaller degree. Also at the interface, Gly251 of the 6/7 loop, Arg267 of the 8/9 loop, and T286 of the 10/11 loop in contrast become more flexible on the fast time scale in the complex (Figure 6a,b).

### **Long-range effects of binding on fast timescale motion**

For a number of residues outside this binding surface for phosphorylated RLKs (Figure 6d), mobility of KI-FHA on the psec to nsec scale changes upon binding the pT868 peptide from the CLV1 RLK. In the loops of the 1/2 and 10/11  $\beta$ -hairpins pointing toward the interface, residues where binding increases the amplitude of psec-nsec motions (drop in  $S^2$ ) are Ala191 and Thr286 (Figure 6a,b). In the loop of the 7/8  $\beta$ -hairpin distant from the interface, binding may have increased the fast motion of Ser256 (Figure 6a). By contrast, pT868 peptide binding appears to have increased rigidity in  $\beta$ -strands of all four of the  $\beta$ -hairpins. This is especially evident in both strands of the 1/2  $\beta$ -hairpin, both strands of the 5/6  $\beta$ -hairpin,  $\beta$ 8 of the 7/8  $\beta$ -hairpin, and more subtly in both strands of 10/11  $\beta$ -hairpin (Figure 6a,b). The apparent changes in  $S^2$  in the highly disordered termini are unlikely to be significant since the termini lack the globular diffusion behavior needed for accurate description by the model-free approach (Figure 6a). The 2/3 loop and 9/10 loop each cross between the two  $\beta$ -sheet layers of the sandwich and are most distant from the binding surface. The peptide binding effect on the 2/3 loop includes mobilization of Ser202 and increased rigidity of Ser203 and Leu206 (Figure 6a). The phosphoThr peptide binding rigidifies Leu275-Asp278 of the remote 9/10 loop.

## Binding effects on flexibility on slow timescale

Exchange broadening suggests motions on the  $\mu$ sec to msec timescale for around 49 residues in the free state and 51 residues in the bound state of KI-FHA, by relaxation dispersion (Figure 1d, Figure 2d) or statistical need for  $R_{\text{ex}}$  term in model-calculations (Figure 7). The slow fluctuations affect a contiguous swath of residues across  $\beta$ 2,  $\beta$ 1,  $\beta$ 11, and  $\beta$ 10 of the six-stranded sheet (appearing left to right in Figure 7, Figure 8). Adjacent to this central swath are other clusters of slowly exchanging residues. At the edge of the five-stranded sheet,  $\beta$ 4 and the 3/4 and 4/5 recognition loops are affected (Figure 7a). At the extreme opposite edge of the six-stranded sheet, residues of  $\beta$ 8, the 7/8 loop, and the 8/9 loop manifest motions on the  $\mu$ sec to msec scale (Figure 7a). Also exchanging are residues in the 1/2 and 10/11 loops near the binding site, the faraway 2/3 and 9/10 loops, and C-terminus. With the CLV1 pT868 peptide present at 3.5-fold excess, the peptide is expected to saturate 98% of KI-FHA based on the  $K_D$  of 20  $\mu$ M. At 98% saturation, Gly265 and Lys268 -Gly270 of the 8/9 peripheral recognition loop and Gly251 of the 6/7 recognition loop have exchange broadening (Figure 4b, Figure 5b, and Figure 7). Asp222, Gly227, and Thr285, with peaks most shifted and broadened during titration with the pT868 peptide (Figure 6a), are too broad in the peptide-saturated state for relaxation fitting.

## Dissociation of CLV1 pT868 peptide from KI-FHA

Intermediate exchange broadening during titration with the pT868 peptide can be exploited to estimate the exchange rate  $k_{\text{ex}}$ . The off-rate  $k_{\text{off}}$  is equivalent to  $k_{\text{ex}}$ , since  $k_{\text{on}}$  is not at all limiting, being of the order of  $\sim 5 \cdot 10^7 \text{ M}^{-1}\text{s}^{-1}$ . The range of  $k_{\text{ex}}$  and  $k_{\text{off}}$  possible is defined by the binding-induced chemical shift changes of Gly227 of  $\Delta v_H = 569 \text{ Hz}$  and Thr285 of  $\Delta v_H = 356 \text{ Hz}$  at 600 MHz; spectral and thermodynamic details will be published elsewhere. Gly227 in the slow-intermediate exchange regime limits  $k_{\text{ex}}$  to the range of  $2.2 * \Delta v_{H,G227} > k_{\text{ex}} > 0.01 * \Delta v_{H,G227}$ , i.e.  $1250 \text{ s}^{-1} > k_{\text{ex}} > 5.7 \text{ s}^{-1}$ . Thr285 in the fast-intermediate exchange regime limits  $k_{\text{ex}}$  to this range:  $100 * \Delta v_{H,T285} > k_{\text{ex}} > \Delta v_{H,T285}$ , i.e.  $35,600 \text{ s}^{-1} > k_{\text{ex}} > 356 \text{ s}^{-1}$ . Combining these two ranges, leads to the conservative estimate that  $k_{\text{off}}$  and  $k_{\text{ex}}$  lie between  $356 \text{ s}^{-1}$  and  $1250 \text{ s}^{-1}$  for the pT868 interaction with KI-FHA. When pT868 is present at 1.5-fold excess conferring 94% saturation of KI-FHA, additional exchange broadening is seen in recognition loops in a comparison of 600 MHz  $R_2$  values at 94% and 98% saturation. This is most evident in the 6/7 loop, followed by the 8/9 loop, the 4/5 loop, and conserved Gly211 (Figure S2 of Supporting Information). A set of model-free results were obtained from relaxation measurements at the 1.5-fold excess of pT868 that results in ~94% saturation of KI-FHA. At 94% saturation relative to 98% saturation, more residues in recognition loops manifest chemical exchange broadening, namely at 6/7 loop residues Met246, Leu249, Asn250, and Thr252, as well as conserved Gly211 and Asp263 and Ser266 of the 8/9 loop (Figures S2, S3 of Supporting Information). Since each of these residues undergo chemical shift changes upon titration with the pT868 peptide (Figure 6a), it appears likely that their broadening at 94% saturation results from the peptide's apparent off-rate of between 350 and  $1250 \text{ s}^{-1}$ .

## DISCUSSION

### Rigidity of residues conserved in the recognition site of FHA domains

Along the edge of the five-stranded  $\beta$ -sheet, conserved Gly211, Arg212 (3/4 loop), Ser226, His229 (4/5 loop), and Asn250 (6/7 loop) (KAPP numbering) are very rigid (Figure 4a, Figure 5a) and critical for forming the phosphoprotein binding site of FHA domains; the arginine, serine, and asparagine contact the phosphoThr of the ligand (3, 5–7, 22, 25, 26). The high rigidity of Gly211 and Arg212 of the 3/4 loop and His229 of the 4/5 loop (Figure 4, Figure 5) is consistent with the structure-stabilizing role proposed for conserved Gly211 and His229 (KAPP numbering) (3). The unique conformation of absolutely conserved Gly211 could be important for positioning Arg212 to interact with the phosphate of the partner. Gly211 and

Arg212 being recessed and largely buried under the interaction surface may confer their rigidity. When the  $S^2$  values of Gly211, Arg212, His229, and Asn250 are interpreted with a diffusion-in-a-cone model of motion (36), the estimated amplitudes of N-H bond reorientations of their psec – nsec have cone semi-angles of  $5^\circ / 2^\circ$ ,  $11^\circ / \text{NA}$ ,  $14^\circ / 13^\circ$ , and  $10^\circ / 4^\circ$ , for the free / bound states respectively. The backbone rigidity of Gly211, Arg212, His229, and Asn250 in the free state decreases energetic costs of loss of configurational entropy upon binding.

### Non-conserved residues of the phosphoprotein-binding surface

Comparison of FHA domain structures of divergent sequence (3,76) suggested that KI-FHA residues Glu224 in the 4/5 loop, Leu249 in the 6/7 loop, and Thr285 in the 10/11 loop may contribute to the key pT+3 site and KI-FHA's distinctive phosphopeptide specificity (7). In yeast Rad53 FHA domains, two more residues appear to contact the pT+3 peptide position (77); the first is equivalent to Gly284 and the second possibly to Thr286 of the 10/11 loop of KAPP. The amide peaks most shifted upon addition of the CLV1 pT868 peptide (with radial  $\Delta\omega_{\text{HN}} > 0.15$  ppm; Figure 6c) suggest a broad phosphoThr peptide recognition surface (Figure 6d). This same broad phosphoprotein (RLK)-binding surface is suggested by Evolutionary Trace analysis and extensive conservation among plant KAPPs on its flanks (7). Most residues of this surface are not, however, widely conserved among FHA domains (7). These non-conserved residues of the recognition loops can be placed in three groups of psec-nsec mobility ( $S^2$ ). The first group is very rigid in both free and pT868-bound states: Lys221, Gly227, Leu249, Trp269, and Gly284. The second group is more flexible in the free state, undergoes restriction in the bound state, and is listed here with estimates of wobbling-in-cone semi-angles (see eq. 5) for free / bound states in parentheses: Val213 ( $18^\circ / 8^\circ$ ), Ser216 ( $35^\circ / 23^\circ$ ), Glu224 ( $28^\circ / 17^\circ$ ), Lys228 ( $34^\circ / 10^\circ$ ), and Ser266 ( $27^\circ / 13^\circ$ ). – These are from solvent-exposed portions of the 3/4, 4/5, and 8/9 loops, with the central 4/5 loop seeming to be in especially intimate contact with the phosphopeptide (Figure 6). The third group at the edge of the recognition surface has increased mobility in the pT868-bound state and is listed with estimated cone semi-angles: Gly251 ( $5^\circ / 16^\circ$ ) of the 6/7 loop, Arg267 ( $21^\circ / 26^\circ$ ) of the 8/9 loop, and Thr286 ( $18^\circ / 25^\circ$ ) of the 10/11 loop.

### Rigidity and role in affinity of residues at the pT+3-site and conserved neighbors

Rigid portions of the phosphoprotein-binding surface could confer affinity to this FHA domain and others. This hypothesis derives from observations on modules binding pTyr peptides: First, the residues of a PTB domain that contact a relevant pTyr peptide are rigid, presumably promoting enthalpically favorable van der Waals contacts (78). Second, in the association of SH2 domains with pTyr peptides, high rigidity is characteristic of sites conferring affinity (12). Clearly, mostly conserved Gly211, Arg212, Ser226, His229, and Asn250 (KAPP numbering) are rigid in the free state (Figure 4) and apparently essential for affinity for receptor-like kinase domains (22). This behavior is consistent with the studies of pTyr peptide recognition. The pT+3 position of peptide ligands is the most important determinant of their affinity for FHA domains (3,79). The non-conserved residues that divergent FHA domains place around the pT+3 site appear to be important for affinity (77). These residues of KI-FHA of KAPP, namely Gly227, Leu249, Gly284, and Thr285 are rigid (Figure 4,Figure 5), consistent with the theme of high rigidity of residues conferring affinity (12,78).

### Significance of pThr peptide binding-dependent flexibility changes remote from RLK binding site

Energetic costs of pThr peptide binding include the loss of conformational entropy of the peptide, at sites in the 3/4, 4/5, and 8/9 recognition loops, at sites in seven  $\beta$ -strands, and in three other loops (1/2, 2/3, and 9/10 loops) (Figure 6a,b). In the free state, there appears to be more nsec-scale motion than in the bound state in all five recognition loops, other loops, and

$\beta$ 8 (Figure 3a, Figure 4c, Figure 5c). Since  $S^2$  is more likely to be underestimated at these locations (74, 75) in the *free* state, the binding-linked increases in rigidity and the entropic costs at these recognition loop residues could be underestimated. The modest increases in rigidity of seven of the 11  $\beta$ -strands may not represent just costs in configurational entropy. The binding-dependent increases in  $S^2$  appear to occur across networks of hydrogen-bonded residues across both  $\beta$ -sheets; one such network includes  $\beta$ 3– $\beta$ 5– $\beta$ 6– $\beta$ 9 while another network includes  $\beta$ 2– $\beta$ 1– $\beta$ 11– $\beta$ 10 (Figure 6b). The increased rigidity seen in both  $\beta$ -sheets suggests the possibility of improved hydrogen-bonding and side chain packing or van der Waals contacts, for a favorable change in the enthalpy of the bound state. This could provide part of the compensating favorable enthalpy driving the association of pThr peptides to KI-FHA (Ding et al., to be published) and to another FHA domain (3). Peptide binding was also observed to rigidify and stabilize an SH3 domain both near and far from the interface(13). Precedent for enthalpic stabilization at long-range was suggested by antibody binding to lysozyme increasing its hydrogen exchange protection to the distal side (80). Sites of binding-enhanced fast motions of KI-FHA seen in the 1/2, 2/3, 6/7, 7/8, 8/9, and 10/11 loops provide favorable increases in conformation entropy to compensate the entropic costs listed above.

### Flexibility of recognition surface in view of breadth of specificity

Several non-conserved residues of the recognition loops are more flexible than average: Ser216 (3/4 loop); Asp222, Glu224, Lys228 (4/5 loop); Ser260, Gly265 - Arg267 (8/9 loop); and Thr286 (10/11 loop). The mobility of these in the free state appears to be on the nsec scale (Figure 3a, Figure 4c). The studies of SH2 domain affinity for pTyr peptides suggest that flexibility correlates well with contact sites of little importance to affinity. -- Mobility on the psec to nsec scale in a recognition surface may decrease the contribution of the mobile groups to affinity by diminishing favorable, strongly distance-dependent van der Waals interactions (12). Flexibility in the phosphoprotein-binding surface of KI-FHA and other FHA domains might confer their breadth of specificity. An example of such flexibility could be the 8/9 loop of KI-FHA of KAPP that distinguishes it from the phosphoprotein-binding surface of other FHA domains. This 8/9 loop of KI-FHA is clearly longer, more solvent-exposed, and mobile on the nsec scale. The prominence and mobility of the 8/9 loop may be important in KAPP's recognition of several RLK targets in plants. Arabidopsis KAPP interacts in a phosphorylation-dependent manner with the following receptor-like kinases critically important to plant development and defense against infection: CLV1 (81), HAESA (82), WAK1 (18), BAK1 (20), SERK1 (21), and FLS2 (19). KAPP being promiscuous enough to attenuate multiple RLK-dependent signaling pathways in plants is supported moreover by KAPP being a single gene product without any known paralog of overlapping function (17). Yet, KAPP failed to interact with a number of kinases tested (17). The psec - nsec mobility of the RLK-binding surface of KI-FHA could foster the limited diversity of RLK partners recognized by KAPP. Perhaps an analogy could be drawn with the cytokine IL-2 where the flexible portion of its ligand-binding surface binds a far greater diversity than does the rigid portion (83). Both the psec-nsec flexibility and binding-inducible  $\mu$ sec to msec mobility of the recognition loops could enable KI-FHA of KAPP to adjust to a minority of the 417 RLKs encoded by the Arabidopsis genome (84).

### Significance of $\mu$ sec to msec fluctuations of 6-stranded $\beta$ -sheet remote from phosphoprotein binding surface

Slow fluctuations of  $\beta$ 2,  $\beta$ 4, and  $\beta$ 8 on the edges of the  $\beta$ -sandwich of KI-FHA may simply result from their solvent exposure. The largest line broadening evidence of fluctuations on the  $\mu$ sec to msec scale occurs at Ser256 of the 7/8 loop (Figure 4b, Figure 5b, Figure 7). Neighboring residues of the 7/8  $\beta$ -hairpin,  $\beta$ 10, and the 9/10 loop also display line broadening evidence of conformational exchange on the  $\mu$ sec to msec scale (Figure 4b, Figure 5b, and Figure 7). This patch of slowly fluctuating residues coincides with a potential interaction

surface predicted by Evolutionary Trace analysis to include many of the same residues of the  $\beta$ 7/8 hairpin,  $\beta$ 10, and the 9/10 loop (7). It is unlikely that a phosphoprotein partner can wrap 80° around from the phosphorecognition surface to this face of FHA domains. The  $\beta$ 7/8 - L9/10 -  $\beta$ 10 patch coincides with the site of segment-swapped dimerization in crystals of the FHA domain of human Chfr (6). (In concentrated solutions of the Chfr FHA domain, the dimer is a minor form relative to the predominant monomer (6).) Perhaps the slow conformational fluctuations of the  $\beta$ 7/8 - L9/10 -  $\beta$ 10 patch facilitate rearrangements such as the straightening of the 7/8  $\beta$ -hairpin of the segment-swapped Chfr FHA dimer. The question has been raised as to whether this independent patch could be a site of dimerization of FHA domains more generally (7). An alternative postulate regarding the patch is that it might play a role in self-recognition of another part of the same protein chain or signaling assembly (7). Slow fluctuations over  $\mu$ sec to msec have been observed at a number of protein-protein interaction sites (27–29, 33, 85).

A swath of contiguous residues across the six-stranded sheet undergo exchange broadening of their amide lines (sheet in foreground of Figure 7). Most of these have interior side chains arrayed in two packed rows across  $\beta$ 2,  $\beta$ 1,  $\beta$ 11, and  $\beta$ 10, plus  $\beta$ 4 of the other sheet (Figure 8). Among these nine side chains, four—Leu184, Ile281, Val289, and Val291— are among a dozen hydrophobic positions characteristic of the folding core of all FHA and SMAD MH2 domains (7). The upper row comprises Leu218 of  $\beta$ 4, Leu194 of  $\beta$ 2, Val186 of  $\beta$ 1, Val289 of  $\beta$ 11, and Ile281 of  $\beta$ 10, progressing left-to-right across Figure 8. The lower row comprises His196 of  $\beta$ 2, Leu184 of  $\beta$ 1, Val291 of  $\beta$ 11, and Asp279 of  $\beta$ 10. The tight packing of this cluster of interior side chains plus the inter-strand hydrogen bonds might coordinate the backbone motions on the  $\mu$ sec to msec scale.

To recap, the conformational fluctuations of the loops on the nsec scale in the free state of KI-FHA are diminished when the pT868 peptide fragment of the CLAVATA1 receptor kinase partner is bound (Figure 3a). A network of residues across the six-stranded  $\beta$ -sheet undergoes conformational fluctuations on the  $\mu$ sec to msec scale independent of binding state (Figure 8). Microsec to msec scale fluctuations are found at in a patch of residues at  $\beta$ 7/8 - L9/10 -  $\beta$ 10 corresponding to the site of crystallization-promoted, segment-swapped dimer formation in the FHA domain of human Chfr (Figure 7). With the phosphoThr peptide bound, more sites in the 6/7 and 8/9 recognition loops display evidence of  $\mu$ sec to msec scale motion (Figure 5, Figure 7). Most of the broadening in the 6/7 loop seems to result from the peptide's off-rate of several hundred per sec. Peptide binding-enhanced rigidity of ten or more residues of recognition loops and adjoining 1/2 loop is compensated by flexibility increases on the fast timescale of six residues on the periphery of the active site (Figure 6). The net overall increase in rigidity of KI-FHA phosphoThr peptide bound, with average  $\Delta S^2=0.04$ , includes increased rigidity at sites in seven  $\beta$ -strands (Figure 6); this suggests the possibility of long-range enhancement of favorable enthalpy to compensate entropic costs of binding. Favorable binding enthalpy is likely from the backbone rigidity at the conserved residues clearly important to affinity and from neighboring residues most likely to contact the key pT+3 position that influences affinity.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank NMRFAM (Madison, WI) for access to the cryoprobe™-equipped Bruker DMX-500. We thank Professors Palmer, Marion, Zuiderweg, Bax, and Nicholson for software used in this study.

This research was supported by NSF grant MCB0111589 to SRV. The Varian Inova 600 and Bruker DRX-500 spectrometers were funded in part by NSF grants DBI0070359 and CHE8908304. NMRFAM is supported by the National Center for Research Resources.

## ABBREVIATIONS

FHA domain	Forkhead-associated domain
HPLC	high pressure liquid chromatography
KAPP	kinase-associated protein phosphatase
LC-ESI-MS	liquid chromatography-electrospray ionization mass spectrometry
NOE	nuclear Overhauser effect
RDC	residual dipolar coupling
RLK	receptor-like kinase of plants
R <sub>1</sub>	spin-lattice or longitudinal relaxation rate
R <sub>2</sub>	spin-spin or transverse relaxation rate
SVD	singular value decomposition

## REFERENCES


1. Durocher D, Jackson SP. The FHA domain. *FEBS Lett* 2002;513:58–66. [PubMed: 11911881]
2. Liao H, Byeon IJ, Tsai MD. Structure and function of a new phosphopeptide-binding domain containing the FHA2 of Rad53. *J. Mol. Biol* 1999;294:1041–1049. [PubMed: 10588905]
3. Durocher D, Taylor IA, Sarbassova D, Haire LF, Westcott SL, Jackson SP, Smerdon SJ, Yaffe MB. The molecular basis of FHA domain:phosphopeptide binding specificity and implications for phospho-dependent signaling mechanisms. *Mol. Cell* 2000;6:1169–1182. [PubMed: 11106755]
4. Liao H, Yuan C, Su MI, Yongkiettrakul S, Qin D, Li H, Byeon IJ, Pei D, Tsai MD. Structure of the FHA1 domain of yeast Rad53 and identification of binding sites for both FHA1 and its target protein Rad9. *J. Mol. Biol* 2000;304:941–951. [PubMed: 11124038]
5. Li J, Williams BL, Haire LF, Goldberg M, Wilker E, Durocher D, Yaffe MB, Jackson SP, Smerdon SJ. Structural and functional versatility of the FHA domain in DNA-damage signaling by the tumor suppressor kinase Chk2. *Mol. Cell* 2002;9:1045–1054. [PubMed: 12049740]
6. Stavridi ES, Huyen Y, Loreto IR, Scolnick DM, Halazonetis TD, Pavletich NP, Jeffrey PD. Crystal Structure of the FHA Domain of the Chfr Mitotic Checkpoint Protein and Its Complex with Tungstate. *Structure* 2002;10:891–899. [PubMed: 12121644]
7. Lee GI, Ding Z, Walker JC, Van Doren SR. NMR structure of the forkhead-associated domain from the Arabidopsis receptor kinase-associated protein phosphatase. *Proc. Natl. Acad. Sci. U S A* 2003;100:11261–11266. [PubMed: 14500786]
8. Li H, Byeon IJ, Ju Y, Tsai MD. Structure of human Ki67 FHA domain and its binding to a phosphoprotein fragment from hNIFK reveal unique recognition sites and new views to the structural basis of FHA domain functions. *J Mol Biol* 2004;335:371–381. [PubMed: 14659764]
9. Ishima R, Torchia DA. Protein dynamics from NMR. *Nat. Struct. Biol* 2000;7:740–743. [PubMed: 10966641]
10. Stone MJ. NMR relaxation studies of the role of conformational entropy in protein stability and ligand binding [Review]. *Acc. Chem. Res* 2001;34:379–388. [PubMed: 11352716]
11. Atkinson RA, Kieffer B. The role of protein motions in molecular recognition: insights from heteronuclear NMR relaxation measurements. *Prog. Nucl. Magn. Reson. Spectrosc* 2004;44:141–187.
12. Kay LE, Muhandiram DR, Wolf G, Shoelson SE, Forman-Kay JD. Correlation between binding and dynamics at SH2 domain interfaces. *Nat. Struct. Biol* 1998;5:156–163. [PubMed: 9461082]

13. Wang C, Pawley NH, Nicholson LK. The role of backbone motions in ligand binding to the c-Src SH3 domain. *J. Mol. Biol* 2001;313:873–887. [PubMed: 11697910]
14. Stone JM, Colinge MA, Smith RD, Horn MA, Walker JC. Interaction of a protein phosphatase with an *Arabidopsis* Serine-Threonine Receptor Kinase. *Science* 1994;266:793–795. [PubMed: 7973632]
15. Stone JM, Trotochaud AE, Walker JC, Clark SE. Control of meristem development by CLAVATA1 receptor kinase and KAPP protein phosphatase interactions. *Plant Physiology* 1998;117:1217–1225. [PubMed: 9701578]
16. Williams RW, Wilson JM, Meyerowitz EM. A possible role for kinase-associated protein phosphatase in the *Arabidopsis* CLAVATA1 signaling pathway. *Proc. Natl. Acad. Sci. USA* 1997;94:10467–10472. [PubMed: 9294234]
17. Braun DM, Stone JM, Walker JC. Interaction of the maize and *Arabidopsis* kinase interaction doamins with a subset of receptor-like protein kinases: implication for transmembrane signaling in plants. *Plant J* 1997;12:83–95. [PubMed: 9263453]
18. Park AR, Cho SK, Yun UJ, Jin MY, Lee SH, Sachetto-Martins G, Park OK. Interaction of the *Arabidopsis* receptor protein kinase Wak1 with a glycine-rich protein, AtGRP-3. *J. Biol. Chem* 2001;276:26688–26693. [PubMed: 11335717]
19. Gómez-Gómez L, Bauer Z, Boller T. Both the extracellular leucine-rich repeat domain and the kinase activity of FLS2 are required for flagellin binding and signaling in *Arabidopsis*. *Plant Cell* 2001;13:1155–1163. [PubMed: 11340188]
20. Li J, Wen JQ, Lease KA, Doke JT, Tax FE, Walker JC. BAK1, an *Arabidopsis* LRR receptor-like protein kinase, interacts with BRI1 and modulates brassinosteroid signaling. *Cell* 2002;110:213–222. [PubMed: 12150929]
21. Shah K, Russinova E, Gadella TW Jr, Willemse J, De Vries SC. The *Arabidopsis* kinase-associated protein phosphatase controls internalization of the somatic embryogenesis receptor kinase 1. *Genes Dev* 2002;16:1707–1720. [PubMed: 12101128]
22. Li J, Smith GP, Walker JC. Kinase interation domain of kinase-associated protein phosphatase, a phosphoptotein-binding domain. *Proc. Natl. Acad. Sci. USA* 1999;96:7821–7826. [PubMed: 10393905]
23. Huse M, Muir TW, Xu L, Chen YG, Kuriyan J, Massague J. The TGF beta receptor activation process: an inhibitor- to substrate-binding switch. *Mol. Cell* 2001;8:671–682. [PubMed: 11583628]
24. Yuan C, Yongkiettrakul S, Byeon IJ, Zhou S, Tsai MD. Solution structures of two FHA1-phosphothreonine peptide complexes provide insight into the structural basis of the ligand specificity of FHA1 from yeast Rad53. *J. Mol. Biol* 2001;314:563–575. [PubMed: 11846567]
25. Wang P, Byeon I-JL, Liao H, Beebe KD, Yongkiettrakul S, Pei D, Tsai M-D. Structure and Specificity of the Interaction between the FHA2 Domain of RAD53 and Phosphotyrosyl Peptides. *J. Mol. Biol* 2000;302:927–940. [PubMed: 10993733]
26. Byeon IJ, Yongkiettrakul S, Tsai MD. Solution structure of the yeast Rad53 FHA2 complexed with a phosphothreonine peptide pTXXL: comparison with the structures of FHA2-pYXL and FHA1-pTXD complexes. *J. Mol. Biol* 2001;314:577–588. [PubMed: 11846568]
27. Wyss DF, Dayie KT, Wagner G. The counterreceptor binding site of human CD2 exhibits an extended surface patch with multiple conformations fluctuating with millisecond to microsecond motions. *Protein Sci* 1997;6:534–542. [PubMed: 9070436]
28. Feher VA, Cavanagh J. Millisecond-timescale motions contribute to the function of the bacterial response regulator protein SpoOF. *Nature* 1999;400:289–293. [PubMed: 10421374]
29. Huang K, Ghose R, Flanagan JM, Prestegard JH. Backbone dynamics of the N-terminal domain in *E. coli* DnaJ determined by <sup>15</sup>N- and <sup>13</sup>CO-relaxation measurements. *Biochemistry* 1999;38:10567–10577. [PubMed: 10441154]
30. Hare BJ, Wyss DF, Osburne MS, Kern PS, Reinhartz EL, Wagner G. Structure, specificity and CDR mobility of a class II restricted single-chain T-cell receptor. *Nat. Struct. Biol* 1999;6:574–581. [PubMed: 10360364]
31. Ye J, Mayer KL, Stone MJ. Backbone dynamics of the human CC-chemokine eotaxin. *J. Biomol. NMR* 1999;15:115–124. [PubMed: 10605085]
32. Lee AL, Kinnear SA, Wand AJ. Redistribution and loss of side chain entropy upon formation of a calmodulin-peptide complex. *Nat. Struct. Biol* 2000;7:72–77. [PubMed: 10625431]

33. Gao G, Semenchenko V, Arumugam S, Van Doren SR. Tissue inhibitor of metalloproteinases-1 undergoes microsecond to millisecond motions at sites of matrix metalloproteinase-induced fit. *J. Mol. Biol* 2000;301:537–552. [PubMed: 10926526]
34. Lipari G, Szabo A. Model-Free Approach to the Interpretation of Nuclear Magnetic Resonance Relaxation in Macromolecules. 1. Theory and Range of Validity. *J. Am. Chem. Soc* 1982;104:4546–4559.
35. Clore GM, Szabo A, Bax A, Kay LE, Driscoll PC, Gronenborn AM. Deviations from the simple two-parameter model-free approach to the interpretation of nitrogen-15 nuclear magnetic relaxation of proteins. *J. Am. Chem. Soc* 1990;112:4989–4991.
36. Lipari G, Szabo A. Model-Free Approach to the Interpretation of Nuclear Magnetic Resonance Relaxation in Macromolecules. 2. Analysis of Experimental Results. *J. Am. Chem. Soc* 1982;104:4559–4570.
37. Bracken C, Carr PA, Cavanagh J, Palmer AG 3rd. Temperature Dependence of Intramolecular Dynamics of the Basic Leucine Zipper of GCN4: Implications for the Entropy of Association with DNA. *J. Mol. Biol* 1999;285:2133–2146. [PubMed: 9925790]
38. Ishima R, Nagayama K. Protein Backbone Dynamics Revealed by Quasi Spectral Density Function Analysis of Amide N-15 Nuclei. *Biochemistry* 1995;34:3162–3171. [PubMed: 7880811]
39. Farrow NA, Zhang O, Szabo A, Torchia DA, Kay LE. Spectral density function mapping using <sup>15</sup>N relaxation data exclusively. *J. Biomol. NMR* 1995;6:153–162. [PubMed: 8589604]
40. Peng JW, Wagner G. Frequency Spectrum of NH Bonds in Eglin C from Spectral Density Mapping at Multiple Fields. *Biochemistry* 1995;34:16733–16752. [PubMed: 8527448]
41. Lefevre J-F, Dayie KT, Peng JW, Wagner G. Internal Mobility in the Partially Folded DNA Binding and Dimerization Domains of GAL4: NMR Analysis of the N-H Spectral Density Functions. *Biochemistry* 1996;35:2674–2686. [PubMed: 8611573]
42. Tolman JR, Flanagan JM, Kennedy MA, Prestegard JH. NMR evidence for slow collective motions in cyanometmyoglobin. *Nat. Struct. Biol* 1997;4:292–297. [comment]. [PubMed: 9095197]
43. Tolman JR, Al-Hashimi HM, Kay LE, Prestegard JH. Structural and dynamic analysis of residual dipolar coupling data for proteins. *J Am Chem Soc* 2001;123:1416–1424. [PubMed: 11456715]
44. Wang L, Pang Y, Holder T, Brender JR, Kurochkin AV, Zuiderweg ER. Functional dynamics in the active site of the ribonuclease binase. *Proc. Natl. Acad. Sci. U S A* 2001;98:7684–7689. [PubMed: 11438724]
45. Stone JM, Trotochaud AE, Walker JC, Clark SE. Control of meristem development by CLAVATA1 receptor kinase and kinase-associated protein phosphatase interaction. *Plant Physiol* 1998;117:1217–1225. [PubMed: 9701578]
46. Lee G, Li J, Walker JC, Van Doren SR. <sup>1</sup>H, <sup>13</sup>C and <sup>15</sup>N resonance assignments of kinase-interacting FHA domain from Arabidopsis phosphatase KAPP. *J. Biomol. NMR* 2003;25:253–254. [PubMed: 12652139]
47. Farrow NA, Muhandiram R, Singer AU, Pascal SM, Kay CM, Gish G, Shoelson SE, Pawson T, Forman-Kay JD, Kay LE. Backbone Dynamics of a Free and a Phosphopeptide-Complexed Src Homology 2 Domain Studied by <sup>15</sup>N Relaxation. *Biochemistry* 1994;33:5984–6003. [PubMed: 7514039]
48. Boyd J, Hommel U, Campbell ID. Influence of cross-correlation between dipolar and anisotropic chemical shift relaxation mechanisms upon longitudinal relaxation rates of <sup>15</sup>N in macromolecules. *Chemical Physics Letters* 1990;175:477–482.
49. Sklenar V, Piotto M, Leppik R, Saudek V. Gradient-Tailored Water Suppression for <sup>1</sup>H-<sup>15</sup>N HSQC Experiments Optimized to Retain Full Sensitivity. *J. Magn. Reson. A* 1993;102:241–245.
50. Barbato G, Ikura M, Kay LE, Pastor RW, Bax A. Backbone dynamics of calmodulin studied by <sup>15</sup>N relaxation using inverse detected two-dimensional NMR spectroscopy: the central helix is flexible. *Biochemistry* 1992;31:5269–5278. [PubMed: 1606151]
51. Loria JP, Rance M, Palmer AG 3rd. A Relaxation-Compensated Carr-Purcell-Meiboom-Gill Sequence for Characterizing Chemical Exchange by NMR Spectroscopy. *J. Am. Chem. Soc* 1999;121:2331–2332.
52. Loria JP, Rance M, Palmer AG 3rd. A TROSY CPMG sequence for characterizing chemical exchange in large proteins. *J. Biomol. NMR* 1999;15:151–155. [PubMed: 10605088]


53. Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J. Biomol. NMR* 1995;6:277–293. [PubMed: 8520220]
54. Goddard, TD.; Kneller, DG. San Francisco: University of California, San Francisco; 2000.
55. Palmer AG 3rd, Rance M, Wright PE. Intramolecular Motions of a Zinc Finger DNA-Binding Domain from Xfin Characterized by Proton-Detected Natural Abundance  $^{13}\text{C}$  Heteronuclear NMR Spectroscopy. *J. Am. Chem. Soc* 1991;113:4371–4380.
56. Dosset P, Hus JC, Blackledge M, Marion D. Efficient analysis of macromolecular rotational diffusion from heteronuclear relaxation data. *J. Biomol. NMR* 2000;16:23–28. [PubMed: 10718609]
57. Pawley NH, Wang C, Koide S, Nicholson LK. An improved method for distinguishing between anisotropic tumbling and chemical exchange in analysis of  $^{15}\text{N}$  relaxation parameters. *J. Biomol. NMR* 2001;20:149–165. [PubMed: 11495246]
58. Tjandra N, Szabo A, Bax A. Protein Backbone Dynamics and  $^{15}\text{N}$  Chemical Shift Anisotropy from Quantitative Measurement of Relaxation Interference Effects. *J. Am. Chem. Soc* 1996;118:6986–6991.
59. Fushman D, Cowburn D. Model-Independent Analysis of  $^{15}\text{N}$  Chemical Shift Anisotropy from NMR Relaxation Data. Ubiquitin as a Test Example. *J. Am. Chem. Soc* 1998;120:7109–7110.
60. Volkman BF, Alam SL, Satterlee JD, Markley JL. Solution structure and backbone dynamics of component IV of Glycerea dibranchiata monomeric hemoglobin-CO. *Biochemistry* 1998;37:10906–10919. [PubMed: 9692983]
61. d'Auvergne EJ, Gooley PR. The use of model selection in the model-free analysis of protein dynamics. *J. Biomol. NMR* 2003;25:25–39. [PubMed: 12566997]
62. Akaike, H. Information theory and an extension of the maximum likelihood principle. Petrov, BN.; Csaki, F., editors. Budapest: Academiai Kiado; 1973. p. 267–281.
63. Larsson G, Martinez G, Schleucher J, Wijmenga SS. Detection of nano-second internal motion and determination of overall tumbling times independent of the time scale of internal motion in proteins from NMR relaxation data. *J. Biomol. NMR* 2003;27:291–312. [PubMed: 14512728]
64. Hansen MR, Hanson P, Pardi A. Filamentous bacteriophage for aligning RNA, DNA, and proteins for measurement of nuclear magnetic resonance dipolar coupling interactions. *Methods Enzymol* 2000;317:220–240. [PubMed: 10829283]
65. Rückert M, Otting G. Alignment of Biological Macromolecules in Novel Nonionic Liquid Crystalline Media for NMR Experiments. *J. Am. Chem. Soc* 2000;122:7793–7797.
66. Oh MH, Ray WK, Huber SC, Asara JM, Gage DA, Clouse SD. Recombinant brassinosteroid insensitive 1 receptor-like kinase autophosphorylates on serine and threonine residues and phosphorylates a conserved peptide motif in vitro. *Plant Physiol* 2000;124:751–766. [PubMed: 11027724]
67. Jin D, Andrec M, Montelione GT, Levy RM. Propagation of experimental uncertainties using the Lipari-Szabo model-free analysis of protein dynamics. *J. Biomol. NMR* 1998;12:471–492. [PubMed: 9862126]
68. Zweckstetter M, Bax A. Prediction of Sterically Induced Alignment in a Dilute Liquid Crystalline Phase: Aid to Protein Structure Determination by NMR. *J Am Chem Soc* 2000;122:3791–3792.
69. Kay LE, Torchia DA, Bax A. Backbone Dynamics of Proteins as Studied by  $^{15}\text{N}$  Inverse Detected Heteronuclear NMR Spectroscopy: Application to Staphylococcal Nuclease. *Biochemistry* 1989;28:8972–8979. [PubMed: 2690953]
70. Blackledge M, Cordier F, Dosset P, Marion D. Precision and Uncertainty in the Characterization of Anisotropic Rotational Diffusion by  $^{15}\text{N}$  Relaxation. *J Am Chem Soc* 1998;120:4538–4539.
71. Dayie KT, Wagner G, Lefevre JF. Theory and practice of nuclear spin relaxation in proteins. *Annual Review of Physical Chemistry* 1996;47:243–282.
72. Fushman D, Weismann R, Thüring H, Rüterjans H. Backbone dynamics of ribonuclease T1 and its complex with 2'GMP studied by two-dimensional heteronuclear NMR spectroscopy. *J. Biomol. NMR* 1994;4:61–78.
73. Buchberger A, Howard MJ, Freund SM, Proctor M, Butler PJ, Fersht AR, Bycroft M. Biophysical characterization of elongin C from *Saccharomyces cerevisiae*. *Biochemistry* 2000;39:11137–11146. [PubMed: 10998253]

74. Tugarinov V, Liang Z, Shapiro YE, Freed JH, Meirovitch E. A structural mode-coupling approach to <sup>15</sup>N NMR relaxation in proteins. *J Am Chem Soc* 2001;123:3055–3063. [PubMed: 11457016]
75. Vugmeyster L, Raleigh DP, Palmer AG 3rd, Vugmeister BE. Beyond the decoupling approximation in the model free approach for the interpretation of NMR relaxation of macromolecules in solution. *J Am Chem Soc* 2003;125:8400–8404. [PubMed: 12837113]
76. Wang P, Byeon IJ, Liao H, Beebe KD, Yongkiettrakul S, Pei D, Tsai MD. II. Structure and specificity of the interaction between the FHA2 domain of Rad53 and phosphotyrosyl peptides. *J. Mol. Biol* 2000;302:927–940. [PubMed: 10993733]
77. Yongkiettrakul S, Byeon IJ, Tsai MD. The ligand specificity of yeast Rad53 FHA domains at the +3 position is determined by nonconserved residues. *Biochemistry* 2004;43:3862–3869. [PubMed: 15049693]
78. Olejniczak ET, Zhou MM, Fesik SW. Changes in the NMR-derived motional parameters of the insulin receptor substrate 1 phosphotyrosine binding domain upon binding to an interleukin 4 receptor phosphopeptide. *Biochemistry* 1997;36:4118–4124. [PubMed: 9100005]
79. Durocher D, Henckel J, Fersht AR, Jackson SP. The FHA domain is a modular phosphopeptide recognition motif. *Mol. Cell* 1999;4:387–394. [PubMed: 10518219]
80. Williams DC Jr, Benjamin DC, Poljak RJ, Rule GS. Global Changes in Amide Hydrogen Exchange Rates for a Protein Antigen in Complex with Three Different Antibodies. *J. Mol. Biol* 1996;257:866–876. [PubMed: 8636987]
81. Trotochaud AE, Hao T, Wu G, Yang Z, Clark SE. The CLAVATA1 Receptor-like kinase requires CLAVATA3 for its assembly into a signaling Complex that includes KAPP and a Rho -related protein. *Plant Cell* 1999;11:393–405. [PubMed: 10072399]
82. Jinn T-L, Stone JM, Walker JC. HAESA, and Arabidopsis leucine-rich repeat receptor kinase, controls floral organ abscission. *Genes & Development* 2000;14:108–117. [PubMed: 10640280]
83. Arkin MR, Randal M, DeLano WL, Hyde J, Luong TN, Oslob JD, Raphael DR, Taylor L, Wang J, McDowell RS, Wells JA, Braisted AC. Binding of small molecules to an adaptive protein-protein interface. *Proc. Natl. Acad. Sci. U S A* 2003;100:1603–1608. [PubMed: 12582206]
84. Shiu SH, Bleecker AB. Receptor-like kinases from Arabidopsis form a monophyletic gene family related to animal receptor kinases. *Proc. Natl. Acad. Sci. USA* 2001;98:10763–10768. [PubMed: 11526204]
85. Berjanskii MV, Riley MI, Van Doren SR. Hsc70-interacting HPD loop of the J domain of polyomavirus T antigens fluctuates in psec to nsec and μsec to msec. *J. Mol. Biol* 2002;321:503–516. [PubMed: 12162962]


**Figure 1.**

$^{15}\text{N}$  relaxation data of KI-FHA measured at 500 and 600 MHz and plotted vs. residue from the KAPP sequence: a)  $^{15}\text{N}\{^1\text{H}\}$  NOE plotted as  $I_{\text{sat}}/I_{\text{nonsat}}$ , b) spin-lattice relaxation rate constant,  $R_1$ , c) spin-spin relaxation rate constant,  $R_2$  and d) relaxation-compensated CPMG measurement of  $\Delta R_2 = R_2(7.5 \text{ ms}) - R_2(2 \text{ ms})$ , where 7.5 and 2 ms are the spacing between  $\pi$  pulses of the CPMG trains. 25 °C was used for a) – c). 22 °C was used for d). Uncertainties were estimated as described in Methods. In b), gray squares mark four cases of poor single exponential fits due to the clear presence of a second, fast,  $T_2$ -like exponential decay; for the four affected residues, the fit of the slow decaying component is shown with black squares.


The locations of  $\beta$ -strands are marked with gray bars in each panel. Filled squares represent 500 MHz data. Open circles represent 600 MHz data.

**Figure 2.**

${}^{15}\text{N}$  relaxation data of KI-FHA, bound to the CLV1 pT868 peptide at 25 °C, and plotted vs. residue number from the KAPP sequence: a)  ${}^{15}\text{N}\{{}^1\text{H}\}$  NOE, b)  $R_1$ , c)  $R_2$  and d)  $\Delta R_2 = R_2(7.5 \text{ ms}) - R_2(2 \text{ ms})$ . 25 °C was used for a) – c). 22 °C was used for d). The peptide was present at 1.5-fold excess, except for the 3.5-fold excess used for  $R_2$  measurements in c) and d) at 600 MHz. The symbol code of Figure 1 is used.


**Figure 3.**

Evidence for nsec backbone motions. a) Reduced spectral density function  $J(\omega_N)$  plotted vs. residue, where  $\omega_N=3.8 \times 10^8$  rad/sec since the relaxation data were collected at 600 MHz. Open squares represent the free state and filled triangles represent the state bound to pT868 from CLV1. b) Sites in free KI-FHA where residual dipolar couplings (RDCs) deviate from those predicted by SVD fits to the NMR structural coordinates of KI-FHA (PDB code 1mzk). Squared deviations of individual  ${}^1D_{\text{NH}}$  measured RDCs from those predicted by SVD fits of all RDCs to the entire backbone are normalized by  $D_a$ , the axial component of the powder pattern distribution of RDCs observed. This quantity is averaged for the ten lowest energy NMR structures of KI-FHA.




**Figure 4.**


Model-free dynamics results for the free state of KI-FHA from fits of  $^{15}\text{N}$  relaxation at two fields, 500 MHz and 600 MHz. a) The generalized order parameter  $S^2$ , b) the line broadening term  $R_{\text{ex}}$  and c) the internal correlation time  $\tau_e$  are plotted vs. sequence position in KI-FHA. Locations of  $\beta$ -strands are marked with bars.

**Figure 5.**

Model-free dynamics results for KI-FHA bound to CLV1 pT868 using  $^{15}\text{N}$  relaxation collected at both 500 MHz and 600 MHz. a) The generalized order parameter  $S^2$ , b) the line broadening term  $R_{\text{ex}}$  and c) the internal correlation time  $\tau_e$  are plotted vs. sequence position in KI-FHA. Locations of  $\beta$ -strands are marked with bars.


**Figure 6.**

Comparison of pT868 peptide-induced changes in KI-FHA flexibility over psec to nsec and in amide NMR peak positions at the interface. a) Changes in the generalized order parameter  $S^2$  upon the binding of the pT868 peptide are plotted vs. residue of KAPP. The locations of  $\beta$ -strands are marked with gray bars. Residues with  $S^2_{\text{bound}} - S^2_{\text{free}} \geq 0.065$ , marked by a dashed line, represent the upper 30-percentile of  $\Delta S^2$  values. b) Sites of  $\Delta S^2$  are colored on the stereo view of the backbone of free KI-FHA with tube width proportional to  $1 - S^2_{\text{free}}$ , i.e. for the free state. In (a), residues rigidified by pT868 peptide binding are colored dark blue where  $S^2_{\text{bound}} - S^2_{\text{free}} > 0.14$  and lighter blue where  $0.14 > \Delta S^2 > 0.065$ . Residues mobilized by pT868 peptide binding are colored red where  $S^2_{\text{bound}} - S^2_{\text{free}} \leq -0.065$ . b) Shifts of amide NMR peaks of KI-FHA upon addition of a 4-fold excess of the CLV1 pT868 peptide are plotted vs. location in KAPP. The radial shift of the peak  $\Delta \omega_{\text{NH}} = (\Delta \omega_{\text{H}}^2 + (\Delta \omega_{\text{H}}/6)^2)^{1/2}$ , where  $\Delta \omega_{\text{H}}$  and  $\Delta \omega_{\text{N}}$  are the changes in  $^1\text{H}$  and  $^{15}\text{N}$  dimensions in ppm. In d), pT868-induced chemical shift changes,  $\Delta \omega_{\text{NH}}$ , are mapped onto the stereo view of the backbone of the free KI-FHA structure with tube width proportional to  $1 - S^2_{\text{bound}}$ , representing the bound state. Residues are colored in red where  $\Delta \omega_{\text{NH}} > 0.35$ , orange where  $0.35 > \Delta \omega_{\text{NH}} > 0.15$ , and gold where  $0.15 > \Delta \omega_{\text{NH}} > 0.067$ . A few of the most shifted residues are labeled. In c) and d), unobservable residues are colored a darker shade of gray.



**Figure 7.**  
 Mapping of sites of  $\mu$ sec to msec scale fluctuations and of nsec scale motions, of backbone amide peaks upon the KI-FHA NMR structure for the free state (a) and pT868-bound state (b). In (a) and (b), residues with  $R_{ex} > 1 \text{ s}^{-1}$  exchange broadening evidence of the slower motions are colored red and those with  $1 \text{ s}^{-1} > R_{ex} > 0 \text{ s}^{-1}$  are colored orange. Residues with apparent  $\tau_e > 0.7$  nsec are marked with blue spheres. Unobservable residues are colored a darker shade of gray. Structural elements with  $R_{ex}$  in free KI-FHA are labeled in (a). In (b), residues are labeled where binding of the pT868 peptide introduces  $R_{ex}$ . The binding surface for phosphoThr peptides or phosphoproteins (receptor-like kinases such as CLV1 in plants) is indicated with red arrows. Residues in the bound state undergoing too much chemical exchange

broadening for relaxation fitting are labeled in pink; these residues exhibit intermediate exchange broadening during titration with the pT868 peptide.

**Figure 8.**

A network of two tiers of packed interior side chains, plotted in blue, where the backbone is affected by conformational exchange detected as line broadening. The color code of the backbone is that of Figure 7a.

**Table 1**

Spectral density functions used to fit relaxation data in the extended Lipari-Szabo model-free approach

Model	Fit parameters	Spectral density function
1	$S^2$	$J(\omega)=2/5\{(\bar{S}^2\tau_m)/[1+(\omega\tau_m)^2]\}$
2	$S^2, \tau_e$	$J(\omega)=2/5\{(\bar{S}^2\tau_m)/[1+(\omega\tau_m)^2]+[(1-\bar{S}^2)\tau]/[1+(\omega\tau)^2]\}$
3	$S^2, R_{ex}$	$J(\omega)=2/5\{(\bar{S}^2\tau_m)/[1+(\omega\tau_m)^2]\}$
4	$S^2, \tau_e, R_{ex}$	$J(\omega)=2/5\{(\bar{S}^2\tau_m)/[1+(\omega\tau_m)^2]+[(1-\bar{S}^2)\tau]/[1+(\omega\tau)^2]\}$
5	$S_s^2, S_f^2, \tau_e$	$J(\omega)=(2S_f^2/5)\{(\bar{S}_s^2\tau_m)/[1+(\omega\tau_m)^2]+[(1-\bar{S}_s^2)\tau]/[1+(\omega\tau)^2]\}$

$\tau = \tau_e \tau_m / (\tau_e + \tau_m)$ , where  $\tau_m$  is the isotropic rotational correlation time and  $\tau_e$  is the effective correlation time for internal motions.  $S^2 = S_f^2 S_s^2$  is the square of the generalized order parameters.  $S_f^2$  and  $S_s^2$  are the squares of the order parameters for the internal motions on the fast and slow time scales.

Average filtered relaxation parameters of KI-FHA, free and bound to pT868.

**Table 2**

state of KI-FHA	Field, Tesla	coarse-filtered A			fine-filtered B			$\tau_m, C$ nsec	$D_{\parallel}/D_{\perp}, C$
		$^{15}\text{N}(\text{H})$	$\bar{R}_{\text{p}, \text{s}^{-1}}$	$R_{2, \text{s}^{-1}}$	$n_B$	$\bar{R}_{\text{p}, \text{s}^{-1}}$	$R_{2, \text{s}^{-1}}$		
Free									
	11.74	83	0.76	2.07	11.6	46	0.76	2.05	11.5
	14.1	80	0.78	1.89	12.6	43	0.79	1.85	12.3
Bound to pT868									
	11.74	57	0.75	1.91	14.4	39	0.72	1.87	14.2
	14.1	84	0.80	1.41	16.9 E	42	0.80	1.35	16.4 E
					(16.2) F				(16.1) F

NOE, R<sub>1</sub>, R<sub>2</sub> and R<sub>2</sub>/R<sub>1</sub> were trimmed prior to averaging.

*A* Coarse filtering employed Normadyn software to remove residues with NOE < 0.65 or with  $T_2 \leq \overline{T}_2 + \sigma_{T_2}$  unless  $T_1 \geq \overline{T}_1 + \sigma_{T_1}$  (57). Number of residues after filtration is *n*.

*B* Fine filtering removed residues poorly fit by simple model-free expressions 1 and 2 (Table 1). Residues with exchange broadening suggested by relaxation dispersion at 600 MHz or statistical model section were removed. Residues suggested to have nsec-scale motions by  $J(\overline{\omega}_N) = 61 \text{MHz} \geq J(\overline{\omega}_N) + \sigma_{J(\overline{\omega}_N)}$  were also removed. Number of residues after filtration is *n*.

*C* Rotational correlation time  $\tau_m$  and the  $D_{\parallel}/D_{\perp}$  ratio were estimated using the fine-filtered set of residues, their 600 MHz R<sub>1</sub> and R<sub>2</sub> relaxation data, Tensor2 (56), and PDB coordinates 1MZK.

*D* Except where noted, KI-FHA was present at 0.53 mM and the peptide at 1.5-fold excess, suggesting 94% peptide saturation of KI-FHA since K<sub>D</sub> is ~20  $\mu\text{M}$ .

*E* These values result from increasing the peptide excess to 3.5-fold, boosting peptide saturation of the 0.33 mM KI-FHA to 98%.

*F* Parenthesized values derive from R<sub>2</sub> measured after dilution of KI-FHA to 0.23 mM with 0.35 mM pT868 peptide. These conditions decreased peptide saturation of KI-FHA to ~88%. Uncertainties in R<sub>2</sub> at 0.23 mM exceed differences from R<sub>2</sub> at higher concentration and saturation.