

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/44588872>

# tRNA Stabilization by Modified Nucleotides

ARTICLE *in* BIOCHEMISTRY · JUNE 2010

Impact Factor: 3.02 · DOI: 10.1021/bi100408z · Source: PubMed

---

CITATIONS

121

---

READS

230

## 2 AUTHORS:



[Yuri Motorin](#)

University of Lorraine

44 PUBLICATIONS 1,479 CITATIONS

[SEE PROFILE](#)



[Mark Helm](#)

Johannes Gutenberg-Universität Mainz

113 PUBLICATIONS 2,735 CITATIONS

[SEE PROFILE](#)

# tRNA Stabilization by Modified Nucleotides<sup>†</sup>

Yuri Motorin<sup>‡</sup> and Mark Helm<sup>\*,§</sup>

<sup>‡</sup>*Laboratoire ARN-RNP Maturation-Structure-Fonction, Enzymologie Moléculaire et Structurale (AREMS), UMR 7214 CNRS-UHP Faculté des Sciences et Techniques, Université Henri Poincaré, Nancy 1, Bld des Aiguillettes, BP 70239, 54506 Vandoeuvre-les-Nancy, France, and* <sup>§</sup>*Institute of Pharmacy and Biochemistry, Johannes Gutenberg-University of Mainz, Staudinger Weg 5, 55099 Mainz, Germany*

*Received March 16, 2010; Revised Manuscript Received May 11, 2010*

**ABSTRACT:** Post-transcriptional ribonucleotide modification is a phenomenon best studied in tRNA, where it occurs most frequently and in great chemical diversity. This paper reviews the intrinsic network of modifications in the structural core of the tRNA, which governs structural flexibility and rigidity to fine-tune the molecule to peak performance and to regulate its steady-state level. Structural effects of RNA modifications range from nanometer-scale rearrangements to subtle restrictions of conformational space on the angstrom scale. Structural stabilization resulting from nucleotide modification results in increased thermal stability and translates into protection against unspecific degradation by bases and nucleases. Several mechanisms of specific degradation of hypomodified tRNA, which were only recently discovered, provide a link between structural and metabolic stability.

With more than 100 distinct chemical structures, the high frequency and chemical diversity of ribonucleotide modifications occurring in natural RNA present a complicated puzzle; clearly, modifications must be of high importance or else this complexity would not have evolved. A pronounced hot spot for sophisticated nucleotide modification is found in the anticodon loop of tRNA, in particular at positions 34 and 37. These are known to optimize the mRNA decoding function by directly affecting codon–anticodon interactions (1). The sophisticated involvement of modifications at the latter positions in codon recognition and related issues has been expertly reviewed elsewhere (2, 3). Many such modifications were shown in early work to be essential, which can be easily rationalized with their important function in the crucial process of accurate translation.

In contrast, the functions of most modifications in the second cluster of modifications, situated in the structural core of tRNA, are ill understood. These modifications are typically unsophisticated with respect to their chemistry: in addition to pseudouridine (Ψ), they consist mostly of methylated nucleotides. Some examples discussed in this review are shown in Figure 1. Although a distinction between structure and function can never be complete, we will treat these chemically simple core modifications as “structural” modifications for the purpose of this work. In tRNAs, which are the most heavily modified and also most investigated RNAs, efforts to clearly define the function of single modifications suffer from the apparent collective nature of the ensemble of modifications. Indeed, the absence of only a single modification remains very frequently without obvious consequence (4), and effects become apparent only when additional modifications or elements of tRNA structure are also missing (5, 6). The common perception in the field (7–10) pictures nucleotide modifications in the tRNA core as modulators of its

structural flexibility, which can rigidify certain areas and render others flexible (11). A central topic of this review is the structural impact of core modifications, and the difficulties in studying such an impact. Because tRNAs exhibit a very conserved and well-defined base pairing pattern, significant structural alterations as a consequence of nucleotide modification are unexpected. Rather, it is typically expected that primary transcripts already form the classical cloverleaf structure and the three-dimensional L shape known from crystallography (12). As has been shown in hundreds of cases with unmodified *in vitro* transcripts, this is indeed the case; most unmodified tRNAs are functional, e.g., in aminoacylation *in vitro* (13–15) or even in *in vitro* translation (16). There are, however, several reported cases in which transcripts fail *in vitro* (17–19), and these are clearly of great interest with respect to post-transcriptional modifications. Here again, it is helpful to distinguish modifications that directly interact with partners of the translation machinery from those that shape and fine-tune the tRNA core structure for peak performance. Recent evidence indicates that tRNAs do not function *in vivo* without post-transcriptional modifications that reinforce the structural tRNA core (5, 6, 20).

As the laws of thermodynamics literally dictate that there must be more than one structure for any given RNA (10), it is advisable to think of one’s favorite structure not as the “correct” structure but rather as the “predominant”, most populated conformation in a given equilibrium. As a direct consequence of modification, we will repeatedly discuss cases of structural equilibrium and alternative structures. Indeed, tRNA participates in a multitude of interactions during its involvement in protein biosynthesis: depending on its binding partner, it has been found in a multitude of structural variations of the L shape (12, 21–24) that require structural flexibility in addition to stability.

In what follows, we will review our knowledge of the stabilizing impact of nucleotide modifications, first on the level of large structural alterations that involve significant rearrangements of Watson–Crick interactions, i.e., nanoscale structural changes, and then touching on the more subtle structural effects that can

<sup>†</sup>M.H. acknowledges funding by the DFG in the frame of the Forschergruppe Dnmt2 (HE 3397/6, FOR 1082) and by the Dr. Gerhard and Martha Röttger Foundation.

<sup>\*</sup>To whom correspondence should be addressed. E-mail: mhelm@uni-mainz.de. Phone: 49-6131-3925731. Fax: 49-6131-3920373.




FIGURE 1: Sites of chemically simple modifications on the Watson–Crick edges and the Hoogsteen edges (purines) or C–H edges (pyrimidines) (162, 163). Single methylation of endocyclic nitrogens of the Watson–Crick edge or double methylation of exocyclic amines ablates the possibility of standard Watson–Crick pairing. Methylations of the 2'-OH group, which may modulate the hydrogen bond pattern of the sugar edge, are also indicated. Modified nucleotides are abbreviated according to common nomenclature (164).

be measured on the angstrom scale. The stabilization of RNA will then be discussed from a thermodynamic point of view, i.e., thermal stability. Finally, we turn to biological consequences for tRNAs lacking structural stabilization, in particular, degradation pathways of unstructured tRNA.

### NANOSCALE REARRANGEMENTS OF tRNA AS AN EFFECT OF NUCLEOTIDE MODIFICATION

Large structural effects in RNA, measured on the nanometer scale, are likely to include the dissolution and re-formation of helical structures. The latter implies alteration on the level of hybridization, and thus of numerous Watson–Crick interactions. Modifications are suitable for altering hydrogen bond interactions on the Watson–Crick edge, the Hoogsteen edge, or the sugar edge of the target nucleotide. Methylations are the most simple and most widespread modifications known to block formation of hydrogen bonds between nucleotides. Accordingly, the few known examples of modification-dependent Watson–Crick rearrangements involve nucleotide methylations. Among the best understood cases is human mitochondrial tRNA<sup>Lys</sup> which adopts a noncanonical extended hairpin structure when unmodified (19, 25). As was shown by single-molecule fluorescence energy transfer studies, this structure is the predominant one in a dynamic equilibrium with the originally expected cloverleaf structure (26). Despite the presence of this correctly folded subpopulation, the unmodified form is not a substrate for aminoacylation by the cognate lysyl-tRNA synthetase (27). Significantly, the introduction of m<sup>1</sup>A at position 9 shifts the dynamic equilibrium in favor of the cloverleaf structure (25, 26, 28) (Figure 2A). Biochemical investigations suggest

that m<sup>1</sup>A formation is an early maturation event (28), which may occur directly after tRNA 5' maturation by RNase P (29). Studies of DNA indicate a penalty of ~6.5 kJ/mol for the introduction of m<sup>1</sup>A into a helical structure, where m<sup>1</sup>A is reported to form a T(*anti*)·m<sup>1</sup>A(*syn*) Hoogsteen base pair with the thymidine residue in the opposing strand (30). In tRNA<sup>Lys</sup>, the methyl group afforded a stabilization of 3–4 kJ/mol of the cloverleaf relative to the extended hairpin structure depending on the magnesium ion concentration (31). A mutational analysis of various transcripts is consistent with the major contribution of this effect coming from the interdiction of a Watson–Crick interaction by the methyl group. Mutations that imitate this effect lead to a predominance of the cloverleaf conformation and to good substrates for aminoacylation (27). Initial studies, conducted by structural probing, had already yielded strong indications of high structural plasticity and of alternative structures in this tRNA. Similar studies revealed structural plasticity in transcripts of several other human mitochondrial tRNAs, including tRNA<sup>Leu(UUR)</sup> and tRNA<sup>Asp</sup> (32–34). For the latter, a noncloverleaf conformation with a strong resemblance to the aforementioned extended hairpin of tRNA<sup>Lys</sup> has been proposed (35) (Figure 2B). Aminoacylation studies in vitro show that transcripts of animal mitochondrial tRNAs are altogether worse substrates than transcripts of canonical tRNAs in the cytosol (33, 34, 36–40). These findings imply that such plasticity is a common characteristic of animal mitochondrial tRNAs and that despite a general conservation of the base pairing in stem regions (41, 42), the minimal structural characteristics of the cloverleaf may frequently be reached only after certain modification steps.




FIGURE 2: Alternative secondary structures related to nucleotide modification of mitochondrial tRNAs. (A) Structural equilibrium of human mitochondrial tRNA<sup>Lys</sup>. The equilibrium is displaced toward the structure on the right by a single methylation on N9 of adenosine 9 as evidenced by structural probing (19, 25) and single-molecule FRET experiments (26). (B) Alternative structures proposed for tRNA<sup>Asp</sup> based on structural probing (35).

There has been speculation (10) that the apparent folding problem of mitochondrial tRNAs may in part be related to the nucleotide bias in the mitochondrial genome and in mitochondrial tRNAs (42). In brief, the “RNA folding problem” is related to the fact that Watson–Crick pairs are quite strong and govern RNA folding in such a predominant way that their low diversity of elemental units (only four different base pairs) makes it difficult for a given sequence to specify a truly unique secondary structure (10). The resulting problem of various alternative structures with similar energy is obviously compounded in sequences with a nucleotide bias, such as RNAs transcribed from mitochondrial genomes, which are rich in A–T pairs. As we have seen in the case of tRNA<sup>Lys</sup>, the introduction of post-transcriptional nucleotide modifications may partially remedy a problem of alternative structures by interdiction of certain Watson–Crick interactions.

Consequently, one might expect similar problems and solutions for many tRNAs that are rich in G and C nucleotides. Indeed, although there is no direct experimental proof so far, circumstantial evidence that suggests that m<sup>2</sup><sub>2</sub>G might have a role in tRNA from thermophiles that is similar to that of m<sup>1</sup>A in

mitochondrial tRNA<sup>Lys</sup> is accumulating. Interestingly, those hydrogen bond donors and acceptors that are involved in the formation of the G–U wobble base pair are left unaltered upon modification of G to m<sup>2</sup><sub>2</sub>G. While its ability to form a Watson–Crick pair with cytosine is ablated, m<sup>2</sup><sub>2</sub>G may still undergo noncanonical interactions involving the N1 and O2 positions and thus form an m<sup>2</sup><sub>2</sub>G–U wobble interaction. Scenarios in which this property might favor the functional cloverleaf structure have been proposed for m<sup>2</sup><sub>2</sub>G at positions 10 (43) and 26 (44).

Large RNA structural alterations on the nanometer scale, which do not involve significant changes in base pairing, may be achieved by changes in the relative reorientation of structural domains. Modifications that may trigger or contribute to such effects are situated in the structural core of the tRNA, where they are thought to stabilize a number of tertiary interactions, which, in turn, govern the interstem angle between the anticodon and acceptor domains among other things. A compilation of biochemical and biophysical data for modified and unmodified tRNA (10, 12, 14, 15, 45–52) indicates strong interplay between magnesium ion coordination and post-transcriptional modification. Apparently, tRNA reaches its functional optimum at physiological concentrations of magnesium ions when fully modified, but the absence of modification can be partially compensated by increasing the magnesium ion concentration. In general, small RNAs have been observed to undergo compaction at increased concentrations of magnesium ions, likely as a general consequence of the increased level of shielding of the phosphates (31, 53). There is evidence that structural core modifications (54), including, e.g., m<sup>5</sup>C40 (51, 55, 56), improve the binding of magnesium ions. As the interstem angle of unmodified tRNA is known to strongly decrease at elevated concentrations of magnesium ions (57–59), nucleotide modifications are expected to promote a decrease in the angle at significantly lower, namely, physiological, concentrations of Mg<sup>2+</sup> ions.

## SUBTLE EFFECTS ON THE ANGSTROM SCALE

Only a few of the plethora of nucleotide modifications exert their influence via blocking of Watson–Crick interactions. Rather, most influence parameters of the local RNA structure and lead to stabilizing effects by restricting conformational flexibility on the angstrom scale. Viewed on that scale, e.g., by nuclear magnetic resonance (NMR), RNA appears as a string of nucleotides whose moieties are constantly and very rapidly changing their relative orientations. This fact is often less obvious in the commonplace presentation of crystal structures, but readily obvious in studies of RNA dynamics by NMR (reviewed in ref 60); it also forms the basis of molecular dynamics (MD) simulations (61, 62).

The structural characterization of a nucleotide unit in an RNA chain includes the sugar pucker and the relative conformation of the sugar and base as central parameters that are readily investigated by NMR (60). In unstructured RNAs, there is an equilibrium between the two predominant sugar conformations, 2′-endo and 3′-endo (Figure 3), which is shifted in favor of the 3′-endo conformation by hybridization to A-form helices. Nucleotide modifications favoring the 3′-endo conformation, such as methylation of the sugar 2′-OH group, pseudouridylation, or thiolation of the pyrimidine O2 atom (Figure 3), will thus stabilize helical conformations (for references, see, e.g., refs (63–65); reviews in refs 60 and 66).

Because NMR experiments are subject to restrictions of RNA size, only a limited number of studies on full-length tRNAs have






FIGURE 3: Local structural impact of chemically simple modifications. Nucleotide conformations are illustrated with  $s^2T$ , a modification typically occurring at position 54 in thermophiles growing above 80 °C: (left) 3'-endo conformation of the ribose and *syn* conformation of the base, (middle) 3'-endo conformation of the ribose and *anti* conformation of the base (most favored in  $s^2T$ ), and (right) 2'-endo conformation of the ribose and *anti* conformation of the base.




FIGURE 4: Coordination of water molecules to N1 of  $\Psi$  as modeled onto the X-ray structure of *E. coli* tRNA<sup>Gln</sup> based on an electron density difference map of modified vs unmodified tRNA<sup>Gln</sup> in a complex with its cognate amino acyl-tRNA synthetase (90). On the right side, the N1H group in  $\Psi$  is encircled, and the coordinated water molecule as well as their presumed hydrogen bonds to the neighboring phosphates is indicated.

been conducted (47, 50, 67, 68). Truncated T-stem loops (69) and truncated anticodon stem loops (ASLs) of ~17 nucleotides are typical model systems, in which subtle to strong effects of various modified nucleotides, including Cm, m<sup>1</sup>G, m<sup>5</sup>C,  $\Psi$ , m<sup>6</sup>A, i<sup>6</sup>A, ms<sup>2</sup>i<sup>6</sup>A, t<sup>6</sup>A, ms<sup>2</sup>t<sup>6</sup>A, mnm<sup>5</sup>s<sup>2</sup>U, mcm<sup>5</sup>s<sup>2</sup>U, and cmo<sup>5</sup>U, have been studied. Although it has been suggested on the basis of tRNA crystal structures (12, 70) that potential long-range effects of the tRNA core structure on the anticodon loop conformation may have been lost, ASLs are, in principle, functional in ribosome binding assays (71). This underscores the biological significance of studies including modified nucleotides in tRNA<sup>Phe</sup> from *Escherichia coli* (72–75) and yeast (76–78), tRNA<sup>Lys</sup> from *E. coli* and humans (79–85), and tRNA<sup>Val</sup> from *E. coli* (86). Of particular interest are studies on the shaping of the *E. coli* tRNA<sup>Phe</sup> ASL by modifications, because in the absence of modifications the characteristic anticodon loop structure is collapsed into a three-nucleotide loop at the expense of an elongated helix featuring U32-A38 and U33-A37 pairs as extra base pairs (72–75). Recent studies revealed an interesting effect of  $\Psi$  residues at positions 32 and 39, which contribute to the native loop structure and disrupt the abnormal U33-A37 base pair. Again, the interplay among magnesium ion binding, modification, and RNA structure appears to be an important feature (12, 72).

Pseudouridine, once named the fifth ribonucleotide (87) because of its abundance, is an isomer of uridine. Formation of  $\Psi$  is an isomerization reaction catalyzed by pseudouridine synthases (Pus), which proceeds by cleavage of the N-glycosidic bond, base rotation, and reconnection to the sugar at C5 in a C–C bond (88). The structural impact of  $\Psi$  has been studied not only in a large number of RNA sequences, including various duplexes (88, 89)

and somewhat more complex anticodon stem–loop structures (ASLs), but also in structurally more complex systems.  $\Psi$  is present in numerous crystal structures of native RNAs, of which one is particularly instructive. Crystal structures of glutaminyl-tRNA synthetase have been obtained in complex with the native tRNA<sup>Gln</sup>, as well as with the unmodified in vitro transcript. An electron density difference map revealed additional electron density near the “signature” endocyclic N1 of every  $\Psi$  present in the structure, which was interpreted as a water molecule whose order stems from coordination to the extra hydrogen bond site (90) (Figure 4). Depending on the exact structural context, this water molecule is thought to undergo additional interactions, in particular with the phosphate on the 5' side of the  $\Psi$  (Figure 4). A coordinated water molecule as a mediator of a stabilizing influence of  $\Psi$  on RNA structure has also been predicted by MD simulations (91, 92), and its presence is furthermore supported by results from advanced NMR techniques. In addition, in modified RNAs containing  $\Psi$  studied to date, the 3-endo sugar pucker conformation predominates (93–95). Both effects are consistent with numerous studies using CD and NMR spectroscopy and UV melting, which consistently reveal a structural stabilization of helices by  $\Psi$  residues, typically identified as an increased melting temperature (88, 89, 93–97).

## THERMOSTABILITY

The structural impact of nucleotide modifications is often evaluated via UV melting experiments, in which certain structural transitions, in particular, melting of helices, are detected as an increase in UV light absorption along a temperature gradient.

Other types of spectroscopy, including NMR, have been used to the same end (64, 66, 98, 99). An increase in the characteristic midpoint temperature of a transition, the so-called melting temperature, is indicative of a structural stabilization. This interpretation is well-grounded in thermodynamics, but interpretation is straightforward only for the melting of simple RNA duplexes (100). In biological interpretations of melting temperatures, it is often tacitly assumed that an increase in thermal stability and the associated structural stabilization are equivalent to increased resistance against unspecific catalytic degradation by general bases or nucleases (101), and thus associated with increased biochemical and metabolic stability. Support for this notion comes from the fact that hydrolysis of the phosphodiester backbone requires a linear “in-line” geometry of the phosphate and the neighboring 2'-OH moiety, which is the attacking nucleophile (102). This geometry is most easily achieved in largely unstructured, single-stranded regions, which can be distinguished from double-stranded and otherwise stable structures by detection of spontaneous hydrolysis in so-called in-line probing experiments (102). Because the 3-endo sugar pucker of helical regions prevents the in-line geometry, it protects the structure from spontaneous cleavage, as do nucleotide modifications that favor the 3-endo conformation or which lend thermal stability to RNA structures in general.

An interesting case in which thermal stabilization of tRNA structure by modifications is equivalent to physiological stabilization is given in thermophilic organisms. Their RNA must obviously retain a functional structure at temperatures as high as that of boiling water. In addition to the denaturation of secondary and tertiary structure, RNA is ordinarily subject to rapid spontaneous in-line-type hydrolysis at these temperatures. Thermophiles have developed several strategies for increasing the stability of tRNA structure (reviewed in ref 103), including the synthesis of polyamines (104, 105) and a high content of G·C base pairs, which is linearly correlated with an increase in melting temperature (66, 106). Interestingly, certain nucleotide modifications, including G<sub>m</sub>, m<sup>2</sup>G<sub>m</sub>, ac<sup>4</sup>C<sub>m</sub>, s<sup>2</sup>T, and m<sup>1</sup>A, are induced at high temperatures (107–110). Biosynthesis of these modifications conferred significant thermal stabilization to thus modified tRNAs (109–111), and null mutants deficient in the synthesis of certain modifications exhibit impeded growth at elevated temperatures (112, 113). Among the best understood modifications from thermophiles with respect to biosynthesis and structural stabilization are G<sub>m</sub>18, s<sup>2</sup>T54, and m<sup>1</sup>A58 (66, 114). These modifications are clustered near or directly at those nucleotides forming the “classical” G18–Ψ55 and G19–C56 interactions between loop D and loop T, implying that they may (together with Ψ55) serve to stabilize this important tertiary interaction. Intriguingly, investigations of G<sub>m</sub>18, which is also known in mesophiles, did not produce convincing evidence of significant tRNA thermal stabilization by this modification (108), and the same has been reported for m<sup>1</sup>A58 (66, 108). However, the presence of m<sup>1</sup>A58 is required for the thiolation of T54 to s<sup>2</sup>T54 in vitro and in vivo (113), and thus, the fact that growth of knockout strains of m<sup>1</sup>A58 was impeded at elevated temperatures (112) was interpreted as being due to the lack of s<sup>2</sup>T54 (113).

The thermophile-specific hypermodification s<sup>2</sup>T54, however, is very effective in increasing the thermal stability of tRNA (66, 114). In a groundbreaking early study, comparison of the melting profiles of purified *Thermus thermophilus* tRNA<sup>Ile</sup><sub>1</sub> containing T or s<sup>2</sup>T at position 54 revealed an increase of 3 °C in thermal stability due to the presence of sulfur in

s<sup>2</sup>T54 (64, 111, 115). Studies of the structure and dynamics of s<sup>2</sup>T as a nucleoside and in the context of the tRNA scaffold have revealed a strong stabilization for the 3'-endo/*anti* conformation (66, 114) (Figure 3). Although residue 54 is not formally part of the T-stem helix as depicted in the cloverleaf structure (e.g., Figure 2), the three-dimensional loop T structure features an extension of the helix, also including T54, which undergoes an additional reverse Hoogsteen interaction with A58/m<sup>1</sup>A58.

## BIOLOGY OF tRNA MODIFICATIONS

Although synthesis and maturation of tRNA vary significantly among organelles, organisms, and kingdoms (116), many common features have been retained during evolution. With only one known exception (117), primary tRNA transcripts, which are frequently polycistronic, are processed at the 5' end by RNase P. This enzyme removes leader sequences and leaves a 5'-phosphate on the tRNA's first nucleotide. Various mechanisms for removal of 3' trailer sequences are known, and the universal 3' end may be added or reconstituted by the CCA-adding enzyme (118–120). Early experiments in eukaryotes (121) suggest that certain modifications already appear at the level of the primary transcript, while some others take place only at the level of almost mature tRNA. It has been proposed that this may reflect the capability of these enzymes to recognize simple RNA structural elements present in the primary transcript (122), although differential expression and catalytic efficiencies of the various modification enzymes may arguably be an important factor. The tRNA maturation process is further complicated by elimination of introns inserted into the anticodon loops of many eukaryotic and archaeal tRNAs (123), and in eukaryotes, yet another layer of complexity is added by the export of partially mature tRNA species to the cytoplasm and possible later retrograde movement to the nucleus (120). The stepwise character of tRNA maturation and modification was clearly demonstrated in eukaryotes (121). Figure 5 (top panel) gives a brief visual outline of this complex maturation process and the implications of tRNA modifications in it. Recent findings suggest that tRNA modification does not passively accompany tRNA maturation but rather actively shapes the unstructured tRNA precursor into the fully functional tRNA having a complex three-dimensional structure. The interdependence of certain tRNA modifications (124–126) strongly indicates a step-by-step nature of at least parts of the whole tRNA maturation process. Indeed, speculation about a “molecular assembly line”, in which the maturing tRNA is handed over from one modification enzyme to the next, has been supported by a certain amount of experimental evidence through the characterization of multienzyme complexes (29, 127, 128), and synthetic lethality mutants between the La protein implicated in tRNA processing and several tRNA modification enzymes, including tRNA:m<sup>2</sup>G26-methyltransferase Trm1 or tRNA:pseudouridine synthase Pus4 (129).

Fully trimmed but incompletely modified tRNA can already partake in protein synthesis. This is obvious from studies on multiple deficiencies of tRNA modification enzymes, which lead to slow growth and translational defects both in bacteria [*E. coli* *trmH trmA truB* mutants (lacking G<sub>m</sub>18, m<sup>5</sup>U54, and Ψ55) (130)] and in yeast (lacking Ψ26–28, Ψ34, Ψ36, Ψ65, and Ψ67 from Pus1 and Ψ55 from Pus4) (131); further examples will be mentioned below. In yeast, this happens despite a certain extent of proofreading before export from the nucleus (132). It follows that systems for further surveillance and degradation act on a population of tRNAs, which are active in the translation




FIGURE 5: Biogenesis and degradation of tRNA. The top panel is a sketch of modification-induced structure acquisition during maturation. Early modification events are thought to be effected by enzymes recognizing partial tRNA features (122). The precursor is processed at the 3' and 5' ends on the basis of its intermediate tRNA structure, and further modifications are added (121), which lend additional stabilization to the tertiary structure. Certain modification enzymes are known to act on the unspliced precursor tRNA (123). In eukaryotes, splicing is effected after partial modification, but before transport from the nucleus to the cytosol. The bottom panel shows a sketch of targeted degradation of hypomodified tRNAs. It is thought that degradation proceeds as a consequence of destabilized three-dimensional interaction between loops D and T $\Psi$  of tRNA, followed by increased accessibility of the three-dimensional tRNA core and extremities to nucleases and poly(A) polymerase.

machinery and were traditionally considered to be shielded from degradation (133).

**Impact of Modifications on tRNA Processing, Stability, and Degradation.** Tight relationships between the integrity of global tRNA structure and stability have become especially apparent in mitochondrial tRNA, possibly because these are less well structured and less extensively modified than their cytosolic counterparts. Certain tRNA-dependent mitochondrial dysfunctions have been described as a consequence of the relative instability of aberrant tRNA species. Nucleotide substitutions observed in pathogenic tRNAs may affect various aspects of tRNA metabolism, like processing (134), modification, or aminoacylation (135, 136). Certain point mutations touch the tRNA's structural scaffold itself (39), while a mutation in the *pus1* gene causes MLASA<sup>1</sup> (myopathy with lactic acidosis and sideroblastic anemia) as a consequence of absent structural  $\Psi$  residues, e.g., at positions 27 and 28 of human mitochondrial tRNA (137, 138). In certain cases, the consequence of mitochondrial tRNA mutations clearly was an intrinsic instability of tRNA species in vivo. For example, mutation in human mitochondrial tRNA<sup>Ile</sup> reduces the stability of this tRNA in vivo and, as a consequence, reduces the level of amino acyl-tRNA. The half-life

of mutant tRNA is considerably reduced both in vivo and in vitro, and the melting temperature was also found to be reduced (139). Similar observations were also made for other point mutants of mitochondrial tRNAs (135, 136).

Recent studies of tRNA degradation pathways in yeasts clearly demonstrated that tRNA stability in vivo depends not only on the tRNA sequence itself but also on its modification profile. Several yeast deletion mutants with multiple deficiencies in tRNA modification exhibit a strong growth phenotype, indicating the importance of these modified residues in vivo.

**Pathways for Degradation of Hypomodified tRNA in Vivo.** Accelerated degradation of hypomodified tRNA species was observed in both bacteria and eukaryotes. Careful characterization of the  $\Delta TrmB$  mutant of the extreme thermophilic eubacterium *T. thermophilus* demonstrated the absence of m<sup>7</sup>G46 in tRNAs in vivo. Unexpectedly, the lack of m<sup>7</sup>G46 also led to hypomodification of Gm18 and m<sup>1</sup>G37. Even if aminoacylation properties of hypomodified tRNA species seem not to be affected, the melting temperature and stability of certain tRNAs in vivo were found to be considerably reduced. As a consequence, the protein synthesis is strongly depressed and the  $\Delta TrmB$  mutant exhibits a strong growth phenotype at 80 °C (140).

Degradation pathways of hypomodified tRNA species are far better studied in lower eukaryotes. Yeast strains deficient in

<sup>1</sup>Abbreviations: MLASA, myopathy with lactic acidosis and sideroblastic anemia; RTD, rapid tRNA decay; tRFs, tRNA fragments or tRNA halves.



Trm6 (or Trm61) tRNA:m<sup>1</sup>A58-methyltransferases display a reduced level of initiator tRNA<sup>Met</sup><sub>i</sub> at elevated temperatures (20, 141). The selective degradation of tRNA<sup>Met</sup><sub>i</sub> lacking m<sup>1</sup>A58 can be attributed to a weak tertiary interaction between loops D and T that is characteristic of tRNA<sup>Met</sup><sub>i</sub> (142). Similar data were also obtained for tRNA<sup>Val(AAC)</sup> in *trm8-Δ trm4-Δ* mutants, which lack m<sup>7</sup>G46 and m<sup>5</sup>C34, -40, -48, and -49 in the respective tRNAs (6). Reduced in vivo levels of tRNA<sup>Ser(CGA)</sup> were observed in yeast strains accumulating both a tRNA<sup>Ser</sup> mutation and deficiencies in Trm2 or Pus4 activities (143). The same observation was also made for yeast in strains lacking Um44 and ac<sup>4</sup>C12 due to deletion of *TRM44* and *TAN1* genes (144). In all studied cases, hypomodified tRNAs are rapidly degraded, but their degradation seems to proceed by two different independent pathways, depending on the cellular compartment.

The first degradation pathway for elimination of destabilized tRNA species in the nucleus is mediated by the TRAMP complex which contains the poly(A) polymerase Trf4, the RNA-binding protein Air2, and the DExH RNA helicase Mtr4p (20, 145). Following polyadenylation, these molecules are degraded by exonuclease Rrp6 and nuclear exosome.

An alternative pathway of tRNA degradation in the cytoplasm ("rapid tRNA decay", abbreviated RTD) is independent of TRF4/RRP6. The existence of an Trf4-independent pathway was initially suggested by analysis of La protein synthetic mutants, and its mechanism was determined by analysis of tRNA<sup>Val(AAC)</sup> degradation in the *trm8-Δ trm4-Δ* strain lacking m<sup>7</sup>G46 and m<sup>5</sup>C34, -40, -48, and -49 (5, 6). This RTD pathway includes the 5'-3' exonucleases Rat1 and Xrn1 and an auxiliary protein Met22. Met22 is probably not directly involved in RNA degradation but rather in cleavage of the pAp intermediate resulting from Rat1 and Xrn1 activity. However, the role of the RTD pathway seems to be more complicated since the aminoacylation level of tRNA<sup>Val(AAC)</sup> is affected by mutations of its components.

Taken together, these observations provide a link between specific pathways of tRNA degradation and general surveillance mechanisms ensuring RNA quality control in the cell.

## CONCLUSIONS AND OUTLOOK

Lessons learned from studying various species of modified RNA and its structural rearrangements (93–95, 146–150) supplement the principal conclusions drawn from tRNA-based investigations. Numerous RNA modifications are present not only in tRNAs but also in most other cellular RNAs, like rRNA, snRNA, snoRNA, and mRNA (7–9). Particularly interesting RNA structural data were obtained for a fragment of 28S rRNA bearing five highly conserved pseudouridine residues (150) and for a short RNA duplex imitating the interaction between the mRNA branch point and U2 snRNA (93–95, 146). The presence of modified residues was also revealed in recent studies of regulatory RNAs (151).

Coming full circle, modifications of the "structural" type would be present in the small fragments of cleaved tRNAs (tRFs or tRNA halves), which were recently discovered in eukaryotic cells (152, 153). These are thought to play a role in regulation of various functions (154, 155) and, possibly, malfunctions such as cancer. Indeed, the differential occurrence in cancer cells of modifications such as G<sub>m</sub> and C<sub>m</sub>, albeit in the anticodon, has been observed in the early going (156–159), and

urinary modified nucleosides are still being considered as cancer markers (160, 161). Little is known to date about the functional and structural importance of these modifications, but undoubtedly, the knowledge accumulated from tRNA studies will have a direct impact on our ability to detect and interpret modifications in newly discovered RNA species.

## REFERENCES

- Weixlbaumer, A., Murphy, F. V. t., Dziergowska, A., Malkiewicz, A., Vendeix, F. A., Agris, P. F., and Ramakrishnan, V. (2007) Mechanism for expanding the decoding capacity of transfer RNAs by modification of uridines. *Nat. Struct. Mol. Biol.* 14, 498–502.
- Agris, P. F. (2008) Bringing order to translation: The contributions of transfer RNA anticodon-domain modifications. *EMBO Rep.* 9, 629–635.
- Agris, P. F. (2004) Decoding the genome: A modified view. *Nucleic Acids Res.* 32, 223–238.
- Björk, G. R. (1995) Genetic dissection of synthesis and function of modified nucleosides in bacterial transfer RNA. *Prog. Nucleic Acid Res. Mol. Biol.* 50, 263–338.
- Chernyakov, I., Whipple, J. M., Kotelawala, L., Grayhack, E. J., and Phizicky, E. M. (2008) Degradation of several hypomodified mature tRNA species in *Saccharomyces cerevisiae* is mediated by Met22 and the 5'-3' exonucleases Rat1 and Xrn1. *Genes Dev.* 22, 1369–1380.
- Alexandrov, A., Chernyakov, I., Gu, W., Hiley, S. L., Hughes, T. R., Grayhack, E. J., and Phizicky, E. M. (2006) Rapid tRNA decay can result from lack of nonessential modifications. *Mol. Cell* 21, 87–96.
- Grosjean, H., and Benne, R. (1998) Modification and Editing of RNA, American Society for Microbiology Press, Washington, DC.
- Grosjean, H., Ed. (2005) Fine-Tuning of RNA Functions by Modification and Editing, Springer, Berlin.
- Grosjean, H., Ed. (2009) DNA and RNA Modification Enzymes: Structure, Mechanism, Function and Evolution, Landes Bioscience, Austin, TX.
- Helm, M. (2006) Post-transcriptional nucleotide modification and alternative folding of RNA. *Nucleic Acids Res.* 34, 721–733.
- Dalluge, J. J., Hashizume, T., Sopchik, A. E., McCloskey, J. A., and Davis, D. R. (1996) Conformational flexibility in RNA: The role of dihydrouridine. *Nucleic Acids Res.* 24, 1073–1079.
- Byrne, R. T., Konevega, A. L., Rodnina, M. V., and Antson, A. A. (2010) The crystal structure of unmodified tRNA<sup>Phe</sup> from *Escherichia coli*. *Nucleic Acids Res.* (in press).
- Saks, M. E., Sampson, J. R., and Abelson, J. N. (1994) The transfer RNA identity problem: A search for rules. *Science* 263, 191–197.
- Sampson, J. R., and Uhlenbeck, O. C. (1988) Biochemical and physical characterization of an unmodified yeast phenylalanine transfer RNA transcribed *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.* 85, 1033–1037.
- Perret, V., Garcia, A., Puglisi, J. D., Grosjean, H., Ebel, J.-P., Florentz, C., and Giegé, R. (1990) Conformation in solution of yeast tRNA<sup>Asp</sup> transcripts deprived of modified nucleotides. *Biochimie* 72, 735–744.
- Harrington, K. M., Nazarenko, I. A., Dix, D. B., Thompson, R. C., and Uhlenbeck, O. C. (1993) In vitro analysis of translational rate and accuracy with an unmodified tRNA. *Biochemistry* 32, 7617–7622.
- Muramatsu, T., Nishikawa, K., Nemoto, F., Kuchino, Y., Nishimura, S., Miyazawa, T., and Yokoyama, S. (1988) Codon and amino-acid specificities of a transfer RNA are both converted by a single post-transcriptional modification. *Nature* 336, 179–181.
- Isel, C., Lanchy, J.-M., Le Grice, S. F. J., Ehresmann, C., Ehresmann, B., and Marquet, R. (1996) Specific initiation and switch to elongation of human immunodeficiency virus type 1 reverse transcription require the post-transcriptional modifications of primer tRNA<sup>Lys</sup>. *EMBO J.* 15, 917–924.
- Helm, M., Brulé, H., Degoul, F., Cepanec, C., Leroux, J. P., Giegé, R., and Florentz, C. (1998) The presence of modified nucleotides is required for cloverleaf folding of a human mitochondrial tRNA. *Nucleic Acids Res.* 26, 1636–1643.
- Kadaba, S., Krueger, A., Trice, T., Krecic, A. M., Hinnebusch, A. G., and Anderson, J. (2004) Nuclear surveillance and degradation of hypomodified initiator tRNA<sup>Met</sup> in *S. cerevisiae*. *Genes Dev.* 18, 1227–1240.
- Moras, D., Comarmond, M. B., Fischer, J., Weiss, R., Thierry, J.-C., Ebel, J.-P., and Giegé, R. (1980) Crystal structure of yeast tRNA<sup>Asp</sup>. *Nature* 288, 669–674.



22. Cavarelli, J., Rees, B., Ruff, M., Thierry, J.-C., and Moras, D. (1993) Yeast tRNA<sup>Asp</sup> recognition by its cognate class II aminoacyl-tRNA synthetase. *Nature* 362, 181–184.
23. Ruff, M., Krishnaswamy, S., Boeglin, M., Poterszman, A., Mitschler, A., Podjarny, A., Rees, B., Thierry, J.-C., and Moras, D. (1991) Class II aminoacyl transfer RNA synthetases: Crystal structure of yeast aspartyl-tRNA synthetase complexed with tRNA<sup>Asp</sup>. *Science* 252, 1682–1689.
24. Ishitani, R., Nureki, O., Nameki, N., Okada, N., Nishimura, S., and Yokoyama, S. (2003) Alternative tertiary structure of tRNA for recognition by a posttranscriptional modification enzyme. *Cell* 113, 383–394.
25. Helm, M., Giegé, R., and Florentz, C. (1999) A Watson-Crick base-pair-disrupting methyl group (m<sup>1</sup>A9) is sufficient for cloverleaf folding of human mitochondrial tRNA<sup>Lys</sup>. *Biochemistry* 38, 13338–13346.
26. Voigts-Hoffmann, F., Hengesbach, M., Kobitski, A. Y., van Aerschot, A., Herdewijn, P., Nienhaus, G. U., and Helm, M. (2007) A methyl group controls conformational equilibrium in human mitochondrial tRNA<sup>Lys</sup>. *J. Am. Chem. Soc.* 129, 13382–13383.
27. Sissler, M., Helm, M., Frugier, M., Giegé, R., and Florentz, C. (2004) Aminoacylation properties of pathology-related human mitochondrial tRNA<sup>Lys</sup> variants. *RNA* 10, 841–853.
28. Helm, M., and Attardi, G. (2004) Nuclear Control of Cloverleaf Structure of Human Mitochondrial tRNA<sup>Lys</sup>. *J. Mol. Biol.* 337, 545–560.
29. Holzmann, J., Frank, P., Löffler, E., Bennett, K. L., Gerner, C., and Rossmann, W. (2008) RNase P without RNA: Identification and functional reconstitution of the human mitochondrial tRNA processing enzyme. *Cell* 135, 462–474.
30. Yang, H., and Lam, S. L. (2009) Effect of 1-methyladenine on thermodynamic stabilities of double-helical DNA structures. *FEBS Lett.* 583, 1548–1553.
31. Kobitski, A. Y., Hengesbach, M., Helm, M., and Nienhaus, G. U. (2008) Sculpting an RNA conformational energy landscape by a methyl group modification: A single-molecule FRET study. *Angew. Chem., Int. Ed.* 47, 4326–4330.
32. Wittenhagen, L. M., and Kelley, S. O. (2003) Impact of disease-related mitochondrial mutations on tRNA structure and function. *Trends Biochem. Sci.* 28, 605–611.
33. Sohm, B., Frugier, M., Brule, H., Olszak, K., Przykorska, A., and Florentz, C. (2003) Towards understanding human mitochondrial leucine aminoacylation identity. *J. Mol. Biol.* 328, 995–1010.
34. Sohm, B., Sissler, M., Park, H., King, M. P., and Florentz, C. (2004) Recognition of human mitochondrial tRNA<sup>Leu(UUR)</sup> by its cognate leucyl-tRNA synthetase. *J. Mol. Biol.* 339, 17–29.
35. Messmer, M., Pütz, J., Suzuki, T., Sauter, C., Sissler, M., and Catherine, F. (2009) Tertiary network in mammalian mitochondrial tRNA<sup>Asp</sup> revealed by solution probing and phylogeny. *Nucleic Acids Res.* 37, 6881–6895.
36. Sissler, M., Lorber, B., Messmer, M., Schaller, A., Pütz, J., and Florentz, C. (2008) Handling mammalian mitochondrial tRNAs and aminoacyl-tRNA synthetases for functional and structural characterization. *Methods* 44, 176–189.
37. Bullard, J. M., Cai, Y. C., Demeler, B., and Spremulli, L. L. (1999) Expression and characterization of a human mitochondrial phenylalanyl-tRNA synthetase. *J. Mol. Biol.* 288, 567–577.
38. Bullard, J. M., Cai, Y. C., and Spremulli, L. L. (2000) Expression and characterization of the human mitochondrial leucyl-tRNA synthetase. *Biochim. Biophys. Acta* 1490, 245–258.
39. Wittenhagen, L. M., Roy, M. D., and Kelley, S. O. (2003) The pathogenic U3271C human mitochondrial tRNA(Leu(UUR)) mutation disrupts a fragile anticodon stem. *Nucleic Acids Res.* 31, 596–601.
40. Messmer, M., Gaudry, A., Sissler, M., and Florentz, C. (2009) Pathology-related mutation A7526G (A9G) helps in the understanding of the 3D structural core of human mitochondrial tRNA<sup>Asp</sup>. *RNA* 15, 1462–1468.
41. Helm, M., Brulé, H., Friede, D., Giegé, R., Pütz, D., and Florentz, C. (2000) Search for characteristic structural features of mammalian mitochondrial tRNAs. *RNA* 6, 1356–1379.
42. Pütz, J., Dupuis, B., Sissler, M., and Florentz, C. (2007) Mamit-tRNA, a database of mammalian mitochondrial tRNA primary and secondary structures. *RNA* 13, 1184–1190.
43. Urbonavicius, J., Armengaud, J., and Grosjean, H. (2006) Identity elements required for enzymatic formation of N2,N2-dimethylguanosine from N2-monomethylated derivative and its possible role in avoiding alternative conformations in archaeal tRNA. *J. Mol. Biol.* 357, 387–399.
44. Steinberg, S., and Cedergren, R. (1995) A correlation between N2-dimethylguanosine presence and alternate tRNA conformers. *RNA* 1, 886–891.
45. Sampson, J. R., DiRenzo, A. B., Behlen, L. S., and Uhlenbeck, O. C. (1990) Role of the tertiary nucleotides in the interaction of yeast phenylalanine tRNA with its cognate synthetase. *Biochemistry* 29, 2523–2532.
46. Hall, K. B., Sampson, J. R., Uhlenbeck, O. C., and Redfield, A. G. (1989) Structure of an unmodified tRNA molecule. *Biochemistry* 28, 5794–5801.
47. Yue, D., Kintanar, A., and Horowitz, J. (1994) Nucleoside modifications stabilize Mg<sup>2+</sup> binding in *Escherichia coli* tRNA<sup>Val</sup>: An imino proton NMR investigation. *Biochemistry* 33, 8905–8911.
48. Serebrov, V., Clarke, R. J., Gross, H. J., and Kisselev, L. (2001) Mg<sup>2+</sup>-induced tRNA folding. *Biochemistry* 40, 6688–6698.
49. Serebrov, V., Vassilenko, K., Kholod, N., Gross, H. J., and Kisselev, L. (1998) Mg<sup>2+</sup> binding and structural stability of mature and in vitro synthesized unmodified *Escherichia coli* tRNA<sup>Phe</sup>. *Nucleic Acids Res.* 26, 2723–2728.
50. Vermeulen, A., McCallum, S. A., and Pardi, A. (2005) Comparison of the global structure and dynamics of native and unmodified tRNA<sup>Val</sup>. *Biochemistry* 44, 6024–6033.
51. Agris, P. F. (1996) The importance of being modified: Roles of modified nucleosides and Mg<sup>2+</sup> in RNA structure and function. *Prog. Nucleic Acid Res. Mol. Biol.* 53, 79–129.
52. Oliva, R., Tramontano, A., and Cavallo, L. (2007) Mg<sup>2+</sup> binding and archaeosine modification stabilize the G15 C48 Levitt base pair in tRNAs. *RNA* 13, 1427–1436.
53. Kobitski, A. Y., Nierth, A., Helm, M., Jaschke, A., and Nienhaus, G. U. (2007) Mg<sup>2+</sup>-dependent folding of a Diels-Alderase ribozyme probed by single-molecule FRET analysis. *Nucleic Acids Res.* 35, 2047–2059.
54. Jones, C. I., Spencer, A. C., Hsu, J. L., Spremulli, L. L., Martinis, S. A., DeRider, M., and Agris, P. F. (2006) A counterintuitive Mg<sup>2+</sup>-dependent and modification-assisted functional folding of mitochondrial tRNAs. *J. Mol. Biol.* 362, 771–786.
55. Chen, Y., Sierzputowska-Gracz, H., Guenther, R., Everett, K., and Agris, P. F. (1993) 5-Methylcytidine is required for cooperative binding of Mg<sup>2+</sup> and a conformational transition at the anticodon stem-loop of yeast phenylalanine tRNA. *Biochemistry* 32, 10249–10253.
56. Dao, V., Guenther, R. H., and Agris, P. F. (1992) The role of 5-methylcytidine in the anticodon arm of yeast tRNA<sup>Phe</sup>: Site-specific Mg<sup>2+</sup> binding and coupled conformational transition in DNA analogs. *Biochemistry* 31, 11012–11019.
57. Friederich, M. W., Gast, F. U., Vacano, E., and Hagerman, P. J. (1995) Determination of the angle between the anticodon and aminoacyl acceptor stems of yeast phenylalanyl tRNA in solution. *Proc. Natl. Acad. Sci. U.S.A.* 92, 4803–4807.
58. Friederich, M. W., and Hagerman, P. J. (1997) The angle between the anticodon and aminoacyl acceptor stems of yeast tRNA<sup>Phe</sup> is strongly modulated by magnesium ions. *Biochemistry* 36, 6090–6099.
59. Friederich, M. W., Vacano, E., and Hagerman, P. J. (1998) Global flexibility of tertiary structure in RNA: Yeast tRNA<sup>Phe</sup> as a model system. *Proc. Natl. Acad. Sci. U.S.A.* 95, 3572–3577.
60. Davis, D. R. (1998) Biophysical and conformational properties of modified nucleosides. In *Modification and Editing of RNA* (Grosjean, H., and Benne, R., Eds.) pp 85–102, American Society for Microbiology Press, Washington, DC.
61. Auffinger, P., Louise-May, S., and Westhof, E. (1995) Multiple molecular dynamics simulations of the anticodon loop of tRNA<sup>Asp</sup> in aqueous solution with counterions. *J. Am. Chem. Soc.* 117, 6720–6727.
62. Auffinger, P., Louise-May, S., and Westhof, E. (1999) Molecular dynamics simulations of solvated yeast tRNA<sup>Asp</sup>. *Biophys. J.* 76, 50–64.
63. Lee, C. H., and Tinoco, I., Jr. (1977) Studies of the conformation of modified dinucleoside phosphates containing 1,N6-ethenoadenosine and 2'-O-methylcytidine by 360-MHz <sup>1</sup>H nuclear magnetic resonance spectroscopy. Investigation of the solution conformations of dinucleoside phosphates. *Biochemistry* 16, 5403–5414.
64. Watanabe, K., Yokoyama, S., Hansske, F., Kasai, H., and Miyazawa, T. (1979) CD and NMR studies on the conformational thermostability of 2-thioribothymidine found in the TΨC loop of thermophile tRNA. *Biochem. Biophys. Res. Commun.* 91, 671–677.
65. Kawai, G., Yamamoto, Y., Kamimura, T., Masegi, T., Sekine, M., Hata, T., Imori, T., Watanabe, T., Miyazawa, T., and Yokoyama, S. (1992) Conformational rigidity of specific pyrimidine residues in

- tRNA arises from posttranscriptional modifications that enhance steric interaction between the base and the 2'-hydroxyl group. *Biochemistry* 31, 1040–1046.
66. Yokoyama, S., Watanabe, K., and Miyazawa, T. (1987) Dynamic structures and functions of transfer ribonucleic acids from extreme thermophiles. *Adv. Biophys.* 23, 115–147.
  67. Tisne, C., Rigourd, M., Marquet, R., Ehresmann, C., and Dardel, F. (2000) NMR and biochemical characterization of recombinant human tRNA<sup>(Lys)</sup><sub>3</sub> expressed in *Escherichia coli*: Identification of posttranscriptional nucleotide modifications required for efficient initiation of HIV-1 reverse transcription. *RNA* 6, 1403–1412.
  68. Chu, W. C., and Horowitz, J. (1989) <sup>19</sup>F NMR of 5-fluorouracil-substituted transfer RNA transcribed in vitro: Resonance assignment of fluorouracil-guanine base pairs. *Nucleic Acids Res.* 17, 7241–7252.
  69. Koshlap, K. M., Guenther, R., Sochacka, E., Malkiewicz, A., and Agris, P. F. (1999) A distinctive RNA fold: The solution structure of an analogue of the yeast tRNA<sup>Phe</sup> TΨC domain. *Biochemistry* 38, 8647–8656.
  70. Benas, P., Bec, G., Keith, G., Marquet, R., Ehresmann, C., Ehresmann, B., and Dumas, P. (2000) The crystal structure of HIV reverse-transcription primer tRNA<sup>(Lys,3)</sup> shows a canonical anticodon loop. *RNA* 6, 1347–1355.
  71. Ashraf, S. S., Sochacka, E., Cain, R., Guenther, R., Malkiewicz, A., and Agris, P. F. (1999) Single atom modification (O → S) of tRNA confers ribosome binding. *RNA* 5, 188–194.
  72. Tworowska, I., and Nikonowicz, E. P. (2006) Base pairing within the  $\psi$ 32, $\psi$ 39-modified anticodon arm of *Escherichia coli* tRNA<sup>Phe</sup>. *J. Am. Chem. Soc.* 128, 15570–15571.
  73. Cabello-Villegas, J., and Nikonowicz, E. P. (2000) Discriminating duplex and hairpin oligonucleotides using chemical shifts: Application to the anticodon stem-loop of *Escherichia coli* tRNA<sup>(Phe)</sup>. *Nucleic Acids Res.* 28, E74.
  74. Cabello-Villegas, J., Tworowska, I., and Nikonowicz, E. P. (2004) Metal ion stabilization of the U-turn of the A37 N6-dimethylallyl-modified anticodon stem-loop of *Escherichia coli* tRNA<sup>Phe</sup>. *Biochemistry* 43, 55–66.
  75. Cabello-Villegas, J., Winkler, M. E., and Nikonowicz, E. P. (2002) Solution conformations of unmodified and A(37)N(6)-dimethylallyl modified anticodon stem-loops of *Escherichia coli* tRNA<sup>(Phe)</sup>. *J. Mol. Biol.* 319, 1015–1034.
  76. Clore, G. M., Gronenborn, A. M., Piper, E. A., McLaughlin, L. W., Graessner, E., and Van Boom, J. H. (1984) The solution structure of a RNA pentadecamer comprising the anticodon loop and stem of yeast tRNA<sup>Phe</sup>. A 500 MHz proton NMR study. *Biochem. J.* 221, 737–751.
  77. Yarian, C. S., Basti, M. M., Cain, R. J., Ansari, G., Guenther, R. H., Sochacka, E., Czerwinska, G., Malkiewicz, A., and Agris, P. F. (1999) Structural and functional roles of the N1- and N3-protons of  $\Psi$  at tRNA's position 39. *Nucleic Acids Res.* 27, 3543–3549.
  78. Stuart, J. W., Koshlap, K. M., Guenther, R., and Agris, P. F. (2003) Naturally-occurring modification restricts the anticodon domain conformational space of tRNA<sup>(Phe)</sup>. *J. Mol. Biol.* 334, 901–918.
  79. Yarian, C., Marszalek, M., Sochacka, E., Malkiewicz, A., Guenther, R., Miskiewicz, A., and Agris, P. F. (2000) Modified nucleoside dependent Watson-Crick and wobble codon binding by tRNA<sup>LysUUU</sup> species. *Biochemistry* 39, 13390–13395.
  80. Agris, P. F., Guenther, R., Ingram, P. C., Basti, M. M., Stuart, J. W., Sochacka, E., and Malkiewicz, A. (1997) Unconventional structure of tRNA<sup>LysUUU</sup> anticodon explains tRNA's role in bacterial and mammalian ribosomal frameshifting and primer selection by HIV-1. *RNA* 3, 420–428.
  81. Bajji, A. C., and Davis, D. R. (2002) Synthesis of the tRNA(Lys,3) anticodon stem-loop domain containing the hypermodified ms<sup>2</sup>t<sup>6</sup>A nucleoside. *J. Org. Chem.* 67, 5352–5358.
  82. Durant, P. C., Bajji, A. C., Sundaram, M., Kumar, R. K., and Davis, D. R. (2005) Structural effects of hypermodified nucleosides in the *Escherichia coli* and human tRNA<sup>Lys</sup> anticodon loop: The effect of nucleosides s2U, mcm5U, mcm5s2U, mnm5s2U, t6A, and ms2t6A. *Biochemistry* 44, 8078–8089.
  83. Stuart, J. W., Gdaniec, Z., Guenther, R., Marszalek, M., Sochacka, E., Malkiewicz, A., and Agris, P. F. (2000) Functional anticodon architecture of human tRNA<sup>Lys3</sup> includes disruption of intraloop hydrogen bonding by the naturally occurring amino acid modification, t6A. *Biochemistry* 39, 13396–13404.
  84. Sundaram, M., Crain, P. F., and Davis, D. R. (2000) Synthesis and characterization of the native anticodon domain of *E. coli* tRNA<sup>(Lys)</sup>: Ssimultaneous incorporation of modified nucleosides mnm(5)s(2)U, t(6)A, and pseudouridine using phosphoramidite chemistry. *J. Org. Chem.* 65, 5609–5614.
  85. Sundaram, M., Durant, P. C., and Davis, D. R. (2000) Hypermodified nucleosides in the anticodon of tRNA<sup>Lys</sup> stabilize a canonical U-turn structure. *Biochemistry* 39, 12575–12584.
  86. Vendex, F. A., Dziergowska, A., Gustilo, E. M., Graham, W. D., Sproat, B., Malkiewicz, A., and Agris, P. F. (2008) Anticodon domain modifications contribute order to tRNA for ribosome-mediated codon binding. *Biochemistry* 47, 6117–6129.
  87. Davis, F. F., and Allen, F. W. (1957) Ribonucleic acids from yeast which contain a fifth nucleotide. *J. Biol. Chem.* 227, 907–915.
  88. Charette, M., and Gray, M. W. (2000) Pseudouridine in RNA: What, where, how, and why. *IUBMB Life* 49, 341–351.
  89. Davis, D. R. (1995) Stabilization of RNA stacking by pseudouridine. *Nucleic Acids Res.* 23, 5020–5026.
  90. Arnez, J. G., and Steitz, T. A. (1994) Crystal structure of unmodified tRNA<sup>Gln</sup> complexed with glutamyl-tRNA synthetase and ATP suggests a possible role for pseudo-uridines in stabilization of RNA structure. *Biochemistry* 33, 7560–7567.
  91. Auffinger, P., and Westhof, E. (1998) Effects of Pseudouridylation on tRNA Hydration and Dynamics. In *Modification and Editing of RNA* (Grosjean, H., and Benne, R., Eds.) pp 103–112, American Society for Microbiology Press, Washington, DC.
  92. Auffinger, P., and Westhof, E. (1998) Simulations of the molecular dynamics of nucleic acids. *Curr. Opin. Struct. Biol.* 8, 227–236.
  93. Newby, M. I., and Greenbaum, N. L. (2001) A conserved pseudouridine modification in eukaryotic U2 snRNA induces a change in branch-site architecture. *RNA* 7, 833–845.
  94. Newby, M. I., and Greenbaum, N. L. (2002) Investigation of Overhauser effects between pseudouridine and water protons in RNA helices. *Proc. Natl. Acad. Sci. U.S.A.* 99, 12697–12702.
  95. Newby, M. I., and Greenbaum, N. L. (2002) Sculpting of the spliceosomal branch site recognition motif by a conserved pseudouridine. *Nat. Struct. Biol.* 9, 958–965.
  96. Meroueh, M., Grohar, P. J., Qiu, J., SantaLucia, J., Jr., Scaringe, S. A., and Chow, C. S. (2000) Unique structural and stabilizing roles for the individual pseudouridine residues in the 1920 region of *Escherichia coli* 23S rRNA. *Nucleic Acids Res.* 28, 2075–2083.
  97. Davis, D. R., and Poulter, C. D. (1991) <sup>1</sup>H-<sup>15</sup>N NMR studies of *Escherichia coli* tRNA(Phe) from hisT mutants: A structural role for pseudouridine. *Biochemistry* 30, 4223–4231.
  98. Davanloo, P., Sprinzl, M., Watanabe, K., Albani, M., and Kersten, H. (1979) Role of ribothymidine in the thermal stability of transfer RNA as monitored by proton magnetic resonance. *Nucleic Acids Res.* 6, 1571–1581.
  99. Kumar, R. K., and Davis, D. R. (1997) Synthesis and studies on the effect of 2-thiouridine and 4-thiouridine on sugar conformation and RNA duplex stability. *Nucleic Acids Res.* 25, 1272–1280.
  100. Puglisi, J. D., and Tinoco, I., Jr. (1989) Absorbance melting curves of RNA. *Methods Enzymol.* 180, 304–325.
  101. Giegé, R., Helm, M., and Florentz, C. (2001) Classical and Novel Chemical Tools for RNA Structure Probing, Chapter 5, pp 71–89, Springer, Berlin.
  102. Wakeman, C. A., and Winkler, W. C. (2009) Analysis of the RNA backbone: Structural analysis of riboswitches by in-line probing and selective 2'-hydroxyl acylation and primer extension. *Methods Mol. Biol.* 540, 173–191.
  103. Grosjean, H., and Oshima, T. (2007) How nucleic acids cope with high temperature. In *Physiology and Biochemistry of Extremophiles* (Gerday, C., and Glansdorff, N., Eds.) pp 39–56, American Society for Microbiology Press, Washington, DC.
  104. Terui, Y., Ohnuma, M., Hiraga, K., Kawashima, E., and Oshima, T. (2005) Stabilization of nucleic acids by unusual polyamines produced by an extreme thermophile, *Thermus thermophilus*. *Biochem. J.* 388, 427–433.
  105. Hayrapetyan, A., Grosjean, H., and Helm, M. (2009) Effect of a quaternary pentamine on RNA stabilization and enzymatic methylation. *Biol. Chem.* 390, 851–861.
  106. Watanabe, K., Oshima, T., Iijima, K., Yamaizumi, Z., and Nishimura, S. (1980) Purification and thermal stability of several amino acid-specific tRNAs from an extreme thermophile, *Thermus thermophilus* HB8. *J. Biochem.* 87, 1–13.
  107. Agris, P. F., Koh, H., and Söll, D. (1973) The effect of growth temperatures on the in vivo ribose methylation of *Bacillus stearothermophilus* transfer RNA. *Arch. Biochem. Biophys.* 154, 277–282.
  108. Kumagai, I., Watanabe, K., and Oshima, T. (1980) Thermally induced biosynthesis of 2'-O-methylguanosine in tRNA from an extreme thermophile *Thermus thermophilus* HB27. *Proc. Natl. Acad. Sci. U.S.A.* 77, 1922–1926.



109. Watanabe, K., Shinma, M., Oshima, T., and Nishimura, S. (1976) Heat-induced stability of tRNA from an extreme thermophile, *Thermus thermophilus*. *Biochem. Biophys. Res. Commun.* 72, 1137–1144.
110. Kowalak, J. A., Dalluge, J. J., McCloskey, J. A., and Stetter, K. O. (1994) The role of posttranscriptional modification in stabilization of transfer RNA from hyperthermophiles. *Biochemistry* 33, 7869–7876.
111. Horie, N., Hara-Yokoyama, M., Yokoyama, S., Watanabe, K., Kuchino, Y., Nishimura, S., and Miyazawa, T. (1985) Two tRNA<sup>Phe</sup> species from an extreme thermophile, *Thermus thermophilus* HB8: Effect of 2-thiolation of ribothymidine on the thermostability of tRNA. *Biochemistry* 24, 5711–5715.
112. Droogmans, L., Roovers, M., Bujnicki, J. M., Tricot, C., Hartsch, T., Stalon, V., and Grosjean, H. (2003) Cloning and characterization of tRNA (m1A58) methyltransferase (TrmI) from *Thermus thermophilus* HB27, a protein required for cell growth at extreme temperatures. *Nucleic Acids Res.* 31, 2148–2156.
113. Shigi, N., Suzuki, T., Terada, T., Shirouzu, M., Yokoyama, S., and Watanabe, K. (2006) Temperature-dependent biosynthesis of 2-thioribothymidine of *Thermus thermophilus* tRNA. *J. Biol. Chem.* 281, 2104–2113.
114. Watanabe, K. (2007) Role of Modified Nucleosides in the Translation Function of tRNAs from Extreme Thermophilic Bacteria and Animal Mitochondria. *Bull. Chem. Soc. Jpn.* 80, 1253–1267.
115. Yamamoto, Y., Yokoyama, S., Miyazawa, T., Watanabe, K., and Higuchi, S. (1983) NMR analyses on the molecular mechanism of the conformational rigidity of 2-thioribothymidine, a modified nucleoside in extreme thermophile tRNAs. *FEBS Lett.* 157, 95–99.
116. Hopper, A. K., and Phizicky, E. M. (2003) tRNA transfers to the limelight. *Genes Dev.* 17, 162–180.
117. Randau, L., Schroder, I., and Söhl, D. (2008) Life without RNase P. *Nature* 453, 120–123.
118. Hartmann, R. K., Gossringer, M., Spath, B., Fischer, S., and Marchfelder, A. (2009) The making of tRNAs and more: RNase P and tRNase Z. *Prog. Nucleic Acid Res. Mol. Biol.* 85, 319–368.
119. Vörtler, S., and Mörl, M. (2010) tRNA-nucleotidyltransferases: Highly unusual RNA polymerases with vital functions. *FEBS Lett.* 584, 297–302.
120. Hopper, A. K., Pai, D. A., and Engelke, D. R. (2010) Cellular dynamics of tRNAs and their genes. *FEBS Lett.* 584, 310–317.
121. Nishikura, K., and De Robertis, E. M. (1981) RNA processing in microinjected *Xenopus* oocytes. Sequential addition of base modifications in the spliced transfer RNA. *J. Mol. Biol.* 145, 405–420.
122. Grosjean, H., Edqvist, J., Straby, K. B., and Giegé, R. (1996) Enzymatic formation of modified nucleosides in tRNA: Dependence on tRNA architecture. *J. Mol. Biol.* 255, 67–85.
123. Grosjean, H., Szweykowska-Kulinska, Z., Motorin, Y., Fasiolo, F., and Simos, G. (1997) Intron-dependent enzymatic formation of modified nucleosides in eukaryotic tRNAs: A review. *Biochimie* 79, 293–302.
124. Jiang, H. Q., Motorin, Y., Jin, Y. X., and Grosjean, H. (1997) Pleiotropic effects of intron removal on base modification pattern of yeast tRNA<sup>Phe</sup>: An in vitro study. *Nucleic Acids Res.* 25, 2694–2701.
125. Pintard, L., Lecoite, F., Bujnicki, J. M., Bonnerot, C., Grosjean, H., and Lapeyre, B. (2002) Trm7p catalyses the formation of two 2'-O-methylribose in yeast tRNA anticodon loop. *EMBO J.* 21, 1811–1820.
126. Rider, L. W., Ottosen, M. B., Gattis, S. G., and Palfey, B. A. (2009) Mechanism of dihydrouridine synthase 2 from yeast and the importance of modifications for efficient tRNA reduction. *J. Biol. Chem.* 284, 10324–10333.
127. Studte, P., Zink, S., Jablonowski, D., Bar, C., von der Haar, T., Tuite, M. F., and Schaffrath, R. (2008) tRNA and protein methylase complexes mediate zymocin toxicity in yeast. *Mol. Microbiol.* 69, 1266–1277.
128. Okada, K., Muneyoshi, Y., Endo, Y., and Hori, H. (2009) Production of yeast (m<sup>2</sup>G10) methyltransferase (Trm11 and Trm112 complex) in a wheat germ cell-free translation system. *Nucleic Acids Symp. Ser.*, 303–304.
129. Copela, L. A., Chakshumathi, G., Sherrer, R. L., and Wolin, S. L. (2006) The La protein functions redundantly with tRNA modification enzymes to ensure tRNA structural stability. *RNA* 12, 644–654.
130. Urbonavicius, J., Durand, J. M., and Björk, G. R. (2002) Three modifications in the D and T arms of tRNA influence translation in *Escherichia coli* and expression of virulence genes in *Shigella flexneri*. *J. Bacteriol.* 184, 5348–5357.
131. Grosshans, H., Lecoite, F., Grosjean, H., Hurt, E., and Simos, G. (2001) Pusp1p-dependent tRNA pseudouridylation becomes essential when tRNA biogenesis is compromised in yeast. *J. Biol. Chem.* 276, 46333–46339.
132. Lund, E., and Dahlberg, J. E. (1998) Proofreading and aminoacylation of tRNAs before export from the nucleus. *Science* 282, 2082–2085.
133. Engelke, D. R., and Hopper, A. K. (2006) Modified view of tRNA: Stability amid sequence diversity. *Mol. Cell* 21, 144–145.
134. Levinger, L., Giegé, R., and Florentz, C. (2003) Pathology-related substitutions in human mitochondrial tRNA<sup>Ile</sup> reduce precursor 3' end processing efficiency in vitro. *Nucleic Acids Res.* 31, 1904–1912.
135. Florentz, C. (2002) Molecular investigations on tRNAs involved in human mitochondrial disorders. *Biosci. Rep.* 22, 81–98.
136. Florentz, C., Sohm, B., Tryoen-Toth, P., Pütz, J., and Sissler, M. (2003) Human mitochondrial tRNAs in health and disease. *Cell. Mol. Life Sci.* 60, 1356–1375.
137. Bykhovskaya, Y., Casas, K., Mengesha, E., Inbal, A., and Fischel-Ghodsian, N. (2004) Missense mutation in pseudouridine synthase 1 (PUS1) causes mitochondrial myopathy and sideroblastic anemia (MLSA). *Am. J. Hum. Genet.* 74, 1303–1308.
138. Patton, J. R., Bykhovskaya, Y., Mengesha, E., Bertolotto, C., and Fischel-Ghodsian, N. (2005) Mitochondrial myopathy and sideroblastic anemia (MLSA): Missense mutation in the pseudouridine synthase 1 (PUS1) gene is associated with the loss of tRNA pseudouridylation. *J. Biol. Chem.* 280, 19823–19828.
139. Yasukawa, T., Hino, N., Suzuki, T., Watanabe, K., Ueda, T., and Ohta, S. (2000) A pathogenic point mutation reduces stability of mitochondrial mutant tRNA(Ile). *Nucleic Acids Res.* 28, 3779–3784.
140. Tomikawa, C., Yokogawa, T., Kanai, T., and Hori, H. (2010) N7-Methylguanine at position 46 (m7G46) in tRNA from *Thermus thermophilus* is required for cell viability at high temperatures through a tRNA modification network. *Nucleic Acids Res.* 38, 942–957.
141. Anderson, J., Phan, L., Cuesta, R., Carlson, B. A., Pak, M., Asano, K., Björk, G. R., Tamame, M., and Hinnebusch, A. G. (1998) The essential Gcd10p-Gcd14p nuclear complex is required for 1-methyladenosine modification and maturation of initiator methionyl-tRNA. *Genes Dev.* 12, 3650–3662.
142. Basavappa, R., and Sigler, P. B. (1991) The 3 Å crystal structure of yeast initiator tRNA: Functional implications in initiator/elongator discrimination. *EMBO J.* 10, 3105–3111.
143. Johansson, M. J., and Byström, A. S. (2002) Dual function of the tRNA(m<sup>5</sup>)U54 methyltransferase in tRNA maturation. *RNA* 8, 324–335.
144. Kotelawala, L., Grayhack, E. J., and Phizicky, E. M. (2008) Identification of yeast tRNA Um(44) 2'-O-methyltransferase (Trm44) and demonstration of a Trm44 role in sustaining levels of specific tRNA<sup>Ser</sup> species. *RNA* 14, 158–169.
145. Kadaba, S., Wang, X., and Anderson, J. T. (2006) Nuclear RNA surveillance in *Saccharomyces cerevisiae*: Trf4p-dependent polyadenylation of nascent hypomethylated tRNA and an aberrant form of 5S rRNA. *RNA* 12, 508–521.
146. Lin, Y., and Kielkopf, C. L. (2008) X-ray structures of U2 snRNA-branchpoint duplexes containing conserved pseudouridines. *Biochemistry* 47, 5503–5514.
147. Höbartner, C., Ebert, M.-O., Jaun, B., and Micura, R. (2002) RNA two-state conformation equilibria and the effect of nucleobase methylation. *Angew. Chem., Int. Ed.* 41, 605–609.
148. Höbartner, C., and Micura, R. (2003) Bistable secondary structures of small RNAs and their structural probing by comparative imino proton NMR spectroscopy. *J. Mol. Biol.* 325, 421–431.
149. Micura, R., and Höbartner, C. (2003) On secondary structure rearrangements and equilibria of small RNAs. *ChemBioChem* 4, 984–990.
150. Sumita, M., Desaulniers, J. P., Chang, Y. C., Chui, H. M., Clos, L., II, and Chow, C. S. (2005) Effects of nucleotide substitution and modification on the stability and structure of helix 69 from 28S rRNA. *RNA* 11, 1420–1429.
151. Zhao, X., Patton, J. R., Davis, S. L., Florence, B., Ames, S. J., and Spanjaard, R. A. (2004) Regulation of nuclear receptor activity by a pseudouridine synthase through posttranscriptional modification of steroid receptor RNA activator. *Mol. Cell* 15, 549–558.
152. Lee, Y. S., Shibata, Y., Malhotra, A., and Dutta, A. (2009) A novel class of small RNAs: tRNA-derived RNA fragments (tRFs). *Genes Dev.* 23, 2639–2649.
153. Thompson, D. M., Lu, C., Green, P. J., and Parker, R. (2008) tRNA cleavage is a conserved response to oxidative stress in eukaryotes. *RNA* 14, 2095–2103.
154. Kawamura, Y., Saito, K., Kin, T., Ono, Y., Asai, K., Sunohara, T., Okada, T. N., Siomi, M. C., and Siomi, H. (2008) *Drosophila*

- endogenous small RNAs bind to Argonaute 2 in somatic cells. *Nature* 453, 793–797.
155. Lau, N. C., Seto, A. G., Kim, J., Kuramochi-Miyagawa, S., Nakano, T., Bartel, D. P., and Kingston, R. E. (2006) Characterization of the piRNA complex from rat testes. *Science* 313, 363–367.
156. Tseng, W. C., Medina, D., and Randerath, K. (1978) Specific inhibition of transfer RNA methylation and modification in tissues of mice treated with 5-fluorouracil. *Cancer Res.* 38, 1250–1257.
157. Randerath, K., Agrawal, H. P., and Randerath, E. (1983) tRNA alterations in cancer. *Recent Results Cancer Res.* 84, 103–120.
158. Randerath, E., Gopalakrishnan, A. S., Gupta, R. C., Agrawal, H. P., and Randerath, K. (1981) Lack of a specific ribose methylation at guanosine 17 in Morris hepatoma 5123D tRNA<sup>Ser1</sup><sub>IGA</sub>. *Cancer Res.* 41, 2863–2867.
159. Kuchino, Y., Borek, E., Grunberger, D., Mushinski, J. F., and Nishimura, S. (1982) Changes of post-transcriptional modification of wye base in tumor-specific tRNA<sup>Phe</sup>. *Nucleic Acids Res.* 10, 6421–6432.
160. Li, H. Y., Wang, S. M., Liu, H. M., Li, J., Han, D., Bu, S. S., and Zhang, M. Z. (2008) Analysis of modified nucleosides in the urine of patients with malignant cancer by liquid chromatography/electrospray ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* 22, 3161–3171.
161. Kellner, S., Burhenne, J., and Helm, M. (2010) Detection of RNA modifications. *RNA Biol* 7, YYY.
162. Leontis, N. B., Stombaugh, J., and Westhof, E. (2002) The non-Watson-Crick base pairs and their associated isostericity matrices. *Nucleic Acids Res.* 30, 3497–3531.
163. Leontis, N. B., and Westhof, E. (2001) Geometric nomenclature and classification of RNA base pairs. *RNA* 7, 499–512.
164. Rozenski, J., Crain, P. F., and McCloskey, J. A. (1999) The RNA Modification Database: 1999 update. *Nucleic Acids Res.* 27, 196–197.