

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5776277>

Cobalamin-Dependent Methionine Synthase: Probing the Role of the Axial Base in Catalysis of Methyl Transfer between Methyltetrahydrofolate and Exogenous Cob(I)alamin or Cob(I)inami...

ARTICLE *in* BIOCHEMISTRY · JANUARY 2004

Impact Factor: 3.02 · DOI: 10.1021/bi035525t · Source: PubMed

CITATIONS

19

READS

54

3 AUTHORS, INCLUDING:



Rowena Green Matthews

University of Michigan

172 PUBLICATIONS 13,394 CITATIONS

SEE PROFILE

SUPPORTING INFORMATION

Cobalamin-dependent Methionine Synthase: Probing the Role of the Axial Base in Catalysis of Methyl Transfer between Methyltetrahydrofolate and Exogenous Cob(I)alamin and Cob(I)inamide

Jeanne Sirovatka Dorweiler, Richard G. Finke and Rowena G. Matthews*

Figure A. Initial rate of methyl transfer from CH₃-H₄folate to cob(I)inamide vs initial concentration of cob(I)inamide measured at 37 ° C. The concentration of MetH(2-649) was 6 μ M, and the concentration of CH₃-H₄folate was 1 mM. Error bars were calculated from a least squares fit to the initial data points. The rate constant calculated from these individual measurements is 183 M⁻¹ s⁻¹.




Figure B. Initial rate of methyl transfer from CH₃-H₄folate to cob(I)inamide vs the concentration of MetH(2-649) measured at 37 °C. The initial concentration of cob(I)inamide was 100 μ M and the concentration of CH₃-H₄folate was 1 mM. Error bars were calculated from a least squares fit to the initial data points. The constant calculated from these individual measurements is 220 M⁻¹ s⁻¹.




Figure C. Initial rate of methyl transfer from methylcobinamide to H₄folate vs the concentration of methylcobinamide measured at 37 °C. The concentration of MetH(2-649) was 6.5 μ M, and the concentration of H₄folate was 4 mM. At methylcobinamide concentrations lower than 200 μ M, the reaction produces a mixture of cob(I)inamide and cob(II)inamide, making the reaction kinetics more difficult to follow. Error bars were calculated from a least squares fit to the initial data points.




Figure D. Initial rate of methyl transfer from methylcobinamide to H₄folate vs the concentration of MetH(2-649) measured at 37 °C. The initial concentration of methylcobinamide was 200 μM , and the concentration of H₄folate was 4 mM. Error bars were calculated from least squares fitting of the initial data points. The rate constant calculated from the individual data points was 3170 $\text{M}^{-1} \text{s}^{-1}$.




Figure E. Effect of pyridine on methylcobalamin:H₄folate (●) and methylcobinamide:H₄folate (■) methyltransferase activities. Assays were performed as described in Materials and Methods. The initial concentrations in the methylcobalamin:H₄folate methyltransferase assay were: MetH(2-649), 7.4 μ M, methylcobalamin, 337 μ M, and H₄ folate, 4mM. The initial concentrations in the methylcobinamide:H₄folate methyltransferase assays were: MetH(2-649), 6.4 μ M; methylcobinamide, 400 μ M; and H₄ folate, 4 mM.




Figure F. Titration of methylcobinamide with pyridine. ~100 μM MeCbi in 50 μM

potassium phosphate buffer, pH 7.2 at 37 ° C was titrated with aliquots of neat pyridine. Spectra were measured in the presence of 0, 25, 50 100, 200 and 400 mM pyridine. The calculated association constant for this titration is $6.2 \pm 0.8 \text{ M}^{-1}$.




Figure G. Corrected methylcobinamide:H₄folate methyltransferase rate constants measured in the presence of pyridine vs. the percent base-on methylcobinamide measured under the same conditions. The uncorrected data for the effect of pyridine on the methylcobinamide:H₄folate methyltransferase assay are taken from Figure E, and corrected for the percent inhibition seen in the methylcobalamin:H₄folate methyltransferase assay data also shown in Figure E. These corrected second order rate constants are plotted against the percent base-on methylcobinamide calculated from the association constant measured in the experiment shown in Figure F.




Figure H. The observed rate constant for methyl transfer from $\text{PhNMe}_3^+\text{Cl}^-$ to cob(I)alamin vs the concentration of $\text{PhNMe}_3^+\text{Cl}^-$. Assays were conducted in 50 mM potassium phosphate buffer, pH 7.2 and 25 ° C. The initial concentration of cob(I)alamin was 100 μM , and values for k_{obs} were determined by dividing the observed initial rate of reaction by the initial concentration of cob(I)alamin. Error bars were calculated from least squares fitting of the initial data points. The second order rate constant calculated from the individual data points is $1.06 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$.




Figure I. The observed rate constant for methyl transfer from $\text{PhNMe}_3^+\text{Cl}^-$ to cob(I)inamide vs the concentration of $\text{PhNMe}_3^+\text{Cl}^-$. The initial concentration of cob(I)inamide was 100 μM , and values for k_{obs} were determined by dividing the observed initial rate of reaction by the initial concentration of cob(I)inamide. Error bars were calculated from least squares fitting of the initial data points. The second order rate constant calculated from the individual values for k_{obs} was $0.94 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$.

