

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/266975443>

Condition-Determined Multicomponent Reactions of 1,3-Dicarbonyl Compounds and Formaldehyde

ARTICLE in ACS COMBINATORIAL SCIENCE · OCTOBER 2014

Impact Factor: 3.03 · DOI: 10.1021/co5001019 · Source: PubMed

CITATION

1

READS

88

5 AUTHORS, INCLUDING:



Changhui Liu

Huazhong University of Science and Technology

10 PUBLICATIONS 20 CITATIONS

SEE PROFILE



Bingbing Lai

Huazhong University of Science and Technology

8 PUBLICATIONS 25 CITATIONS

SEE PROFILE



Amir Taheri

Huazhong University of Science and Technology

15 PUBLICATIONS 75 CITATIONS

SEE PROFILE



Yanlong Gu

Huazhong University of Science and Technology

87 PUBLICATIONS 2,946 CITATIONS

SEE PROFILE


Condition-Determined Multicomponent Reactions of 1,3-Dicarbonyl Compounds and Formaldehyde

Changhui Liu,^{†,§} Meng Shen,^{†,§} Bingbing Lai,[†] Amir Taheri,[†] and Yanlong Gu^{*,†,‡}

[†]Key Laboratory for Large-Format Battery Materials and System, Ministry of Education, School of Chemistry and Chemical Engineering, Huazhong University of Science and Technology, 430074, Wuhan, China

[‡]State Key Laboratory for Oxo Synthesis and Selective Oxidation, Lanzhou Institute of Chemical Physics, Lanzhou, 730000 People's Republic of China

S Supporting Information



ABSTRACT: By means of changing the reaction parameters, different products could be generated selectively starting from the same combination of substrates involving 1,3-dicarbonyl compounds and formaldehyde. This strategy enabled us to access diverse molecules without changing both starting material and reactor, maximizing thus the multifunctionality of the synthetic system. For example, starting from a 1,3-dicarbonyl compound, formaldehyde and 1,1-diphenylethylene, two kinds of products could be selectively formed including (i) a densely substituted dihydropyran and (ii) a C2-cinnamyl substituted 1,3-dicarbonyl compound. A one-pot three-component reaction of phenacylpyridinium salt, 1,3-dicarbonyl compound, and formaldehyde was also investigated, which produced either 2,4-diacyl-2,3-dihydrofuran or 2,4-diacyl-2-hydroxymethyl-2,3-dihydrofuran in good to excellent yield.

KEYWORDS: multicomponent reactions, diversity-oriented synthesis, condition-determined MCR, combinatorial chemistry

INTRODUCTION

Multicomponent reactions (MCRs) are convergent reactions of three or more starting materials, which have emerged as an efficient method for rapidly generating complex molecules with diverse functional substituents.¹ MCRs have often been used to establish expedient and ecofriendly chemical methods for the discovery of new chemical entities required by pharmaceutical and agrochemical industries.² Most MCRs were established by a reaction sequence involving (i) generation of an active intermediate through a reaction of the first two or three components and (ii) trapping of the intermediate with the same or another component. The generated intermediates generally have a very high reactivity, which enabled us to construct new molecular scaffolds sometimes. Therefore, most of the research interests focus on either the exploration of a suitable trapping reagent or derivatization of the intermediate with the hope of establishing a new reaction sequence.³ However, there is a perceived challenge in the face of the ever increasing demand for novel medicinally active compounds. This forced us to think how to

maximize the efficiency of establishing molecule libraries for biological screening.

Control of the reaction selectivity, for example, chemo-, stereo-, and regioselectivity, is one of the most important objectives of organic chemistry.⁴ Many different reaction parameters such as temperature, pressure, solvent, and catalyst type, and other factors can be utilized to modulate the selectivity of organic reactions. Because three or more substrates are involved in a MCR, it is conceivable that by carefully manipulating the reaction parameters, it might be possible to establish two or more MCRs with the same combination of substrates. This strategy can increase the number of MCRs without increasing the number of substrates. Previously, a few reports have disclosed some individual examples of the synthesis of different products from the same substrates.⁵ It offered an effective means to us for enriching the diversity of

Received: July 3, 2014

Revised: September 18, 2014


the MCR product libraries, which in turn facilitates biological screening.

We were attracted by the unique advantages of this strategy and started a research program on this topic some time ago. To utilize this strategy, we have to find a suitable intermediate, which not only has a high reactivity but also is amenable to diversified reaction modes, allowing us to trap it in different reaction pathways. Recently, the Knoevenagel reaction of 1,3-dicarbonyl compounds and formaldehyde has been used to create MCRs.⁶ The generated 2-methylene-1,3-dicarbonyl intermediate not only acts as an *oxo*-diene in Diels–Alder reaction but also serves as a Michael acceptor in conjunction with some Michael donors, favoring thus construction of many MCRs.⁷ We were attracted by the multifunctionality of this intermediate and started our MCR investigation with a combination of 1,3-dicarbonyl compounds and formaldehyde.

RESULTS AND DISCUSSION

Initially, a three-component reaction of 1,1-diphenylethylene **1a**, acetoacetone **2a**, and formaldehyde was investigated. As shown in Scheme 1, when formalin was used as HCHO source,

Scheme 1. Three-Component Reaction of 1a, 2a, and Formaldehyde



a dihydropyran **3a** was obtained in 75% yield after 5 h of reaction at 80 °C in acetonitrile. The reaction is very clean, and the unreacted 1,1-diphenylethylene can be fully recovered. Interestingly, when paraformaldehyde was used as the HCHO

source, a different compound, **4a** was obtained in 80% yield in the presence of toluenesulfonic acid (PTSA) at 60 °C. These results imply that the source of HCHO and the reaction conditions played key roles in controlling the reaction selectivity.

These results also gave us impetus to investigate the reaction mechanism. It is well-known that **3a** was formed through a tandem Knoevenagel/*oxo*-Diels–Alder reaction pathway, in which **1a** acted as a dienophile to trap the generated 3-methylene-2,4-pentadione (intermediate **I**, Figure 1).⁸ In order to shed light on the mechanism for the formation of **4a**, several control experiments were then carried out. First, although the Prins cyclization product of **1a** and paraformaldehyde, **5a**, could be formed with the aid of PTSA catalyst, it cannot be converted into **4a** under the reaction conditions (Scheme 2). Because **3a** could be also detected during the

Scheme 2. Control Experiments for Understanding the Mechanism of 4a Formation





Figure 1. Proposed mechanism for the formations of 3a and 4a.




Figure 2. Progress of a PTSA-catalyzed reaction of **1a**, **2a**, and paraformaldehyde monitored by ^1H NMR.

reaction forming **4a** (Figure 2), we therefore treated **3a** with PTSA in acetonitrile. After 5 h of reaction at 80 °C, **4a** was formed in 80% of yield. However, this result is insufficient to lead us to draw a conclusion for the formation of **4a** because kinetic investigation of the reaction between **1a**, **2a**, and paraformaldehyde revealed that no significant accumulation of **3a** was observed during the reaction (Figure S2, Supporting Information). In addition, monitoring of the reaction progress by means of ^1H NMR demonstrated that (i) intermediate **I** was generated quickly in the first 30 min of the reaction and then its concentration gradually decreased and (ii) the formation of **4a** occurred in the beginning of the reaction and lasted all 12 h as the concentration of **4a** increased gradually during the reaction. All these results led us to deduce that **4a** might be formed through a direct Michael reaction of the intermediate **I** and **1a**. Because the isolation of pure intermediate **I** is not possible, methyl vinyl ketone **6a** was therefore used as a Michael acceptor, which has a relatively lower reactivity than the intermediate **I**. As shown in Scheme 2, the expected product **7a** was obtained in 80% of yield. This result implies that **4a** might be formed through a tandem Knoevenagel/Michael reaction pathway. Incidentally, because Knoevenagel/oxo-Diels–Alder reaction is a noncatalytic reaction sequence, formation of **3a** is inevitable during the synthesis of **4a**. A Knoevenagel/oxo-Diels–Alder/ring-opening reaction sequence may be also operative for the formation of **4a** (Figure 1). The ring-opening reaction pathway is able to convert **3a** into **4a**, ensuring thus a good selectivity of **4a**.

The PTSA/acetonitrile system was successfully used to establish the three-component reactions of a wide range of 1,3-dicarbonyl compounds, $(\text{HCHO})_n$, and 1,1-diarylethylenes, and the results are shown in Figure 3. Many linear β -ketoesters or 1,3-diketones reacted readily with **1a** and paraformaldehyde, affording the corresponding products in generally excellent yields. Cyclopropyl and methoxy groups are tolerable in this system (**4g**). A secondary β -ketoamide can also be used uneventfully (**4i**). Some other 1,1-diphenylethylene derivatives

could also be used. Particularly, a diarylethylene with thienyl group participated readily in this reaction as well (**4l**). It is significant to note that **1a** could be replaced by 1,1-diphenylethanol, which is less-expensive compared with **1a**, in this reaction. This offered a cost-effective alternative route to access **4a**-type products (Scheme 3). It should be noted also that the same products in Figure 3 could be synthesized by many reported methods, most of which involve the use of harsh conditions and expensive reagents and suffer from the lack of simplicity and also the yields and selectivities reported are sometimes far from satisfactory.⁹ Therefore, the present three-component reaction opened a simple and effective route to access these compounds. However, attempts to use normal 1-arylethylenes, such as 4-methylstyrene and α -methylstyrene, as substrates in the PTSA/acetonitrile system were in vain. The reactions suffered from a lack of selectivity as messy mixtures were formed in these cases. By the same token, formaldehyde cannot be replaced by other aliphatic or aromatic aldehydes in this reaction.

The above-mentioned results demonstrated that the development of condition-determined MCRs based on a combination of a 1,3-dicarbonyl compound and formaldehyde is indeed possible. Encouraged by these results, we then investigated the condensation reaction of *N*-phenacylpyridinium bromide **7a**, 1,3-cyclohexanedione **2b**, and formaldehyde, which can hopefully produce a 2,4-diacyl-2,3-dihydrofuran derivative, **8a** through a cascade Knoevenagel/[4 + 1] annulation reaction under appropriate conditions.¹⁰ The reaction was also triggered by a Knoevenagel condensation of **2b** with formaldehyde, which generated a 2-methylene-1,3-cyclohexanedione intermediate (**II**) that can be trapped by phenacylpyridinium salt through [4 + 1] annulation reaction (Figure 4). As shown in Table 1, a product was indeed formed in the presence of an inorganic base, $\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$, in DMSO; however, it was the hydroxymethylation product of the expected one, **8a'**. Because compound **8a** was also detected at the end of the reaction, we therefore deduced that **8a'** might be formed through a cascade





Figure 3. PTSA-catalyzed three-component reaction of 1,1-diarylethylene, 1,3-dicarbonyl compounds, and $(\text{HCHO})_n$.

Scheme 3. PTSA-Catalyzed Three-Component Reaction of 1,1-Diphenylethanol, 2a, and Paraformaldehyde



166 Knoevenagel/[4 + 1] annulation/hydroxymethylation reaction
 167 (Figure 4). Indeed, treatment of **8a** in DMSO in the presence
 168 of paraformaldehyde resulted in an evident formation of **8a'**
 169 (Scheme 4). To our great delight, the quasi-four-component
 170 reaction was found to be very efficient, and the yield of **8a'**
 171 reached 83% after 4 h of reaction at 80 °C (entry 1). This
 172 observation encouraged us to scrutinize the effects of reac-
 173 tion parameters including base, solvent, and reaction temper-
 174 ature. No or only trace amount of product was obtained with
 175 inorganic bases, such as $\text{K}_3\text{PO}_4 \cdot 3\text{H}_2\text{O}$ and K_2CO_3 (entries 2
 176 and 3). Organic bases like NEt_3 and DBU were also ineffective

for this reaction (entries 4 and 5). Among different solvents 177
 tested in the reaction, DMSO clearly stood out, producing **8a'** 178
 with the highest yield, with DMF and acetonitrile in a distant 179
 second place (ca. 40% yields). PEG400, ionic liquid [BMIm]- 180
 BF_4 , and water resulted in significantly lower efficiency of the 181
 reaction (entries 8 to 10). Ratio of **7a/2b/HCHO** can also 182
 significantly affect the yield of **8a'**, and the best is **7a/2b/** 183
 $\text{HCHO} = 1.0/2.0/2.5$. Poor yields were obtained with much 184
 excess of **2b** or HCHO , which might result from an extensive 185
 formation of a byproduct through Knoevenagel/Michael 186
 reaction of **2b** and HCHO (entries 11 and 12). Interestingly, 187
 when ratio of the **7a/2b/HCHO** was changed to 1.0/1.5/2.0, 188
8a was produced as a major product, and **8a'** was formed only 189
 in tiny amounts (entry 13). These results imply that substrate 190
 ratio has a subtle influence on the reaction selectivity, and 191
 amounts of **2b** and formaldehyde are both important to 192
 determine the reaction selectivity. It offered us a possible means 193
 to control the reaction selectivity by tuning the reaction param- 194
 eters. It should be noted that, in all the previous reports on 195
 Knoevenagel/[4 + 1] annulation sequential reaction of 196





Figure 4. Proposed mechanism for the formation of **8a** and **8a'**.

Table 1. Three-Component Reaction of *N*-Phenacylpyridinium Bromide, Acetylacetone, and Formaldehyde under Different Conditions^a




entry	base	solvent	ratio of 7a/ 2b/HCHO	temp (°C)	yield (%)	
					8a	8a'
1	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	80	<5	83
2	K ₃ PO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	80	<5	<5
3	K ₂ CO ₃	DMSO	1.0/2.0/2.5	80	0	0
4	Et ₃ N	DMSO	1.0/2.0/2.5	80	8	5
5	DBU	DMSO	1.0/2.0/2.5	80	0	0
6	K ₂ HPO ₄ ·3H ₂ O	DMF	1.0/2.0/2.5	80	<5	36
7	K ₂ HPO ₄ ·3H ₂ O	CH ₃ CN	1.0/2.0/2.5	80	<5	39
8	K ₂ HPO ₄ ·3H ₂ O	PEG400	1.0/2.0/2.5	80	<5	<5
9	K ₂ HPO ₄ ·3H ₂ O	[BMIIm]BF ₄	1.0/2.0/2.5	80	9	<5
10	K ₂ HPO ₄ ·3H ₂ O	H ₂ O	1.0/2.0/2.5	80	8	<5
11	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/1.0/2.0	80	<5	67
12	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.5/3.0	80	9	22
13	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/1.5/2.0	80	80	<5
14	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	50	0	5
15	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	100	5	10
16 ^b	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	80	24	51
17 ^c	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	80	<5	11
18 ^d	K ₂ HPO ₄ ·3H ₂ O	DMSO	1.0/2.0/2.5	80	0	0

^aConditions: 1a, 1.0 mmol; paraformaldehyde was used as HCHO source; solvent 1.0 mL; reaction time 4 h. ^bReaction time 2 h.

^cAqueous solution of formaldehyde was used as HCHO source.

^dTrioxymethylene was used as HCHO source.

Scheme 4. Hydroxymethylation of 8a to 8a'




phenacylpyridinium salt, the use of aromatic aldehyde is mandatory in order to facilitate control of the reaction selectivity.¹¹ The present synthesis of 8a-like 2,3-dihydrofurans represents the first example of using nonaromatic aldehyde as substrate. Additionally, the reaction was also affected by temperature and reaction time, and the maximum yield of 8a' was obtained at 80 °C after 4 h of reaction (entries 14 to 16). It is worthwhile to note that, under the optimal conditions, efforts to replace paraformaldehyde with either formalin (37 wt %) or trioxymethylene were in vain (entries 17 and 18).

We also probed the scope of the reaction with respect to both the pyridinium bromide and the 1,3-dicarbonyl compounds. As evidenced by the results in Table 2, *N*-phenacylpyridinium bromides with both electron-donating and moderately electron-withdrawing groups smoothly reacted with 2b, producing 2-hydroxymethylated 2,3-dihydrofuran derivatives in generally


good yields (entries 1–5). By decreasing the ratio of 7/2/213 HCHO, we are able to suppress the hydroxymethylation. 214 Particularly, when *N*-(4-methoxyphenacyl)pyridinium bromide 215 was used, yield of the tandem Knoevenagel/[4 + 1] annulation 216 product, 8e, reached 95% with the ratio of 7/2/HCHO = 1.0/ 217 1.5/2.0. However, increasing the ratio to 1.0/2.0/2.5 was in vain 218 to obtain its hydroxymethylated counterpart, 8e'. In this case, 219 high excess of paraformaldehyde has to be used in order to get a 220 good yield of 8e' (entry 4). Acetoacetone 2a reacted readily with 221 7a and formaldehyde; however, extra effort has to be paid to 222 control the reaction selectivity because change of the substrate 223 ratio cannot alter significantly the product distribution. Addition 224 of solvent amount of xylene, which constructed a biphasic system 225 along with DMSO, proved to be an effective way to suppress the 226 hydroxymethylation reaction of 8g (entry 6). In order to get 8g', 227 the reaction has to be performed at 30 °C. Fortunately, when the 228 other *N*-phenacylpyridinium bromide derivatives were used to 229 react with 2a, it was quite easy to control the reaction 230 selectivity. In the presence of a large excess of paraformaldehyde, 231 the hydroxymethylated product will be preferentially 232 formed as usual, whereas the major products are the non- 233 hydroxymethylated 2,3-dihydrofurans when the ratio of 7/2b/ 234 HCHO is 1.0/2.0/2.5 (entries 7–14). This strategy is particularly 235 effective for tuning the selectivity of condensation between *N*-(4- 236 phenylphenacyl)pyridinium bromide, 2a, and formaldehyde. Both 237 hydroxymethylated and nonhydroxymethylated products could be 238 obtained in more than 90% yields in this case (entry 10). 1-(2- 239 Naphthoylethyl)pyridinium bromide also proved to be an 240 eligible substrate that reacted smoothly with either 2b or 2a, 241 providing both hydroxymethylated and nonhydroxymethylated 242 products in good yields (entries 5 and 11). It should be noted that 243 the OH group in the phenacylpyridinium salt can be delivered 244 uneventfully (entry 12). This facilitates further conversions of 245 the obtained 2,3-dihydrofurans. A heterocyclic group, such as 246 thienyl, is also tolerable in the present reaction (entry 13). 247 Reactions with β -ketoesters also proceeded very well, and the 248 products succeeded the ester moieties without any damage 249 (entries 14–17). The ether fragment in 2-methoxyethyl 250 acetoacetate is also tolerable. Due to an insusceptibility of the 251 reaction toward the change of the substrate ratio, the DMSO/ 252 xylene biphasic system was employed when methyl isobutyr- 253 ylate and 2-methoxyethyl acetoacetate were used to react 254 with 7a (entries 16 and 17). It should be noted that when an 255 aqueous solution of acetaldehyde was used instead of 256 paraformaldehyde, no expected substituted dihydrofuran 257 derivative was formed. 258

Because the hydroxymethylated products contain some 259 reactive groups, we suspected that these molecules might be 260 susceptible under acidic conditions. As we expected, treatment 261 of 8g' in ethanol in the presence of Sc(OTf)₃ resulted in 262 selective formation of diphenyl derivative 9a (Scheme 5). The 263 existence of the hydroxyl group in 8g' proved to be crucial for 264 rendering this reaction possible because no reaction was ob- 265 served when 8g was used as substrate under the identical con- 266 ditions. The initial step of the reaction might be an intra- 267 molecular ring-opening and ring-closing reaction of 8g' with 268 the aid of acid catalyst, which generated an epoxide inter- 269 mediate (IV). The following ring-opening of IV with ethanol 270 produced an intermediate V that underwent an intramolecular 271 aldol reaction¹² and subsequent retro-Claisen condensation¹³ 272 to form the final product 9a. This reaction not only displayed 273 an interesting reaction sequence but also offered us the first 274

Table 2. Substrate Scope of Three-Component Reaction of *N*-Phenacylpyridinium Bromides, 1,3-Dicarbonyl Compounds, and Paraformaldehyde^a




7




2

$(\text{HCHO})_n$
 $\xrightarrow[\text{DMSO, 80 } ^\circ\text{C, 4 h}]{\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O (2.0 equiv)}}$



8b-8r



8b'-8r'




































entry	1/2/HCHO	major product	yield (%) ^b	1/2/HCHO	major product	Yield (%) ^b
1	1.0/1.5/2.0		8b 74 (12)	1.0/2.0/2.5		8b' 60 (14)
2	1.0/1.5/2.0		8c 62 (9)	1.0/2.0/2.5		8c' 68 (13)
3	1.0/1.5/2.0		8d 70 (8)	1.0/2.0/2.5		8d' 70 (10)
4	1.0/1.5/2.0		8e 95 (< 1)	1.0/2.0/7.0		8e' 71 (19)
5	1.0/1.5/2.0		8f 86 (7)	1.0/2.0/7.0		8f' 76 (14)
6	1.0/2.0/2.5		8g 82 ^c (6)	1.0/2.0/2.5		8g' 85 ^d (7)
7	1.0/2.0/2.5		8h 69 (16)	1.0/2.0/7.0		8h' 70 (11)
8	1.0/2.0/2.5		8i 73 (8)	1.0/2.0/7.0		8i' 51 (9)
9	1.0/2.0/2.5		8j 94 (< 5)	1.0/2.0/7.0		8j' 93 (< 5)
10	1.0/2.0/2.5		8k 72 (12)	1.0/2.0/7.0		8k' 57 (18)
11	1.0/2.0/2.5		8l 65 (17)	1.0/2.0/7.0		8l' 62 (15)
12	1.0/2.0/2.5		8m 75 (11)	1.0/2.0/7.0		8m' 50 (13)

Table 2. continued

entry	1/2/HCHO	major product	yield (%) ^b	1/2/HCHO	major product	Yield (%) ^b
13	1.0/2.0/2.5		8n 81 (10)	1.0/2.0/7.0		8n' 61 (16)
14	1.0/2.0/2.5		8o 49 (8)	1.0/2.0/7.0		8o' 54 (< 5)
15	1.0/2.0/2.5		8p 60 (14)	1.0/2.0/7.0		8p' 71 (7)
16	1.0/2.0/2.5		8q 50 ^c (9)	1.0/2.0/2.5		8q' 76 (13)
17	1.0/2.0/2.5		8r 50 ^c (11)	1.0/2.0/2.5		8r' 97 (< 1)

^aConditions: *N*-Phenacylpyridinium bromide 0.5 mmol; DMSO 1.0 mL; K₂HPO₄·3H₂O 1.0 mmol; 80 °C, 4 h. ^bValue in parentheses is the yield of the minor product. ^cXylene was added. ^dReaction performed at 30 °C.

Scheme 5. Conversion of 8g' to 9a



example that can produce aromatic ether from five-member ring heterocycles without oxidation.¹⁴

CONCLUSION

Some condition-determined MCRs of 1,3-dicarbonyl compounds and formaldehyde were reported. Reaction of a 1,3-dicarbonyl compound, formaldehyde, and 1,1-diphenylethylene produced either a densely substituted 3,4-dihydropyran or a C2-cinnamyl substituted 1,3-dicarbonyl compound. A pseudo-four-component reaction of *N*-phenacylpyridinium bromide, 1,3-dicarbonyl compound, and formaldehyde was also developed, which involved a hitherto unreported Knoevenagel/[4 + 1] annulation/hydroxymethylation reaction sequence. All these examples demonstrated that the concept of condition-determined MCR is indeed useful for divergence-oriented organic synthesis.

EXPERIMENTAL SECTION

General. Melting points were determined by microscopic melting point meter and were uncorrected. IR spectra were recorded on a FT-IR, Bruker (EQUINOX 55), using KBr pellets or neat liquid technology. ¹H and ¹³C NMR spectra were recorded on a Bruker AV-400. Chemical shifts are expressed in ppm relative to Me₄Si in solvent. All chemicals used were of reagent grade and were used as received without further purification. All reactions were conducted in a 10 mL V-type flask equipped with triangle magnetic stirring.

Reaction of 1,1-Diarylethylene, 1,3-Dicarbonyl Compounds, and (HCHO)_n. In a typical reaction, the 1,3-dicarbonyl compound (0.2 mmol) was mixed with paraformaldehyde (0.2 mmol), 1,1-diarylethylene (0.25 mmol), and PTSA (0.02 mmol, 3.8 mg, 10% mol) in acetonitrile (1.0 mL). The mixture was then stirred at 60 °C for 12 h. After reaction, the mixture was cooled to room temperature, and the product was obtained by isolation with preparative TLC (eluting solution, petroleum ether/ethyl acetate = 5/1 (v/v)). Tests for substrate scope were all performed with an analogous procedure.

Three-Component Reaction of *N*-Phenacylpyridinium Bromides, 1,3-Dicarbonyl Compounds, and (HCHO)_n. *N*-Phenacylpyridinium bromide (0.25 mmol) was mixed with the 1,3-dicarbonyl compound (0.375 mmol), and paraformaldehyde (0.5 mmol). The mixture was then stirred at 80 °C for 4 h. After reaction, the mixture was cooled to room temperature, and the product 2,4-diacyl-2,3-dihydrofuran derivative was obtained by isolation with preparative TLC (eluting solution, petroleum ether/ethyl acetate = 10/1 (v/v)). Tests for substrate scope were all performed with an analogous procedure. The hydroxymethylation product was obtained by only changing the ratio of *N*-phenacylpyridinium bromide, 1,3-dicarbonyl compound, and paraformaldehyde to 1.0/2.0/2.5.

Synthesis of 9a from 8g'. Compound 8g' (52 mg, 0.2 mmol) and Sc(OTf)₃ (10 mg, 10% mol) was added to ethanol

(1 mL), and the mixture was then stirred at 80 °C for 4 h. After reaction, the product **9a** was obtained by isolation with preparative TLC (eluting solution, petroleum ether/ethyl acetate = 20/1 (v/v)) with yield of 61%.

■ ASSOCIATED CONTENT

● Supporting Information

Additional experimental details, reaction progress of the three-component reaction of **1a**, **2a**, and paraformaldehyde, characterization data of new compounds, and NMR spectra of compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Phone: (0)86-(0)27-87 54 37 32. Fax: (0)86-(0)27-87 54 45 32. E-mail: klgyl@hust.edu.cn.

Author Contributions

[§]Changhui Liu and Meng Shen contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors thank for National Natural Science Foundation of China for the financial support (Grants 21173089 and 21373093) and also for the Analytical and Testing Center of HUST. Chutian Scholar Program of the Hubei Provincial Government and the Cooperative Innovation Center of Hubei Province are also acknowledged. This work is also supported by the Fundamental Research Funds for the Central Universities of China (Grant 2014ZZGH019).

■ REFERENCES

(1) (a) Toure, B. B.; Hall, D. G. Natural product synthesis using multicomponent reaction strategies. *Chem. Rev.* **2009**, 109 (9), 4439–4486. (b) Sunderhaus, J. D.; Martin, S. F. Applications of multicomponent reactions to the synthesis of diverse heterocyclic scaffolds. *Chem.—Eur. J.* **2009**, 15 (6), 1300–1308. (c) Ganem, B. Strategies for innovation in multicomponent reaction design. *Acc. Chem. Res.* **2009**, 42 (3), 463–472. (d) Jiang, B.; Rajale, T.; Wever, W.; Tu, S.-J.; Li, G. Multicomponent reactions for the synthesis of heterocycles. *Chem.—Asian J.* **2010**, 5 (11), 2318–2335. (e) Yu, J.; Shi, F.; Gong, L.-Z. Brønsted-acid-catalyzed asymmetric multicomponent reactions for the facile synthesis of highly enantioenriched structurally diverse nitrogenous heterocycles. *Acc. Chem. Res.* **2011**, 44 (11), 1156–1171. (f) de Graaff, C.; Ruijter, E.; Orru, R. V. A. Recent developments in asymmetric multicomponent reactions. *Chem. Soc. Rev.* **2012**, 41 (10), 3969–4009. (2) (a) Ramazani, A.; Kazemizadeh, A. R. Preparation of stabilized phosphorus ylides via multicomponent reactions and their synthetic applications. *Curr. Org. Chem.* **2011**, 15 (23), 3986–4020. (b) Bonne, D.; Coquerel, Y.; Constantieux, T.; Rodriguez, J. I. 3- Dicarbonyl compounds in stereoselective domino and multicomponent reactions. *Tetrahedron: Asymmetry* **2010**, 21 (9–10), 1085–1109. (c) Willy, B.; Mueller, T. J. J. Multicomponent heterocycle syntheses via catalytic generation of alkynes. *Curr. Org. Chem.* **2009**, 13 (18), 1777–1790. (d) Isambert, N.; Lavila, R. Heterocycles as key substrates in multicomponent reactions: The fast lane towards molecular complexity. *Chem.—Eur. J.* **2008**, 14 (28), 8444–8454. (e) Cioc, R. C.; Ruijter, E.; Orru, R. V. A. Multicomponent reactions: Advanced tools for sustainable organic synthesis. *Green Chem.* **2014**, 16 (6), 2958–2975. (3) (a) Ruijter, E.; Scheffelaar, R.; Orru, R. V. A. Multicomponent reaction design in the quest for molecular complexity and diversity. *Angew. Chem., Int. Ed.* **2011**, 50, 6234–6246. (b) Bhunia, A.; Biju, A. T. Employing arynes in transition-metal- free, N- heterocycle-initiated

multicomponent reactions. *Synlett* **2014**, 25 (5), 608–614. (c) See also the *Chemical Society Reviews* issue on the Rapid formation of molecular complexity in organic synthesis: *Chem. Soc. Rev.* **2009**, 38, 2969–3276.

(4) See some examples: (a) Bocokić, V.; Kalkan, A.; Lutz, M.; Spek, A. L.; Gryko, D. T.; Reek, J. N. H. Capsule-controlled selectivity of a rhodium hydroformylation catalyst. *Nat. Commun.* **2013**, 4, 2670. (b) Xu, X.; Zhou, J.; Yang, L.; Hu, W. Selectivity control in enantioselective four-component reactions of aryl diazoacetates with alcohols, aldehydes and amines: An efficient approach to synthesizing chiral β -amino- α -hydroxyesters. *Chem. Commun.* **2008**, 6564–6566. (c) Shawali, A. S. Chemoselectivity in 1,3-Dipolar Cycloaddition Reactions of Nitrilimines with Multifunctionalized Dipolarophiles. *Curr. Org. Chem.* **2014**, 18, 598–614. (d) Ji, J.; Zhang, X.; Zhu, Y.; Qian, Y.; Zhou, J.; Yang, L.; Hu, W. Diastereoselectivity switch in cooperatively catalyzed three-component reactions of an aryldiazoacetate, an alcohol, and a β,γ -unsaturated α -keto ester. *J. Org. Chem.* **2011**, 76, 5821–5824. (e) Shawali, A. S.; Abdelhamid, A. O. Synthesis of spiro-heterocycles via 1,3-dipolar cycloadditions of nitrilimines to exoheterocyclic enones. site-, regio- and stereo-selectivities overview. *Curr. Org. Chem.* **2012**, 16, 2673–2689.

(5) See some recent examples: (a) Gao, H.; Sun, J.; Yan, C.-G. Selective synthesis of functionalized spiro[indoline-3,2'-pyridines] and spiro[indoline-3,4'-pyridines] by lewis acid catalyzed reactions of acetylenedicarboxylate, arylamines, and isatins. *J. Org. Chem.* **2014**, 79, 4131–4136. (b) Zhou, F.; Liu, X.; Zhang, N.; Liang, Y.; Zhang, R.; Xin, X.; Dong, D. Copper-catalyzed three-component reaction: Solvent-controlled regioselective synthesis of 4-amino- and 6-amino-2-iminopyridines. *Org. Lett.* **2013**, 15, 5786–5789. (c) Jaegli, S.; Dufour, J.; Wei, H.-L.; Piou, T.; Duan, X.-H.; Vors, J.-P.; Neuville, L.; Zhu, J. Palladium-catalyzed carbo-heterofunctionalization of alkenes for the synthesis of oxindoles and spirooxindoles. *Org. Lett.* **2010**, 12, 4498–4501. (d) Li, J.; Wang, N.; Li, C.; Jia, X. Multicomponent reaction to construct spirocyclic oxindoles with a Michael (triple Michael)/cyclization cascade sequence as the key step. *Chem.—Eur. J.* **2012**, 18, 9645–9650. (f) Chebanov, V. A.; Saraev, V. E.; Desenko, S. M.; Chernenko, V. N.; Knyazeva, I. V.; Groth, U.; Glasnov, T. N.; Kappe, C. O. Tuning of chemo- and regioselectivities in multicomponent condensations of 5-aminopyrazoles, dimedone, and aldehydes. *J. Org. Chem.* **2008**, 73, 5110–5118. (g) Elders, N.; Ruijter, E.; de Kanter, F. J. J.; Groen, M. B.; Orru, R. V. A. Selective formation of 2-imidazolines and 2-substituted oxazoles by using a three-component reaction. *Chem.—Eur. J.* **2008**, 14, 4961–4973. (h) Karapetyan, G.; Dang, T. T.; Sher, M.; Ghochikyan, T. V.; Saghyan, A.; Langer, P. Diversity-oriented synthesis of functionalized phenols by regioselective [3 + 3] cyclocondensations of 1,3-bis(silyloxy)-1,3-butadienes with 3-alkoxy-2-en-1-ones and related substrates. *Curr. Org. Chem.* **2012**, 16, 557–565. (i) He, H.; Sharif, M.; Neumann, H.; Beller, M.; Wu, X.-F. A convenient palladium-catalyzed carbonylative synthesis of 4(3H)-quinazolinones from 2-bromoformanilides and organo nitros with $\text{Mo}(\text{CO})_6$ as a multiple promoter. *Green Chem.* **2014**, 16, 3763–3767. (j) Chen, J.; Neumann, H.; Beller, M.; Wu, X.-F. Palladium-catalyzed synthesis of isoindoloquinazolinones via dicarbonylation of 1,2-dibromoarenes. *Org. Biomol. Chem.* **2014**, 12, 5835–5838. (k) Chen, J.; Natta, K.; Spannenberg, A.; Neumann, H.; Beller, M.; Wu, X.-F. Efficient palladium-catalyzed double carbonylation of *o*-dibromobenzenes: Synthesis of thalidomide. *Org. Biomol. Chem.* **2014**, 12, 5578–5581. (l) Li, H.-Q.; Li, W.-F.; Spannenberg, A.; Baumann, W.; Neumann, H.; Beller, M.; Wu, X.-F. A Novel Domino Synthesis of Quinazolinones by Palladium-Catalyzed Double Carbonylation. *Chem.—Eur. J.* **2014**, 20, 8541–8544. (m) Chen, J.; Natta, K.; Spannenberg, A.; Neumann, H.; Langer, P.; Beller, M.; Wu, X.-F. Base-Controlled Selectivity in the Synthesis of Linear and Angular Fused Quinazolinones by a Palladium-Catalyzed Carbonylation/Nucleophilic Aromatic Substitution Sequence. *Angew. Chem., Int. Ed.* **2014**, 53, 7579–7583. (n) He, L.; Li, H.-Q.; Neumann, H.; Beller, M.; Wu, X.-F. Highly Efficient Four-Component Synthesis of 4(3H)-Quinazolinones: Palladium-

- 454 Catalyzed Carbonylative Coupling Reactions. *Angew. Chem., Int. Ed.*
455 **2014**, 53, 1420–1424.
- 456 (6) (a) Gu, Y.; de Sousa, R.; Frapper, G.; Bachmann, C.; Barrault, J.;
457 Jérôme, F. Catalyst-free aqueous multicomponent domino reactions
458 from formaldehyde and 1,3-dicarbonyl derivatives. *Green Chem.* **2009**,
459 *11*, 1968–1972. (b) Gu, Y.; Barrault, J.; Jérôme, F. Trapping of active
460 methylene intermediates with alkenes, indoles or thiols: towards highly
461 selective multicomponent reactions. *Adv. Synth. Catal.* **2009**, 351,
462 3269–3278. (c) Tan, J.-N.; Li, H.; Gu, Y. Water mediated trapping of
463 active methylene intermediates generated by IBX-induced oxidation of
464 Baylis–Hillman adducts with nucleophiles. *Green Chem.* **2010**, 12,
465 1772–1782. (d) Li, M.; Chen, C.; He, F.; Gu, Y. Multicomponent
466 reactions of 1,3-cyclohexanediones and formaldehyde in glycerol:
467 stabilization of paraformaldehyde in glycerol resulted from using
468 dimedone as substrate. *Adv. Synth. Catal.* **2010**, 352, 519–530.
469 (e) Tan, J.-N.; Li, M.; Gu, Y. Multicomponent reactions of 1,3-
470 disubstituted 5-pyrazolones and formaldehyde in environmentally
471 benign solvent systems and their variations with more fundamental
472 substrates. *Green Chem.* **2010**, 12, 908–914.
- 473 (7) (a) Kumar, A.; Kumar, M.; Gupta, M. K. An efficient
474 organocatalyzed multicomponent synthesis of diarylmethanes via
475 Mannich type Friedel–Crafts reaction. *Tetrahedron Lett.* **2009**, 50,
476 7024–7027. (b) Kumar, A.; Sharma, S.; Maurya, R. A. A novel multi-
477 component reaction of indole, formaldehyde, and tertiary aromatic
478 amines. *Tetrahedron Lett.* **2009**, 50, 5937–5940. (c) Ferreira, S. B.;
479 Gonzaga, D. T. G.; de Carvalho da Silva, F.; de Lima Araújo, K. G.;
480 Ferreira, V. F. Synthesis of new *o*-quinone methides from β -lapachone
481 analogues. *Synlett* **2011**, 1623–1625. (d) Massa, A. Recent advances in
482 the chemistry of active methylene compounds. *Curr. Org. Chem.* **2012**,
483 *16*, 2159–2159. (e) Di Mola, A.; Palombi, L.; Massa, A. Active
484 methylene compounds in the synthesis of 3-substituted isobenzofur-
485 anones, isoindolinones and related compounds. *Curr. Org. Chem.*
486 **2012**, 16, 2302–2320.
- 487 (8) Frapper, G.; Bachmann, C.; Gu, Y.; de Sousa, R. C.; Jérôme, F.
488 Mechanisms of the Knoevenagel hetero Diels–Alder sequence in
489 multicomponent reactions to dihydropyrans: experimental and
490 theoretical investigations into the role of water. *Phys. Chem. Chem.*
491 *Phys.* **2011**, 13, 628–636.
- 492 (9) (a) Safi, M.; Sinou, D. Palladium(0)-catalyzed substitution of
493 allylic substrates in a two-phase aqueous-organic medium. *Tetrahedron*
494 *Lett.* **1991**, 32, 2025–2028. (b) Maity, A. K.; Chatterjee, P. N.; Roy, S.
495 Multimetallic Ir-Sn₃-catalyzed substitution reaction of π -activated
496 alcohols with carbon and heteroatom nucleophiles. *Tetrahedron*
497 **2013**, 69, 924–956. (c) Wahl, B.; Giboulot, S.; Mortreux, A.;
498 Castanet, Y.; Sauthier, M.; Liron, F.; Poli, G. Straightforward synthesis
499 of allylated keto esters: The Palladium-catalysed haloketone
500 alkoxycarbonylation/allylation domino reaction. *Adv. Synth. Catal.*
501 **2012**, 354, 1077–1083. (d) Rueping, M.; Nachtsheim, B. J.; Kuenkel,
502 A. Efficient metal-catalyzed direct benzylation and allylic alkylation of
503 2,4-pentanediones. *Org. Lett.* **2007**, 9, 825–828.
- 504 (10) (a) Chuang, C.-P.; Chen, K.-P.; Hsu, Y.-L.; Tsai, A.-I.; Liu, S.-T.
505 α -Nitro carbonyl compounds in the synthesis of 2,3-dihydrofurans.
506 *Tetrahedron* **2008**, 64, 7511–7516. (b) Arai, S.; Nakayama, K.; Suzuki,
507 Y.; Hatano, K.-I.; Shioiri, T. Stereoselective synthesis of dihydrofurans
508 under phase-transfer catalyzed conditions. *Tetrahedron Lett.* **1998**, 39,
509 9739–9742. (c) Chuang, C. P.; Tsai, A.-I. Pyridinium ylides in the
510 synthesis of 2,3-dihydrofurans. *Synthesis* **2006**, 675–679. (d) Induma-
511 thi, S.; Perumal, S.; Anbananthan, N. A facile eco-friendly three-
512 component protocol for the regio- and stereoselective synthesis of
513 functionalized *trans*-dihydrofuro[3,2-*c*]-quinolin-4(2*H*)-ones. *Green*
514 *Chem.* **2012**, 14, 3361–3367. (e) Gunasekaran, P.; Balamurugan, K.;
515 Sivakumar, S.; Perumal, S.; Menéndez, J. C.; Almansour, A. I. Domino
516 reactions in water: Diastereoselective synthesis of densely function-
517 alized indolyl dihydrofuran derivatives. *Green Chem.* **2012**, 14, 750–
518 757. (f) Chuang, C.-P.; Chen, K.-P. *N*-Phenacylpyridinium bromides
519 in the one-pot synthesis of 2,3-dihydrofurans. *Tetrahedron* **2012**, 68,
520 1401–1406. (g) Wang, Q.-F.; Hou, H.; Hui, L.; Yan, C.-G.
521 Diastereoselective synthesis of *trans*-2,3-dihydrofurans with pyridinium
522 ylide assisted tandem reaction. *J. Org. Chem.* **2009**, 74, 7403–7406.
- (h) Yang, Z.; Fan, M.; Mu, R.; Liu, W.; Liang, Y. A facile synthesis of
523 highly functionalized dihydrofurans based on 1,4-diazabicyclo[2.2.2]-
524 octane (DABCO) catalyzed reaction of halides with enones. *Tetrahedron*
525 **2005**, 61, 9140–9146. (i) Vitale, P.; Scilimati, A. Five-
526 membered ring heterocycles by reacting enolates with dipoles. *Curr.*
527 *Org. Chem.* **2013**, 17, 1986–2000.
- (11) Literature survey stated that while sulfonium ylide and
529 imidazolium ylide enabled the use of aliphatic aldehyde, the reactions
530 with pyridium ylide are still not amenable to aliphatic aldehyde:
531 (a) Jiang, Y.; Ma, D. Synthesis of enantiopure substituted
532 dihydrofurans via the reaction of (*S*)-glyceraldehyde acetone- or
533 Garner aldehyde acetone-derivative enones with sulfonium ylides.
534 *Tetrahedron: Asymmetry* **2002**, 13, 1033–1038. (b) Kumar, A.;
535 Srivastava, S.; Gupta, G. Cascade [4 + 1] annulation *via* more
536 environmentally friendly nitrogen ylides in water: synthesis of bicyclic
537 and tricyclic fused dihydrofurans. *Green Chem.* **2012**, 14, 3269–3272.
538
- (12) (a) Chen, L.; Luo, S.; Li, J.; Li, X.; Cheng, J.-P. Organocatalytic
539 kinetic resolution *via* intramolecular aldol reactions: Enantioselective
540 synthesis of both enantiomers of chiral cyclohexenones. *Org. Biomol.*
541 *Chem.* **2010**, 8, 2627–2632. (b) Jin, T.; Yamamoto, Y. Gold-catalyzed
542 intramolecular carbocyclization of alkynyl ketones leading to highly
543 substituted cyclic enones. *Org. Lett.* **2007**, 9, 5259–5262.
- (13) Biswas, S.; Maiti, S.; Jana, U. An efficient iron-catalyzed carbon–
545 carbon single-bond cleavage via Retro-Claisen Condensation: a mild
546 and convenient approach to synthesize a variety of esters or ketones.
547 *Eur. J. Org. Chem.* **2010**, 2861–2866.
- (14) Synthesis of aromatic ether from non-aromatic precursor has
549 just been established by means of oxidative condensation of alcohol
550 and 2-cyclohexenone: Simon, M.-O.; Girard, S. A.; Li, C.-J. Catalytic
551 aerobic synthesis of aromatic ethers from non-aromatic precursors.
552 *Angew. Chem., Int. Ed.* **2012**, 51, 7537–7540.