

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51554048>

Discovery of a Cytokinin Deaminase

ARTICLE in ACS CHEMICAL BIOLOGY · AUGUST 2011

Impact Factor: 5.33 · DOI: 10.1021/cb200198c · Source: PubMed

CITATIONS

8

READS

36

4 AUTHORS:



Alissa M Goble

Texas A&M University

3 PUBLICATIONS 19 CITATIONS

SEE PROFILE



Hao Fan

Agency for Science, Technology and Research...

32 PUBLICATIONS 989 CITATIONS

SEE PROFILE



Andrej Sali

University of California, San Francisco

358 PUBLICATIONS 40,377 CITATIONS

SEE PROFILE



Frank Raushel

Texas A&M University

312 PUBLICATIONS 9,454 CITATIONS

SEE PROFILE

Discovery of a Cytokinin Deaminase


Alissa M. Goble,[†] Hao Fan,[‡] Andrej Sali,^{*,‡} and Frank M. Raushel^{*,†}

[†]Department of Chemistry, Texas A&M University, P.O. Box 30012, College Station, Texas 77843-3012, United States

[‡]Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, 1700 Fourth Street, San Francisco, California 94158, United States

S Supporting Information

ABSTRACT: An enzyme of unknown function within the amidohydrolase superfamily was discovered to catalyze the hydrolysis of *N*-6-substituted adenine derivatives, several of which are cytokinins. Cytokinins are a common type of plant hormone and *N*-6-substituted adenines are also found as modifications to tRNA. Patl2390, from *Pseudoalteromonas atlantica* T6c, was shown to hydrolytically deaminate *N*-6-isopentenyladenine to hypoxanthine and isopentenylamine with a k_{cat}/K_m of $1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$. Additional substrates include *N*-6-benzyl adenine, *cis*- and *trans*-zeatin, kinetin, *O*-6-methylguanine, *N*-6-butyladenine, *N*-6-methyladenine, *N,N*-dimethyladenine, 6-methoxypurine, 6-chloropurine, and 6-thiomethylpurine. This enzyme does not catalyze the deamination of adenine or adenosine. A comparative model of Patl2390 was computed using the three-dimensional crystal structure of Pa0148 (PDB code 3PAO) as a structural template, and docking was used to refine the model to accommodate experimentally identified substrates. This is the first identification of an enzyme that will hydrolyze an *N*-6-substituted side chain larger than methylamine from adenine.



With the rapidly increasing volume of protein sequence data available, a need has been created for a comprehensive strategy for analyzing and annotating newly sequenced genomes. Clusters of orthologous groups (COG) were created to aid in the assignment of function for related groups of proteins. Within the amidohydrolase superfamily (AHS) there are 24 clusters of orthologous groups as defined by NCBI. Members of the AHS may have either a mononuclear or binuclear metal center embedded within a $(\beta/\alpha)_8$ -barrel structural fold with metal binding residues found at the C-terminal ends of β -strands 1, 4, 5, 6, and 8.¹ The metal center is responsible for activating a hydrolytic water molecule for nucleophilic attack on amino acids, sugars, nucleic acids, and organophosphate esters. Three of these clusters of enzymes are known to catalyze aromatic deamination reactions: cog1001, cog0402, and cog1816. One of these clusters, cog1816, contains nearly 500 bacterial proteins that are currently annotated as adenosine deaminases.² The prototypical adenine deaminase (ADE) and *N*-6-methyladenine deaminase (6-MAD) are members of cog1001.³ Enzymes that deaminate guanine, cytosine, *S*-adenosylhomocysteine, thiomethyl adenosine, *N*-formimino-L-glutamate, and 8-oxoguanine are found in cog0402.⁴

A sequence similarity network for cog1816 at an E-value cutoff of 10^{-70} is presented in Figure 1.^{5,6} Enzymes with experimentally confirmed adenosine deaminase activity belong to Group 5 of cog1816, and enzymes recently identified as adenine deaminases belong to Group 3.⁷ The authentic adenine and adenosine deaminases of cog1816, in addition to the other known deaminases from cog0402, share an invariant HxxE motif at the C-terminal end of β -strand 5. Eleven proteins at the periphery of Group 3 have an HxD motif instead of HxxE at this position in the protein sequence. These proteins (Patl2390 from *Pseudoalteromonas atlantica*,





Figure 1. Sequence similarity network created using Cytoscape⁶ of cog1816 from the amidohydrolase superfamily. Each node in the network represents a single sequence, and each edge (depicted as lines) represents the pairwise connection between two sequences at a BLAST E-value of better than 1×10^{-70} . Lengths of edges are not significant, except for tightly clustered groups, which are more closely related than sequences with only a few connections. Group 5 contains the *E. coli* adenosine deaminase. The yellow dots in Group 3 represent proteins that will deaminate adenine. The proteins represented by the pink dots at the periphery of Group 3 are characterized in this paper.

gi|109898705; Leum0809 from *Leuconostoc mesenteroides*, gi|116617920; Lsa0086 from *Lactobacillus sakei*, gi|81427699; Cja0578 from *Cellvibrio japonicus*, gi|192359911; MADE1015570 from *Alteromonas macleodii*, gi|332142506; Jden_1580 from *Jonesia denitrificans* DSM 20603, gi|256832804; Glaag_1976 from *Glaciecola agarilytica* 4H-3-7+YE-5, gi|332173667; ambt_10110 from *Alteromonas* sp. SN2, gi|332993493; Sros_2140 from *Streptosporangium roseum* DSM 43021, gi|271963671; Sked_12620 from *Sanguibacter keddiei* DSM 10542, gi|269794583; and LBU_0899

Received: June 20, 2011

Accepted: August 8, 2011

Scheme 1



from *Lactobacillus delbrueckii* subsp. *Bulgaricus* 2038, gi|325125754) are highlighted in pink in Figure 1. These 11 proteins originate from organisms with an authentic adenine deaminase from either cog1001 or Group 3 of cog1816. On the basis of these observations, we predicted that the 11 proteins at the periphery of Group 3 from cog1816 will not deaminate adenine but will deaminate an adenine-like substrate.

Substrate Discovery. We established the substrate profile for one of the proteins at the periphery of Group 3 from cog1816; Patl2390 from *P. atlantica* T6c, an agarolytic bacterium.⁸ The gene for Patl2390 was cloned and expressed, and the resultant protein purified to homogeneity. The purified protein contained 0.6 equiv of Zn^{2+} per monomer. The substrate profile for Patl2390 was determined by monitoring the changes in absorbance after the addition of enzyme to a small library of modified purines, including adenosine, adenine, *N*-6-methyladenine, and *N*-6-isopentenyladenine. On the basis of the results of the initial screen, a larger collection of 6-substituted purines was assayed, including 6-chloropurine, 6-thiomethylpurine, *N,N*-dimethyladenine, *trans*-zeatin, kinetin, benzyladenine, 6-methoxypurine, *N*-butyladenine, *O*-6-methylguanine, and *cis*-zeatin. The structures of the most active compounds are presented in Scheme 1. The enzyme exhibited a complete lack of activity (<0.005%) for the deamination of adenosine or adenine. Enzymatic conversion of substrate to product was verified by UV–vis spectroscopy and mass spectrometry. A coupled assay with alcohol dehydrogenase was used to detect the formation of methanol from 6-methoxypurine.⁹ The kinetic constants for the reactions of compounds 1–12 catalyzed by Patl2390 are presented in Table 1. The best substrate for Patl2390 is *N*-6-isopentenyladenine with a $k_{\text{cat}}/K_{\text{m}}$ of $1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$.

Structure of Patl2390. The determination of the three-dimensional structure of Patl2390 by X-ray crystallography has thus far failed. A comparative model was therefore constructed for Patl2390 using the most closely related structure in the PDB database as the template. This protein, Pa0148 from *Pseudomonas aeruginosa* PAO1 (PDB id 3OU8) is an adenine deaminase from Group 3 of cog1816 and shares a 45% sequence identity to Patl2390. A protein sequence alignment of Patl2390 and Pa0148, along with four proteins predicted to

Table 1. Kinetic Constants for Patl2390

substrate	k_{cat} (s^{-1})	K_{m} (μM)	$k_{\text{cat}}/K_{\text{m}}$ ($\text{M}^{-1} \text{s}^{-1}$)
1	7.3 ± 0.1	0.6 ± 0.1	$1.2 (0.2) \times 10^7$
2	8.5 ± 0.9	86 ± 7	$9.9 (0.3) \times 10^4$
3	10.1 ± 0.5	9.2 ± 0.9	$1.1 (0.1) \times 10^6$
4	10.2 ± 0.6	9 ± 2	$1.1 (0.3) \times 10^6$
5	5.6 ± 0.5	9 ± 2	$6.2 (0.2) \times 10^5$
6	6.5 ± 0.4	50 ± 6	$1.3 (0.2) \times 10^5$
7	13.1 ± 0.6	10 ± 1	$1.3 (0.1) \times 10^6$
8	11.4 ± 0.6	1.5 ± 0.3	$8 (1) \times 10^6$
9	11.6 ± 0.6	7 ± 1	$1.6 (0.3) \times 10^6$
10	13 ± 1	35 ± 5	$3.6 (0.5) \times 10^5$
11			$3.2 (0.3) \times 10^4$
12	13 ± 1	38 ± 5	$3.3 (0.4) \times 10^5$

have the same substrate profile as Patl2390, is presented in Supplementary Figure S1. The crystal structure of Pa0148 with adenine bound (PDB id 3PAO) was used as the template for comparative modeling by satisfaction of spatial restraints, as implemented in Modeler-9v8.¹⁰ In the resulting model of Patl2390, most residues in the proximity of adenine are conserved relative to 3PAO, except for Asp203 and Phe257. The backbone and side chain conformations of segments near the two variable residues (Cys202 to Gln206, Pro252 to Ser261) were refined in the presence of adenine by Modeler for loops and by PLOP for side chains.¹¹ The high-energy intermediate (HEI)^{12,13} library that contains ~22,500 different intermediate forms of 4207 KEGG^{14,15} molecules was docked to this refined model of Patl2390 by DOCK 3.5.54.^{16,17} Six known substrates that were included in the docking library (*N*-6-benzyladenine, *cis*-zeatin, kinetin, *N*-6-isopentenyladenine, *trans*-zeatin, *N*-6-methyladenine) were ranked 56, 60, 64, 88, 147, and 328, respectively. The docking poses of these six substrates are similar to the binding mode of adenine in 3PAO. For *N*-6-benzyladenine, the activated hydroxide attacks C-6 on the *re* face of the purine ring, forming a tetrahedral intermediate with *R*-stereochemistry. Asp203 is hydrogen bonded to protonated N-1 of the purine ring.





Figure 2. Structural model of the active site of Patl2390 with *N*-6-benzyladenine.

In the comparative model, residues that coordinate the mononuclear metal center (His21, His23, His201, and Asp282) and residues that apparently are involved in substrate recognition (Tyr68, Asp105, Ser174, and Asp283) superimpose with the aligned residues from the structural template. A deletion of two Patl2390 residues in the sequence ²⁵²PISNR--FV²⁵⁸ compared with the structural template changes the second half of a two-turn helix in Pa0148 into a more flexible loop conformation containing residue Phe257. This deletion opens up a hydrophobic pocket that enables the binding of substrates with branched alkyl chains as well as aromatic substituents into the active site. Pi-stacking interactions are possible between the substituents attached to *N*-6 of the substrate with the side chains of Phe64 and Phe257.

Another region that is different in the structural model of Patl2390, compared to the template, is in the active site following β -strand 5. Instead of the HxxE motif found in Pa0148, Patl2390 has a two-residue insertion with the sequence HxDxxQ at the C-terminal of β -strand 5. The invariant histidine in Patl2390 (His227) overlaps in the structural model with the corresponding histidine in Pa0148 (His196). The highly conserved glutamate in the HxxE motif is functionally quite critical, since this residue has been shown to be essential for the delivery of a proton to *N*-1 during the deamination reaction in enzymes from the AHS that react with adenosine, adenine, and guanine.^{1,18} The glutamine in the HxDxxQ motif in Patl2390 would be unable to function as a proton donor. The active site of the comparative model of Patl2390 with *N*-6-benzyladenine bound in the active site is presented in Figure 2.

Since Patl2390 does not have an appropriately positioned glutamate after β -strand 5, residues that were likely to fill this function were identified from the initial comparative model. Asp203 from the HxDxxQ motif was mutated to an asparagine (D203N), and Gln206 was mutated to glutamate (Q206E). In addition to these two residues, a conserved glutamate after β -strand 4 that is in the vicinity of the active site was mutated to glutamine and alanine (E176Q and E176A). E176Q was insoluble, but E176A could be purified to homogeneity. The kinetic constants for the E176A mutant using *N*-6-benzyladenine as a substrate were as follows: $k_{\text{cat}} = 0.013 \text{ s}^{-1}$, $K_{\text{m}} = 4.9 \text{ }\mu\text{M}$, and $k_{\text{cat}}/K_{\text{m}} = 2600 \text{ M}^{-1} \text{ s}^{-1}$. The value of $k_{\text{cat}}/K_{\text{m}}$ has been reduced by nearly 4 orders of magnitude. The Q206E mutant exhibited a substantial reduction in the values of the kinetic constants relative to the wild type enzyme. The values of k_{cat} , K_{m} , and $k_{\text{cat}}/K_{\text{m}}$ for

Scheme 2



this mutant are 0.11 s^{-1} , $10 \text{ }\mu\text{M}$, and $1.1 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$, respectively. We were unable to detect any activity ($<0.001\%$ of wild-type enzyme) when Asp203 was mutated to an asparagine. On the basis of these results and the docking of known substrates into the active site of Patl2390, we conclude that Asp203 functions to deliver a proton to *N*-1 of the substrate during catalysis.

The substrates identified for Patl2390 are known as cytokinins. These compounds promote cell growth and division in plants.¹⁹ They are produced by nearly all plants and have been isolated from red algae.²⁰ Similar compounds are also found as modified adenosines in serine and phenylalanine tRNA.^{21–23} Naturally occurring cytokinins include *N*-6-benzyladenine, kinetin, *cis*-zeatin, *trans*-zeatin, and *N*-6-isopentenyladenine. *N*-6-Isopentenyladenine, as well as *cis*- and *trans*-zeatin, are believed to originate from dimethylallyl pyrophosphate transfer to AMP.²¹ The effects of cytokinins on plants have been well-studied, but the metabolism of these compounds has received less attention.²⁴ A cytokinin oxidase from *Zea mays* catalyzes the oxidative cleavage of *N*-6-isopentenyl or zeatin side chains of both the adenine and nucleoside forms.²⁵

The bacterium *P. atlantica* T6c was first isolated in association with marine red algae, which are known to contain agar in their cell walls and to produce cytokinins.²⁰ *P. atlantica* T6c is able to degrade the cell walls of red algae through production of an extracellular agarase, which enables the bacteria to acquire nutrients from the algae.²⁶ The enzyme identified here may function in salvaging of nutrients from algae through the conversion of cytokinins into hypoxanthine, which can then be incorporated into the purine salvage pathway. The reaction catalyzed by Patl2390 with isopentenyl adenine is presented in Scheme 2. The ability to remove a methoxy group from *O*-6-methylguanine may also indicate a role of this enzyme in DNA repair.²⁷

Strategy for Functional Annotation. Current estimates put the percent of unknown, uncertain, or incorrect annotations of bacterial genes at more than 30%.²⁸ For this investigation the NCBI database was used to construct a sequence similarity diagram for cog1816 from the amidohydrolase superfamily, which identified 11 proteins as having a high sequence identity to authentic adenine deaminases of Group 3 but missing some key catalytic residues. The genes for these proteins are located in the vicinity of genes for other enzymes involved in purine metabolism, including the experimentally verified adenine deaminases. Using adenine as a structural scaffold, a small library of compounds was assembled to ultimately identify the substrate profile for Patl2390. We demonstrated that Patl2390 is capable of accepting cytokinins and several modified purines as substrates. No catalytic activity toward adenine or adenosine was observed. The construction of a comparative structural model for Patl2390 allowed for the identification of the structural rationale for the preferential hydrolysis of large substituents attached to C-6 of the purine base.

METHODS

Materials. All chemicals were purchased from Sigma unless otherwise stated. 6-Methoxypurine was obtained from Tokyo Chemical Industry Co. *N*-6-Methyladenine was procured from Spectrum. *N*-6-Butyladenine was bought from Ryan Scientific. *cis*-Zeatin was ordered from MP Biomedicals, and zeatin riboside was acquired from Carbo-synth. *O*-6-Methylguanine was obtained from LT Pharmatech.

Cloning and Purification of Patl2390 from *Pseudoalteromonas atlantica* T6c. Patl2390 was cloned from *P. atlantica* T6c genomic DNA (ATCC) using restriction sites for *Bam*HI and *Eco*RI, digested, and ligated into a pET-30a(+) vector. The cloned gene fragment was sequenced to verify fidelity of the PCR amplification. Patl2390 in a pET-30a(+) vector was overexpressed in BL21(DE3) cells (Novagen) and purified by affinity chromatography using a HISTRap column (GE Healthcare). Additional information is available in Supporting Information.

Activity Screens. Patl2390 (10 nM) was incubated for 16 h with adenine, 2,6-diaminopurine, 6-mercaptopurine, 6-methylthiopurine, 6-chloropurine, 6-methylpurine, *N*-6-isopentyl adenine, isoguanine, *N*-6-methyladenine, zeatin, zeatin riboside, 4-aminopyrimidine, 4,6-diaminopyrimidine, 2,4-diaminopyrimidine, 7-methyladenine, toxypyrimidine, cytosine, 5-hydroxymethylcytosine, 2-chloroadenine, and 2-dimethylaminoadenine. Substrate concentration was 80 μ M for all screens. Activity was monitored by changes in absorbance between 240 and 300 nm on a SpectraMax384Plus spectrophotometer (Molecular Devices).

Measurement of Enzymatic Activity. Assays were conducted with 3–200 μ M substrate. Dechlorination of 6-chloropurine was monitored by an increase in absorbance at 250 nm. Formation of hypoxanthine from *N*-6-isopentenyladenine, *N*-6-methyladenine, *trans*-zeatin, benzyladenine, 6-methoxypurine, and *N*-6-butyladenine was monitored at 270 nm. Decreases in absorbance at 275 nm were used to monitor formation of hypoxanthine from *cis*-zeatin and *N*, *N*-dimethyladenine. Conversion of 6-methylthiopurine and kinetin were monitored by decreases in absorbance at 290 and 274 nm, respectively. Formation of guanine from *O*-6-methylguanine was monitored at 255 nm. Difference extinction coefficients were calculated by subtracting the extinction coefficient of the product from the extinction coefficient for the substrate for 6-methylthiopurine ($\Delta\epsilon_{290} = 16500 \text{ M}^{-1} \text{ cm}^{-1}$), *N*-6-methyladenine ($\Delta\epsilon_{270} = 15000 \text{ M}^{-1} \text{ cm}^{-1}$), *N*-6-isopentenyladenine ($\Delta\epsilon_{270} = 11700 \text{ M}^{-1} \text{ cm}^{-1}$), *N,N*-dimethyladenine ($\Delta\epsilon_{275} = 18300 \text{ M}^{-1} \text{ cm}^{-1}$), *trans*-zeatin ($\Delta\epsilon_{270} = 17700 \text{ M}^{-1} \text{ cm}^{-1}$), kinetin ($\Delta\epsilon_{274} = 8900 \text{ M}^{-1} \text{ cm}^{-1}$), *N*-6-benzyladenine ($\Delta\epsilon_{270} = 12300 \text{ M}^{-1} \text{ cm}^{-1}$), *N*-6-butyladenine ($\Delta\epsilon_{270} = 12600 \text{ M}^{-1} \text{ cm}^{-1}$), and *cis*-zeatin ($\Delta\epsilon_{275} = 12000 \text{ M}^{-1} \text{ cm}^{-1}$). For 6-chloropurine ($\Delta\epsilon_{250} = 9600 \text{ M}^{-1} \text{ cm}^{-1}$), *O*-6-methylguanine ($\Delta\epsilon_{255} = 3200 \text{ M}^{-1} \text{ cm}^{-1}$), and 6-methoxypurine ($\Delta\epsilon_{270} = 1937 \text{ M}^{-1} \text{ cm}^{-1}$), difference extinction coefficients were calculated by subtracting the extinction coefficient for the substrate from the extinction coefficient for the product. Production of the corresponding alkylamine side chains and hypoxanthine were confirmed by mass spectrometry for the reactions of *N*-6-benzyladenine, *N*-6-isopentenyladenine, *N,N*-dimethyladenine, *N*-6-methyladenine, *N*-6-butyladenine, *cis*- and *trans*-zeatin, 6-chloropurine, 6-methylthiopurine, and kinetin. Production of methanol from 6-methoxypurine by 420 nM Patl2390 was detected using alcohol dehydrogenase (4 mg mL⁻¹) and monitoring the conversion of NAD⁺ to NADH at 340 nm.⁹ Removal of methanol from *O*-6-methylguanine was confirmed using a coupled assay with guanine deaminase to produce the demethylated and deaminated product, xanthine.

Mutation of E176Q, E176A, D203N, and Q206E. Single-site mutations were constructed using the standard QuickChange PCR protocol according to the manufacturer's instructions. The mutants were expressed and purified using the protocol used for wild-type Patl2390.

Metal Analysis. Metal content of the proteins was determined by ICP-MS.²⁹ Protein samples for ICP-MS were digested with HNO₃ by refluxing for ~45 min to prevent protein precipitation during the measurement. Protein concentration was adjusted to ~1.0 μ M with 1% (v/v) HNO₃.

Data Analysis. Sequence alignments were created using ClustalW at <http://www.compbio.dundee.ac.uk/JalviewWS/services/ClustalWS>. Steady-state kinetic data were analyzed using Softmax Pro version 5.4. Kinetic parameters were determined by fitting the data to eq 1 using the nonlinear least-squares fitting program in SigmaPlot 9.0, where *A* is the substrate concentration, *K*_m is the Michaelis constant, *v* is the velocity of the reaction, and *k*_{cat} is the turnover number.

$$v/E_t = k_{\text{cat}}A/(K_m + A) \quad (1)$$

Model Building and Computational Docking. The sequence alignment between Patl2390 and Pa0148 was computed by MUSCLE (Multiple Sequence Comparison by Log-Expectation).³⁰ The original alignment was manually adjusted, so that the active site residue Asp203 in Patl2390 was aligned to Glu199 in Pa0148. The crystal structure of Pa0148 with adenine bound (PDB id 3PAO) was used as the template. A total of 500 comparative models were generated with the standard "automodel" class in Modeler-9v8.¹⁰ The model with the best DOPE³¹ score was selected for refinement of backbone and side chain conformations of residues Cys202 to Gln206 and Pro252 to Ser261, using the "loopmodel" class in Modeler. Side chains of residues in these two loops were subsequently refined using the "side chain prediction" protocol in PLOP,³² in the presence of adenine. This refinement resulted in two representative models of Patl2390, in which (1) residues Cys202 to Gln206 were refined with respect to the original model and (2) residues Cys202 to Gln206 and Pro252 to Ser261 were refined with respect to the original model. The high-energy intermediate (HEI)^{12,13} library of KEGG^{14,15} molecules was docked to both refined models of Patl2390 by DOCK 3.5.54.^{16,17} The second model yielding a higher enrichment for six known substrates (*N*-6-benzyladenine, *cis*-zeatin, kinetin, *N*-6-isopentenyladenine, *trans*-zeatin, *N*-6-methyladenine) was selected to represent the binding mode of *N*-6-benzyladenine in Patl2390 (Figure 2).

ASSOCIATED CONTENT

S Supporting Information. Protein sequence alignment of Patl2390 with five other proteins (Figure S1) and additional information on the cloning and purification of Patl2390. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: sali@salilab.org; raushel@tamu.edu.

ACKNOWLEDGMENT

This work was supported in part by the National Institutes of Health (GM 71790, GM 54762, and GM 93342) and the Robert A. Welch Foundation (A-840).

ABBREVIATIONS

ADE, adenine deaminase; AHS, amidohydrolase superfamily; ICP-MS, inductively coupled plasma mass spectrometry

REFERENCES

- (1) Seibert, C. M., and Raushel, F. M. (2005) Structural and catalytic diversity within the amidohydrolase superfamily. *Biochemistry* 44, 6383–6391.

- (2) Ho, M. C., Cassera, M. B., Madrid, D. C., Ting, L. M., Tyler, P. C., Kim, K., Almo, S. C., and Schramm, V. L. (2009) Structural and metabolic specificity of methylthioformycin for malarial adenosine deaminases. *Biochemistry* 48, 9618–9626.
- (3) Kamat, S. S., Fan, H., Sauder, J. M., Burley, S. K., Shoichet, B. K., Sali, A., and Raushel, F. M. (2011) Enzymatic deamination of the epigenetic base N-6-methyladenine. *J. Am. Chem. Soc.* 133, 2080–2083.
- (4) Hall, R. S., Agarwal, R., Hitchcock, D., Sauder, J. M., Burley, S. K., Swaminathan, S., and Raushel, F. M. (2010) Discovery and structure determination of the orphan enzyme isoxanthopterin deaminase. *Biochemistry* 49, 4374–4382.
- (5) Atkinson, H. J., Morris, J. H., Ferrin, T. E., and Babbitt, P. C. (2009) Using sequence similarity networks for visualization of relationships across diverse protein superfamilies. *PLoS One* 4, e4345.
- (6) Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 13, 2498–2504.
- (7) Goble, A. M., Zhang, Z., Sauder, J. M., Burley, S. K., Swaminathan, S., and Raushel, F. M. (2011) Pa0148 from *Pseudomonas aeruginosa* catalyzes the deamination of adenine. *Biochemistry* 50, 6589–6597.
- (8) Yaphe, W. (1957) The use of agarase from *Pseudomonas atlantica* in the identification of agar in marine algae (Rhodophyceae). *Can. J. Microbiol.* 3, 987–993.
- (9) Bergmeyer, H. U., Ed. (1974) *Methods of Enzymatic Analysis*, Academic Press, New York, NY.
- (10) Sali, A., and Blundell, T. L. (1993) Comparative protein modeling by satisfaction of spatial restraints. *J. Mol. Biol.* 234, 779–815.
- (11) Tubbs, J. L., Latypov, V., Kanugula, S., Butt, A., Melikishvili, M., Kraehenbuehl, R., Fleck, O., Marriott, A., Watson, A. J., Verbeek, B., McGown, G., Thorncroft, M., Santibanez-Koref, M. F., Millington, C., Arvai, A. S., Kroeger, M. D., Peterson, L. A., Williams, D. M., Fried, M. G., Margison, G. P., Pegg, A. E., and Tainer, J. A. (2009) Flipping of alkylated DNA damage bridges base and nucleotide excision repair. *Nature* 459, 808–813.
- (12) Hermann, J. C., Ghanem, E., Li, Y. C., Raushel, F. M., Irwin, J. J., and Shoichet, B. K. (2006) Predicting substrates by docking high-energy intermediates to enzyme structures. *J. Am. Chem. Soc.* 128, 15882–15891.
- (13) Hermann, J. C., Marti-Arbona, R., Fedorov, A. A., Fedorov, E., Almo, S. C., Shoichet, B. K., and Raushel, F. M. (2007) Structure-based activity prediction for an enzyme of unknown function. *Nature* 448, 775–U772.
- (14) Kanehisa, M., and Goto, S. (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.* 28, 27–30.
- (15) Kanehisa, M., Goto, S., Hattori, M., Aoki-Kinoshita, K. F., Itoh, M., Kawashima, S., Katayama, T., Araki, M., and Hirakawa, M. (2006) From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Res.* 34, D354–D357.
- (16) Lorber, D. M., and Shoichet, B. K. (2005) Hierarchical docking of databases of multiple ligand conformations. *Curr. Top. Med. Chem. Sci.* 739–749.
- (17) Meng, E. C., Shoichet, B. K., and Kuntz, I. D. (1992) Automated docking with grid-based energy evaluation. *J. Comput. Chem.* 13, 505–524.
- (18) Mohamedali, K. A., Kurz, L. C., and Rudolph, F. B. (1996) Site-directed mutagenesis of active site glutamate-217 in mouse adenosine deaminase. *Biochemistry* 35, 1672–1680.
- (19) Skoog, F., Hamzi, H. Q., Szwedkow, Am, Leonard, N. J., Carraway, K. L., Fujii, T., Helgeson, J. P., and Loeppky, R. N. (1967) Cytokinins - Structure/Activity Relationships. *Phytochemistry* 6, 1169.
- (20) Yokoya, N. S., Stirk, W. A., van Staden, J., Novak, O., Tureckova, V., Pencik, A., and Strnad, M. (2010) Endogenous cytokinins, auxins, and abscisic acid in red algae from Brazil. *J. Phycol.* 46, 1198–1205.
- (21) Kakimoto, T. (2003) Biosynthesis of cytokinins. *J. Plant Res.* 116, 233–239.
- (22) Uziel, M., and Gassen, H. G. (1969) Phenylalanine transfer ribonucleic acid from *Escherichia coli* B. Isolation and characterization of oligonucleotides from ribonuclease T-1 and ribonuclease A hydrolysates. *Biochemistry* 8, 1643–1655.
- (23) Nishimura, S., Yamada, Y., and Ishikura, H. (1969) The presence of 2-methylthio-N6-(delta-2-isopentenyl) adenosine in serine and phenylalanine transfer RNAs from *Escherichia coli*. *Biochim. Biophys. Acta* 179, 517–520.
- (24) Chen, C., Smith, O. C., and McChesney, J. (1975) Biosynthesis and cytokinin activity of 8-hydroxy and 2,8-dihydroxy derivatives of zeatin and N-6(increment-2-isopentenyl)adenine. *Biochemistry* 14, 3088–3093.
- (25) Whitty, C. D., and Hall, R. H. (1974) A cytokinin oxidase in *Zea mays*. *Can. J. Biochem.* 52, 789–799.
- (26) Holmstrom, C., and Kjelleberg, S. (1999) Marine Pseudoalteromonas species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiol. Ecol.* 30, 285–293.
- (27) Morita, R., Nakagawa, N., Kuramitsu, S., and Masui, R. (2008) An O6-methylguanine-DNA methyltransferase-like protein from *Thermus thermophilus* interacts with a nucleotide excision repair protein. *J. Biochem.* 144, 267–277.
- (28) Schnoes, A. M., Brown, S. D., Dodevski, I., and Babbitt, P. C. (2009) Annotation error in public databases: misannotation of molecular function in enzyme superfamilies. *PLoS Comput. Biol.* 5, e1000605.
- (29) Hall, R. S., Xiang, D. F., Xu, C., and Raushel, F. M. (2007) N-Acetyl-D-glucosamine-6-phosphate deacetylase: substrate activation via a single divalent metal ion. *Biochemistry* 46, 7942–7952.
- (30) Edgar, R. C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797.
- (31) Shen, M. Y., and Sali, A. (2006) Statistical potential for assessment and prediction of protein structures. *Protein Sci.* 15, 2507–2524.
- (32) Sherman, W., Day, T., Jacobson, M. P., Friesner, R. A., and Farid, R. (2006) Novel procedure for modeling ligand/receptor induced fit effects. *J. Med. Chem.* 49, 534–553.