

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/99049>

Please be advised that this information was generated on 2016-01-31 and may be subject to change.

leads to deactivation of the system. GPC analysis of the polymer indicated  $M_n$  and  $M_w$  values of 21 000 and 47 900 daltons (Da), respectively, and a DSC curve for the polymer exhibited no detectable exotherm prior to the onset of crystalline melting at 131.6 °C. In contrast, the reactions of  $(\text{Ph}_3\text{SiO})_3\text{VO}$  and  $(n\text{-PrO})_3\text{VO}$  under identical conditions gave no polyethylene; the only observable reaction (by  $^{51}\text{V}$  NMR) was metathesis of alkyl ligands on Al for alkoxide ligands on V.<sup>7d,8</sup> We have not extensively explored the use of other cocatalysts with **2**, but we have observed similar activities with  $\text{Et}_3\text{Al}$  and somewhat lower activities with  $(\text{Me}_3\text{SiCH}_2)_3\text{Al}$ .

A survey of the catalyst's reactivity indicates that other olefins can also be polymerized or copolymerized, although not as efficiently as ethylene. For example, the reaction of **2** and  $\text{Me}_3\text{Al}$  (3 equiv) in propylene (25 °C, ~8 atm, 3 h) gave a small amount (~125 turnovers) of atactic polypropylene<sup>9</sup> ( $MW < 10\,000$ ). Similarly, the copolymerization of ethylene (1%) in neat propylene produced small amounts (~350 turnovers) of copolymer, which contained 5–10% propylene (by  $^{13}\text{C}$  NMR).<sup>9</sup> One olefin that is polymerized well by our catalyst is 1,3-butadiene; an ampule containing 50 mL of neat butadiene (bp -4.5 °C) completely solidifies within 30 min to create a partial vacuum inside the ampule.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy<sup>10</sup> indicated that the resulting product was >95% *trans*-1,4-polybutadiene.<sup>11</sup>

The activity of the catalyst is sensitive to the amount of  $\text{Me}_3\text{Al}$  used. As shown in Figure 1, polyethylene production (1 atm, 25 °C, 3 h) is maximized when approximately 3 equiv of  $\text{Me}_3\text{Al}$  is used as cocatalyst. Large excesses of  $\text{Me}_3\text{Al}$  lead to complete deactivation of the catalyst. Both observations are contrary to the behavior of typical "soluble catalysts"<sup>12</sup> prepared from trialkylvanadates, which typically require large excesses of alkyl-aluminum reagents (15–500 equiv) to initiate olefin polymerization.<sup>13</sup> In light of the narrow polydispersity measured for our polyethylene sample (2.28), these results suggest that a well-defined catalyst is being formed<sup>14</sup> and that this catalyst is fundamentally different from these conventional soluble V catalysts.

The presence and equilibration of both **2** and **3** in the starting solution greatly complicate mechanistic studies, but several polymerization reactions performed at -30 °C, where the equilibration of **2** and **3** is negligibly slow, indicate that the active catalyst is derived from the reaction of **2** with  $\text{Me}_3\text{Al}$ . Specifically, the addition of  $\text{Me}_3\text{Al}$  to an ethylene-saturated solution of **2** and **3** (~10:90 by  $^{51}\text{V}$  NMR) at -30 °C initiates ethylene polymerization by slowly consuming **2**, but does not affect the amount of **3** in solution. Furthermore, ethylene polymerization is only initiated when the  $^{51}\text{V}$  NMR resonance for **2** is initially present.

The active polymerization catalyst(s) in our system is (are) currently not known, and it would be inappropriate to speculate about its identity without additional data.<sup>15</sup> It is, however,

important to note that this POMSS-based catalyst is capable of polymerizing olefins when stoichiometrically similar complexes (e.g.,  $(\text{Ph}_3\text{SiO})_3\text{VO}$  and  $(n\text{-PrO})_3\text{VO}$ ) show little or no reactivity. Since our understanding of surface catalysis is both based upon and inherently limited by known reaction chemistry of solution complexes, the unique chemistry of V-containing POMSS may provide new insights into the chemistry of silica-supported vanadate catalysts. Efforts to elucidate the active polymerization catalyst(s) in this interesting system are currently in progress.

**Acknowledgment.** These studies were supported by the National Science Foundation (CHE-8703016 and CHE-9011593) and an NSF Presidential Young Investigator Award (CHE-8657262). Acknowledgement is also made to the Chevron Research Company and to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research (22376-AC3). F.J.F. gratefully acknowledges financial support from an Alfred P. Sloan Foundation Research Fellowship.

(15) A referee has suggested that a V(IV) species might be the active catalyst. This is clearly a reasonable hypothesis, but we have prepared  $[(\text{C}_6\text{H}_{11})_2\text{Si}_7\text{O}_{12}\text{V}(\text{CH}_2\text{SiMe}_3)]$  and found that it does not initiate ethylene polymerization. The synthesis of this complex and the results from our other efforts to elucidate the identity of the polymerization catalyst(s) will be reported in a subsequent article.

## 2D Nuclear Magnetic Resonance Study of the Structure of the Fullerene $\text{C}_{70}$

Robert D. Johnson,\* Gerard Meijer,<sup>1</sup> Jesse R. Salem, and Donald S. Bethune

IBM Research Division, Almaden Research Center  
650 Harry Road, San Jose, California 95120-6099

Received January 10, 1991

Geodesic structures were developed by R. Buckminster Fuller<sup>2</sup> on the basis of his insight into their structural economy and stability. In 1985, similar considerations led Smalley, Kroto, et al.<sup>3</sup> to propose that  $\text{C}_{60}$ , observed in carbon cluster beam experiments,<sup>3,4</sup> possessed the geometry of a soccer ball, and they named it "Buckminsterfullerene". Their prediction of the stability of this molecule, as well as other fullerenes, has been stunningly verified by the recent synthesis of macroscopic amounts of  $\text{C}_{60}$  and  $\text{C}_{70}$ .<sup>5-8</sup> This development has sparked intense research activity in the production and characterization of these materials.<sup>8-18</sup> Raman<sup>9,18</sup>

(8) C.f.: (a) Lachowicz, A.; Thiele, K. H. *Z. Anorg. Allg. Chem.* **1977**, *431*, 88–94. (b) Thiele, K. H.; Adler, B.; Grahlert, H.; Lachowicz, A. *Z. Anorg. Allg. Chem.* **1974**, *403*, 279–88.

(9) Determined by  $^{13}\text{C}$  NMR spectroscopy: (a) Tonelli, A. E. *NMR Spectroscopy and Polymer Microstructure*; VCH Publishers: New York, 1989 and references cited therein. (b) Cheng, H. N. *Polym. Bull.* **1985**, *14*, 347–54. (c) Randall, J. C., Ed. *NMR and Macromolecules: Sequence, Dynamic, and Domain Structure*; ACS Symposium Series 246; American Chemical Society: Washington, DC, 1984 and references cited therein. (d) Randall, J. C. *Polymer Sequence Determination; Carbon-13 NMR Method*; Academic Press: New York, 1977 and references cited therein.

(10) (a) Elgert, K. F.; Quack, G.; Stutz, B. *Polymer* **1976**, *16*, 154–6. (b) Santee, E. R.; Mochel, V. D.; Morton, M. J. *Polym. Sci., Polym. Lett. Ed.* **1973**, *11*, 453–5. (c) Santee, E. R.; Chang, R.; Morton, M. J. *Polym. Sci., Polym. Lett. Ed.* **1973**, *11*, 449–52. (d) Mochel, V. D. *J. Polym. Sci., Part A-1* **1972**, *10*, 1009–18.

(11) Neither pure **2** nor pure  $\text{Me}_3\text{Al}$  reacts with 1,3-butadiene, and only trivial amounts of 1,3-butadiene are polymerized by  $(\text{Ph}_3\text{SiO})_3\text{VO}/\text{Me}_3\text{Al}$  or  $(n\text{-PrO})_3\text{VO}/\text{Me}_3\text{Al}$  under identical conditions.

(12) The term "soluble" has been used quite loosely to describe catalyst solutions that appear to remain homogeneous.<sup>13b</sup> In many instances, serious questions can be raised regarding the apparent homogeneity of the catalyst.

(13) (a) Shuke, J.; Dingshing, Y. *Polym. J.* **1985**, *17*, 899–907. (b) Christman, D. L. *J. Polym. Sci., Part A-1* **1972**, *10*, 471–87. (c) Uchida, Y.; Furuhata, K.; Ishiwatari, H. *Bull. Chem. Soc. Jpn.* **1971**, *44*, 1118–21.

(14) Peebles, L. H. *Molecular Weight Distributions in Polymers*; Wiley-Interscience Publishers: New York, 1971; Chapter 2.

(1) Present address: Department of Molecular and Laser Physics, Catholic University of Nijmegen, Toernooiveld, 6525 ED Nijmegen, The Netherlands.

(2) Fuller, R. B. *Ideas and Integrity*; Collier Books: New York, 1963.

(3) Kroto, H. W.; Heath, J. R.; O'Brien, S. C.; Curl, R. F.; Smalley, R. E. *Nature* **1985**, *318*, 162–163.

(4) Rohlfing, E. A.; Cox, D. M.; Kaldor, A. *J. Chem. Phys.* **1984**, *81*, 3322.

(5) Krätschmer, W.; Fostiropoulos, K.; Huffman, D. R. *Chem. Phys. Lett.* **1990**, *170*, 167.

(6) Krätschmer, W.; Lamb, L. D.; Fostiropoulos, K.; Huffman, D. R. *Nature* **1990**, *347*, 354.

(7) Meijer, G.; Bethune, D. S. *J. Chem. Phys.* **1990**, *93*, 7800.

(8) Taylor, R.; Hare, J. P.; Abdul-Sada, A. K.; Kroto, H. W. *J. Chem. Soc., Chem. Commun.* **1990**, 1423.


(9) Bethune, D. S.; Meijer, G.; Tang, W. C.; Rosen, H. J. *Chem. Phys. Lett.* **1990**, *174*, 219.

(10) Johnson, R. D.; Meijer, G.; Bethune, D. S. *J. Am. Chem. Soc.* **1990**, *112*, 8983.

(11) Ajie, H.; Alvarez, M. M.; Anz, S. J.; Beck, R. D.; Diederich, F.; Fostiropoulos, K.; Huffman, D. R.; Krätschmer, W.; Rubin, Y.; Schriver, K. E.; Sensharma, D.; Whetten, R. L. *J. Phys. Chem.* **1990**, *94*, 8630–8633.

(12) Haufler, R. E.; Conceicao, J.; Chibante, L. P. F.; Chai, Y.; Byrne, N. E.; Flanagan, S.; Haley, M. M.; O'Brien, S. C.; Pan, C.; Xiao, Z.; Billups, W. E.; Ciufolini, M. A.; Hauge, R. H.; Margrave, J. L.; Wilson, L. J.; Curl, R. F.; Smalley, R. E. *J. Phys. Chem.* **1990**, *94*, 8634–8636.

(13) Wilson, R. J.; Meijer, G.; Bethune, D. S.; Johnson, R. D.; Chambliss, D. D.; de Vries, M. S.; Hunziker, H. E.; Wendt, H. R. *Nature* **1990**, *348*, 621–622.



**Figure 1.** Upper trace: aromatic region of the  $^{13}\text{C}$  NMR spectrum of  $\sim 300\ \mu\text{g}$  of  $\text{C}_{70}/\text{C}_{60}$  in toluene- $d_8$  without  $^1\text{H}$  decoupling, obtained with a Bruker AM-500 NMR spectrometer operating at 125.7 MHz at 25  $^\circ\text{C}$ . Lower spectrum: the 2D NMR "INADEQUATE" spectrum of  $\text{C}_{70}$ . Doublets (circled) are obtained at a common double quantum frequency for the two resonances of a bonded carbon pair. The connectivity obtained is linear and highlighted by lines; the asymmetrical intensity string of 10:10:20:20:10 allows the assignment of polar end cap carbon a and the remaining carbons. The spectrum was obtained in the presence of  $\text{Cr}(\text{ac})_3$ , by using the pulse sequence of Mareci and Freeman<sup>25</sup> with a refocusing delay of 5 ms, a sweep width of 13 000 Hz digitized into 8K-work datasets for the horizontal dimension, and a sweep width of 7000 Hz for the double quantum dimension digitized into 512 blocks, with 512 scans/block.

and infrared<sup>5,6,17,18</sup> spectroscopy support the proposed  $\text{C}_{60}$  soccer ball structure, and the single-resonance  $^{13}\text{C}$  NMR spectrum of  $\text{C}_{60}$  is strong evidence for the icosahedral symmetry of this molecule.<sup>8,10</sup>

NMR spectroscopy is a powerful probe of chemical structure.<sup>19,20</sup>  $^{13}\text{C}$  NMR analysis of fullerenes probes structure, torsional strain, and bonding in these molecules. The  $^{13}\text{C}$  NMR spectrum of  $\text{C}_{60}$  is a single line at 143 ppm (in both liquids<sup>8,10</sup> and solids<sup>15,16</sup>), showing that all carbons in the molecule are chemically equivalent. The resonance position for  $\text{C}_{60}$  is in excellent accord with similar carbons<sup>19</sup> in azulene (140.2 ppm), fluorene (141.6 and 143.2 ppm), and 3,5,8-trimethylaceptylene (146.8 ppm); this is consistent with the soccer ball structure for  $\text{C}_{60}$  (Figure 1).

The 1D  $^{13}\text{C}$  NMR spectrum for  $\text{C}_{70}$ , shown in the upper trace of Figure 1, consists of five lines with intensities in the ratio 10:20:10:20:10, as reported by Taylor et al.,<sup>8</sup> which strongly supports the  $D_{5h}$   $\text{C}_{70}$  structure<sup>21</sup> shown in Figure 1. This structure has five chemically distinct kinds of carbon atoms and is similar to the  $\text{C}_{60}$  structure with the insertion of 10 carbons forming a belt around the molecule. Characterization of the bonding of  $\text{C}_{70}$  is important for understanding its structure, and assignment of

the NMR lines is critical for their interpretation. An assignment of the resonances of  $\text{C}_{70}$  was proposed by Taylor et al.,<sup>8</sup> based on model compounds and relative strain in the molecule. However, unequivocal experimental assignment of the lines is clearly desirable.

In this communication we show that it is possible to map the bonding connectivities of  $\text{C}_{70}$  and obtain definitive resonance assignments by the 2D "INADEQUATE" NMR experiment.<sup>22-25</sup> This experiment correlates the  $^{13}\text{C}$  NMR line of a carbon to that of its bonded neighbor by excitation of a shared double quantum coherence via scalar  $J$  coupling, thus yielding the carbon connectivity map of the molecule.  $\text{C}_{70}$  samples enriched to  $\sim 20\%$   $^{13}\text{C}$  were used to increase the probability of adjacent  $^{13}\text{C}$  atoms, reducing the data acquisition time to a feasible 4 days despite low solubility. The enriched clusters were prepared by using cored carbon rods loaded with amorphous  $^{13}\text{C}$  powder (Cambridge Isotopes) in an arc fullerene generator.<sup>6,12,26</sup>

The 2D INADEQUATE NMR spectrum of  $\text{C}_{70}$  is shown in Figure 1. For this experiment, two bonded carbons share a double quantum frequency in the vertical dimension, and peaks occur at the two respective chemical shifts in the horizontal dimension, allowing the correlation to be made. In addition, each peak will be split into a doublet by the relevant  $^1J_{\text{CC}}$  coupling constant. The  $D_{5h}$  structure of  $\text{C}_{70}$  suggests a linear connectivity of the five carbon types, with the polar carbons (labeled a) representing one end of the connectivity and the belt carbons (e) representing the other. The 2D spectrum reveals the four connectivities, showing a single string of connected resonances, with respective intensities 10:10:20:20:10, solidly supporting the  $D_{5h}$  structure. The crucial connectivity is obtained between the two intensity 10 lines at 150.8 ppm and 147.8 ppm, forcing the assignment of the line at 150.8 ppm to the polar end cap carbon a, and hence the remaining assignments. We note that the resonances correlating a to b are the least intense, as only one out of the three bonds of an end cap carbon a connects to a carbon b. In contrast, the belt carbons e at 130.8 ppm are each bonded to two type-d carbons at 144.4 ppm, giving cross peaks twice as intense as those revealing the a-b connectivity. The small separation between resonances b and c gives rise to second-order effects in the cross peaks, reducing the outer line of each multiplet.<sup>27</sup> These results confirm the assignments of ref 8.

Taylor et al.<sup>8</sup> noted that the  $\text{C}_{70}$  chemical shifts indicated torsional strain. Belt carbon e resonating at 130.8 ppm can be contrasted to similar carbons in benzo[a]pyrene (125.5 and 123.8 ppm);<sup>28</sup> the downfield position of e is consistent with torsional strain (e.g., the bridgehead carbons in paracyclophane<sup>19</sup> resonate at 140.4 ppm). However, while carbon a has a similar structural environment to those in  $\text{C}_{60}$ , it resonates 7 ppm further downfield. The origin of this relative shift is not clear, but may be due to differences in ring currents or aromaticity between the two molecules.<sup>29,30</sup>

One-bond carbon-carbon coupling constants ( $^1J_{\text{CC}}$ ) in polycyclic aromatic compounds generally range from 53 to 63 Hz.<sup>31</sup> It has been concluded that the size of the coupling is related to the s character of the bond, and a linear correlation between  $^1J_{\text{CC}}$  and decreasing bond lengths or increasing  $\pi$ -bond orders has been

(14) Wragg, J. L.; Chamberlain, J. E.; White, H. W.; Krätschmer, W.; Huffman, D. R. *Nature* **1990**, *348*, 623-624.

(15) Yannoni, C. S.; Johnson, R. D.; Meijer, G.; Bethune, D. S.; Salem, J. R. *J. Phys. Chem.* **1991**, *95*, 9-10.

(16) Tycko, R.; Haddon, R. C.; Dabbagh, G.; Glarum, S. H.; Douglass, D. C.; Muijsce, A. M. *J. Phys. Chem.* **1991**, *95*, 518-520.

(17) Cox, D. M.; Behal, S.; Disko, M.; Gorun, S.; Greaney, M.; Hsu, C. S.; Kollin, E.; Millar, J.; Robbins, W.; Sherwood, R.; Tindall, P. *J. Am. Chem. Soc.*, in press.

(18) Bethune, D. S.; Meijer, G.; Tang, W. C.; Rosen, H.; Golden, W. G.; Seki, H.; Brown, C. A.; de Vries, M. S. *Chem. Phys. Lett.*, in press.

(19) Kalinowski, H.; Berger, S.; Braun, S. *Carbon-13 NMR Spectroscopy*; Wiley and Sons: New York, 1988.

(20) Ernst, R. R.; Bodenhausen, G.; Wokaun, A. *Principles of NMR in One and Two Dimensions*; Oxford University Press: Oxford, 1987.

(21) Heath, J. R.; O'Brien, S. C.; Zhang, Q.; Liu, Y.; Curl, R. F.; Kroto, H. W.; Zhang, Q.; Tittel, F. K.; Smalley, R. E. *J. Am. Chem. Soc.* **1985**, *107*, 7779.

(22) Johnson, R. D.; Yannoni, C. S.; Salem, J. R.; Meijer, G.; Bethune, D. S. Partially presented at the Material Research Society, Special Session on Buckminsterfullerene, Boston, MA, Nov 29, 1990 (in press).

(23) Bax, A.; Freeman, R.; Kempell, S. P. *J. Am. Chem. Soc.* **1980**, *102*, 4849-4851.

(24) Bax, A.; Freeman, R.; Frenkiel, T. A.; Levitt, M. H. *J. Magn. Reson.* **1981**, *43*, 478-483.

(25) Mareci, T. H.; Freeman, R. *J. Magn. Reson.* **1982**, *48*, 158-163.

(26) Krätschmer, W.; Sorg, N.; Huffman, D. R. *Surf. Sci.* **1985**, *156*, 814.

(27) Pople, J. A.; Schneider, W. G.; Bernstein, H. J. *High Resolution NMR*; McGraw-Hill: New York, 1959.

(28) Unkefer, C. J.; London, R. E.; Waley, T. W.; Daub, G. H. *J. Am. Chem. Soc.* **1983**, *105*, 733.

(29) Haddon, R. C.; Schneemeyer, L. F.; Waszczak, J. V.; Tycko, R.; Dabbagh, G.; Muller, A. J.; Muijsce, A. M.; Rosseinsky, M.; Zahurak, S. M.; Thiel, F. A.; Raghavachari, K.; Elser, V., submitted to *Nature*.

(30) Ruoff, R. S.; Beach, D.; Cuomo, J.; McGuire, T.; Whetten, R. L.; Diederich, F., submitted to *J. Phys. Chem.*

(31) Hansen, P. E. *Org. Magn. Reson.* **1979**, *12*, 109-142.

suggested by several groups for polycyclic aromatic compounds.<sup>28,31-34</sup> A value of  $^1J_{9,10}$  of 45 Hz for azulene was interpreted as evidence for this bond being weak.<sup>35</sup> For C<sub>70</sub>, the 2D spectrum gives the  $^1J_{CC}$  values:  $^1J_{a,b} = 68$ ,  $^1J_{b,c} = 55$ ,  $^1J_{c,d} = 55$ , and  $^1J_{d,e} = 62$  Hz. These values indicate that the four bonds have substantial s character and  $\pi$ -bond order. Bonds d-e and a-b fuse six-membered rings and may be compared with  $^1J_{9,10}$  in 1-methyl- and 2-methylnaphthalene at 52 and 53 Hz, respectively.<sup>33</sup> In analogy with cyclopropane derivatives,<sup>36,37</sup> the larger value of  $^1J_{a,b}$  may arise from both carbons in bond a-b belonging to five-membered rings, whose bonds have greater p character due to smaller internal angles. This should increase the s character of the a-b bond and hence the coupling constant; this effect should be less for bond d-e, as this bond has only carbon d in a five-membered ring. We note that the large values for  $^1J_{CC}$  we report are evidence against proposed structures for fullerenes involving three-membered rings,<sup>38</sup> as by analogy with cyclopropane derivatives<sup>36,37</sup> these rings would be expected to have markedly small coupling constants.

The 2D NMR spectrum of C<sub>70</sub> yields bonding topology, coupling constants, and a definitive assignment of the  $^{13}\text{C}$  NMR spectrum. The bonding topology and coupling constants solidly support the "rugby ball"  $D_{5h}$  structure for this molecule. The resonance assignments confirm those previously proposed.<sup>8</sup> The  $^1J_{CC}$  values are relevant to investigations of reactivity and bonding in fullerenes.

**Acknowledgment.** We gratefully acknowledge stimulating discussions with C. S. Yannoni, F. A. L. Anet, G. M. Wallraff, and C. G. Wade, and we thank C. R. Moylan for a critical reading of the manuscript.

(32) Marshall, J. L.; Ihrig, A. M.; Miller, D. E. *J. Magn. Reson.* **1974**, *16*, 439.

(33) Berger, S. *Org. Magn. Reson.* **1984**, *22*, 47-51.

(34) Hansen, P. E.; Poulsen, O. K.; Berg, A. *Org. Magn. Reson.* **1979**, *12*, 43-49.

(35) Berger, S.; Zeller, K. *J. Org. Chem.* **1984**, *49*, 3725-3728.

(36) Weigert, J. A.; Roberts, J. D. *J. Am. Chem. Soc.* **1972**, *94*, 6021.

(37) Lippert, E.; Prigge, H. *Ber. Bunsen-Ges. Phys. Chem.* **1963**, *67*, 415.

(38) Shibuya, T.-I.; Yoshitani, M. *Chem. Phys. Lett.* **1987**, *137*, 13-16.

## Evidence from EXAFS for a Copper Cluster in the Metalloregulatory Protein CUP2 from Yeast

Kent H. Nakagawa,<sup>†</sup> Carla Inouye,<sup>‡</sup> Britt Hedman,<sup>§</sup>  
Michael Karin,<sup>‡</sup> Thomas D. Tullius,<sup>||</sup> and  
Keith O. Hodgson<sup>\*†</sup>

Department of Chemistry, Stanford University  
Stanford, California 94305-5080


Department of Pharmacology, M-036, School of Medicine  
University of California, San Diego  
La Jolla, California 92093

Stanford Synchrotron Radiation Laboratory  
Stanford University, SLAC, Bin 69, P.O. Box 4349  
Stanford, California 94309


Department of Chemistry, The Johns Hopkins University  
Baltimore, Maryland 21218

Received December 10, 1990

Expression of yeast metallothionein, which binds copper specifically, is regulated by the protein CUP2 (also known as ACE1).<sup>1,2</sup> CUP2 itself is activated for binding to DNA by



**Figure 1.** EXAFS data for CUP2. The solid line is the raw data. The dashed line is the data obtained by Fourier transforming the data into  $R$  space, applying a filter from 0.80 to 3.35 Å, and back-transforming. The dotted line is the calculated EXAFS from two-shell fits described in the text and reported in Table I.



**Figure 2.** Comparison of the Fourier transform on filtered EXAFS data ( $R = 0.80$ – $3.35$  Å,  $k$  range of  $3.0$ – $10.0$  Å<sup>-1</sup>) from CUP2 (solid line) with transforms of a one-shell fit containing S (dashed line) and a two-shell fit containing S and Cu (dotted line). Notice the excellent agreement between the fit and data for the first shell in both cases. However, while the second peak is not reproduced by a single S shell, it is well reproduced with the presence of a second shell of Cu.

copper(I).<sup>3</sup> Yeast metallothionein contains a cluster of eight copper(I) ions bridged by thiolate ligands that are likely provided by the 12 cysteines of the protein.<sup>4</sup> How copper is bound to CUP2 is unknown, however. Since stimulation by copper(I) of CUP2 binding to DNA is a cooperative process,<sup>5</sup> and the DNA binding domain of CUP2 contains 12 cysteines,<sup>3,6</sup> the presence of a copper cluster in CUP2 is also likely. Here we report that Cu K-edge extended X-ray absorption fine structure (EXAFS) gives strong evidence that the coppers bound to CUP2 are sulfur-coordinated and in close proximity to each other, most likely bridged by thiolate sulfurs. The Cu K-edge X-ray absorption edge structure demonstrates that the coppers in CUP2 are in the +1 oxidation state and furthermore indicates that their electronic environment is closest to 3-fold coordination.

Cu K-edge X-ray absorption spectra were collected at the Stanford Synchrotron Radiation Laboratory on wiggler beam line 4-2 (unfocused) under dedicated ring conditions (3.0 GeV, 70–90 mA) using a Si(220) double-crystal monochromator. Protein<sup>7</sup>

(1) Thiele, D. *J. Mol. Cell. Biol.* **1988**, *8*, 2745-2752.

(2) Welch, J.; Fogel, S.; Buchman, C.; Karin, M. *EMBO J.* **1989**, *8*, 255-260.

(3) Buchman, C.; Skroch, P.; Welch, J.; Fogel, S.; Karin, M. *Mol. Cell. Biol.* **1989**, *9*, 4091-4095.

(4) George, G. N.; Byrd, J.; Winge, D. R. *J. Biol. Chem.* **1988**, *263*, 8199-8203.

(5) Fürst, P.; Hamer, D. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, *86*, 5267-5271.

(6) Fürst, P.; Hu, S.; Hackett, R.; Hamer, D. *Cell* **1988**, *55*, 705-717.

<sup>†</sup> Department of Chemistry, Stanford, University.

<sup>‡</sup> Department of Pharmacology, M-036, School of Medicine, University of California, San Diego.

<sup>§</sup> Stanford Synchrotron Radiation Laboratory, Stanford University.

<sup>||</sup> Department of Chemistry, The Johns Hopkins University.