

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/244438028>

Through-Space Polar. π . Effects on the Acidity and Hydrogen-Bonding Capacity of Carboxylic Acids

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · JUNE 1994

Impact Factor: 12.11 · DOI: 10.1021/ja00092a053

CITATIONS

67

READS

70

2 AUTHORS:



chao-tsen Chen

National Taiwan University

71 PUBLICATIONS 2,104 CITATIONS

SEE PROFILE



Jay Siegel

Tianjin University

298 PUBLICATIONS 8,104 CITATIONS

SEE PROFILE

Communications to the Editor

Through-Space Polar- π Effects on the Acidity and Hydrogen-Bonding Capacity of Carboxylic Acids

Chao-Tsen Chen and Jay S. Siegel*


Department of Chemistry
University of California at San Diego
La Jolla, California 92093-0358

Received December 13, 1993

Acidity and hydrogen bonding capacity are crucial factors in molecular recognition¹ and biocatalysis.² The ability to design and create biomimetic catalysts depends on an understanding of the way these properties change as a function of environment. Environmental control of functional group reactivity is a principal mechanism by which the chemistry of active site residues is modified.³ Given that the interaction between two simple phenyl derivatives contains a significant component that is transmitted electrostatically and through space (polar- π),⁴ the question arises of whether this polar- π interaction can be used to affect the properties of a functional group like a carboxylic acid. This report presents our findings for a series of 2,6-bis(*p*-X-phenyl)benzoic acids (DPBA).

Benzoic acids, with flanking ortho phenyl rings, were chosen to maximize the polar- π effect by symmetry and proximity.⁵ The flanking phenyl groups sandwich the carboxyl group, excluding solvent from the faces,^{6,7} and the symmetry of the system amplifies the effect of substitution. Substituents are held distal to the reactive site to minimize local steric effects.

The series 1-9 was prepared by the Hart reaction of the desired X-phenyl Grignard with 2,6-dichloriodobenzene; the resultant anion was quenched with CO₂ (Scheme 1).⁸ Measurement of the pK_a values was done by titrating a solution (1 \times 10⁻³ M) of the acid in 80% (w/w) methylcellulose/water with a standardized potassium hydroxide solution (0.1 M) using a potentiometric microtitration apparatus.^{6,9,10} The titration data were fitted using a nonlinear least squares regression analysis. Two runs were averaged to produce the reported pK_a (Table 1). Binding constants for 1, 6, and 7 with 9-ethyladenine (9-EA) were determined in dry chloroform-*d* by ¹H NMR (at 500 MHz) from a series of solutions of the acid (2 \times 10⁻⁴ M) mixed with various proportions of 9-ethyladenine (2 \times 10⁻⁴ to 2.5 \times 10⁻² M). The values for the


Scheme 1^a

^a (a) Mg, THF; (b) 2,6-dichloriodobenzene; (c) CO₂; (d) prepared as cyclic acetal, deprotected with acidic wet silica in dichloromethane.

Table 1. Values for pK_a and Binding Constants with 9-EA for 1-9

compd	X	σ_p^a	pK _a ^b	K _b (M ⁻¹) ^c
1	<i>p</i> -OMe	-0.28	6.61	830
2	<i>p</i> -Me	-0.14	6.50	
3	H	0.00	6.39	
4	<i>p</i> -F	0.15	6.14	
5	<i>p</i> -Cl	0.24	5.97	
6	<i>p</i> -Br	0.26	5.98	1560
7	<i>p</i> -C(O)Me	0.47	5.87	1930
8	<i>m</i> -OMe	0.10	6.61	
9	<i>m</i> -C(O)Me	0.36	5.87	

^a See ref 12. ^b Measured by microtitration in 80% methyl cellulose/water. ^c Measured by ¹H NMR titration in chloroform-*d*.

Figure 1. Plot of $\Sigma\sigma_{para}$ vs pK_a for Acids 1-9.

binding constants were derived by nonlinear least squares analysis of the data between 20% and 80% saturation.¹¹

A plot of pK_a vs $\Sigma\sigma_{para}$ ¹² for 1-7 shows a strong linear relationship with a ρ value of 1.1 (Figure 1).¹⁰ Substituted benzoic acid, *cis*-cinnamic acid, 2-phenylbenzoic acid (2-PBA), and 4-phenylbenzoic acid in the same solvent display ρ values of 1.7, 0.86, 0.62, and 0.43, respectively.^{7,10,13} The through-space nature of the interaction is demonstrated by the pK_a values of the meta-substituted methoxy and acetyl compounds. Within experimental

(1) (a) Hamilton, A. D. *J. Chem. Educ.* **1990**, *67*, 821. (b) Neder, K. M.; Whitlock, H. W. *J. Am. Chem. Soc.* **1990**, *112*, 9412.

(2) Jenks, W. P. *Catalysis in Chemistry and Enzymology*; McGraw-Hill: New York, 1969.

(3) Fersht, A. R. *Enzyme Structure and Mechanism*, 2nd ed.; Freeman: New York, 1985.

(4) (a) Cozzi, F.; Cinquini, M.; Annunziata, R.; Siegel, J. S. *J. Am. Chem. Soc.* **1993**, *115*, 5330. (b) Cozzi, F.; Cinquini, M.; Annunziata, R.; Dwyer, T.; Siegel, J. S. *J. Am. Chem. Soc.* **1992**, *114*, 5729. (c) Petti, M. A.; Sheppard, T. J.; Barans, R. E., Jr.; Dougherty, D. A. *J. Am. Chem. Soc.* **1988**, *110*, 6825.

(5) Empirical force field and semiempirical calculations using the AM1 Hamiltonian predict the structure of 1 to have the flanking phenyl groups and the best plane of the carboxyl group perpendicular to the central phenyl ring.

(6) This reduces the interference from solvent effects speculated to cause problems in systems like 2-phenylbenzoic acid.

(7) Bowden, K.; Hojati, J. *Chem. Soc., Perkin Trans. 2* **1990**, 1201.

(8) (a) Du, C.-J. F.; Hart, H.; Ng, K.-K. D. *J. Org. Chem.* **1986**, *51*, 3162. (b) All compounds have been chromatographically and spectroscopically analyzed for identity and purity.

(9) This solvent and procedure have been used extensively by Simon¹⁰ and Bowden.⁷ Thus a general wide-range analysis of these results in an appropriate thermodynamic context is possible.

(10) (a) Simon, W.; Mörikofer, A.; Heilbronner, E. *Helv. Chim. Acta* **1957**, *40*, 1918. (b) Simon, W.; Lyssy, G. H.; Mörikofer, A.; Heilbronner, E. *Zusammenstellung von Scheinbaren Dissoziationskonstanten im Lösungsmittelsystem Methylcellulose/Wasser*; Juris-Verlag: Zürich, 1959, Vol. 1.

(11) (a) Schneider, H.-J.; Dürr, H. In *Frontiers in Supramolecular Organic Chemistry and Photochemistry*; VCH: Weinheim, 1990. (b) Connors, K. A. In *Binding Constants*; John Wiley & Sons, Inc.: New York, 1987. (c) *Eureka: The Solver*, version 1.01, Borland International, 1987.

(12) Chapman, N. B.; Shorter, J. *Correlation Analysis in Chemistry*; Plenum Press: New York, 1978.

(13) Fuchs, R.; Bloomfield, J. J. *J. Org. Chem.* **1966**, *31*, 3423.

error, there is no difference between the pK_a values of meta- and para-substituted compounds; therefore, when evaluated at their σ_{meta} values, these data deviate greatly from the fitted line.

There are four transmission mechanisms normally observed for aromatic ring substituents: resonance, π -steric, polarization (through bond or space), and field.¹⁴ Because conjugation between the central ring and both the substituted phenyl and the carboxyl groups is conformationally prohibited, resonance is excluded. When invoked, the π -steric effect is independent of the electronic character of the substituent, and is therefore not applicable here because of the direct and significant relationship between pK_a and σ . The result that para- and meta-substituted acids give the same pK_a also eliminates this possibility. The data are most consistent with the polar- π effect being transmitted through space with strong field and π -polarization components.¹⁵

In the case of para-substituted benzoic acids, pK_a correlates linearly with K_b for 9-EA.¹⁶ The 9-EA/DPBA complex shows a similar relationship between $\Sigma\sigma$ and K_b . For the three compounds studied, 1, 6, and 7, K_b increased from 1 to 7 as pK_a decreased, but with a larger slope than for the benzoic acids (1430 vs 520 $\text{M}^{-1} \sigma^{-1}$). In addition, although the pK_a for 2-PBA (6.47) is comparable to 1, the K_b (270 M^{-1}) of 2-PBA^{11a} is roughly $1/3$ that for 1. Thus, there is something more than hydrogen bonding governing the strength of complexation.

Shifts in the ^1H NMR spectra during titration provide some insight into the geometry of complexation.¹⁷ The amino protons of 9-EA shift downfield, consistent with participation in a hydrogen bond. The ortho aryl protons on the flanking rings and H-2 and H-8 of 9-EA move upfield, indicating that 9-EA sits between the flanking rings in a range close enough to feel the ring current. Two possible arrangements still exist for the complex, one with Watson-Crick and one with Hoogsteen type geometries, however, the magnitude of the shift at H-2 vs H-8 (ca. 250 vs 50 Hz) favors the Watson-Crick option.¹⁸ Thus, a tight complex with close aryl-heterocycle interactions is the proposed picture (Figure 2).

(14) (a) Reynolds, W. F. *Prog. Phys. Org. Chem.* **1983**, *14*, 165. (b) Topsom, R. D. *Prog. Phys. Org. Chem.* **1981**, *12*, 1.

(15) By analogy, these data confirm a direct through-space mechanism for the *cis*-cinnamic acids and 2-PBA. For a related study on 3-phenylnorbornene-2-carboxylic acids, see: Beugelmans-Verrier, M.; Nicolas, L.; Gaudemer, A.; Parello, J. *Tetrahedron Lett.* **1976**, 361.

(16) Zimmerman, S. C. *Top. Curr. Chem.* **1993**, *165*, 71.

(17) Separate titrations were done using either component as titrant (guest). Shift data are interpreted from shifts of the species acting as host.




Figure 2. Proposed general complex of 9-EA with a generic DPBA.

The interaction energy between stacked phenyls has been shown to vary as a function of the electronic demand of the substituent. Therefore it seems to be a reasonable extension to assume that a secondary interaction of this type is leading to the enhanced sensitivity of K_b to substitution.¹⁹

From our intuitive understanding of the distance dependence of chemical forces it is natural to assume that through-bond effects should fall off rapidly with distance compared to simple electrostatic effects.²⁰ Nonetheless, discussion still remains as to the nature of how substituent effects are transmitted.²¹ The terphenyl acids studied here offer a simple and interpretable model. The trends revealed by complexation and acidity experiments lead to the conclusion that chemical intuition is quite accurate here and that through-space effects should become an integral part of our repertoire in the design of molecular receptors and catalysts and in the rationalization of their mode of action.²²

Acknowledgment. This work was supported by the National Science Foundation (CHE-9307582), the NATO Collaborative Research Programme, and the Alfred P. Sloan Foundation.

(18) This is a best guess based on chemical shifts; a more rigorous analysis including NOE and 2-D NMR would be more conclusive.

(19) In a related context Jorgensen has discussed the role of secondary dipole interactions on the stability of hydrogen bond complexes. See: Jorgensen, W. L.; Pranata, J. *J. Am. Chem. Soc.* **1990**, *112*, 2008.

(20) (a) Israelachvili, J. N. *Intermolecular and Surface Forces*; Academic: New York, 1985. (b) Bowden, K.; Chapman, N. B.; Shorter, J. *Can. J. Chem.* **1964**, *42*, 1979.

(21) Bowden, K. *J. Chim. Phys.* **1992**, *89*, 1647.

(22) Such effects are already being designed into "concave reagents" see: Luning, U.; Baumstark, R.; Wagnick, C.; Muller, M.; Schyja, W.; Gerst, M.; Gelbert, M. *Pure Appl. Chem.* **1993**, *65*, 527.