

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/236620710>

# Hydrolysis of Monomethyl-, Dimethyl-, and Trimethyltin(IV) Cations in Fairly Concentrated Aqueous Solutions at $I = 1 \text{ mol } 3 \text{ L}^{-1}$ ( $\text{NaNO}_3$ ) and $T = 298.15 \text{ K}$ . Evidence for the Predominan...

ARTICLE *in* JOURNAL OF CHEMICAL & ENGINEERING DATA · APRIL 2011

Impact Factor: 2.04 · DOI: 10.1021/jje101069y

---

CITATIONS

5

---

READS

48

## 5 AUTHORS, INCLUDING:



**Concetta De Stefano**

Università degli Studi di Messina

226 PUBLICATIONS 3,057 CITATIONS

SEE PROFILE



**Antonio Gianguzza**

Università degli Studi di Palermo

162 PUBLICATIONS 1,948 CITATIONS

SEE PROFILE

Hydrolysis of Monomethyl-, Dimethyl-, and Trimethyltin(IV) Cations in Fairly Concentrated Aqueous Solutions at  $I = 1 \text{ mol} \cdot \text{L}^{-1}$  ( $\text{NaNO}_3$ ) and  $T = 298.15 \text{ K}$ . Evidence for the Predominance of Polynuclear SpeciesRosalia M. Cigala,<sup>†</sup> Concetta De Stefano,<sup>†</sup> Alba Giacalone,<sup>‡</sup> Antonio Gianguzza,<sup>‡</sup> and Silvio Sammartano<sup>\*,†</sup><sup>†</sup>Dipartimento di Chimica Inorganica, Chimica Analitica e Chimica Fisica, Università di Messina, Viale Ferdinando Stagno d'Alcontres 31, I-98166, Messina (Vill. S. Agata), Italy<sup>‡</sup>Dipartimento di Chimica "Stanislao Cannizzaro", Università di Palermo, Viale delle Scienze, I-90128 Palermo, Italy

**ABSTRACT:** The hydrolysis of methyltin(IV) cations at fairly high concentrations was investigated to evaluate the formation of polynuclear species in aqueous solution. The hydrolysis of monomethyltin(IV), dimethyltin(IV), and trimethyltin(IV) was studied by potentiometry at  $T = 298.15 \text{ K}$  and at  $I = 1 \text{ mol} \cdot \text{L}^{-1}$  in  $\text{NaNO}_3$  aqueous solutions. The results obtained gave evidence for the formation of the following polynuclear species, in addition to the mononuclear species already reported, which were also considered in the models proposed for the three systems investigated:  $[(\text{CH}_3)_n\text{Sn}(\text{OH})_3]^0$ ,  $[(\text{CH}_3)_n\text{Sn}(\text{OH})_4]^-$ ,  $[(\text{CH}_3)_n\text{Sn}_2(\text{OH})_4]^{2+}$ ,  $[(\text{CH}_3)_n\text{Sn}_2(\text{OH})_5]^+$ ,  $[(\text{CH}_3)_n\text{Sn}_2(\text{OH})_7]^-$ ,  $[(\text{CH}_3)_n\text{Sn}_3(\text{OH})_5]^{4+}$ ,  $[(\text{CH}_3)_n\text{Sn}_3(\text{OH})_7]^{2+}$ ,  $[(\text{CH}_3)_n\text{Sn}_5(\text{OH})_9]^{6+}$ , and  $[(\text{CH}_3)_n\text{Sn}_5(\text{OH})_{11}]^{4+}$  for the monomethyltin(IV) system;  $[(\text{CH}_3)_2\text{Sn}(\text{OH})]^+$ ,  $[(\text{CH}_3)_2\text{Sn}(\text{OH})_2]^0$ ,  $[(\text{CH}_3)_2\text{Sn}(\text{OH})_3]^-$ ,  $[(\text{CH}_3)_2\text{Sn}_2(\text{OH})_2]^{2+}$ ,  $[(\text{CH}_3)_2\text{Sn}_2(\text{OH})_3]^+$ ,  $[(\text{CH}_3)_2\text{Sn}_3(\text{OH})_4]^{2+}$ , and  $[(\text{CH}_3)_2\text{Sn}_4(\text{OH})_6]^{2+}$  for the dimethyltin(IV) system;  $[(\text{CH}_3)_3\text{Sn}(\text{OH})]^0$ ,  $[(\text{CH}_3)_3\text{Sn}(\text{OH})_2]^-$ ,  $[(\text{CH}_3)_3\text{Sn}_2(\text{OH})]^+$ ,  $[(\text{CH}_3)_3\text{Sn}_2(\text{OH})_2]^0$ , and  $[(\text{CH}_3)_3\text{Sn}_2(\text{OH})_4]^{2-}$  for the trimethyltin(IV) system. The formation constants  $\beta_{pq}^{\text{OH}}$  (relative to the equilibrium  $p(\text{CH}_3)_n\text{Sn}^{(4-n)+} + q\text{OH}^- = ((\text{CH}_3)_n\text{Sn})_p(\text{OH})_q^{(p(4-n)-q)}$ ) can be expressed as a function of the cation charge  $z$  ( $z = 4 - n$ ) by the general equation:  $\log \beta_{pq}^{\text{OH}} / (p + q) = 1 + (2.50 \pm 0.05)z$ , that allows the calculation of rough values of  $\beta_{pq}^{\text{OH}}$  for any  $(p, q)$  couple. The reliability of the proposed speciation models is discussed. As expected for a tricharged cation, by increasing the concentration of  $(\text{CH}_3)_3\text{Sn}^{3+}$ , the yield of polynuclear species becomes predominant; this behavior is less marked for  $(\text{CH}_3)_2\text{Sn}^{2+}$  and  $(\text{CH}_3)_3\text{Sn}^{+}$ .

## ■ INTRODUCTION

Organotin(IV) compounds are used as active components in many industrial formulations for different purposes.<sup>1–3</sup> The most important applications are as pesticides in agriculture,<sup>4,5</sup> as antifouling preservatives in coatings for wood and stones and in paints for ships (see ref 6 and refs therein), as additives for thermal stabilization in the plastic industry,<sup>7</sup> and as catalysts for polyurethane and foams production<sup>8,9</sup> in the glass industry.<sup>10</sup> The wide distribution into the environment of organotin(IV) compounds, whose toxicity is well-recognized,<sup>11–13</sup> is a reason of great concern, also considering their high stability and, as a consequence, their accumulation in biota.<sup>14–20</sup> In spite of the policy adopted by many countries to reduce the use of organotin(IV) compounds in industrial applications,<sup>21</sup> especially as additives in antifouling paints,<sup>22</sup> these compounds, owing to their persistence, are still present in aquatic ecosystems.<sup>23–26</sup> Therefore, it is important to keep under control the environmental distribution and fate of organotins, as well as their reactivity in natural fluids. As a consequence of the acidic character of the organotin(IV) cations, hydrolysis is the most important reaction undertaken by this class of compounds in aqueous solution, often leading to the formation of polynuclear hydroxo-complexes in solution  $(\text{R}_n\text{Sn})_p(\text{OH})_q^{(p(4-n)-q)}$  with  $p > 1$ , depending on the organotin(IV) cation concentration. The hydrolysis of organotin(IV) cations has been subject of investigation for more than fifty years by different research groups,<sup>27–46</sup> using different techniques. To give a contribution to the speciation of organotin(IV) compounds in natural waters, the hydrolysis of the mono-,<sup>47</sup>

di-,<sup>48–50</sup> and trimethyltin(IV)<sup>50–52</sup> cations has also been investigated in our laboratories in different ionic media and at different ionic strengths and temperatures. Moreover, some databases report quantitative data on the hydrolysis of organotin(IV) compounds,<sup>53–55</sup> and some recommended hydrolysis constants were given by Baes and Mesmer.<sup>56</sup>

Recently we were involved in the study of mixed metal hydrolytic species formation, in aqueous solution.<sup>57,58</sup> In these investigations, the use of rather high metal concentrations was necessary to evidence mixing effects, in particular the enhancement of hydrolysis. The results obtained in our previous studies show that the deviation of experimental data from those calculated, by considering only the hydrolysis constants determined for the single metal systems, is in general fairly small. Therefore, constants as precise as possible, determined under the same conditions for both single and mixed systems, are needed. We are now investigating some mixed metal systems that include the methyltin(IV) cation  $[(\text{CH}_3)_n\text{Sn}^{(4-n)+}]$ . For this reason we need accurate values of hydrolysis constants for these cations determined at fairly high metal concentrations.

Special Issue: John M. Prausnitz Festschrift

Received: October 21, 2010

Accepted: January 29, 2011

Published: February 22, 2011

**Table 1. Experimental Conditions for Measurements at  $I = (1.00 \pm 0.05) \text{ mol} \cdot \text{L}^{-1}$  and  $T = 298.15 \text{ K}$** 

methyltin(IV)	HNO <sub>3</sub>	pH range	runs <sup>a</sup>
mmol · L <sup>-1</sup>	mmol · L <sup>-1</sup>		
	(CH <sub>3</sub> ) <sub>3</sub> Sn <sup>3+</sup>		
20–80	10–20	1–11	14
	(CH <sub>3</sub> ) <sub>2</sub> Sn <sup>2+</sup>		
20–80	10–15	1.5–7	13
	(CH <sub>3</sub> ) <sub>3</sub> Sn <sup>+</sup>		
60–85	5–10	2–12	13

<sup>a</sup> 40 to 70 titration points per run.


## EXPERIMENTAL SECTION

**Reagents and Materials.** All reagents were of analytical grade, and the solutions were always prepared using CO<sub>2</sub>-free freshly prepared distilled water ( $R \geq 18 \text{ M}\Omega \cdot \text{cm}$ ). Nitric acid and sodium hydroxide solutions were prepared by diluting concentrated solutions (Riedel-de Haën), and standardized by sodium carbonate (Aldrich) and potassium hydrogen phthalate (Fluka), respectively. Sodium nitrate (Aldrich), used as background salt, was always dried in a oven at 383.15 K before use. Monomethyltin(IV), dimethyltin(IV), and trimethyltin(IV), as chloride salts (by Aldrich), with purities of (> 97, 98, and 98) %, respectively, were used without further purification.

**Safety Considerations.** The effects of methyltin compounds' toxicity are skin and eye irritation, inflammation of the bile duct, hepatotoxicity, and neurotoxicity. Trimethyltin chloride, the most toxic, causes neuronal necrosis. Its ingestion or inhalation can cause lethal poisoning. Certain dialkyltin compounds have been shown to cause adverse effects on cell-mediated immunity, specifically on the T cell lymphocyte. Experimental studies have failed to reveal any evidence of carcinogenicity, mutagenicity, or teratogenicity. Direct contact with the skin must be avoided. Precautions and adequate clothing are absolutely necessary when manipulating the reagent. Disposal of all tin-containing waste from the experiment should be done in accordance with facility guidelines.

**Apparatus and Procedure.** Potentiometric titrations were carried out by using an apparatus consisting of a combination Orion-Ross 8172 glass electrode connected to a Crison pH2002 potentiometer coupled with a Metrohm mod. 665 automatic titrant dispenser. The equipment was connected to a personal computer, and a homemade computer program was used to control titrant delivery and data acquisition and to check for emf stability. The estimated precision of this system was  $\pm 0.15 \text{ mV}$  and  $\pm 0.003 \text{ mL}$  for electromotive force (emf) and titrant volume readings, respectively. All titrations were carried out at  $T = (298.15 \pm 0.10) \text{ K}$ , by using a HAAKE F3 digital thermocryostat, under magnetic stirring and bubbling purified presaturated N<sub>2</sub> through the solution to exclude O<sub>2</sub> and CO<sub>2</sub> inside the solution.

To determine the glass electrode formal potential ( $E^\circ$ ) and, when necessary, the junction potential ( $E_j = j_a[\text{H}^+]$ ), independent titrations of acidic solutions (nitric acid) were performed with standard sodium hydroxide solutions in the same experimental conditions of ionic strength and temperature as the systems under study. The free hydrogen ion concentration scale




**Figure 1.** Distribution diagram of (CH<sub>3</sub>)<sub>3</sub>Sn<sup>3+</sup>: (a)  $c(\text{CH}_3)_3\text{Sn}^{3+} = 1 \text{ mmol} \cdot \text{L}^{-1}$ : 1 (CH<sub>3</sub>)<sub>3</sub>Sn<sup>3+</sup> free; 2 (1,1); 3 (1,2); 4 (1,3); 5 (1,4); 6 (2,5); (b)  $c(\text{CH}_3)_3\text{Sn}^{3+} = 80 \text{ mmol} \cdot \text{L}^{-1}$ : 1 (1,3); 2 (1,4); 3 (2,4); 4 (2,5); 5 (2,7); 6 (3,5); 7 (3,7); 8 (5,9); 9 (5,11); in NaNO<sub>3</sub>  $I = 1.00 \text{ mol} \cdot \text{L}^{-1}$  at  $T = 298.15 \text{ K}$ .

was used for all measurements ( $\text{pH} \equiv -\log[\text{H}^+]$ ). A solution volume of 25 mL containing monomethyltin(IV) trichloride, dimethyltin(IV) dichloride, or trimethyltin(IV) chloride, nitric acid, and sodium nitrate to adjust the ionic strength to the desired values ( $1 \text{ mol} \cdot \text{L}^{-1}$ ), was titrated with standard CO<sub>2</sub>-free sodium hydroxide solutions up to pH ~ 7 for dimethyltin(IV) and up to ~12 for the monomethyltin(IV) and trimethyltin(IV) cations, respectively. The experimental conditions used are reported in Table 1.


From separate titrations of strong acid in NaNO<sub>3</sub> media, the value  $-\log K_w = 13.73 \pm 0.02$  at  $I = (1.00 \pm 0.05) \text{ mol} \cdot \text{L}^{-1}$  and  $T = 298.15 \text{ K}$  was obtained.

**Calculations.** To determine all of the parameters of an acid–base titration (analytical reagent concentrations, electrode formal potential  $E^\circ$ , junction potential coefficient  $j_a$ ) and to calculate complex formation constants (and  $\log K_w$  values), the computer programs ESAB2M and BSTAC were used.<sup>59,60</sup> The ES4ECI<sup>60</sup> computer program was used to draw speciation diagrams and to compute the formation percentage of the species.

Hydrolysis constants are expressed according to the equilibrium 1:



and  $((\text{CH}_3)_n\text{Sn})_p(\text{OH})_q^{(p(4-n)-q)}$  species formation constants are given by the equilibrium 2:



**Table 2.** Summary of Tested Speciation Models for the Hydrolysis of Monomethyltin(IV) [ $I = (1.00 \pm 0.05) \text{ mol} \cdot \text{L}^{-1}$ ,  $T = 298.15 \pm 0.10 \text{ K}$ ]

$\log \beta_{pq}^a$	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
1,2	-	-	-	$-2.840(8)^b$	-	-	-
1,3	$-9.912(5)^b$	$-9.948(4)^b$	$-9.961(5)^b$	$-9.925(5)$	$10.567(13)^b$	$-9.702(18)^b$	$-9.767(15)^b$
1,4	$-22.52(4)$	-	$-21.915(6)$	$-21.888(5)$	-	-	$-21.854(17)$
2,4	$-3.811(9)$	$-3.824(8)$	$-3.816(7)$	-	$-3.656(10)$	$-3.036(13)$	$-3.094(10)$
2,5	$-8.103(7)$	$-8.151(7)$	$-8.154(6)$	$-8.093(7)$	$-9.01(20)$	$-7.762(27)$	$-7.860(23)$
2,7	$-30.65(2)$	$-30.477(8)$	-	-	$-31.425(24)$	$-30.146(32)$	-
3,5	$-1.887(12)$	$-1.891(11)$	$-1.871(10)$	$-1.873(13)$	$-1.548(10)$	$-1.227(16)$	$-1.307(11)$
3,7	$-8.226(7)$	$-8.272(8)$	$-8.272(7)$	$-8.252(8)$	$-8.845(20)$	$-7.652(23)$	$-7.765(18)$
5,9	$-2.725(10)$	$-2.789(11)$	$-2.788(9)$	$-2.845(12)$	-	$-3.57(11)$	-
5,11	$-9.163(11)$	$-9.232(12)$	$-9.229(9)$	$-9.290(13)$	$-10.084(20)$	-	-
$\sigma^2/\sigma_0^{2c}$	1.000	1.100	1.147	1.200	1.565	4.83	5.16

<sup>a</sup>  $\beta_{pq}$  refers to the equilibrium  $p(\text{CH}_3)_3\text{Sn}^{3+} + q\text{H}_2\text{O} = [((\text{CH}_3)_3\text{Sn})_p(\text{OH})_q]^{(3p-q)} + q\text{H}^+$ . <sup>b</sup> Standard deviation in the last figure(s) in parentheses. <sup>c</sup>  $\sigma^2 =$  variance in the fit;  $\sigma_0^2 =$  variance in the fit for the best speciation model.

## RESULTS

**Hydrolysis of  $(\text{CH}_3)_3\text{Sn}^{3+}$ .** The hydrolysis of the monomethyltin(IV) cation in dilute aqueous solutions ( $\leq 1 \text{ mmol} \cdot \text{L}^{-1}$ ) includes the formation of the mononuclear species (in parentheses the  $\log \beta_{pq}$  at infinite dilution and  $T = 298.15 \text{ K}$ , from refs 47 and 50):




and the polynuclear species:



As shown in Figure 1a, this cation undergoes strong hydrolysis, over the whole pH range 2 to 12, and at millimolar concentrations, the formation of polynuclear species is quite significant in the pH range 4 to 8. When higher concentrations are considered, as in our experiments [ $(20 \text{ to } 80) \text{ mmol} \cdot \text{L}^{-1}$ ], other polynuclear species are formed, namely,  $(p,q) = (2,4), (2,7), (3,5), (3,7), (5,9)$ , and  $(5,11)$ . Several speciation models were tested: in Table 2 seven models are summarized (other intermediate combinations of species were rejected by the computer program). The best model was chosen on the basis of lowest overall variance. The speciation diagram drawn in Figure 1b shows high yields for polynuclear species over the whole pH range 2 to 12, and in this range the amount of free cation is negligible.


**Hydrolysis of  $(\text{CH}_3)_2\text{Sn}^{2+}$ .** The hydrolysis of the dimethyltin(IV) cation in dilute aqueous solutions ( $\leq 1 \text{ mmol} \cdot \text{L}^{-1}$ ) includes the formation of the mononuclear species (in parentheses the  $\log \beta_{pq}$  at infinite dilution and  $T = 298.15 \text{ K}$ , from refs 48 to 50):



**Figure 2.** Distribution diagram of  $(\text{CH}_3)_2\text{Sn}^{2+}$ : (a)  $c(\text{CH}_3)_2\text{Sn}^{2+} = 1 \text{ mmol} \cdot \text{L}^{-1}$ : 1  $(\text{CH}_3)_2\text{Sn}^{2+}$  free; 2 (1,1); 3 (1,2); 4 (1,3); 5 (2,2); 6 (2,3); (b)  $c(\text{CH}_3)_2\text{Sn}^{2+} = 80 \text{ mmol} \cdot \text{L}^{-1}$ : 1  $(\text{CH}_3)_2\text{Sn}^{2+}$  free; 2 (1,1); 3 (1,2); 4 (1,3); 5 (2,2); 6 (2,3); 7 (3,4); 8 (4,6); in  $\text{NaNO}_3$   $I = 1.00 \text{ mol} \cdot \text{L}^{-1}$  at  $T = 298.15 \text{ K}$ .



and the polynuclear species:




This cation undergoes fairly strong hydrolysis, as shown in Figure 2a, over the pH range 2 to 12, and at millimolar concentrations the formation of polynuclear species is always

**Table 3. Summary of Tested Speciation Models for the Hydrolysis of Dimethyltin(IV) [ $I = (1.00 \pm 0.05) \text{ mol} \cdot \text{L}^{-1}$ ,  $T = (298.15 \pm 0.10) \text{ K}$ ]**

$\log \beta_{pq}^a$	Model 1	Model 2	Model 3	Model 6	Model 5	Model 4
1,1	-3.185(2) <sup>b</sup>	-3.222(3) <sup>b</sup>	-3.248(3)	-3.181(6) <sup>b</sup>	-3.155(6) <sup>b</sup>	-3.179(3) <sup>b</sup>
1,2	-8.459(3)	-8.473(4)	-8.525(4)	-8.442(9)	-8.466(11)	-8.430(4)
1,3	-15.939(18)	-	-	-	-15.582(20)	-
2,2	-5.293(4)	-5.255(4)	-5.210(4)	-5.341(10)	-5.284(11)	-5.331(4)
2,3	-9.449(7)	-9.646(10)	-9.660(9)	-9.513(11)	-9.312(10)	-9.493(5)
3,4	-11.548(12)	-11.358(9)	-11.358(6)	-11.328(27)	-	-11.454(8)
4,6	-18.65(5)	-17.872(24)	-17.721(15)	-	-	-
4,7	-	-	-	-23.89(12) <sup>d</sup>	-	-
5,10	-	-	-36.426(24) <sup>d</sup>	-	-	-
$\sigma^2/\sigma_0^{2c}$	1.000	1.177	1.232	1.895	1.705	1.303

<sup>a</sup>  $\beta_{pq}$  refers to the equilibrium  $p(\text{CH}_3)_2\text{Sn}^{2+} + q\text{H}_2\text{O} = [((\text{CH}_3)_2\text{Sn})_p(\text{OH})_q]^{(2p-q)} + q\text{H}^+$ . <sup>b</sup> Standard deviation in the last figure(s) in parentheses. <sup>c</sup>  $\sigma^2$  = variance in the fit;  $\sigma_0^2$  = variance in the fit for the best speciation model. <sup>d</sup> Very low formation percentages observed for this species.



**Figure 3.** Distribution diagram of  $(\text{CH}_3)_3\text{Sn}^+$ : (a)  $c(\text{CH}_3)_3\text{Sn}^+ = 1 \text{ mmol} \cdot \text{L}^{-1}$ : 1  $(\text{CH}_3)_3\text{Sn}^+$  free; 2 (1,1); 3 (1,2); 4 (2,1); (b)  $c(\text{CH}_3)_3\text{Sn}^+ = 80 \text{ mmol} \cdot \text{L}^{-1}$ : 1  $(\text{CH}_3)_3\text{Sn}^+$  free; 2 (1,1); 3 (1,2); 4 (2,1); 5 (2,2); 6 (2,4); in  $\text{NaNO}_3$   $I = 1.00 \text{ mol} \cdot \text{L}^{-1}$  at  $T = 298.15 \text{ K}$ .

lower than 0.05 (fraction of hydrolyzed cation). When higher concentrations are considered, as in our experiments [ $(20 \text{ to } 80) \text{ mmol} \cdot \text{L}^{-1}$ ], other polynuclear species are formed, namely,  $(p,q) = (3,4)$  and  $(4,6)$ . As for the monomethyl derivative, several speciation models were tested: in Table 3 six models are considered (other intermediate combinations of species were rejected by the computer program), and the best model was chosen on the basis of lowest variance. The speciation diagram ( $80 \text{ mmol} \cdot \text{L}^{-1}$ ) drawn in Figure 2b shows that high yields of polynuclear species are obtained, in the pH range 3 to 6. Note that for  $(\text{CH}_3)_2\text{Sn}^{2+}$  a shorter pH range was considered since at  $\text{pH} > 7$  the solubility is lower than  $10 \text{ mmol} \cdot \text{L}^{-1}$ . In the pH range 4 to 7 the dimethyltin(IV) cation is fully hydrolyzed.

**Table 4. Summary of Tested Speciation Models for the Hydrolysis of Trimethyltin(IV) [ $I = (1.00 \pm 0.05) \text{ mol} \cdot \text{L}^{-1}$ ,  $T = (298.15 \pm 0.10) \text{ K}$ ]**

$\log \beta_{pq}^a$	Model 2	Model 1a <sup>d</sup>	Model 1	Model 2a <sup>d</sup>
1,1	-6.250(6) <sup>b</sup>	-6.280(16) <sup>b</sup>	-6.240(5) <sup>b</sup>	-6.177(5) <sup>b</sup>
1,2	-18.655(14)	-	-18.968(30)	-
2,1	-4.958(9)	-4.969(20)	-4.932(7)	-5.156(10)
2,2	-11.81(8)	-11.75 (6)	-11.75	-
2,4	-	-	-35.726(24)	-
$\sigma^2/\sigma_0^{2c}$	1.150	1.000	1.000	1.179

<sup>a</sup>  $\beta_{pq}$  refers to the equilibrium  $p(\text{CH}_3)_3\text{Sn}^+ + q\text{H}_2\text{O} = [((\text{CH}_3)_3\text{Sn})_p(\text{OH})_q]^{(p-q)} + q\text{H}^+$ . <sup>b</sup> Standard deviation in the last figure(s) in parentheses. <sup>c</sup>  $\sigma^2$  = variance in the fit;  $\sigma_0^2$  = variance in the fit for the best speciation model. <sup>d</sup> pH range < 8.

**Hydrolysis of  $(\text{CH}_3)_3\text{Sn}^+$ .** The hydrolysis of the trimethyltin(IV) cation in dilute aqueous solutions ( $\leq 10 \text{ mmol} \cdot \text{L}^{-1}$ ) includes the formation of the mononuclear species (in parentheses the  $\log \beta_{pq}$  at infinite dilution and  $T = 298.15 \text{ K}$ , from refs 50 to 52):



and the dinuclear species:



This cation undergoes fairly strong hydrolysis, as shown in Figure 3a, at  $\text{pH} > 4.5$ , and at millimolar concentrations the formation of polynuclear species is fairly negligible. When higher concentrations are considered, as in our experiments [ $(60 \text{ to } 85) \text{ mmol} \cdot \text{L}^{-1}$ ], other polynuclear species are formed, namely,  $(p,q) = (2,2)$  and  $(2,4)$ . Also in this case, several speciation models were tested: in Table 4 four models are considered (other intermediate combinations of species were rejected by the computer program). The best model was chosen on the basis of lowest variance. The speciation diagram drawn in Figure 3b shows that at these concentrations high yields of polynuclear species are obtained, and at  $\text{pH} > 6.5$  the trimethyltin(IV) cation is fully hydrolyzed.



## DISCUSSION

**Reliability of the Proposed Models.** As above reported, the best model was chosen on the basis of a lower variance ratio (see Tables 2 to 4). Nevertheless, it must be emphasized that the reliability of quite complex hydrolysis models, for large pH ranges, can be different in acidic or in alkaline or even in small parts of these regions.

Models 1 to 4 reported in Table 2 for monomethyltin(IV) hydrolysis show statistical parameters that are not very different (even if the differences are significant at the 95 % level), and these models differ mainly for the presence/absence of species formed in the very acidic or alkaline ranges. Moreover, the analysis of residuals shows a small but significant worsening of the least-squares fit in these ranges. On the basis of this evidence we are able to define the reliability ranges:

pH 4.5 to 10 = good

pH 2 to 4.5 and 10 to 12 = fairly good

The model proposed for the dimethyltin(IV) cation is more robust and does not show worsening of the fit at low pH values, but the uncertainty for the equilibrium constant of  $[(\text{CH}_3)_2\text{Sn}(\text{OH})_3]^-$  species is significantly higher with respect to those of other species. In this case the reliability ranges are:

pH 2 to 6 = good


pH 6 to 7 = fairly good

Analogous observations for the hydrolysis of the trimethyltin(IV) cation allow definition of the reliability ranges:

pH 5 to 10 = good

pH 2 to 5 and 10 to 12 = fairly good

**Relevance of Polynuclear Species.** Quite high yields of polynuclear species are observed for the three methyl derivatives with, as expected, the trend:




and, in several conditions, these species are predominant, in particular for the monomethyltin(IV) cation. The mean ratio  $q/p$  of polynuclear species follows the opposite trend, and this indicates that the tendency to form these species is favored by the possibility to form hydroxo bridges. Moreover, we can observe that most of the  $q/p$  (for  $p > 1$ ) values are lower than the charge of methyltin(IV) cations, that is,  $q/p < z$ , while for mononuclear species we observe the maximum values  $q/p = z + 1$ . This means that in the highly alkaline region mononuclear species become predominant. For example,  $(\text{CH}_3)_3\text{Sn}^{3+}$  is characterized by a crossing pH point where the fraction of mononuclear species becomes higher than that of polynuclear ones, and this point is a positive function of the analytical cation concentration [ $\text{pH} \sim 8.4$ ,  $(0.5 \text{ to } 1) \text{ mmol} \cdot \text{L}^{-1}$ ;  $\text{pH} = 10$ ,  $20 \text{ mmol} \cdot \text{L}^{-1}$  and  $\text{pH} = 10.5$ ,  $80 \text{ mmol} \cdot \text{L}^{-1}$ ]. The formation of hydrolysis products with  $p \gg 1$  has been recognized for some 2+ cations<sup>56</sup> and, in particular for 3+ cations, such as  $\text{Al}^{3+}$  and  $\text{Fe}^{3+}$ .<sup>61–64</sup>

**Empirical Relationships.** Another way to report equilibrium constants for the formation of hydrolytic species is given by eq 2

**Table 5. Formation Constants of  $(\text{CH}_3)_n\text{Sn}^{(4-n)+}$  Species [ $I = (1.00 \pm 0.05) \text{ mol} \cdot \text{L}^{-1}$ ,  $T = (298.15 \pm 0.10) \text{ K}$ ]**

$pq$	$\log \beta_{pq}^{\text{OH}}$		
	$(\text{CH}_3)_3\text{Sn}^{3+}$	$(\text{CH}_3)_2\text{Sn}^{2+}$	$(\text{CH}_3)_3\text{Sn}^{+}$
1,1		$10.54 \pm 0.02^a$	$7.49 \pm 0.02^a$
1,2		$19.00 \pm 0.03$	$8.49 \pm 0.07$
1,3	$31.28 \pm 0.04^a$	$25.25 \pm 0.05$	
1,4	$32.40 \pm 0.09$		
2,1			$8.80 \pm 0.02$
2,2		$22.17 \pm 0.03$	$15.71 \pm 0.16$
2,3		$31.74 \pm 0.04$	
2,4	$51.11 \pm 0.04$		$19.19 \pm 0.06$
2,5	$60.55 \pm 0.05$		
2,7	$65.46 \pm 0.07$		
3,4		$43.37 \pm 0.05$	
3,5	$66.76 \pm 0.05$		
3,7	$87.88 \pm 0.05$		
4,6		$63.73 \pm 0.11$	
5,9	$120.84 \pm 0.06$		
5,11	$141.87 \pm 0.07$		

<sup>a</sup>  $\pm 95\%$  C.I.



**Figure 4.** Experimental vs calculated  $\log \beta_{pq}^{\text{OH}}$  values, according to eq 4.  $\circ$ , monomethyltin(IV);  $\triangle$ , dimethyltin(IV);  $\nabla$ , trimethyltin(IV).

where the binding of  $\text{OH}^-$  by the cation is considered ( $\beta_{pq}^{\text{OH}}$ ). In Table 5, we give these formation constants for the mono-, di-, and trimethyl cation derivatives. The analysis of these data reveals that  $\log \beta^{\text{OH}}$  is a fairly linear function of the sum of the stoichiometric coefficients  $p$  and  $q$  of eq 2, according to the relationship:

$$\log \beta_{pq}^{\text{OH}} = A(p + q) \quad (3)$$

Moreover,  $A$  can be expressed as a function of the cation charge  $z$  [ $(\text{CH}_3)_n\text{Sn}^{(4-n)+}$ ,  $z = 4 - n$ ]:

$$A = 1 + Bz$$

and therefore  $\log \beta_{pq}^{\text{OH}}$  values for all methyltin(IV) derivatives can be expressed by the single parameter:

$$B = 2.50 \pm 0.05$$

**Table 6.** Literature Data for the Formation of  $(\text{CH}_3)_n\text{Sn}^{(4-n)+}-\text{OH}^-$  species at  $I = 1 \text{ mol}\cdot\text{L}^{-1}$  and  $T = 298.15 \text{ K}$ 

cation	species	$\log \beta^{\text{OH}}$			ref
		NaCl	NaNO <sub>3</sub>	NaClO <sub>4</sub>	
$(\text{CH}_3)_3\text{Sn}^{3+}$	(1,2)	23.51	23.47		50
	(1,3)	31.89	31.88		50
	(1,4)	34.33	34.41		50
	(2,5)	59.93	59.61		50
$(\text{CH}_3)_2\text{Sn}^{2+}$	(1,1)	9.98	10.58	10.68	53
	(1,2)	18.15	18.98	19.02	53
	(1,3)	20.9	21.7	21.6	53
	(2,2)	21.8	22.4	22.5	53
$(\text{CH}_3)_3\text{Sn}^{3+}$	(2,3)	30.1	31.6	31.8	53
	(1,1)	7.44	7.49		53

In Figure 4, we show  $\log \beta_{pq}^{\text{OH}} (\text{exp})$  vs  $\log \beta_{pq}^{\text{OH}} (\text{calcd})$ , where one can see a quite good linearity ( $R = 0.995$ ). The general equation:

$$\log \beta_{pq}^{\text{OH}} / (p + q) = 1 + 2.50z \quad (4)$$

allows the calculation of rough values of  $\beta_{pq}^{\text{OH}}$  for any  $(p,q)$  couple, that, in turn, can be useful in determining the extra stability of mixed hydrolytic species.<sup>57,58</sup> In fact, to calculate the quantitative parameters for the tendency to form mixed metal species, it is necessary to know the formation constants of some homopolynuclear species often not experimentally accessible.<sup>57,58</sup> Equations with more than one parameter can be obtained, such as that proposed by Brown et al.<sup>63</sup> for the hydrolysis of  $\text{Al}^{3+}$  ( $-\log \beta_{pq} = 5.73 - 3.6p + 4.64q$ ). As an example, for the polynuclear species of  $(\text{CH}_3)_3\text{Sn}^{3+}$ , we have  $-\log \beta_{pq} = 3.33 - 5.78p + 3.16q$ .

## FINAL REMARKS

To show the effect of the supporting electrolyte, we report in Table 6 some literature data in different salts at  $1 \text{ mol}\cdot\text{L}^{-1}$  aqueous solutions. The differences are generally quite small, even if there is a systematic lowering for  $\log \beta_{pq}^{\text{OH}}$  determined in NaCl. This means that the methyltin(IV) cations form very weak ion pairs with chloride ions. Attempts were already made to calculate the formation constants for these ion pairs,<sup>47,48,50,51</sup> with quite small constants resulting. Therefore we can affirm that the amount of  $\text{Cl}^-$ , deriving from methyltin(IV) chlorides, does not seriously affect the equilibrium constants reported in this work.

A comparison with literature data can be done only for the di- and trimethyl derivatives (data for the monomethyl derivative come from these laboratories). Formation constants reported in Tables 5 and 6 are in fairly good accordance, also taking into account that in the present investigation many other polynuclear species were found. These species are predominant, in particular for  $(\text{CH}_3)_3\text{Sn}^{3+}$ , over a large pH range, as shown in Figure 2. This result, together with the possibility of using the general empirical eq 4, is quite useful for the analysis of mixed metal systems. The formation of large amounts of homopolynuclear species, in particular those with  $p > 2$ , allows the prediction of the formation of heteropolynuclear species in mixed solutions containing different kinds of alkyltin(IV) compounds and other highly charged metal ions.

## AUTHOR INFORMATION

### Corresponding Author

\*Silvio Sammartano. E-mail: ssammartano@unime.it. Fax: +39090392827.

### Funding Sources

We thank the Universities of Messina and Palermo for partial financial support.

## REFERENCES

- (1) Davies, A. G. *Organotin Chemistry*; Wiley-VCH Verlag: Weinheim, 2004.
- (2) Champ, M. A.; Seligman, P. F. *Organotin. Environmental fate and effects*; Chapman & Hall: London, 1996.
- (3) Cima, F.; Craig, P. J.; Harrington, C. Organotin compounds in the environment. In *Organometallic compounds in the environment*, 2nd ed.; Craig, P. J., Ed.; John Wiley & Sons: Chichester, U.K., 2003; pp 101–157.
- (4) Crowe, A. J. Organotin compounds in agriculture since 1980. Part. I. Fungicidal, bactericidal and herbicidal properties. *Appl. Organomet. Chem.* **1987**, *1*, 143–156.
- (5) Shukla, S. K.; Tiwari, V. K.; Rani, S.; Ravi, K.; Tewari, I. C. Studies on insecticidal and pesticidal activity of some organotin compounds. *Int. J. Agric. Sci.* **2010**, *2*, 5–10.
- (6) Champ, M. A.; Seligman, P. F. An introduction to organotin compounds and their use in antifouling coatings. In *Organotin. Environmental fate and effects*; Champ, M. A., Seligman, P. F., Eds.; Chapman & Hall: London, 1996; Chapter 1, pp 1–27.
- (7) Ayrey, G.; Hau, S. Y.; Poller, R. C. The use of organotin compounds in the thermal stabilisation of poly(vinylchloride). VI. An assessment of the reactive importance of HCl scavenging, exchange and addition reactions. *J. Polym. Sci.* **1984**, *22*, 2871–2886.
- (8) Jousseau, B.; Noiret, N.; Pereyre, M.; Saux, A.; Frances, J. M. Air activated organotin catalysts for silicone curing and polyurethane preparation. *Organometallics* **1994**, *13* (3), 1034–1038.
- (9) Gaina, C.; Gaina, V.; Cristea, M. Poly(urethane-urea) varnishes containing tributyltin groups. *J. Inorg. Organomet. Polym.* **2009**, *19*, 157–165.
- (10) Le Bourhis, E. *Glass: mechanics and technology*; Wiley-WCH, Verlag: Weinheim, 2008.
- (11) Fent, K. Ecotoxicology of organotin compounds. *Crit. Rev. Toxicol.* **1996**, *26*, 1–117.
- (12) Arakawa, Y.; Wada, O. Biological properties of alkyltin compounds. In *Metal ions in biological systems*; Sigel, H., Sigel, A., Eds.; Marcel Dekker Inc.: New York, 1993; pp 101–137.
- (13) Fent, K. Ecotoxicology of organotin compounds. *Crit. Rev. Toxicol.* **1996**, *26*, 3–117.
- (14) Mizukawa, H.; Takahashi, S.; Nakayama, K.; Sudo, A.; Tanabe, S. Contamination and Accumulation Feature of Organotin Compounds in Common Cormorants (*Phalacrocorax carbo*) from Lake Biwa, Japan. *Interdiscip. Stud. Environ. Chem.— Environ. Res. Asia* **2009**, 153–161.
- (15) Suzuki, T.; Yamamoto, I.; Yamada, H.; Kaniwa, N.; Kondo, K.; Murayama, M. Accumulation, Metabolism, and Depuration of Organotin Compounds in the Marine Mussels *Mytilus graynus* and *Mytilus edulis* under Natural Conditions. *J. Agric. Food Chem.* **1998**, *46*, 304–313.
- (16) Kannan, K.; Corsolini, S.; Focardi, S. Accumulation Pattern of Butyltin Compounds in Dolphin, Tuna and Shark Collected from Italian Coastal Waters. *Environ. Contam. Toxicol.* **1995**, *31*, 19–23.
- (17) Meng-Pei, H.; Shiu-Mei, L. Accumulation of organotin compounds in Pacific oysters, *Crassostrea gigas*, collected from aquaculture sites in Taiwan. *Sci. Total Environ.* **2003**, *313*, 41–48.
- (18) Iwata, H.; Tanabe, S.; Tatsukawa, R. Bioaccumulation of butyltin compounds in marine mammals: the specific tissue distribution and composition. *Appl. Organomet. Chem.* **1997**, *11*, 257–264.
- (19) El Hassani, L. H.; Frenich, A. G.; Martinez Vidal, J. L.; Muros, M. J. S.; Benajiba, M. H. Study of the accumulation of tributyltin and

triphenyltin compounds and their main metabolites in the sea bass, *Dicentrarchus labrax*, under laboratory conditions. *Sci. Total Environ.* **2005**, 348, 191–198.

(20) Strand, J.; Jacobsen, J. A. Accumulation and trophic transfer of organotins in a marine food web from the Danish coastal waters. *Sci. Total Environ.* **2005**, 350, 72–85.

(21) Champ, M. A. A review of organotin regulatory strategies, pending actions, related costs and benefits. *Sci. Total Environ.* **2000**, 258, 21–71.

(22) Omae, I. Organotin antifouling paints and their alternatives. *Appl. Organomet. Chem.* **2003**, 17, 81–105.

(23) Chau, Y. K.; Maguire, R. J.; Brown, M.; Yang, F.; Batchelor, S. P. Occurrence of organotin compounds in the Canadian aquatic environment five years after the regulation of antifouling uses of tributyltin. *Water Qual. Res. J. Can.* **1997**, 32, 453–521.

(24) Diez, S.; Abalos, M.; Bayona, J. M. Organotin contamination in sediments from the Western Mediterranean enclosures following 10 years of TBT regulation. *Water Res.* **2002**, 36, 905–918.

(25) Gibson, C. P.; Wilson, S. P. Imposex still evident in eastern Australia 10 years after tributyltin restrictions. *Mar. Environ. Res.* **2003**, 55, 101–112.

(26) Murai, R.; Takahashi, S.; Tanabe, S.; Takeuchi, I. Status of butyltin pollution along the coasts of western Japan in 2001, 11 years after partial restrictions on the usage of tributyltin. *Mar. Pollut. Bull.* **2005**, 51, 940–949.

(27) Tobias, R. S.; Ogrins, I.; Nevett, B. A. Studies on the mono- and polynuclear hydroxo complexes of dimethyltin(IV) ion in aqueous solution. *Inorg. Chem.* **1962**, 1, 638–646.

(28) Janssen, M. J.; Luijten, J. G. Investigations on organotins compounds. XVIII The basicity of triorganotin hydroxides. *Rec. Trav. Chim.* **1963**, 85, 1008–1014.

(29) McGrady, M. M.; Tobias, R. S. Raman, Infrared and Nuclear Magnetic Resonance Spectroscopic studies on aqueous solutions of dimethyltin(IV) compounds: structure and bonding of the aquo dimethyltin ion. *Inorg. Chem.* **1964**, 3, 1157–1163.

(30) Tobias, R. S.; Yasuda, M. Studies on the soluble intermediates in the hydrolysis of dimethyltin dichloride. *Can. J. Chem.* **1964**, 42, 781–791.

(31) Tobias, R. S.; Freidline, C. E. Raman spectroscopic and e.m.f. studies on aqueous solutions of the trans-tetrahydroxodimethylstannate(IV) ion. The four acid dissociation constants of dimethyltin(IV) aquo ion. *Inorg. Chem.* **1965**, 4, 215–220.

(32) Van den Bergue, E. V.; Van der Kelen, G. P. On the NMR spectra of methyltintrichloride in solution. *Bull. Soc. Chim. Belges.* **1965**, 74, 479–480.

(33) Tobias, R. S. Bonded organometallic cations in aqueous solutions and crystals. *Organometal. Chem. Rev.* **1966**, 1, 93–129.

(34) Luijten, J. G. A. Investigations on organotin compounds. XXII. Hydrolysis products of alkyltin trichlorides. *Recueil.* **1966**, 85, 873–877.

(35) Tobias, R. S.; Farrer, H.; Hughes, M.; Nevett, B. A. Hydrolysis of the aquo ions  $R_3Sn^+$  and  $R_2Sn^{2+}$ : steric effects on the dissociation of aquo acids. *Inorg. Chem.* **1966**, 5, 2052–2055.

(36) Asso, M.; Carpeni, G. Recherches sur le point isohydrique et les équilibres acido-basique de condensation ou association en chimie. XXVII. Les hydroxocomplexes des sels organo stanniques  $(CH_3)_3SnCl$  et  $(C_2H_5)_2SnCl$ , en solutions aqueuses, à 25 °C et à différentes forces ioniques. *Can. J. Chem.* **1968**, 46, 1795–1802.

(37) Devaud, M. Etude en solution hydroalcoolique de l'ethyltrichloroétain. *J. Chim. Phys.* **1969**, 66, 302–312.

(38) Devaud, M. Etude potentiométrique en milieu hydroalcoolique du trichlorure de triperchlorate d'éthylétain. *J. Chim. Phys.* **1979**, 67, 270–278.

(39) Devaud, M. Comportement de l'ethyltrichlorostannane en milieu hydroalcoolique basique. *J. Chim. Phys.* **1972**, 69, 460–469.

(40) Blunden, S. J.; Smith, P. J.; Gillies, D. G. An investigation of the hydrolysis products of mono alkyltin trichlorides by  $^{119}Sn$  Mossbauer and  $^1H$  and  $^{119}Sn$  NMR spectroscopy. *Inorg. Chim. Acta* **1982**, 60, 105–109.

(41) Arena, G.; Purrello, R.; Rizzarelli, E.; Gianguzza, A.; Pellerito, L. Thermodynamics of hydroxo complex formation of dialkyltin(IV) ions in aqueous solution. *J. Chem. Soc., Dalton Trans.* **1989**, 773–777.

(42) Blunden, S. J.; Hill, R. An investigation of the base hydrolysis of methyl- and butyl-tin trichloride in aqueous solution by  $^1H$  and  $^{119}Sn$  NMR spectroscopy. *Inorg. Chim. Acta* **1990**, 177, 219–223.

(43) Cunningham, D.; McManus, J.; Hynes, M. J. Nuclear magnetic resonance studies and structural investigations of the chemistry of organotin compounds. *J. Organomet. Chem.* **1990**, 393, 69–82.

(44) Hynes, M. J.; Keely, J. M.; McManus, J. Investigation of  $[Sn(CH_3)_3(H_2O)_2]^+$  in aqueous solution by Tin-119 Nuclear Magnetic Resonance Spectroscopy. *J. Chem. Soc., Dalton Trans.* **1991**, 3427–3429.

(45) Barbieri, R.; Silvestri, A. The hydrolysis of  $Me_2Sn^{IV}$  and  $Me_3Sn^{IV}$  moieties monitored through  $^{119}Sn$  Mössbauer spectroscopy. *Inorg. Chim. Acta* **1991**, 188, 95–98.

(46) Takahashi, T.; Natsume, N.; Koshino, S.; Funahashi, Y.; Takagi, D. Speciation of trimethyltin(IV): hydrolysis, complexation equilibria, and structures of trimethyltin(IV) ion in aqueous solution. *Can. J. Chem.* **1997**, 75, 1084–1092.

(47) De Stefano, C.; Foti, C.; Gianguzza, A.; Marrone, F.; Sammartano, S. Hydrolysis of methyltin(IV) trichloride in aqueous NaCl and NaNO<sub>3</sub> solution at different ionic strengths and temperatures. *Appl. Organomet. Chem.* **1999**, 13, 805–811.

(48) De Stefano, C.; Foti, C.; Gianguzza, A.; Martino, M.; Pellerito, L.; Sammartano, S. The Hydrolysis of  $(CH_3)_2Sn^{2+}$  in different ionic media. Salt Effects and Complex Formation. *J. Chem. Eng. Data* **1996**, 41, 511–515.

(49) Foti, C.; Gianguzza, A.; Millero, F. J.; Sammartano, S. The speciation of  $(CH_3)_2Sn^{2+}$  in electrolyte solution containing the major components of natural waters. *Aquat. Geochem.* **1999**, 5, 381–398.

(50) Foti, C.; Gianguzza, A.; Piazzese, D.; Trifiletti, G. Inorganic speciation of organotin(IV) cations in natural waters with particular references to seawater. *Chem. Spec. Bioavail.* **2000**, 12, 41–52.

(51) Cannizzaro, V.; Foti, C.; Gianguzza, A.; Marrone, F. Hydrolysis of trimethyltin(IV) cation in NaNO<sub>3</sub> and NaCl aqueous media at different temperatures and ionic strengths. *Ann. Chim.* **1998**, 88, 45–54.

(52) De Stefano, C.; Foti, C.; Gianguzza, A.; Millero, F. J.; Sammartano, S. Hydrolysis of  $(CH_3)_3Sn^+$  in various salt media. *J. Solution Chem.* **1999**, 28 (7), 959–972.

(53) Martell, A. E.; Smith, R. M.; Motekaitis, R. J. *NIST Critically selected stability constants of metal complexes database*, 8.0; National Institute of Standard and Technology: Gaithersburg, MD, 2004.

(54) Pettit, L.; Powell, K. J. *The IUPAC Stability Constants Database*; Academic Software: Otley, U.K., 2001.

(55) May, P. J.; Murray, K. *Joint Expert Speciation System (JESS Primer)*; Murdoch University: Perth, Western Australia, 2000.

(56) Baes, C. F.; Mesmer, R. E. *The Hydrolysis of Cations*; Wiley: New York, 1976.

(57) Crea, F.; Milea, D.; Sammartano, S. Enhancement of hydrolysis through the formation of mixed hetero-metal species. *Talanta* **2005**, 65, 229–238.

(58) Crea, F.; Milea, D.; Sammartano, S. Enhancement of hydrolysis through the formation of mixed hetero-metal species: dioxouranium(VI)-cadmium(II) mixtures. *Ann. Chim. (Rome)* **2005**, 95, 767–778.

(59) De Stefano, C.; Princi, P.; Rigano, C.; Sammartano, S. Computer Analysis of Equilibrium Data in Solution. ESAB2M: An Improved Version of the ESAB Program. *Ann. Chim. (Rome)* **1987**, 77, 43–675.

(60) De Stefano, C.; Mineo, P.; Rigano, C.; Sammartano, S. Ionic strength dependence of formation constants. XVII. The calculation of equilibrium concentrations and formation constants. *Ann. Chim. (Rome)* **1993**, 83, 243–277.

(61) Ciavatta, L.; Grimaldi, M. On the hydrolysis of the iron(III) ion,  $Fe^{3+}$ , in perchlorate media. *J. Inorg. Nucl. Chem.* **1975**, 37, 163–169.

(62) Daniele, P. G.; Rigano, C.; Sammartano, S.; Zelano, V. Ionic strength dependence of formation constants—XVIII. The hydrolysis of iron(III) in aqueous KNO<sub>3</sub> solutions. *Talanta* **1994**, 41, 1577–1582.



(63) Brown, P. L.; Sylva, R. N.; Batley, G. E.; Ellis, J. The hydrolysis of metal ions. Part 8. Aluminium(III). *J. Chem. Soc., Dalton Trans.* **1985**, 1967–1970.

(64) Cigala, R. M.; De Stefano, C.; Giacalone, A.; Gianguzza, A. The hydrolysis of  $\text{Al}^{3+}$  in fairly concentrated solutions (20 to 200  $\text{mmol L}^{-1}$ ) at  $I = 1 \text{ mol L}^{-1}$  ( $\text{NaNO}_3$ ), in the acidic pH range, at different temperatures. *Chem. Spec. Bioavail.*, DOI: 10.3184/095422911X12971889727358.