

J Org Chem. Author manuscript; available in PMC 2009 September 5.

Published in final edited form as:

J Org Chem. 2008 September 5; 73(17): 6928–6931. doi:10.1021/jo801181d.

A Practical and Efficient Route for the Highly Enantioselective Synthesis of Mexiletine Analogues and Novel β -Thiophenoxy and Pyridyl Ethers

Kun Huang, Margarita Ortiz-Marciales^{*}, Viatcheslav Stepanenko, Melvin De Jesús, and Wildeliz Correa

Department of Chemistry, University of Puerto Rico-Humacao, CUH Station, Humacao, Puerto Rico 00791

Abstract

$$R = Me, Aryl$$

A practical and efficient procedure for the enantioselective synthesis of mexiletine analogues with use of 10% of spiroborate ester $\bf 6$ as chirality transfer agent is presented. A variety of mexiletine analogues were prepared in good yield with excellent enantioselectivities (91–97% ee) from readily available starting materials. The developed methodology was also successfully applied for the synthesis of novel β -amino ethers containing thiophenyl and pyridyl fragments.

Mexiletine (Figure 1) is an effective sodium channel blocker used as an antiarrhythmic and analgesic oral drug. 1,2 Structure-activity studies in vivo 3 and in vitro 4 of pharmacologically active mexiletine indicate that its (-)-(R) enantiomer binds preferentially to the cardiac sodium channels. In addition, (-)-(R)-mexiletine is also more active than (+)-(S)-mexiletine on native skeletal muscular fibers. 1 b, 5 The use of mexiletine as a racemate in the treatment of neuromuscular disorders is limited due to its possible side effects. 6 The optically active mexiletine analogue (R)-2 (Figure 1) is 27-fold more potent than (R)-mexiletine in producing a tonic block and 23-fold more potent in condition of high frequency of stimulation (phasic block). 6 Recently, racemate 3 was established as a novel potent blocker of voltage-gated K⁺ channels by using structure-based virtual screening in conjunction with electrophysiological assays in rat hippocampal neurons. 7

The preparation of mexiletine enantiomers has been reported previously by several groups. Generally, the methods involved resolution of racemic intermediates, ⁸ enzymatic hydrolysis of an *N*-acyl derivative, ⁹ or using a stereospecific, four-step procedure, in 7.2% overall yield. ^{8b} Flippin and co-workers ¹⁰ reported a convenient procedure for the preparation of stereoisomers of mexiletine, but the scope of products was limited by the availability of chiral substrates and expensive chromium tricarbonyl complexes of aryl halides: hence, some amines were provided in the racemate form. Although Franchini's group ⁶ synthesized the stereoisomers of mexiletine analogues, the use of 2-phenyloxirane as chiral source restricted

the range of mexiletine analogues. Furthermore, the procedure is also controlled by the regiospecificity of the ring-opening reaction, ¹¹ and the possible racemization of the benzylic carbon by the substitution of the alkoxy group by the amine. ¹² Hence, a practical and efficient route for the synthesis of highly enantiopure mexiletine analogues is highly desirable.

The asymmetric reduction of oxime ethers with borane-based catalysts offers a facile and direct approach to obtain enantioenriched primary amines; however, more than an stoichiometric amount of in situ prepared oxazaborolidine has been employed to obtain a high degree of enantioselectivity. $^{13a-n}$ Our recent success in the borane-mediated asymmetric reduction of O-benzyl oximes, using a truly catalytic amount of the air and moisture stable spiroborate esters indicated in Figure $2^{,13o,p}$ prompted us to investigate their application for the enantioselective synthesis of β -amino ethers of biological interest.

The synthesis of **12a**, as outlined in Scheme 1, was selected as a model protocol. The aryloxy acetophenone **9a** was prepared from the 2-chloro- or 2-bromoacetophenone with different bases (NaOH, NaH, and K_2CO_3). After optimization, it was found that treatment of 2-bromoacetophenone with K_2CO_3 (1.5 equiv) and 2,6-dimethylphenol (1.5 equiv) in DMF at room temperature for 24 h gave the desired ketone in 81% yield. The oxime was then prepared by addition of NH₂OH · HCl and pyridine at 0 °C. ¹⁴ Since the enantiofacial selectivity in the reduction of C=N bonds depends not only on the chirality's transfer agent but also on the E/Z isomeric purity, ¹³ the main (Z) isomer was easily obtained by simple recrystallization. The pure (Z)-benzyl oxime ether **11a** was obtained in high yield from (Z)-oxime with use of NaH and benzyl bromide in DMF at -30 °C.

Initially, we screened the spiroborates presented in Figure 2 for the reduction of *O*-benzyl oxime **11a.** The reactivity of catalysts **5**, **7**, and **8** was rather low at 0 °C. The complete conversion was achieved at room temperature (22 °C) affording 57%, 81%, and 82% ee, respectively, (entries 2, 5, and 6, Table 1). Spiroborate ester **6** provided the best enantioselectivities (Table 1, entries 3 and 4).

Under the optimized conditions, a variety of O-benzyl oximes were prepared and reduced in dioxane employing 10% spiroborate ester $\bf 6$ and 4 equiv of BH $_3$ ·THF at 0 °C. Table 2 illustrates the results for various O-benzyl oximes that were reduced to the enantiopure amines in 91–97% ee in good to excellent isolated yield. Noteworthy, (S)-mexiletine was prepared in 94% ee and 84% chemical yield (entry 2), illustrating, clearly, the catalyst ability to differentiate between the methyl and the alkoxy moiety in the borane reduction. Generally, substituents on each aryl group influenced slightly the enantioselectivities (entries 3–11). For example, O-benzyl oximes bearing both electron-withdrawing and electron-donating aryl groups provided higher enantioselectivities (entry 3 and 6). The absolute configuration of products was determined by comparing the optical rotations with the corresponding known compounds.

Our current efforts in the design of novel neuronal nicotinic acetylcholine receptors (nAChRs) agonists led us to target enantiopure arylamino ethers as therapeutic agents for the treatment of CNS and peripheral nervous system disorders. 15,16 Within the past ten years, a series of selective agonist of human neuronal nAChRs, such as Epibatidine (13), 17 A-85380 (14), 18 ABT-594 (15), 19 A-84543 (16), 20 and SIB-1553A (17) 21 (Figure 3), has been found. Additionally, compounds containing fluorine are widely applied in the fields of agricultural, medicinal, and material chemistry, 22 especially fluorinated aromatics which are widely found in many modern drugs.

On the basis of related structural features present in nAChRs agonists, highly enantiopure benzylic primary amines or fluorinated aryl groups with β -thiophenoxy or pyridyloxy groups of biological significance were prepared by using the previous procedure in good yield with excellent enantioselectivity (94–97% ee), as shown in Table 3.

In summary, the preparation of chiral mexiletine analogues, in addition to novel β -thiophenoxy and pyridyl ethers, was achieved by a facile four-step synthesis in good yield and excellent enantioselectivities. The highly enantioselective reduction of benzyloxime α -ethers, efficiently catalyzed by 10% of the novel spiroborate ester $\mathbf{6}$, is applicable for a wide range of other bioactive compounds. Considering the convenience and efficiency of this route, this methodology should open new avenues for further research.

Experimental Section

General Procedure for the Preparation of Ketones: 2-(2, 6-Dimethylphenoxy)-1-phenylethanone³ (9a)

To a 100 mL round-bottomed flask charged with a magnetic stirrer, 2-bromoacetophenone (1.99 g, 10 mmol), 2,6-dimethylphenol (15 mmol), and K_2CO_3 (15 mmol) was added 20 mL of DMF and the resulting mixture was stirred for 24 h at room temperature. The reaction was quenched with 50 mL of H_2O and the aqueous phase was extracted with diethyl ether (3 × 40 mL). The combined organic phases were washed with 2 N NaOH (2 × 20 mL) to remove the excess of phenol, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel/hexane:ethyl acetate (5:1), as a yellow solid: mp 57–59 °C (lit.6 mp 60–61 °C); yield 81% (1.94 g); 1H NMR (400 MHz, CDCl₃) δ 2.23 (s, 6H), 5.02 (s, 2H), 6.89 (m, 1H), 6.97 (m, 2H), 7.41 (m, 2H), 7.52 (m, 1H), 7.90 (m, 2H); ^{13}C NMR (100 MHz, CDCl₃) δ 16.3, 74.5, 124.5, 127.9, 128.8, 129.0, 130.8, 133.7, 134.7, 155.7, 194.1; GC-MS m/z 240.2 (M⁺).

General Procedure for the Asymmetric Reduction of *O*-Benzyl Oximes: (*R*)-2-(2,6-Dimethylphenoxy)-1-phenylethylamine (12a)

To a 50 mL reaction tube with catalyst 6 (33 mg, 0.1 mmol) under N₂ was added 10 mL of anhydrous dioxane and 4 mL of BH₃ · THF (1 M in THF) in one portion. After the mixture was stirred for 1 h at room temperature, the clear solution was cooled at 0 °C and the benzyl oxime (1 mmol) in 5 mL of dioxane was added dropwise during 1 h with a syringe pump. The resulting mixture was stirred at 0 °C until the conversion was complete in about 48 h. The reaction was quenched with 6 N HCl and basified with 6 N NaOH. The reaction mixture was extracted with ether and the combined organic phases were washed with brine, dried over anhydrous MgSO₄, and concentrated under reduced pressure. The residue was purified by flash column chromatography and purified by silica gel/hexane:ethyl acetate (2:1) column chromatography as a colorless oil: yield 84% (101 mg); 97% ee; $[\alpha]_D$ 20 –28.5 (c 1.2, CHCl₃) [lit⁶ –14 (c 2, MeOH)], % ee not given; ¹H NMR (400 MHz, CDCl₃) δ 1.99 (s, 2H), 2.33 (s, 6H), 3.93 (m, 2H), 4.51 (m, 1H), 6.99 (m, 1H), 7.07 (m, 2H), 7.35–7.44 (m, 3H), 7.52 (m, 2H); 13 C NMR (100 MHz, CDCl₃) δ 16.4, 56.4, 77.8, 124.0, 126.0, 127.6, 128.5, 128.9, 130.9, 142.3, 155.5; GC-MS m/z 241.1 (M⁺). The Enantiomeric excess was determined by HPLC for the acetyl derivative with use of a Chiralcel OD-H column (9:1 hexane:2-propanol), 0.8 mL/min, 254 nm, minor enantiomer $t_R = 11.3 \text{ min}$, major enantiomer $t_R = 16.2 \text{ min}$.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support by the National Institutes of Health through their MBRS (GM 08216) and INBRE (NC P20 RR-016470) grants is greatly appreciated. We express our special gratitude to the NSF-MRI (01–07) and NIH-MBRS programs to have made possible the acquisition of a 400 MHz NMR spectrometer.

References

(a) De Luca A, Natuzzi F, Desaphy J-F, Loni G, Lentini G, Franchini C, Tortorella V, Conte Camerino D. Mol Pharmacol 2000;57:268–277. [PubMed: 10648636] (b) De Luca A, Natuzzi F, Duranti A, Lentini G, Franchini C, Tortorella V, Conte Camerino D. Naunyn-Schmiedeberg's Arch Pharmacol 1997;356:777–787. (c) De Luca A, Pierno S, Natuzzi F, Franchini C, Duranti A, Lentini G, Tortorella V, Jockush H, Conte Camerino D. J Pharmacol Exp Ther 1997;282:93–100. [PubMed: 9223544] (d) Fenster PE, Comes KA. Pharmacotherapy 1986;6:1–9. [PubMed: 3513138]

- (a) Rüdel R, Lehmann-Horn F. Physiol Rev 1985;65:310–356. [PubMed: 2580324] (b) Cannon SC. Trends Neurosci 1996;19:3–10. [PubMed: 8787138] (c) Lehmann-Horn F, Rüdel R. Rev Physiol Biochem Pharmacol 1996;128:195–268. [PubMed: 8791722] (d) Jackson CE, Bason RJ, Ptacek LJ. Muscle Nerve 1994;17:763–768. [PubMed: 8008003]
- 3. Turgeon J, Uprichard ACG, Belanger PM, Harron DWC, Grech-Belanger OJ. Pharm Pharmacol 1991;43:630–635.
- 4. Hill RJ, Duff HJ, Sheldon RS. Mol Pharmacol 1988;34:659–663. [PubMed: 2848186]
- DeLuca A, Natuzzi F, Lentini G, Franchini C, Tortorella V, Conte Camerino D. Naunyn-Schmiedeberg's Arch Pharmacol 1995;352:653–661.
- Franchini C, Carocci A, Catalano AA, Cavalluzzi MM, Corbo F, Lentini G, Scilimati A, Scilimati A, Tortorella P, Camerino DC, Luca A. J Med Chem 2003;46:5238–5248. [PubMed: 14613326]
- Liu H, Gao ZB, Yao Z, Zheng S, Li Y, Zhu W, Tan X, Luo X, Shen J, Chen K, Hu GY, Jiang H. J Med Chem 2007;50:83–93. [PubMed: 17201412]
- (a) Turgeon J, Uprichard ACG, Belanger PM, Harron DWG, Grech-Belanger O. J Pharm Pharmacol 1991;43:630–635. [PubMed: 1685521] (b) Franchini C, Cellucci C, Corbo F, Lentini G, Scilimati A, Tortorella V, Stasi F. Chirality 1994;6:590–595.
- 9. Phillips, GT.; Shears, JH. U.K. Patent GB 2246774 A1. 1992.
- 10. Loughhead DG, Flippin LA, Weikert RJ. J Org Chem 1999;64:3373–3375. [PubMed: 11674449]
- 11. (a) Chem J, Shum W. Tetrahedron Lett 1995;36:2379–2382. (b) Fan RH, Hou XL. J Org Chem 2003;68:726–730. [PubMed: 12558391]
- 12. (a) Allen AD, Kanagasabapathy VM, Tidewell TT. J Am Chem Soc 1985;107:4513–4519. (b) Richard JP, Jencks WP. J Am Chem Soc 1984;106:1383–1396. (c) Crossland RK, Servis KL. J Org Chem 1970;35:3195–3196. (d) Uenishi J, Takagi T, Ueno T, Hiraoka T, Yonemitsu O, Tsukube H. Synlett 1999:41–44.
- 13. Houben-Weyl. Methods of Organic Chemistry, Workbench. In: Helmchen, G.; Hoffmann, RW.; Mulzer, J.; Schaumann, E., editors. Stereoselective Synthesis. Vol. E 21. Vol. 7. Georg Thieme Verlag; S New York: 1996. p. 4224-4231. (b) Glushkov VA, Tolstikov AG. Russ Chem Rev 2004;73:581–608. (c) Itsuno S, Sakurai Y, Shimizu K, Ito K. J Chem Soc, Perkin Trans 1 1990:1859– 1863. (d) Bolm C, Felder M. Synlett 1994:655-666. (e) Cho BT, Ryu MH. Bull Korean Chem Soc 1994;15:191–192. (f) Lantos I, Flisak J, Liu L, Matsunoka R, Mendelson W, Stevenson D, Tubman K, Tucker L, Zhang WY, Adams J, Sorenson M, Garigipati R, Erhardt K, Ross S. J Org Chem 1997;62:5358-5391. (g) Inoue T, Sato D, Komura K, Itsuno S. Tetrahedron Lett 1999;40:5379-5382. (h) Itsuno S, Matsumoto T, Sato D, Inoue T. J Org Chem 2000;65:5879–5881. [PubMed: 10970344] (i) Sailes HE, Watts JP, Whiting A. J Chem Soc, Perkin Trans 1 2000:3362–3374. (j) Fontaine E, Namane C, Menevrol J, Geslin M, Serva L, Roussev E, Tissandie S, Maftouh M, Roger P. Tetrahedron: Asymmetry 2001;12:2185–2189. (k) Krzeminski MP, Zaidlewicz M. Tetrahedron: Asymmetry 2003;14:1463–1466. (1) Sakito Y, Yoneyoshi Y, Suzukamo G. Tetrahedron Lett 1988;29:223-224. (m) Itsuno S, Nakano M, Miyazaki K, Masuda H, Ito K. J Chem Soc, Perkin Trans 1 1985:2039–2044. (n) Itsuno S, Sakurai Y, Ito K, Hirao A, Nakahama S. Bull Chem Soc Jpn 1987;60:395–396. (o) Chu YB, Shan ZX, Liu DJ, Sun NN. J Org Chem 2006;71:3998–4001. [PubMed: 16674084] (p) Huang XG, Ortiz-Marciales M, Huang K, Stepanenko V, Merced FG, Ayala AM, Correa W, De-Jesus M. Org Lett 2007;9:1793-1795. [PubMed: 17397177] (q) Huang K, Ortiz-Marciales M, Merced FG, Meléndez HJ, Correa W, De Jesús M. J Org Chem 2008;73:4017–4026. [PubMed: 18447392]
- 14. Benjahad A, Croisy M, Monneret C, Bisagni E, Mabire D, Coupa S, Poncelet A, Csoka I, Guillemont J, Meyer C, Andries K, Pauwels R, Bethune MP, Himmel DM, Das K, Arnold E, Nguyen CH, David S, Grierson DS. J Med Chem 2005;48:1948–1964. [PubMed: 15771439]

- 15. Lloyd GK, Williams M. J Pharmacol Exp Ther 2000;292:461-467. [PubMed: 10640281]
- 16. Karlin A. Nat Rev Neurosci 2002;3:102–114. [PubMed: 11836518]
- 17. Carroll FI, Lee JR, Navarro HA, Ma W, Brieaddy LE, Abraham P, Damaj MI, Martin BR. J Med Chem 2002;45:4755–4761. [PubMed: 12361403]
- 18. (a) Abreo MA, Lin NH, Garvey DS, Gunn DE, Hettinger AM, Wasicak JT, Pavlik PA, Martin YC, Donnelly-Roberts DL, Anderson DJ, Sullivan JP, Williams M, Arneric SP, Holladay MW. J Med Chem 1996;39:817–825. [PubMed: 8632405] (b) Sullivan JP, Donnelly-Roberts D, Briggs CA, Anderson DJ, Gopalakrishnan M, Piattoni-Kaplan M, Campbell JE, McKenna DG, Molinari E, Hettinger AM, Garvey DS, Wasicak JT, Holladay MW, Williams M, Arneric SP. Neuropharmacology 1996;35:725–734. [PubMed: 8887981]
- (a) Bannon AW, Decker MW, Holladay MW, Curzon P, Donnelly-Roberts D, Puttfarcken PS, Bitner RS, Diaz A, Dickenson AH, Porsolt RD, Williams M, Arneric SP. Science 1998;279:77–81.
 [PubMed: 9417028] (b) Holladay MW, Wasicak JT, He Y, Ryther KB, Bannon AW, Buckley MJ, Kim DB, Decker MW, Anderson DJ, Campbell JE, Kuntzweiler TA, Donnelly-Roberts DL, Piattoni-Kaplan M, Briggs CA, Williams M, Arneric SP. J Med Chem 1998;41:407–412. [PubMed: 9484491] (c) Abreo MA, Lin NH, Garvey DS, Gunn DE, Hettinger AM, Wasicak JT, Pavlik PA, Martin YC, Donnelly-Roberts DL, Anderson DJ, Sullivan JP, Williams M, Arneric SP, Holladay MW. J Med Chem 1996;39:817–825. [PubMed: 8632405]
- 20. Tonder JE, Olesen PH. Curr Med Chem 2001;8:651-674. [PubMed: 11281847]
- Vernier JM, El-Abdellaoui H, Holsenback H, Cosford NDP, Bleicher L, Barker G, Bontempi B, Chavez-Noriega L, Menzaghi F, Rao TS, Reid R, Sacaan AI, Suto C, Washburn M, Lloyd GK, McDonald IA. J Med Chem 1999;42:1684–1686. [PubMed: 10346920]
- 22. Banks, RE.; Smart, BE.; Tatlow, JC. Organofluorine Chemistry, Principle, and Commercial Application. Plenum Press; New York: 1994. Hiyama, T. Organofluorine Compounds, Chemistry and Application. Springer; Berlin, Germany: 2000. Filler, R.; Kobayashi, Y.; Yagupolskii, LM. Organofluorine Chemistry in Medicinal Chemistry and Biomedical Applications. Elsevier; Amsterdam, The Netherlands: 1993. Ojima, I.; McCarthy, JR.; Welch, JT. Biomedical Frontiers of Fluorine Chemistry. American Chemistry Society; Washington, DC: 1996.
- 23. (a) Stepanenko V, Ortiz-Marciales M, De Jesús M, Correa W, Vázquez C, Ortiz L, Guzmán I, De la Cruz W. Tetrahedron: Asymmetry 2007;18:2738–2745. (b) Alexakis A, Frutos JC, Mutti S, Mangeney P. J Org Chem 1992;57:1224–1237. (c) Alexakis A, Frutos JC, Mutti S, Mangeney P. J Org Chem 1994;59:3326–3334.

Figure 1. Mexiletine and examples of its more potent analogues.

Figure 2. Spiroborate esters derived from nonracemic aminoalcohols and ethylene glycol.

Figure 3. Neuronal nicotinic acetylcholine receptors agonist.

Scheme 1. Synthetic Route to 12a

Table 1 Screening of Different Catalysts for the Reduction of 11a

	_p (%) ee	70(R)	57(R)	97(R)	99(R)	81(S)	82(S)
NH ₂	yield (%)	87	40	84	92	08	76
	cat. (equiv)	0.10	0.10	0.10	0.25	0.10	0.10
Cat BH ₃ ·THF Dioxane	temp (°C)	0	22	0	0	22	22
N OBn 11a	cat.	4	S	9	9	7	80
	entry	1	2	3	4	5	9

 $^{\it a}$ Determined by chiral HPLC (Chiralcel OD-H column) of acetyl derivatives.

 $[\alpha]^{20}_{D}$ 12 entry NH_2 -28(1.2 1 2 3 -19(1.9 entry

Huang et al. Page 12

12

 $[a]^{20}_{D}$

-47(1.5 4 d -36(1.2 5 -17(1.0 6 7 -43(1.2 -47(2.4 8

 $[\alpha]^{20}_{D}$ 12 entry 9 -38(1.2)-36(1.0 10 11 -25(1.2

^aCHCl3 as solvent.

 $[^]b\mathrm{Determined}$ by amine on chiral HPLC (Chiralcel OD-H column).

 $^{^{}c}$ Determined by acetyl derivative on Chiralcel OD-H column.

 $d_{\mbox{\scriptsize Determined}}$ by GC of a cetyl derivative on Chiral column (CP-Chirasil-DexCB).

entry	12
1	NH ₂ S
2	F NH ₂ S O
3	F NH₂
	S_OBn
	F
4	NH ₂
5	o NH ₂ O N CI p
	ν

^aCHCl3 as solvent.

 $[^]b{\rm Determined}$ by chiral HPLC (Chiralcel OD-H column).

 $^{^{}c}$ Determined by 31 P NMR analysis of the phosphonate derivative. 23