

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/230821179>

# Conformational Distribution of Gas-phase Glycerol

ARTICLE in THE JOURNAL OF PHYSICAL CHEMISTRY A · NOVEMBER 2000

Impact Factor: 2.69 · DOI: 10.1021/jp002677e

CITATIONS

30

READS

20

6 AUTHORS, INCLUDING:



**Riccardo Chelli**

University of Florence

89 PUBLICATIONS 1,779 CITATIONS

SEE PROFILE



**Francesco Luigi Gervasio**

University College London

94 PUBLICATIONS 3,121 CITATIONS

SEE PROFILE



**Piero Procacci**

University of Florence

117 PUBLICATIONS 2,103 CITATIONS

SEE PROFILE



**Gianni Cardini**

University of Florence

171 PUBLICATIONS 2,373 CITATIONS

SEE PROFILE

# Conformational Distribution of Gas-phase Glycerol

Riccardo Chelli, Francesco L. Gervasio, Cristina Gellini, Piero Procacci,\* Gianni Cardini, and Vincenzo Schettino

Università di Firenze, Dipartimento di Chimica, Via Gino Capponi 9, 50121 Firenze, Italy,  
and European Laboratory for Nonlinear Spectroscopy (LENs), Largo E. Fermi 2, 50125 Florence, Italy

Received: July 26, 2000; In Final Form: September 20, 2000

Combining ab initio and statistical mechanics calculations we have determined the conformational distribution of gas-phase glycerol at different temperatures. The obtained results are consistent with infrared spectroscopy and electron diffraction measurements and are in excellent agreement with previous molecular dynamics simulation data.

In a recent paper<sup>1</sup> we have calculated the infrared absorption of various conformers of glycerol using density functional theory. These conformers are shown in Figure 1. The ab initio data were used in a fitting procedure of the experimental infrared spectrum of gas-phase glycerol.<sup>2</sup> The results indicated that, at 498 K, glycerol is present as a mixture of two conformers, namely  $\alpha\alpha$  and  $\alpha\gamma$ . Unfortunately the fit was not able to give quantitative results because of the large computational error (see discussion in ref 1). However, our conclusions were qualitatively similar to those obtained with electron diffraction measurements<sup>3</sup> and with molecular dynamics simulations<sup>4</sup> using empirical potential models.

At variance with these conclusions, the best fit to the supersonic jet rotational spectrum of a gas sample at 423 K was obtained assuming a distribution of  $\gamma\gamma$  and  $\alpha\gamma$  conformers.<sup>5</sup> Although the independent experimental measurements of the vibrational infrared and rotational microwave spectra refer to comparable thermodynamic conditions, they give contradictory indications on the conformational distribution that cannot be explained on the basis of the temperature difference. In view of this discrepancy, we thought it useful to search for an additional independent evidence for the conformational composition of gas-phase glycerol.

In this paper we report on a quantitatively reliable estimate of the conformational distribution of glycerol in gas phase, by computing the molecular partition function and the equilibrium constants using the results of accurate ab initio data.

In gas-phase glycerol, for any pair of conformational species  $I$  and  $J$ , the following conformational equilibrium holds



In the hypothesis of an ideal mixture, the equilibrium constant  $K_{IJ}$  can be determined from the knowledge of the molecular partition functions<sup>6</sup>

$$\frac{N_J}{N_I} = \frac{d_J q_J}{d_I q_I} = K_{IJ} \quad (2)$$

where  $N_J$  and  $N_I$  are the molar concentrations. The  $d_I$  and  $d_J$  factors correspond to the structural degeneracy of the conform-

ers, namely to the number of different conformational enantiomers. For all of the considered conformers (see Figure 1)  $d = 2$ , except for  $\alpha\alpha 3$  ( $d = 1$ ) because of the presence of a symmetry plane. In the Born–Oppenheimer approximation and neglecting vibro-rotational coupling, the molecular partition function can be factorized into its translational, rotational, vibrational, electronic, and nuclear parts, i.e.,  $q = q_{\text{trans}} q_{\text{rot}} q_{\text{vib}} q_{\text{elec}} q_{\text{nucl}}$ . The translational and nuclear partition functions are identical for all of the species and therefore they are irrelevant for the equilibrium constant of eq 2. The rotational partition function for an asymmetric top such as any glycerol conformer  $I$  is given by<sup>6</sup>

$$q_{\text{rot}}^{(I)} = \pi^{1/2} \left( \frac{8\pi^2}{\beta h^2} \right)^{3/2} (I_A^{(I)} I_B^{(I)} I_C^{(I)})^{1/2} \quad (3)$$

where  $I_A^{(I)}$ ,  $I_B^{(I)}$ , and  $I_C^{(I)}$  are the principal moments of inertia,  $h$  is the Planck constant and  $\beta = 1/(k_B T)$ . The vibrational partition function is given by

$$q_{\text{vib}}^{(I)} = \prod_j \frac{\exp(-\beta h \nu_j^{(I)}/2)}{1 - \exp(-\beta h \nu_j^{(I)})} \quad (4)$$

where  $j$  goes over the 36 vibrational frequencies  $\nu_j^{(I)}$ . Finally the electronic partition function is given by

$$q_{\text{elec}}^{(I)} = \exp(-\beta E_g^{(I)}) \quad (5)$$

where  $E_g^{(I)}$  is the (non degenerate) ground-state electronic energy of the  $I$ th conformer. Using the factorization property of the molecular partition function, the equilibrium constant  $K_{IJ}$  (eq 2) can be written as

$$K_{IJ} = K_{\text{rot}} K_{\text{vib}} K_{\text{elec}} = \frac{q_{\text{rot}}^{(J)} q_{\text{vib}}^{(J)} q_{\text{elec}}^{(J)}}{q_{\text{rot}}^{(I)} q_{\text{vib}}^{(I)} q_{\text{elec}}^{(I)}} \quad (6)$$

Using the ab initio data relative to the various conformers (all the ab initio data were obtained by the Gaussian98 package<sup>7</sup>), i.e., inertia moments, vibrational frequencies, and ground-state electronic energies, the gas-phase partition functions (eqs 3–5) can be calculated. From these, via eq 6, the

\* To whom correspondence should be addressed. E-mail: procacci@chim.unifi.it




Figure 1. Structures of the glycerol conformers.

chemical equilibrium constants between the various species, and hence the conformational distribution, are determined.

At the B3-LYP/6-31G(d) level of theory,<sup>8–10</sup> inertia moments and vibrational frequencies within few percent from the experimental counterparts are generally obtained.<sup>11–16</sup> The size of the basis set, however, is not sufficient to obtain reliable ground state *relative* electronic energies.<sup>17</sup> To minimize the error on the *ab initio* energies, it has been suggested<sup>17</sup> to optimize the structure at the B3-LYP/6-31G(d) level and then to compute the ground-state electronic energy using a single-point calculation with a much larger basis set, such as 6-311+G(3df,2p). Using this procedure, the average errors on molecular properties, such as atomization energies or first ionization potentials, were found to be<sup>17</sup> comparable to or even smaller than those obtained with very expensive correlated methods such as MP2<sup>18</sup> with a large basis set.

The electronic energies of the conformers of Figure 1 computed with B3-LYP using the small 6-31G(d) and the large 6-311+G(3df,2p) basis sets are reported in Table 1. All structures were optimized with the small basis set. As already found in ours<sup>1</sup> and in other *ab initio* calculations,<sup>19</sup>  $\gamma\gamma 1$  turns out to be the most stable conformer. However, as it can be seen from the table, the 6-31G(d) basis set gives larger energy gaps between the  $\gamma\gamma 1$  and other conformers with respect to the 6-311+G(3df,2p) basis set. For instance, the energy difference

TABLE 1: Electronic Energies (in  $\text{kJ mol}^{-1}$ ) of the Glycerol Conformers Calculated at the B3-LYP/6-31G(d) and B3-LYP/6-311+G(3df,2p) Level of Theory<sup>a</sup>

	6-311+G(3df,2p)	6-31G(d)
$\alpha\alpha 1$	3.13	13.59
$\alpha\alpha 2$	2.10	14.73
$\alpha\alpha 3$	7.54	17.14
$\alpha\gamma 1$	1.87	8.30
$\alpha\gamma 2$	0.46	9.32
$\alpha\gamma 3$	5.67	12.85
$\alpha\gamma 4$	4.10	13.65
$\alpha\gamma 5$	5.77	14.92
$\gamma\gamma 2$	2.59	8.91
$\alpha\beta$	13.22	26.30
$\beta\beta$	11.74	23.03
$\beta\gamma$	11.19	20.40

<sup>a</sup>  $\gamma\gamma 1$  conformer is taken as reference for the energy values of all the other conformers.

$\gamma\gamma 1$ – $\alpha\alpha 2$  reduces from 14.7 to 2.1  $\text{kJ mol}^{-1}$  going from the 6-31G(d) to the 6-311+G(3df,2p) basis set. In Table 2 we report the ratios at 498 K between the various molecular partition functions (vibrational, rotational, and electronic) of the most stable  $\gamma\gamma 1$  conformer and those corresponding to the other conformers (the vibrational and rotational partition functions have been computed using the data obtained with the small basis set; the computed vibrational frequencies have been rescaled

**TABLE 2: Electronic ( $K_{\text{elec}} = q_{\text{elec}}^{(\gamma\gamma 1)}/q_{\text{elec}}^{(I)}$ ), Vibrational ( $K_{\text{vib}} = q_{\text{vib}}^{(\gamma\gamma 1)}/q_{\text{vib}}^{(I)}$ ) and Rotational ( $K_{\text{rot}} = q_{\text{rot}}^{(\gamma\gamma 1)}/q_{\text{rot}}^{(I)}$ ) Contributions to the Constant  $K_{I\gamma\gamma 1} = K_{\text{elec}}K_{\text{vib}}K_{\text{rot}}$  of the Conformational Equilibrium  $I \rightleftharpoons \gamma\gamma 1$  at 498 K**

I	$K_{\text{elec}}$	$K_{\text{vib}}$	$K_{\text{rot}}$	$K_{I\gamma\gamma 1}$
$\alpha\alpha 1$	2.13	0.19	0.88	0.35
$\alpha\alpha 2$	1.66	0.13	0.89	0.19
$\alpha\alpha 3$	6.18	0.23	0.86	1.22
$\alpha\gamma 1$	1.57	0.38	0.90	0.54
$\alpha\gamma 2$	1.12	0.27	0.90	0.27
$\alpha\gamma 3$	3.94	0.34	0.89	1.19
$\alpha\gamma 4$	2.69	0.22	0.89	0.54
$\alpha\gamma 5$	4.03	0.22	0.88	0.76
$\gamma\gamma 2$	1.87	0.38	0.98	0.69
$\alpha\beta$	24.37	0.11	0.80	2.13
$\beta\beta$	17.05	0.15	0.84	2.13
$\beta\gamma$	14.93	0.09	0.89	1.17

**TABLE 3: Conformational Distributions (%) of Glycerol Conformers at Various Temperatures<sup>a</sup>**

	298 K	423 K	498 K
$\gamma\gamma 1$	5.79	4.93	4.50
$\gamma\gamma 2$	6.30	6.59	6.52
$\alpha\alpha 1$	12.30	12.82	12.75
$\alpha\alpha 2$	27.29	24.53	23.22
$\alpha\alpha 3$	0.89	1.55	1.84
$\alpha\gamma 1$	9.25	8.78	8.40
$\alpha\gamma 2$	23.56	18.56	16.67
$\alpha\gamma 3$	2.29	3.37	3.78
$\alpha\gamma 4$	6.69	8.02	8.38
$\alpha\gamma 5$	3.69	5.27	5.88
$\alpha\beta$	0.44	1.41	2.11
$\beta\beta$	0.52	1.48	2.11
$\beta\gamma$	0.94	2.69	3.85
$\alpha\alpha$	40.49	38.90	37.81
$\alpha\gamma$	45.50	44.01	43.10
$\gamma\gamma$	12.09	11.52	11.02

<sup>a</sup>  $\gamma\gamma$ ,  $\alpha\gamma$  and  $\alpha\alpha$  are the total concentrations obtained by summing up the contributions of the identical backbone conformers.

by the factor 0.928 as proposed by Rauhut and Pulay<sup>13</sup>). The product of these ratios ( $K_{I\gamma\gamma 1}$  in Table 2) yields the concentration of the  $\gamma\gamma 1$  conformer relative to that of the  $I$ th conformer. According to ab initio calculations,  $\gamma\gamma 1$  is the most stable conformer, hence its presence in the gas phase should be favored with respect to all of the other conformers. However  $\gamma\gamma 1$  has a very “rigid” and “compact” structure with correspondingly small vibrational and rotational partition functions compared to more open and less rigid forms such as  $\alpha\alpha$  or  $\alpha\gamma$ . This behavior is confirmed by  $K_{\text{elec}}$ ,  $K_{\text{rot}}$ , and  $K_{\text{vib}}$  (see Table 2) which are systematically greater than one ( $K_{\text{elec}}$ ) and less than one ( $K_{\text{rot}}$  and  $K_{\text{vib}}$ ). The overall balance is in favor of the  $\alpha\alpha$  and  $\alpha\gamma$  forms (see relative  $K_{I\gamma\gamma 1}$  values in Table 2). The conformational distributions at the temperatures of the infrared<sup>1,2</sup> (498 K) and microwave<sup>5</sup> (423 K) experiments, and at room temperature, are shown in Table 3. The  $\alpha\alpha$  and  $\alpha\gamma$  backbone conformers have similar concentrations (40–45%) with contributions mainly due to  $\alpha\alpha 1$ ,  $\alpha\alpha 2$ , and  $\alpha\gamma 2$ . The remaining  $\alpha\alpha$  and  $\alpha\gamma$  conformers have too high electronic energy (see Table 1) for contributing significantly to the conformational distribution. The  $\gamma\gamma$  backbone conformer has a concentration of about 11–12% with contribu-

tions from both  $\gamma\gamma 1$  and  $\gamma\gamma 2$ . The  $\alpha\beta$ ,  $\beta\beta$  and  $\beta\gamma$  have concentrations of few units of percentage because of their large electronic energy (see Table 1).

The present results add further support to the conclusions of our previous studies of the infrared spectra<sup>1</sup> and those of the electron diffraction experiments.<sup>3</sup> It can also be remarked that the overall concentrations, as well as their small temperature dependence, are in excellent agreement with those reported in a previous molecular dynamics simulation of the gas-phase glycerol.<sup>4</sup> The results of the present work are still at variance with those of Maccaferri et al.,<sup>5</sup> confirming, as suggested in ref 1, that the supersonic jet expansion in their experiment did not produce an equilibrium distribution.

**Acknowledgment.** This work was supported by the Italian Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST), by the Consiglio Nazionale delle Ricerche (CNR) and by the European Union (contract No. ERBFMGECT950017). We are grateful to Raffaele Guru Della Valle for stimulating discussions.

## References and Notes

- (1) Chelli, R.; Gervasio, F. L.; Gellini, C.; Procacci, P.; Cardini, G.; Schettino, V. *J. Phys. Chem. A* **2000**, *104*, 5351.
- (2) NIST Mass Spec Data Center, S. E. Stein, director, “IR and Mass Spectra” in NIST Chemistry WebBook, NIST Standard Reference Database Number 69, Eds. W. G. Mallard and P. J. Linstrom, November 1998, National Institute of Standards and Technology, Gaithersburg MD, 20899 (<http://webbook.nist.gov>), 1998.
- (3) Bastiansen, O. *Acta Chem. Scand.* **1949**, *3*, 415.
- (4) Chelli, R.; Procacci, P.; Cardini, G.; Califano, S. *Phys. Chem. Chem. Phys.* **1999**, *1*, 879.
- (5) Maccaferri, G.; Caminati, W.; Favero, P. G. *J. Chem. Soc., Faraday Trans.* **1997**, *93*, 4115.
- (6) McQuarrie, D. A. *Statistical Mechanics*; HarperCollins, New York NY, 1976.
- (7) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Zakrzewski, V. G.; J. A. Montgomery, J.; Stratmann, R. E.; Burant, J. C.; Dapprich, S.; Millam, J. M.; Daniels, A. D.; Kudin, K. N.; Strain, M. C.; Farkas, O.; Tomasi, J.; Barone, V.; Cossi, M.; Cammi, R.; Mennucci, B.; Pomelli, C.; Adamo, C.; Clifford, S.; Ochterski, J.; Petersson, G. A.; Ayala, P. Y.; Cui, Q.; Morokuma, K.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Cioslowski, J.; Ortiz, J. V.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Gonzalez, C.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Andres, J. L.; Gonzalez, C.; Head-Gordon, M.; Replogle, E. S.; Pople, J. A. *Gaussian 98, Revision A.5 1998*, Gaussian, Inc., Pittsburgh, PA.
- (8) Becke, A. D. *Phys. Rev. A* **1988**, *33*, 3098.
- (9) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.
- (10) Hariharan, P. C.; Pople, J. A. *Theor. Chem. Acta* **1973**, *28*, 213.
- (11) Schettino, V.; Gervasio, F. L.; Cardini, G.; Salvi, P. R. *J. Chem. Phys.* **1999**, *110*, 3241.
- (12) Gervasio, F. L.; Cardini, G.; Salvi, P. R.; Schettino, V. *J. Phys. Chem. A* **1998**, *102*, 2131.
- (13) Rauhut, G.; Pulay, P. *J. Phys. Chem.* **1995**, *99*, 3093.
- (14) Scott, A. P.; Radom, L. *J. Phys. Chem.* **1996**, *100*, 16502.
- (15) Wong, M. W. *Chem. Phys. Lett.* **1996**, *256*, 391.
- (16) Gellini, C.; Salvi, P. R.; Vogel, E. *J. Phys. Chem. A* **2000**, *104*, 3110.
- (17) Foresman, J. B.; Frisch, A. *Exploring Chemistry with Electronic Structure Methods*; Gaussian Inc.: Pittsburgh, PA, 1996.
- (18) Möller, C.; Plesset, M. *Phys. Rev.* **1934**, *46*, 618.
- (19) Teppen, B. J.; Cao, M.; Frey, R. F.; van Alsenoy, C.; Miller, D. M.; Schäfer, L. *J. Mol. Struct. (THEOCHEM)* **1994**, *314*, 169.