

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231527667>

Synthesis and Chemistry of Titanacyclopentane and Titanacyclopropane Rings Supported by Aryloxi Ligand

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · SEPTEMBER 1997

Impact Factor: 12.11 · DOI: 10.1021/ja970515z

CITATIONS

100

READS

31

6 AUTHORS, INCLUDING:



Phillip E. Fanwick

Purdue University

641 PUBLICATIONS 11,269 CITATIONS

SEE PROFILE

Synthesis and Chemistry of Titanacyclopentane and Titanacyclopropane Rings Supported by Aryloxy Ligation

Matthew G. Thorn, John E. Hill, Steven A. Waratuke, Eric S. Johnson, Phillip E. Fanwick, and Ian P. Rothwell*

Contribution from the Department of Chemistry, 1393 Brown Building, Purdue University, West Lafayette, Indiana 47907-1393

Received February 18, 1997[®]

Abstract: Treatment of the titanacyclopentadiene compound $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{C}_4\text{Et}_4)]$ (**3**) ($\text{OC}_6\text{H}_3\text{Ph}_2-2,6 = 2,6$ -diphenylphenoxide) with olefins leads to the formation of a variety of stable titanacyclopentane derivatives along with one equivalent of substituted 1,3-cyclohexadiene. The structural and spectroscopic properties of the ethylene product $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{CH}_2)_4]$ (**4**) show a ground state titanacyclopentane structure, but facile fragmentation on the NMR time scale to form a bis(ethylene) complex has been detected. The substituted products $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{C}_4\text{H}_6\text{R}_2)]$ ($\text{R} = \text{Me}$, **5**; Et , **6**; Ph , **7**) formed from α -olefins $\text{RCH}=\text{CH}_2$ exist as a mixture of regio- and stereoisomers in hydrocarbon solution. Analysis of a crystal obtained from solutions of **7** showed a *trans*-2,5-diphenyl-titanacyclopentane ring to be present in the solid state. Alternative routes to the titanacyclopentane compounds involve treatment of the dichlorides $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\text{Cl}_2]$ (**1**) or $[\text{Ti}(\text{OC}_6\text{HPh}_4-2,3,5,6)_2\text{Cl}_2]$ (**2**) with either sodium amalgam (2 Na per Ti) or 2 equiv of Bu^nLi in the presence of the substrate olefin. Using these conditions the titanacyclopentane compounds $[(\text{ArO})_2\text{Ti}\{\text{CH}_2\text{CH}(\text{C}_4\text{H}_8)\text{CHCH}_2\}]$ ($\text{ArO} = \text{OC}_6\text{H}_3\text{Ph}_2-2,6$, **10**; $\text{OC}_6\text{HPh}_4-2,3,5,6$, **11**) can be obtained by intramolecular coupling of 1,7-octadiene. The spectroscopic properties of **10** and **11** as well as a single-crystal X-ray diffraction analysis of **11** show an exclusive *trans* stereochemistry is present. Thermolysis of **10** or **11** in the presence of excess 1,7-octadiene leads to the catalytic formation of 2-(methylmethylene)cyclohexane (80%) along with *E,Z* isomers of 2,6-octadiene (20%). A kinetic study shows the reaction to be zero order in diene with activation parameters, $\Delta H^\ddagger = +18.7(5) \text{ kcal mol}^{-1}$ and $\Delta S^\ddagger = -26(5) \text{ eu}$. The diphenyltitanacyclopentane **7** will catalyze the dimerization of styrene to *trans*-1,3-diphenylbut-1-ene followed by isomerization to 1,3-diphenylbut-2-ene. This result shows that although a 2,5-diphenyl regiochemistry was observed in the solid state, styrene dimerization occurs via the 2,4-diphenyltitanacyclopentane intermediate. The facile fragmentation of these titanacyclopentane compounds accounts for the products observed in a number of reactions. Addition of phosphine donor ligands (*L*) leads to a series of titanacyclopropane compounds $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\eta^2\text{-CHR}=\text{CH}_2)(\text{L})]$ ($\text{R} = \text{H}$, **14**; Me , **15**; Et , **16**; Ph , **17**) along with 1 equiv of olefin. The solid-state structure of the ethylene complex **14** shows the C_2H_4 unit lies approximately coplanar with the $\text{Ti}-\text{PMe}_3$ bond. This structure is not only maintained in solution but slow olefin rotation is observed on the NMR time scale. In the case of the α -olefin products, two isomers are detected by ^1H , ^{13}C , and ^{31}P NMR spectroscopy. Addition of $\text{Ph}_2\text{C}=\text{O}$ or $\text{PhCH}=\text{NR}$ ($\text{R} = \text{Ph}$, CH_2Ph) to the titanacyclopentane and titanacyclopropane compounds leads to different products depending upon the reagent and reaction conditions. These can be classified as 2-oxa(aza)titanacycloheptanes, 2-oxa(aza)titanacyclopentanes, 2,5-dioxa(diaza)titanacyclopentanes, and examples of 2-oxatitanacyclopropane (η^2 -ketone) and 2,7-dioxatitanacycloheptane compounds. The 2-azatitanacyclopentane compounds $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{(\text{PhCH}_2)\text{NCH}(\text{Ph})\text{CH}_2\text{CH}_2\}]$ (**30**) and *trans*- $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{(\text{Ph})\text{NCH}(\text{Ph})\text{CH}_2\text{CH}(\text{Ph})\}]$ (**31**) react with alkynes to produce the corresponding 2-azatitanacyclopent-4-ene which hydrolyze to produce a stoichiometric equivalent of allylamine. Reaction of **30** with benzonitrile produces the 2,5-diazatitanacyclopent-2-ene $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{N}=\text{CPhCHPhNR})]$ (**35**) along with ethylene.

Introduction

The organometallic chemistry of the group 4 metals continues to be dominated by the use of cyclopentadiene ligation.¹ The metallocene dichlorides first discovered by Wilkinson et al.² represent an important starting material for a range of stoichiometric and catalytic transformations. A few significant developments have been the utilization of bulky cyclopentadiene ligation,³ the isolation and study of cationic alkyl derivatives,⁴ the chemistry of group 4 metal–ligand multiple bonds,^{5,6} and

the evolution of chiral metallocenes for carrying out asymmetric synthesis.⁷ The success of cyclopentadiene-based group 4 metal chemistry has led a number of research groups to attempt parallel chemistry utilizing “metallocene equivalents”.⁸ This approach hinges on the much analyzed σ^2, π^4 nature of the interaction of Cp-based ligation with transition metals.⁸ The use of isolobal

[®] Abstract published in *Advance ACS Abstracts*, September 1, 1997.

(1) *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: New York, 1995; Vol. 4.

(2) (a) Wilkinson, G.; Pauson, P. L.; Birmingham, J. L.; Cotton, F. A. *J. Am. Chem. Soc.* **1953**, *75*, 15. (b) Wilkinson, G.; Birmingham, J. L. *J. Am. Chem. Soc.* **1954**, *76*, 4281.

(3) (a) Bercaw, J. E.; Marvich, R. H.; Bell, L. G.; Brintzinger, H. H. *J. Am. Chem. Soc.* **1972**, *94*, 1219. (b) Threlkel, R. S.; Bercaw, J. E. *J. Organomet. Chem.* **1977**, *136*, 1.


(4) (a) Jordan, R. F. *Adv. Organomet. Chem.* **1991**, *32*, 325. (b) Bochmann, M. *J. Chem. Soc., Dalton Trans.* **1996**, 255. (c) Brintzinger, H. H.; Fischer, D.; Mulhaupt, R.; Rieger, B.; Waymouth, R. M. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1143. (d) Marks, T. J. *Acc. Chem. Res.* **1992**, *25*, 57. (e) Grubbs, R. H.; Coates, G. W. *Acc. Chem. Res.* **1996**, *29*, 85.

(5) Lee, S. Y.; Bergman, R. G. *J. Am. Chem. Soc.* **1996**, *118*, 6396 and references therein.

(6) Howard, W. A.; Parkin, G. *J. Am. Chem. Soc.* **1994**, *116*, 606 and references therein.

(7) (a) Willoughby, C. A.; Buchwald, S. L. *J. Am. Chem. Soc.* **1994**, *116*, 8952. (b) Willoughby, C. A.; Buchwald, S. L. *J. Am. Chem. Soc.* **1994**, *116*, 11703. (c) Verdaguer, X.; Lange, U. E. W.; Reding, M. T.; Buchwald, S. L. *J. Am. Chem. Soc.* **1996**, *118*, 6784. (d) Hicks, F. A.; Buchwald, S. L. *J. Am. Chem. Soc.* **1996**, *118*, 11688.

Scheme 1




ligands such as the formally dianionic alkylimido group⁹ leads to analogous chemistry based upon [Cp(RN)M] (M = group 5) and [(RN)₂M] (M = group 6) units.⁸

The alkoxide ligand, whose transition metal chemistry was originally explored by the Bradley and Chisholm groups is a formally monoanionic ligand that can also bond to the metal in a σ^2, π^4 fashion.¹⁰ This analogy was exploited by Wolczanski et al. in employing ligands such as tritox (Bu_3CO) and silox (Bu_3SiO) whose cylindrical shape and cone angle mimic that of the cyclopentadiene ligand.¹¹ Recently Sato et al. have utilized isopropoxide derivatives of titanium as important synthons in organic chemistry.¹² In our group we have focused on sterically demanding, *ortho*-substituted aryloxy ligation to support organometallic chemistry on early d-block metals.¹³ In this paper we report on the chemistry of titanacyclic rings supported by aryloxy ligation with emphasis on those derived from olefin substrates. A number of parallels between this aryloxy chemistry and known metallocene chemistry are analyzed. Some aspects of this work have been communicated.¹⁴

Results and Discussion

Synthesis and Characterization of Titanacyclopentane Compounds. The dichloride compounds $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\text{Cl}_2]$ (**1**)¹⁵ and $[\text{Ti}(\text{OC}_6\text{HPh}_4-2,3,5,6)_2\text{Cl}_2]$ (**2**) can be obtained in high yield by simple treatment of TiCl_4 with two equivalents of the parent phenol in hydrocarbon solvents (Scheme 1). Previously we have shown that sodium amalgam reduction of hydrocarbon solutions of **1** in the presence of 3-hexyne leads to the titanacyclopentadiene compound $[\text{Ti}(\text{OC}_6\text{H}_3\text{Pr}_2-2,6)_2(\text{C}_4\text{-Et}_4)]$ (**3**) in high yield (Scheme 1).¹⁶ Solutions of **3** in benzene, react rapidly with excess ethylene, propene, 1-butene, and

Scheme 2



styrene to produce solutions containing titanacyclopentane derivatives **4–7** along with 1 equiv of 1,3-cyclohexadiene (Scheme 2). The regio- and stereochemistry as well as the catalytic formation of the 1,3-cyclohexadiene products is discussed in detail in a subsequent paper.¹⁷ The titanacyclopentane compounds **4–6** can be readily isolated pure from these reactions mixtures. A superior synthesis of the diphenyl compound **7** involves sodium amalgam reduction of **1** in benzene solvent in the presence of styrene (Scheme 3). During the course of this reaction an intermediate, intense purple color indicated the formation of the d^1-d^1 dimer $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\mu\text{-Cl})_2]$ (**8**) which can be isolated pure by simply adding one equivalent of Na/Hg to benzene solutions of **1**. The solid-state structure of **8** has been reported and shows terminal phenoxide ligands and a short Ti-Ti distance of 2.9827(7).¹⁸ Attempts to utilize the 2,3,5,6-tetraphenylphenoxide compound **2** by sodium amalgam reduction in the presence of alkynes or olefins is hampered by the formation of the deep blue, sparingly soluble d^1-d^1 compound $[\text{Ti}(\text{OC}_6\text{HPh}_4-2,3,5,6)_2(\mu\text{-Cl})_2]$ (**9**) (Scheme 3). The low solubility of **9** hinders further reduction to form metallacyclic compounds.

Either by addition of 1,7-octadiene to **3**, or reduction of **1** in the presence of 1,7-octadiene, the titanabicyclic derivative **10** can be isolated. As an alternative to sodium amalgam reduction methods we have also explored the reaction of Bu^nLi with the dichlorides **1** and **2**. The addition of 2 equiv of Bu^nLi to Cp_2MCl_2 (the Negishi method)¹⁹ has been utilized to generate a “metallocene equivalent” in the presence of suitable organic substrates. We find that treatment of **1** or **2** with 2 equiv Bu^nLi

(8) Gibson, V. C. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 1565.

(9) Wigley, D. E. *Prog. Inorg. Chem.* **1994**, *42*, 239.

(10) (a) Bradley, D. C.; Mehrotra, R. C.; Gaur, D. P. *Metal Alkoxides*; Academic Press: New York, 1978. (b) Chisholm, M. H. *Chemtracts* **1992**, *4*, 273.

(11) Wolczanski, P. T. *Polyhedron* **1995**, *14*, 3335.

(12) (a) Nakagawa, T.; Kasatkin, A.; Sato, F. *Tetrahedron Lett.* **1995**, *36*, 3207. (b) Kasatkin, A.; Nakagawa, T.; Okamoto, S.; Sato, F. *J. Am. Chem. Soc.* **1995**, *117*, 3881. (c) Okamoto, S.; Kasatkin, A.; Zubaidha, P. K.; Sato, F. *J. Am. Chem. Soc.* **1996**, *118*, 2208. (d) Suzuki, K.; Urabe, H.; Sato, F. *J. Am. Chem. Soc.* **1996**, *118*, 8729. (e) Urabe, H.; Sato, F. *J. Org. Chem.* **1996**, *61*, 6756.

(13) (a) Rothwell, I. P. *J. Chem. Soc., Chem. Commun.*, in press. (b) Rothwell, I. P. *Acc. Chem. Res.* **1988**, *21*, 153.

(14) (a) Hill, J. E.; Fanwick, P. E.; Rothwell, I. P. *Organometallics* **1991**, *10*, 15. (b) Hill, J. E.; Fanwick, P. E.; Rothwell, I. P. *Organometallics* **1992**, *11*, 1771.

(15) Dilworth, J. R.; Hanich, J.; Krestel, M. J. *Organomet. Chem.* **1986**, *315*, 9.


(16) Hill, J. E.; Balaich, G. J.; Fanwick, P. E.; Rothwell, I. P. *Organometallics* **1993**, *12*, 2911.

(17) (a) Johnson, E. S.; Balaich, G. J.; Rothwell, I. P. *J. Am. Chem. Soc.*, in press. (b) Balaich, G. J.; Rothwell, I. P. *J. Am. Chem. Soc.* **1993**, *115*, 1581.

(18) Hill, J. E.; Fanwick, P. E.; Rothwell, I. P. *Polyhedron* **1990**, *9*, 1617.

(19) (a) Negishi, E.; Takahashi, T. *Acc. Chem. Res.* **1994**, *27*, 1277. (b) Negishi, E.; Cederbaum, F. E.; Takahashi, T. *Tetrahedron Lett.* **1986**, *27*, 2829. (c) Negishi, E.; Holmes, S. J.; Tour, J. M.; Miller, J. A.; Cederbaum, F. E.; Swanson, D. R.; Takahashi, T. *J. Am. Chem. Soc.* **1989**, *111*, 3336. (d) Kondakov, D.; Negishi, E. *J. Chem. Soc., Chem. Commun.* **1996**, 963.

Scheme 3



Li in benzene solvent will generate titanacyclopentadiene and titanacyclopentane compounds directly as detected by ^1H NMR spectroscopy (Scheme 3). In most cases, however, isolated yields were low due to difficulty in separating the products from the reaction mixture. One exception is the treatment of tetraphenylphenoxide **2** with Bu^nLi in the presence of 1,7-octadiene. Addition of only 1 equiv of Bu^nLi to benzene solutions of **2** lead to formation of deep blue **9**. When 2 equiv of Bu^nLi are added with rapid agitation, only a small amount of **9** is produced along with a benzene solution of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,3,5,6)_2\{\text{CH}_2\text{CH}(\text{C}_4\text{H}_8)\text{CHCH}_2\}]$ (**11**) (Scheme 3). Filtration, removal of solvent, and addition of toluene was found to lead to crystallization of **11** in moderate yield as a toluene solvate.

The solution NMR spectroscopic properties of the titanacyclopentane compounds **4**–**7**, **10**, and **11** are of interest.²⁰ In the ^{13}C NMR spectra, signals for the $\text{Ti}-\text{C}(\alpha)$ and $\text{Ti}-\text{C}(\beta)$ carbon atoms display $^1J(^{13}\text{C}-^1\text{H})$ coupling constants close to those expected for sp^3 -hybridized carbon atoms. Furthermore, the $\text{Ti}-\text{C}(\alpha)$ chemical shifts are in the range typical for titanium alkyl compounds containing aryloxy ligation. This data supports their proposed structure and is inconsistent with a bis(olefin) formulation for the compounds. There is, however, evidence in the ^1H NMR spectrum of titanacyclopentane **4** for facile, reversible fragmentation into a “bis(ethylene)” species.²¹ In the presence of 1 atm of ethylene, solutions of **4** in $\text{C}_6\text{D}_5\text{CD}_3$ are stable at temperatures up to 80 °C. At ambient temperatures the $\alpha\text{-CH}_2$ and $\beta\text{-CH}_2$ methylene protons appear as well resolved signals (Figure 1). The excess ethylene appears as a sharp singlet at a chemical shift almost identical for C_2H_4 simply dissolved in $\text{C}_6\text{D}_5\text{CD}_3$. Upon raising the temperature





Figure 1. ^1H NMR (200 MHz) spectra of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{CH}_2)_4]$ (**4**) and ethylene in $\text{C}_6\text{D}_5\text{CD}_3$ solvent (* indicates protio impurity) over the temperature range -30 to $+60$ °C.

Scheme 4



of the solution, broadening of the methylene proton signals occurs and at $+60$ °C collapse into the baseline occurs. The signal due to the free ethylene begins to broaden at $+40$ °C. From the coalescence temperature we estimate the activation energy (ΔG^\ddagger) for exchange of the α - and $\beta\text{-CH}_2$ methylene groups within the titanacyclopentane **4** to be $15.9(5)$ kcal mol^{-1} at 55 °C. Simulation of the spectra using a program that accommodates a three-site exchange shows that the rate of exchange of methylene groups within **4** (160 s^{-1} at 55 °C) exceeds the rate of exchange of these groups with free ethylene (40 s^{-1} at 55 °C). Accurate simulation of the spectra was not possible due to the variation of the ethylene intensity with temperature (probably reflecting changes in solubility) and the fact that the methylene signals are in reality unresolved multiplets.²² Furthermore, at temperatures above 60 °C **4** is converted into a new organometallic compound (*vide infra*). We interpret these gross changes as indicating that not only fragmentation of **4** into a “bis(ethylene)” species occurs (Scheme 4) but that exchange of free and coordinated ethylene also occurs. The most plausible pathway for the intermolecular

(20) For leading references to metallacyclopentane compounds, see: (a) Knight, K. S.; Wang, D.; Waymouth R. M.; Ziller, J. *J. Am. Chem. Soc.* **1994**, *116*, 1845. (b) Nugent W. A.; Taber, D. F. *J. Am. Chem. Soc.* **1989**, *111*, 6435. (c) Taber, D. F.; Louey, J. P.; Wang, Y.; Nugent, W. A.; Dixon, D. A.; Harlow, R. L. *J. Am. Chem. Soc.* **1994**, *116*, 9457.

(21) (a) Grubbs, R. H.; Miyashita, A. *J. Chem. Soc., Chem. Commun.* **1977**, 864. (b) Grubbs, R. H.; Miyashita, A. *J. Am. Chem. Soc.* **1978**, *100*, 1300.

(22) For a discussion of the problems of interpreting NMR exchange rates, see: Green, M. L. H.; Wong, L.-L.; Sella, A. *Organometallics* **1992**, *11*, 2660.

The titanacyclopentane compounds **4** and **7** and the titanabicyclic derivative **11** have been subjected to single-crystal X-ray diffraction analysis (Figures 2–4 and Tables 1–3). In all cases the formulation of these compounds was confirmed and in the case of **11**, the *trans* stereochemistry was also established




Figure 2. Molecular structure of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_{2-2,6})_2(\text{CH}_2)_4]$ (**4**).




Figure 3. Molecular structure of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_{2,6})_2\{\text{CH}(\text{Ph})\text{CH}_2\text{CH}_2\text{CH}(\text{Ph})\}]$ (**7**).




Figure 4. Molecular structure of $[\text{Ti}(\text{OC}_6\text{HPh}_4\text{-2,3,5,6})_2\{\text{CH}_2\text{CH}(\text{C}_4\text{H}_8)\text{CHCH}_2\}]$ (**10**).

(Figure 4). A number of attempts at determining the crystal structure of a sample of **7** were unsuccessful. Eventually a data set was obtained on a crystal which was solved showing the presence of the *trans*-2,5-diphenyl-substituted titanacyclopentane

(23) Cohen, S. A.; Auburn, P. R.; Bercaw, J. E. . *J. Am. Chem. Soc.* **1983**, *105*, 1136.

(24) Takahashi, T.; Fujimori, T.; Seki, T.; Saburi, M.; Uchida, Y.; Rousset, C. J.; Negishi, E. *J. Chem. Soc., Chem. Commun.* **1990**, 182.

(25) Warner, B. P.; Davis, W. M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1994**, *116*, 5471.

(26) (a) Schock, L. E.; Marks, T. J. *J. Am. Chem. Soc.* **1988**, *110*, 7701.
(b) Schmidt, J. R.; Duggan, D. M. *Inorg. Chem.* **1981**, *20*, 318.

(27) (a) Gell, K. I.; Schwartz, J. J. *J. Am. Chem. Soc.* **1981**, *103*, 2687.
(b) Akita, M.; Yasuda, H.; Yamamoto, H.; Nakamura, A. *Polyhedron* **1991**, *10*, 1.

(28) Swanson, D. R.; Rousset, C. J.; Negishi, E.; Takahashi, T.; Seki, T.; Saburi, M.; Uchida, Y. *J. Org. Chem.* **1989**, *54*, 3521.

Table 1. Selected Bond Distances (Å) and Angles (deg) for [Ti(OC₆H₃Ph₂-2,6)₂(C₄H₈)] (**4**)

Molecule 1			
Ti(1)–O(110)	1.799(1)	C(11)–C(12)	1.510(3)
Ti(1)–O(120)	1.804(1)	C(12)–C(13)	1.511(4)
Ti(1)–C(11)	2.087(2)	C(13)–C(14)	1.517(4)
Ti(1)–C(14)	2.084(2)		
O(110)–Ti(1)–O(120)	132.62(6)	Ti(1)–C(11)–C(12)	106.6(1)
O(110)–Ti(1)–C(11)	104.96(7)	Ti(1)–C(14)–C(13)	106.3(1)
O(110)–Ti(1)–C(14)	108.12(7)	C(11)–C(12)–C(13)	109.4(2)
C(11)–Ti(1)–C(14)	85.59(9)	C(12)–C(13)–C(14)	109.8(2)
Ti(1)–O(110)–C(111)	157.9(1)	Ti(1)–O(120)–C(121)	155.7(1)
Molecule 2			
Ti(2)–O(210)	1.802(1)	C(21)–C(22)	1.520(3)
Ti(2)–O(220)	1.817(1)	C(22)–C(23)	1.512(4)
Ti(2)–C(21)	2.084(2)	C(23)–C(24)	1.512(4)
Ti(2)–C(24)	2.084(2)		
O(210)–Ti(2)–O(220)	135.05(6)	Ti(2)–C(21)–C(22)	106.1(1)
O(210)–Ti(2)–C(21)	107.84(7)	Ti(2)–C(24)–C(23)	105.4(1)
O(210)–Ti(2)–C(24)	104.23(7)	C(21)–C(22)–C(23)	109.8(2)
C(21)–Ti(2)–C(24)	86.22(8)	C(22)–C(23)–C(24)	109.6(2)
Ti(2)–O(210)–C(211)	159.1(1)	Ti(2)–O(220)–C(221)	151.3(1)

Table 2. Selected Bond Distances (Å) and Angles (deg) for [Ti(OC₆H₃Ph₂-2,6)₂{CH(Ph)CH₂CH₂CH(Ph)}] (**7**)

Ti–O(5)	1.801(3)	Ti–O(6)	1.804(3)
T–C(1)	2.103(6)	Ti–C(4)	2.115(5)
C(1)–C(2)	1.537(7)	C(2)–C(3)	1.516(8)
C(3)–C(4)	1.546(8)		
O(5)–Ti–O(6)	126.1(1)	C(1)–Ti–C(4)	85.6(2)
Ti–C(1)–C(2)	97.7(4)	Ti–C(4)–C(3)	105.6(4)
C(1)–C(2)–C(3)	110.0(5)	C(2)–C(3)–C(4)	111.6(5)
Ti–O(5)–C(51)	157.4(3)	Ti–O(6)–C(61)	161.1(3)

Table 3. Selected Bond Distances (Å) and Angles (deg) for [Ti(OC₆HPh₄-2,3,5,6)₂{CH₂CH(C₄H₈)CHCH₂}] (**11**)

Ti–O(10)	1.802(3)	Ti–C(1)	2.061(5)
C(1)–C(21)	1.49(2)	C(21)–C(21)	1.65(3)
O(10)–Ti–O(10)	135.0(2)	O(10)–Ti–C(1)	106.6(2)
O(10)–Ti–C(1)'	105.2(2)	C(1)–Ti–C(1)	88.5(3)
Ti–O(10)–C(11)	149.4(3)	Ti–C(1)–C(21)	105.0(8)
C(1)–C(21)–C(21)	106.5(8)		



ring (Figure 3). In all three titanacyclopentane derivatives the Ti–O and Ti–C distances are within the ranges typical of aryloxo and alkyl groups bound to 4-coordinate Ti(IV).²⁹

The titanacyclopentane compounds **4–7** react with excess H₂O to produce a mixture of organic products along with 2 equiv of phenol (Scheme 5). For the unsubstituted derivative **4**, butane was detected along with significant amounts of ethylene and ethane (GC analysis and ¹H NMR). In the case of the diphenyl derivative **7** an almost quantitative conversion to a 50/50 mixture of styrene and ethylbenzene was observed. It, therefore, appears that water acts as a simple Lewis donor ligand as well as a protic source and induces the fragmentation of the titanacyclopentane prior to protonation of the bound olefin (Scheme 5). Hydrolysis of titanabicyclo **10** and **11** yields *trans*-1,2-dimethylcyclohexane.

During an attempt to obtain crystals of the dimethyltitanacyclopentane complex **5** by slow cooling of a hexane solution, the accidental introduction of trace amounts of moisture led to the formation of crystals of an oxo complex **12** (Scheme 5). The molecular structure of **12** (Figure 5, Table 4) shows that it has originated by reactions of 2 equiv of **5** with 1 equiv of water. A similar type of reaction occurs between metallocene dialkyls and traces of water.³⁰ The protonation of one of the Ti–C bonds

(29) Smith, G. D.; Fanwick, P. E.; Rothwell, I. P. *Inorg. Chem.* **1990**, 29, 3221.

(30) Hunter, W. E.; Nrcir, D. C.; Bynum, R. V.; Penttila, R. A.; Atwood, J. L. *Organometallics* **1983**, 2, 750.

**Figure 5.** Molecular structure of [{Ti(OC₆H₃Ph₂-2,6)₂(CH₂CHMeCHMe₂)}₂(μ-O)] (**12**).**Scheme 5****Table 4.** Selected Bond Distances (Å) and Angles (deg) for [{Ti(OC₆H₃Ph₂-2,6)₂(CH₂CHMeCHMe₂)}₂(μ-O)] (**12**)

Ti–O(1)	1.793(3)	Ti–O(2)	1.795(2)
Ti–O(B)	1.801(1)	Ti–C(1)	2.087(5)
O(1)–Ti–O(2)	119.4(1)	O(1)–Ti–O(B)	110.2(1)
O(1)–Ti–C(1)	100.0(2)	O(2)–Ti–O(B)	113.46(9)
O(2)–Ti–C(1)	107.2(2)	O(B)–Ti–C(1)	104.5(2)
Ti–O(B)–Ti	152.3(1)	Ti–O(1)–C(11)	160.7(2)
Ti–O(2)–C(21)	167.6(3)	Ti–C(1)–C(2)	129.7(5)

in **5** leads to an alkyl group which gives insight into the regiochemistry of the original titanacyclopentane ring. Clearly the alkyl group in **12** can only be formed from the 3,4-dimethyltitanacyclopentane ring, although the same group can be formed from either the *cis* or *trans* stereoisomers.

Thermal Stability of Titanacyclopentanes: Catalytic Cyclization of 1,7-Octadiene and Dimerization of Styrene. When a solution of titanacyclopentane **7** in C₆H₆ is heated at 100 °C in the presence of excess styrene, catalytic dimerization occurs to initially produce *trans*-1,3-diphenylbut-1-ene (Scheme 6).³¹ This product presumably arises via a 2,4-diphenyltitanacyclopentane intermediate by abstraction of a hydrogen from the 3-position of the ring. The *trans*-1,3-diphenylbut-1-ene is





Figure 6. Plot showing the appearance with time of 2-(methylmethylene)cyclohexane within a C_6D_6 solution containing $[Ti(OC_6H_3Ph_2-2,6)_2\{CH_2CH(C_4H_8)CHCH_2\}]$ (**10**, 0.0345 mmol) and 1,7-octadiene (0.977 mmol) heated at 118 °C.

Scheme 6



subsequently isomerized within the reaction mixture to produce 1,3-diphenylbut-2-ene. This result shows that although structural studies demonstrate the presence of the *trans*-2,5-diphenyl derivative in the solid state, thermal reactivity in solution proceeds via the 2,4-diphenyl isomer.

Titanacyclopentane **10** catalytically converts 1,7-octadiene into a mixture of 2-methyl-methylenecyclohexane^{31–34} (80% by GC) and 2,6-octadiene (predominantly E,E by ¹³C NMR,³⁵ Scheme 6). Previous work by Nakamura et al. has shown that low valent titanocene systems are excellent catalysts for the internalization of terminal olefins and nonconjugated dienes.³⁵ Recent work by Negishi et al. has shown a zirconocene isomerization of α,ω -dienes to produce metal bound 1,3-diene structures.³⁶ Attempts to use **10** as a catalyst for the cyclization of 1,6-heptadiene and 1,8-nonadiene led mainly to olefin isomerization. A kinetic study of the cyclization of 1,7-octadiene showed the reaction to be zero order in [diene]. The catalyst **10** was observed (NMR) to be the predominant Ti species in solution (as long as there is excess 1,7-octadiene present) and remains active for a significant number of turnovers (Figure 6, Table 5). Activation parameters were obtained for the formation of 2-(methylmethylene)cyclohexane over a 40° temperature range (Figure 7).

(31) Dimerization of olefins and cyclization of α,ω -dienes via metalla-cyclopentanes, see: (a) McLain, S. J.; Wood, C. D.; Schrock, R. R. *J. Am. Chem. Soc.* **1978**, *101*, 4558. (b) Smith, G.; McLain, S. J.; Schrock, R. R. *J. Organomet. Chem.* **1980**, *202*, 269.

(32) Linear dimerization of dienes, see: Christoffers, J.; Bergman, R. G. *J. Am. Chem. Soc.* **1996**, *118*, 6422 and references therein.

(33) For an alternative selective cyclization of α,ω -dienes catalyzed by scandium hydrides, see: Piers, W. E.; Shapiro, P. J.; Bunel, E. E.; Bercaw, J. E. *Synlett* **1990**, 74.

(34) For references to the ring closing metathesis of α,ω -dienes, see: Zuercher, W. J.; Hashimoto, M.; Grubbs, R. H. *J. Am. Chem. Soc.* **1996**, *118*, 6634 and references therein.

(35) Akita, M.; Yasuda, H.; Nagasuna, K.; Nakamura, A. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 554.

(36) Negishi, E.; Maye, J. P.; Choueiry, D. *Tetrahedron* **1995**, *51*, 4447.

Table 5. First-Order Rate Constants for the Cyclization of 1,7-Octadiene Catalyzed by **10**

<i>T</i> , °C	$10^3 k$, s ⁻¹	equiv of Ti ⁻¹ h ⁻¹
94	0.102	0.366
108	0.217	0.78
118	0.487	1.75
134	1.28	4.60





Figure 7. Plot of $\ln(k/T)$ vs $1/T$ for the cyclization of 1,7-octadiene into 2-(methylmethylene)cyclohexane (data in Table 5).


Scheme 7



A number of distinct mechanistic pathways for these olefin dimerization/cyclization reactions can be envisaged proceeding via the titanacyclopentane intermediate. The two most reasonable differ as to whether the β -hydrogen abstraction occurs initially by the metal center to generate an intermediate hydride or in a concerted fashion by the adjacent α -carbon center (Scheme 7). This latter process corresponds to an intramolecular σ -bond metathesis reaction, a pathway that is common for ligand cyclometallation at d^0 metal centers.³⁷ These reactions are directly related to the breakdown of compounds such as $[Cp_2M-$

(37) Rothwell, I. P. In *Selective Hydrocarbon Activation*; Davis, J. A., Watson, P. L., Liebman, J. F., Greenberg, A., Eds; VCH Publishers: New York, 1990; pp 43–75.

Scheme 8




(X)(R)] ($M = \text{Zr, Hf}$) in which the X group contains at least one β -CH bond. Mechanistic studies by Buchwald et al. on the formation of thioaldehyde complexes led to the conclusion that the reactions proceed via a concerted pathway (Scheme 7).³⁸ Particularly informative were a large negative entropy of activation and a primary deuterium kinetic isotope effect. A substituent effect study yielded a Hammett plot with $\rho = +0.39$, indicating positive charge build up on the hydrogen being transferred in the transition state.³⁸

The kinetic study of the cyclization of 1,7-octadiene yielded a value of $\Delta S^\ddagger = -26(5)$ eu (Figure 7), consistent with the highly ordered transition state required in the concerted pathway. However, when an equimolar mixture of $\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$ and $\text{C}_6\text{D}_5\text{CD}=\text{CD}_2$ was heated with titanacyclopentane **7** at 100 °C in C_6D_6 , scrambling of both α - and β -olefinic H/D atoms was observed (^1H and ^{13}C NMR) at a rate much faster than dimerization. The catalyst for this exchange process is presumably a titanium hydride species which undergoes reversible insertion of styrene monomers to produce both α - and β -phenethyl groups. It is possible that solutions of **7** contain such a hydride species as a contaminant. An alternative explanation is that the scrambling is catalyzed by the hydride generated by β -hydrogen abstraction by the metal center. If this was indeed the case it would imply that this species was sufficiently long-lived to carry out multiple insertions with free styrene.

The thermal instability of **4** is also mechanistically significant. When solutions of **4** and excess ethylene in C_6D_6 are heated at 80 °C, almost quantitative conversion to the titanacycloheptene complex **13** occurs (Scheme 8). This complex has been previously isolated by addition of 1,3-butadiene to **4**.³⁹ Small amounts of 1-butene and ethane were also present in solution (^1H NMR). The thermal decomposition of metallacyclopentanes of the group 4 metals has been studied by many researchers dating back to the pioneering studies of Whitesides et al.⁴⁰ This reactivity parallels exactly that reported by Erker et al. for the zirconacyclopentane $[\text{Cp}_2\text{Zr}(\text{CH}_2)_4]$, where ethylene dimerization was curtailed by formation of a zirconacycloheptene complex.⁴¹ The ethylene complex $[\text{Cp}^*\text{Ti}(\text{C}_2\text{H}_4)]$ was demonstrated by Cohen and Bercaw to be a catalyst for the conversion of ethylene

Scheme 9



to 1,3-butadiene and ethane.²³ These reactions are believed to proceed via intermediates formed by β -hydrogen abstraction by the metal center. The resulting 3-butenyl hydride can then insert an extra equivalent of ethylene and eliminate ethane with formation of a 1,3-butadiene complex which in the case of **4** would produce **13** (Scheme 8). An alternative pathway proposed by Erker et al. proceeds via a titanium dihydride intermediate.⁴¹

The above observations, therefore, imply that the catalytic dimerization of styrene by **7** and cyclization of 1,7-octadiene by **10** proceed via metal hydride intermediates and not by way of a concerted pathway. The zero-order dependence on olefin is consistent with either rate-determining β -hydrogen abstraction and relatively fast subsequent steps or alternatively reversible β -hydrogen with a rate determining elimination step. This latter scenario could not, however, involve olefin in the rate-determining step. A pathway which is consistent with these mechanistic findings and avoids the intermediacy of a highly unsaturated species $[\text{Ti}(\text{OAr})_2]$ is shown for ethylene in Scheme 8. Rate-determining β -hydrogen abstraction is followed by insertion of an equivalent of olefin to generate a dialkyl intermediate. In the case of ethylene, ethane elimination eventually leads to **13**, whereas in the case of styrene and 1,7-octadiene elimination of product generates a monoolefin species which can rapidly regenerate a titanacyclopentane (Scheme 8).

Synthesis and Characterization of Titanacyclopentane Compounds. The titanacyclopentane compounds **4–7** react rapidly in C_6D_6 solution with either PMe_3 or PMe_2Ph to produce deep-purple solutions containing 1 equiv of olefin and new organometallic compounds. These new compounds are formulated as titanacyclopentane (monoolefin) species containing a single phosphine ligand (Scheme 9). In the case of the propene-, butene- and styrene-derived complexes, a mixture of two isomers is observed in solution by ^1H , ^{13}C , and ^{31}P NMR spectroscopy. The presence of two, nonequivalent Ti–C resonances in the ^{13}C NMR spectrum of the ethylene complex (**14**) indicates a structure for the molecule in which the C–C unit is coplanar, as opposed to perpendicular, to the Ti–P bond. This situation was confirmed by a single-crystal X-ray diffraction analysis of **14a** (Figure 8, Table 6). The two isomers present for **15–17** are due to the olefin substituents being positioned either proximal or distal to the phosphine ligand. This reactivity, the structures of compounds **15–17** as well as the

(38) Buchwald, S. L.; Nielsen, R. B. *J. Am. Chem. Soc.* **1988**, *110*, 3175.

(39) Balaich, G. J.; Hill, J. E.; Waratuke, S. A.; Fanwick, P. E.; Rothwell, I. P. *Organometallics* **1995**, *14*, 656.

(40) McDermott, J. X.; Wilson, M. E.; Whitesides, G. M. *J. Am. Chem. Soc.* **1976**, *98*, 6529.

(41) Erker, G.; Engel, K.; Dorf, U.; Atwood, J. L.; Hunter, W. E. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 914.




Figure 8. Molecular structure of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2\text{-2,6})_2(\text{CH}_2\text{CH}_2)(\text{PMe}_3)]$ (**14a**).

Table 6. Selected Bond Distances (Å) and Angles (deg) for $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2\text{-2,6})_2(\eta^2\text{-CH}_2\text{CH}_2)(\text{PMe}_3)]$ (**14a**)

Ti–P(1)	2.5633(5)	Ti–O(2)	1.835(1)
Ti–O(3)	1.838(1)	Ti–C(41)	2.110(2)
Ti–C(42)	2.148(2)	C(41)–C(42)	1.425(3)
P(1)–Ti–O(2)	93.39(4)	P(1)–Ti–O(3)	98.39(4)
P(1)–Ti–C(41)	122.67(5)	P(1)–Ti–C(42)	83.97(5)
O(2)–Ti–O(3)	140.73(5)	O(2)–Ti–C(41)	100.76(7)
O(2)–Ti–C(42)	100.76(7)	O(3)–Ti–C(41)	103.86(7)
O(3)–Ti–C(42)	107.33(7)	C(41)–Ti–C(42)	39.09(7)
Ti–O(2)–C(21)	160.9(1)	Ti–O(3)–C(31)	161.3(1)

Table 7. Structural Parameters for Selected Group 4 Metallocyclopropane Compounds

compound	M–C, (Å)	C–C, (Å)	ref
$\text{H}_2\text{C}=\text{CH}_2$		1.337(2)	
$[\text{Cp}^*_2\text{Ti}(\text{CH}_2\text{CH}_2)]$	2.160(4)	1.438(5)	22
$[(\text{ArO})_2\text{Ti}(\text{CH}_2\text{CH}_2)(\text{PMe}_3)]$ (14a)	2.110(2)	1.425(3)	a
	2.148(2)		
$[\text{Cp}_2\text{Zr}(\text{CH}_2\text{CH}_2)(\text{PMe}_3)]$	2.354(3)	1.449(6)	42
	2.332(4)		
$[\text{Cp}_2\text{Zr}(\text{CH}_2\text{CHPh})(\text{PMe}_3)]$	2.35(1)	1.46(2)	42
	2.35(2)		
$[\text{Cp}_2\text{Hf}(\text{CH}_2\text{CMe}_2)(\text{PMe}_3)]$	2.316(8)	1.46(10)	43b
	2.368(9)		
$\{(\text{Et}_3\text{P})_2\text{Cl}_3\text{Zr}\}(\text{CH}_2\text{CH}_2)\{(\text{Et}_3\text{P})_2\text{Cl}_3\text{Zr}\}$	2.42(2)	1.69(3)	44
	2.44(2)		

^a This work.

lack of facile olefin rotation is similar to that reported for group 4 analogous containing cyclopentadiene ligation.^{28,42–44}

The most interesting feature of the solid-state structure of **14** pertains to the structural parameters of the titanacyclopentane ring. In Table 7 are collected selected parameters of related molecules for comparison.


It can be seen that although there is a pronounced elongation of the ethylene C–C bond upon coordination to the titanium

(42) (a) Takahashi, T.; Murakami, M.; Kunishige, M.; Saburi, M.; Uchida, Y.; Kozawa, K.; Uchida, T.; Swanson, D. R.; Negishi, E. *Chem. Lett.* **1989**, 761.

(43) Binger, P.; Muller, P.; Benn, R.; Rufinska, A.; Gabor, B.; Kruger, C.; Betz, P. *Chem. Ber.* **1989**, 122, 1035.

(44) (a) Fisher, R. A.; Buchwald, S. L. *Organometallics* **1990**, 9, 871. (b) Buchwald, S. L.; Kreutzer, K. A.; Fisher, R. A. *J. Am. Chem. Soc.* **1990**, 112, 4600.

Scheme 10



center in **14a**, the lengthening is slightly less than found in corresponding mononuclear metallocene derivatives of Ti, Zr, and Hf (Table 7). The bonding of the ethylene moiety in the species $[\text{Cl}_2\text{Ti}(\text{CH}_2\text{CH}_2)]$ has been theoretically analyzed.⁴⁶

Synthesis and Characterization of Oxa- and Azatitanacyclic Compounds. We have probed the reactivity of the isolated titanacyclopentane and titanacyclopentane species toward ketones and imine reagents. Initially focusing upon the reaction with benzophenone, $\text{Ph}_2\text{C}=\text{O}$, by varying the reaction conditions we have been able to isolate and characterize oxatitanacyclopentane, -pentane, and -heptane as well as dioxatitanacyclopentane and -heptane species. The titanacyclopentane **4** will react with $\text{Ph}_2\text{C}=\text{O}$ in the presence of excess ethylene to produce the oxatitanacycloheptane complex **18**. In the absence of excess ethylene in solution, **4** produces a mixture of not only **18** but also oxatitanacyclopentane **19** and dioxatitanacycloheptane **20** (Scheme 10). These latter two compounds are also synthesized by addition of 1 or 2 equiv of $\text{Ph}_2\text{C}=\text{O}$ to $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2\text{-2,6})_2(\eta^2\text{-CH}_2\text{CH}_2)(\text{PMe}_3)]$ **14** (Scheme 10). Cohen and Bercaw demonstrated the clean coupling of η^2 -bound ethylene with acetaldehyde to produce an oxatitanacyclopentane ring.²³ Some related reactions include coupling of η^2 -cyclobutene with acetone^{44a} and the intramolecular coupling of ene–one substrates.⁴⁷ In contrast, addition of $\text{Ph}_2\text{C}=\text{O}$ to the titanabicyclo **10** was found to produce the oxatitanacycloheptane **21** along with small amounts of the dioxatitanacyclopentane (benzopinacolate) **22**. The addition of $\text{Ph}_2\text{C}=\text{O}$ to the diphenyltitanacyclopentane **7** is found, even in the presence of excess styrene, to yield a mixture of oxatitanacyclopentane **23**, dioxatitanacycloheptane **24**, and dioxatitanacyclopentane **22**. In

(45) Cotton, F. A.; Kibala, P. A. *Polyhedron* **1987**, 6, 645.

(46) Steigerwald, M. L.; Goddard, W. A. *J. Am. Chem. Soc.* **1985**, 107, 5027.

(47) (a) Kablaoui, N. M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1996**, 118, 3182. (b) Hewlett, D. F.; Whitby, R. J. *J. Chem. Soc., Chem. Commun.* **1990**, 1684.




Figure 9. Molecular structure of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2\text{-}2,6)_2(\text{OCPh}_2)(\text{PMe}_3)]$ (**25**).


Table 8. Selected Bond Distances (Å) and Angles (deg) for $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2\text{-}2,6)_2(\eta^2\text{-OCPh}_2)(\text{PMe}_3)]$ (**25**)

Ti—P(4)	2.592(3)	Ti—O(1)	1.849(5)
Ti—O(2)	1.839(5)	Ti—O(3)	1.817(4)
Ti—C(1)	2.150(7)	C(1)—O(3)	1.397(8)
P(4)—Ti—O(1)	127.0(2)	P(4)—Ti—O(2)	88.6(2)
P(4)—Ti—O(3)	105.5(2)	P(4)—Ti—C(1)	90.9(2)
O(1)—Ti—O(2)	110.1(2)	O(1)—Ti—O(3)	109.0(2)
O(1)—Ti—C(1)	40.0(2)	O(2)—Ti—O(3)	115.8(2)
O(2)—Ti—C(1)	133.0(3)	O(3)—Ti—C(1)	109.5(2)
Ti—O(2)—C(21)	167.8(5)	Ti—O(3)—C(31)	159.2(4)

product **23**, NMR spectroscopy shows that the ketone has coupled with the methylene carbon of the styrene ligand, leading to the 3,3,5-triphenyl-2-oxa-titanacyclopentane regioisomer. This reaction has also been interrogated utilizing labeled $\text{Ph}_2^{13}\text{C}=\text{O}$, allowing not only confirmation of the ^{13}C NMR chemical shifts of the carbon, but also to confirm that the reagent benzophenone is incorporated into only the three products shown (Scheme 10).

The addition of $\text{Ph}_2\text{C}=\text{O}$ to the propene complex **15** followed by vacuum removal of volatile products, allows the isolation of the oxatitanacyclopropane (η^2 -ketone) complex **25** (Scheme 11). If the propene is not removed, or if excess propene is added to a solution of **25**, the oxatitanacyclopentane **26** is produced along with free PMe_3 . The η^2 -ketone complex **25** has been subjected to single crystal X-ray diffraction analysis (Figure 9, Table 8). Although a large number of aldehyde and ketone derivatives of the group 4 metals are known, they are typically bimolecular with bridging ketonic functions.⁴⁸ There are, however, mononuclear ketene species which are directly related to **25**.⁴⁹ The structural parameters for **25** are interesting, showing a distinct elongation of the C—O bond, 1.397(8), compared to a value of 1.23 in $\text{Ph}_2\text{C}=\text{O}$.⁵⁰ The Ti—O(3)

Scheme 11



distance of 1.817(4) is slightly shorter than the distances to the aryloxy oxygen atoms. These parameters indicate strong π -back-bonding into the π^* ($\text{C}=\text{O}$) orbital, and strongly support the oxatitanacyclopropane resonance picture.

Related reactivity is observed in the reactions involving the imine reagents $\text{PhCH}=\text{NR}$ ($\text{R} = \text{Ph}, \text{CH}_2\text{Ph}$). The titanabicyclo **10** produces the azatitanacycloheptane **27** ($\text{R} = \text{CH}_2\text{Ph}$, stereochemistry undetermined), whereas benzylideneaniline was observed to produce 1,7-octadiene and a new species believed to be diazatitanacyclopentane **28** (Scheme 11) based upon spectroscopic data. The diphenyltitanacyclopentane **7** reacts with $\text{PhCH}=\text{NPh}$ to produce the azatitanacyclopentane **29** in high yield. Analysis of the ^1H NMR spectrum of **29** as well as the labeled compound obtained using $\text{Ph}^{13}\text{CH}=\text{NPh}$, shows a *trans* 2,3,5-triphenyl isomer to be present. We justify the single regio- and stereochemistry for **29** on steric grounds. The *trans* stereochemistry allows the phenyl rings at the C-3 and C-5 positions to occupy pseudoaxial and pseudoequatorial positions respectively (Scheme 11). The axial positioning at C-3 removes conflict with the N-Ph group while the preference for an equatorial site for the phenyl group at C-5 will help avoid clashes with the bulky 2,6-diphenylphenoxide ligands which lie above and below the plane of the azatitanacyclopentane.

Reaction of the ethylene derivative **14** with $\text{PhCH}=\text{NCH}_2\text{Ph}$ produces the azatitanacyclopentane **30** and 1 equiv of PMe_3 (Scheme 12). In contrast the propene complex forms the diazatitanacyclopentane **31** along with propene. The ^{13}C NMR spectrum of **31** shows only one Ti—O—C(aryloxy) resonance consistent with a *trans* stereochemistry as shown (Scheme 12). The azatitanacyclopentanes **29** and **30** react with alkynes to displace olefin and generate azatitanacyclopent-4-enes (Scheme 12). The regiochemistry of these organometallic products is readily confirmed by NMR methods and hydrolysis is found to produce the corresponding allylic amines (Scheme 13). The regio- and stereochemistry of the coupling of alkynes and imines at zirconocene metal centers has been extensively investigated by Buchwald et al.⁵¹ The displacement of ethylene from zirconacyclopentene rings has been shown to be a synthetically useful reaction.⁵²


(48) (a) Erker, G. *Acc. Chem. Res.* **1984**, *17*, 103. (b) Kropp, K.; Skibbe, V.; Erker, G.; Kruger, C. *J. Am. Chem. Soc.* **1983**, *105*, 3353. (c) Erker, G.; Hoffmann, U.; Zwetler, R.; Betz, P.; Kruger, C. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 630. (d) Erker, G.; Czisch, P.; Schlund, R.; Angermund, K.; Kruger, C. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 364. (e) Stella, S.; Floriani, C. *J. Chem. Soc., Chem. Commun.* **1986**, 1053. (f) Erker, G.; Dorf, U.; Czisch, P.; Petersen, J. L. *Organometallics* **1986**, *5*, 668. (g) Fachinetti, G.; Biran, C.; Floriani, C.; Villa, A. C.; Guastini, C. *Inorg. Chem.* **1978**, *17*, 2995.

(49) (a) Ho, S. C. H.; Straus, D. A.; Armantrout, J.; Schaefer, W. P.; Grubbs, R. H. *J. Am. Chem. Soc.* **1984**, *106*, 2210. (b) Meinhardt, J. D.; Santarsiero, B. D.; Grubbs, R. H. *J. Am. Chem. Soc.* **1986**, *108*, 3318.


(50) Fleischer, E. B.; Sung, N.; Hawkinson, S. *J. Phys. Chem.* **1968**, *72*, 4311.

(51) (a) Buchwald, S. L.; Watson, B. T.; Wannamaker, M. W.; Dewan, J. C. *J. Am. Chem. Soc.* **1989**, *111*, 4486. (b) Buchwald, S. L.; Wannamaker, M. W.; Watson, B. T. *J. Am. Chem. Soc.* **1989**, *111*, 776.

Scheme 12



Scheme 13



The azatitanacyclopentane compound **30** reacts with one equivalent of benzonitrile to produce one equivalent of ethylene and a new titanium compound **35** (Scheme 14). The new compound is formulated as a 2,5-diazatitanacyclopent-2-ene derivative on the basis of spectroscopic data and a single crystal diffraction study of **35** (Figure 10, Table 9). The molecular structure of **35** shows a planar metallacycle ring, in contrast to the nonplanar situation found for 2,5-diazatitanacyclopent-3-ene species. Solutions of **35** were found to be converted over days into 2,5-diazatitanacyclopent-3-ene (ene-diamide) **36**, which was obtained as an impure liquid, but found to crystallize as its pyridine adduct **37** (Scheme 14). The isomerization of 2-aza-metallacyclopentene rings has literature precedence.^{23,53}

Conclusions

Good synthetic routes to a variety of titanacyclopentane and titanacyclopentane species for use in organometallic synthesis

(52) Takahashi, T.; Kageyama, M.; Denisov, V.; Hara, R.; Negishi, E. *Tetrahedron Lett.* **1993**, 34, 687.

(53) Strickler, J. R.; Wigley, D. E. *Organometallics* **1990**, 9, 1605 and references therein.




Figure 10. Molecular structure of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NCPHCHPhN}(\text{CH}_2\text{Ph})\}]$ (**35**).

Scheme 14




Table 9. Selected Bond Distances (Å) and Angles (deg) for $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NCPHCHPhN}(\text{CH}_2\text{Ph})\}]$ (**35**)

Ti—O(5)	1.816(2)	Ti—N(4)	1.886(2)
Ti—O(6)	1.829(2)	Ti—N(1)	1.910(2)
N(1)—C(2)	1.277(3)	C(2)—C(3)	1.529(4)
N(4)—C(3)	1.466(3)		
O(5)—Ti—O(6)	124.10(8)	O(5)—Ti—N(1)	113.61(9)
O(5)—Ti—N(4)	108.71(9)	O(6)—Ti—N(1)	114.72(9)
O(6)—Ti—N(4)	104.27(9)	N(4)—Ti—N(1)	81.9(1)
Ti—O(5)—C(51)	161.5(2)	Ti—O(6)—C(61)	151.5(2)

have been developed. The use of sterically demanding aryloxide groups has been found to generate chemistry which is both different and complementary to that developed with cyclopentadiene ligation. The solution spectroscopic properties and reactivity of the titanacyclopentane rings can only be accounted for by facile fragmentation into a bis(olefin) intermediate. The synthetic utility of both types of metallacycles has been probed with ketones and imines and shown to lead to a wide variety of new oxa and aza-titanacycles.

Experimental Section

All manipulations were carried out under N_2 using a Vacuum Atmospheres dry box and conventional Schlenk techniques. All solvents were dried by distillation over sodium/benzophenone under a nitrogen atmosphere. ^1H and ^{13}C NMR spectra were recorded using a Varian Gemini 200 MHz instrument. Microanalytical, mass spectral, and X-ray crystallographic data were acquired through Purdue in-house facilities. The compounds $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\text{Cl}_2]$ (**1**)¹⁵ and $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{C}_4\text{Et}_4)]$ (**3**)¹⁶ were obtained by published procedures. Synthetic details can be found in the Supporting Information.

[Ti(OC₆HPh₄-2,3,5,6,2-Cl₂) (2). Anal. Calcd for TiO₂Cl₂C₆₆H₄₈, 2·C₆H₆: C, 79.92; H, 4.88; Cl, 7.15. Found: C, 79.97; H, 4.70; Cl, 7.21. ¹H NMR (C₆D₆, 30 °C): δ 6.50–7.50 (aromatics).

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂)₂] (4). Anal. Calcd for TiC₄₀H₃₄O₂: C, 80.80; H, 5.76. Found: C, 80.69; H, 6.02. ¹H NMR (C₆D₆, 30 °C): δ 6.90–7.41 (aromatics); 1.98 (broad), 1.60 (broad, CH₂). ¹³C NMR (C₆D₆, 30 °C): δ 160.3 (Ti–O–C); 89.7 [¹J(¹³C–¹H) = 127.4 Hz, TiCH₂]; 31.1 [¹J(¹³C–¹H) = 127.7 Hz, TiCH₂CH₂].

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CHMeCHMeCH₂)] (5). Anal. Calcd for TiC₄₂H₃₀O₂: C, 81.02; H, 6.15. Found: C, 81.31; H, 6.56. ¹H NMR (C₆D₆, 30 °C): *trans* isomer δ 6.88–7.48 (aromatic); 2.04 (m, CHMe); 1.88 (t, 10.6 Hz), 1.24 (dd, TiCH₂, 10.6 Hz, 4.1 Hz); 0.93 (d, CHMe, 5.4 Hz). ¹³C NMR (C₆D₆, 30 °C): *trans* isomer δ 160.1 (Ti–O–C); 98.0 [¹J(¹³C–¹H) = 129.2 Hz, TiCH₂]; 44.8 [¹J(¹³C–¹H) = 126.2 Hz, CHMe]; 23.3 (CHMe).

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CHEtCHEtCH₂)] (6). Anal. Calcd for TiC₄₄H₄₂O₂: C, 81.22; H, 6.51. Found: C, 81.54; H, 6.91. ¹H NMR (C₆D₆, 30 °C): *trans* isomer δ 6.88–7.8 (aromatics); 2.03 (m, CHEt); 1.85 (t), 1.22 (m, TiCH₂); 1.22 (m, CH₂CH₃); 0.60 (t, CH₂CH₃). ¹³C NMR (C₆D₆, 30 °C): *trans* isomer δ 160.2 (Ti–O–C); 95.0 [¹J(¹³C–¹H) = 126.5 Hz, TiCH₂]; 48.7 [¹J(¹³C–¹H) = 126.8 Hz, CHEt]; 28.9 (CH₂CH₃); 18.1 (CH₂CH₃).

[Ti(OC₆H₃Ph₂-2,6,2)(C₄H₆(Ph)₂)] (7). Anal. Calcd for TiC₅₂H₄₂O₂: C, 83.57; H, 5.64. Found: C, 83.63; H, 5.67. ¹H NMR (C₆D₆, 30 °C): δ 6.20–7.41 (aromatics); 3.30 (m), 2.77 (broad m), 2.00 (broad m), 1.30 (m). ¹³C NMR (C₆D₆, 30 °C): δ 160.3 (Ti–O–C); 107.0 (broad, TiCHPh).

Dimerization of Styrene Catalyzed by [Ti(OC₆H₃Ph₂-2,6,2)(C₄H₆(Ph)₂)] (7). In a typical procedure 0.10 g (0.15 mmol) of [Ti(OC₆H₃Ph₂-2,6,2)Cl₂] (1) was dissolved in 5 mL of benzene. To this solution was added 2.1 equiv of BuⁿLi (0.12 mL, 0.31 mmol) and 25 equiv of styrene (0.42 mL, 3.64 mmol). This reaction mixture was then heated at approximately 100 °C initially producing *trans*-1,3-diphenylbut-1-ene. ¹H NMR (C₆D₆, 30 °C): δ 6.80–7.40 (aromatics); 6.33 [AB, ³J(¹H–¹H) = 16.0 Hz, *trans* CH=CH]; 3.42 (m, CHPhCH₃); 1.33 (d, CH₃). ¹³C NMR (C₆D₆, 30 °C): δ 135.1, 128.4 (CHPh=CH); 42.5 (CHPhCH₃); 21.2 (CH₃). HRMS: calcd for C₁₆H₁₆ 208.1252, found 208.1251. Upon further heating this initial product was isomerized to 1,3-diphenylbut-2-ene. ¹H NMR (C₆D₆, 30 °C): δ 6.80–7.40 (aromatics); 5.89 [t, ³J(¹H–¹H) = 7.3 Hz, CH]; 3.30 (d, CH₂Ph); 1.86 (s, CH₃).

[Ti(OC₆H₃Ph₂-2,6,2)(μ-Cl)]₂ (8). Anal. Calcd for Ti₂C₇₂H₅₂O₄Cl₂: C, 75.34; H, 4.57; Cl, 6.18. Found: C, 81.54; H, 4.85; Cl, 6.29. The X-ray crystal structure of 8 has been previously published.

[Ti(OC₆HPh₄-2,3,5,6,2)(μ-Cl)]₂ (9). Anal. Calcd for TiO₄-Cl₂C₁₂₀H₈₄: C, 82.05; H, 4.82; Cl, 4.09. Found: C, 81.75; H, 5.05; Cl, 4.12.

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CH(C₄H₈)CHCH₂)] (10). Anal. Calcd for TiC₄₄H₄₀O₂: C, 81.47; H, 6.22. Found: C, 81.56; H, 6.41. ¹H NMR (C₆D₆, 30 °C): δ 6.90–7.42 (aromatics); 0.4–2.0 (aliphatics). ¹³C NMR (C₆D₆, 30 °C): δ 160.2 (Ti–O–C); 95.9 [¹J(¹³C–¹H) = 128.0 Hz, TiCH₂]; 46.8 [¹J(¹³C–¹H) = 129.0 Hz, TiCH₂CH]; 37.7 [¹J(¹³C–¹H) = 124.5 Hz, TiCH₂CHCH₂]; 27.0 [¹J(¹³C–¹H) = 124.5 Hz, TiCH₂CHCH₂CH₂]. The product of hydrolysis of 10 was identified as *trans*-1,2-dimethylcyclohexane by comparison of the ¹H NMR spectrum and GC trace to that of an authentic sample.

[Ti(OC₆HPh₄-2,3,5,6,2)(CH₂CH(C₄H₈)CHCH₂)] (11). To a stirred mixture of [Ti(OC₆HPh₄-2,3,5,6,2)Cl₂] (2, 3.00 g, 3.03 mmol) and 1,7-octadiene (0.90 mL, 6.06 mmol, two fold excess) in benzene (10 mL) was slowly added BuⁿLi (6.06 mmol) in hexane (2.5 M solution). The resulting brown suspension was filtered and the solvent removed under vacuum. The oily residue was dissolved in a minimum of toluene and allowed to stand whereupon orange crystals of the toluene solvate 3b·2C₇H₈ were deposited. The yield of 3b obtained in this fashion has been found to vary from 15–75%. Satisfactory microanalytical data on 3b could not be obtained presumably due to loss of toluene solvate.

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CHMeCMe₂)]₂(μ-O) (12). Anal. Calcd for Ti₂C₈₄H₆₆O₃: C, 79.99; H, 6.07. Found: C, 80.35; H, 6.58. ¹H NMR (C₆D₆, 30 °C): δ 6.83–7.39 (aromatics); 0.42–1.48 (aliphatics). ¹³C NMR (C₆D₆, 30 °C): δ 160.7 (Ti–O–C); 99.6, 99.2 (TiCH₂); 42.4, 41.8 (CHMe); 35.9, 35.7 (CHMe₂); 20.6, 20.5 (CHMe); 17.8, 17.5, 17.5, 17.3 (CHMe₂).

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CH₂(PMe₃))] (14a). Anal. Calcd for TiC₄₁H₃₉O₂P: C, 76.63; H, 6.12; P, 4.82. Found: C, 76.99; H, 6.14; P, 4.58. ¹H NMR (C₆D₆, 30 °C): δ 6.87–7.36 (aromatics); 1.72 [td, ³J(¹H–³¹P) = 6.8 Hz, ³J(¹H–¹H) = 12 Hz CH₂ adjacent to PMe₃]; 0.51 [t, ³J(¹H–³¹P) < 1 Hz, CH₂ away from PMe₃]; 0.22 (d, PMe₃). ¹³C NMR (C₆D₆, 30 °C): δ 160.0 (Ti–O–C); 78.0 [d, ¹J(¹³C–¹H) = 147.6 Hz, ²J(¹³C–³¹P) = 5.9 Hz, CH₂ adjacent to PMe₃]; 67.0 [¹J(¹³C–¹H) = 149.6 Hz, CH₂ away from PMe₃]; 12.5 (d, PMe₃). ³¹P NMR (C₆D₆, 30 °C): δ 2.43 (PMe₃).

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CH₂(PMe₂Ph))] (14b). ¹H NMR (C₆D₆, 30 °C): δ 6.80–7.27 (aromatics); 1.70 (m), 0.5 (m, CH₂CH₂); 0.45 (d, PMe₂Ph). ¹³C NMR (C₆D₆, 30 °C): δ 160.0 (Ti–O–C); 79.4 [d, ²J(¹³C–³¹P) = 2.8 Hz, CH₂ adjacent to PMe₂Ph]; 68.8 (CH₂ away from PMe₂Ph); 11.0 (d, PMe₂Ph).

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CHMe)(PMe₃)] (15a). Anal. Calcd for TiC₄₂H₄₁O₂P: C, 76.82; H, 6.29; P, 4.72. Found: C, 76.83; H, 6.46; P, 4.98. ¹H NMR (C₆D₆, 30 °C): major isomer δ 6.8–7.6 (aromatics); 1.61 (m), 1.48 (m, CH₂); 1.43 (d, 8.0 Hz, CHMe); 0.53 (m, CHMe); 0.14 (d, PMe₃); minor isomer δ 2.32 (m, CHMe); 1.04 (d, 8.0 Hz, CHMe); 0.90 (m), 0.44 (m, CH₂); 0.02 (d, PMe₃). ³¹C NMR (C₆D₆, 30 °C) major isomer δ 80.7 [d, ¹J(¹³C–¹H) = 148.1 Hz, ²J(¹³C–³¹P) = 5.5 Hz, CH₂]; 76.2 [¹J(¹³C–¹H) = 148.1 Hz, CHMe]; 22.6 (CHMe); 12.5 (PMe₃); minor isomer δ 89.2 [d, ¹J(¹³C–¹H) = 148.3 Hz, ²J(¹³C–³¹P) = 6.4 Hz, CHMe]; 74.0 [¹J(¹³C–¹H) = 146.5 Hz, CH₂]; 20.9 (CHMe); 12.8 (PMe₃). ³¹P NMR (C₆D₆, 30 °C): major isomer δ 2.19 (PMe₃); minor isomer δ 1.40 (PMe₃).

[Ti(OC₆H₃Ph₂-2,6,2)(CH₂CHMe)(PMe₂Ph)] (15b). Anal. Calcd for TiC₄₇H₄₃O₂P: C, 78.54; H, 6.03; P, 4.31. Found: C, 75.23; H, 5.88; P, 3.02. ¹H NMR (C₆D₆, 30 °C): major isomer δ 6.73–7.59 (aromatics); 1.46 (d, 6.2 Hz, CHMe); 1.63 (m), 1.47 (m, CH₂); 0.41 (m, CHMe); 0.64 (d), 0.26 (d, 5.8 Hz, PMe₂Ph); minor isomer δ 2.23 (m, CHMe); 1.13 (d, 6.4 Hz, CHMe); 1.00 (m), 0.4 (m, CH₂); 0.36 (d, 6.4 Hz, PMe₂Ph). ¹³C NMR (C₆D₆, 30 °C): major isomer δ 81.8 [d, ¹J(¹³C–¹H) = 146.2 Hz, ²J(¹³C–³¹P) = 2.6 Hz, CH₂]; 77.5 [¹J(¹³C–¹H) = 148.3 Hz, CHMe]; 22.6 (CHMe); minor isomer δ 90.0 (CH₂); 76.2 (CHMe). ³¹P NMR (C₆D₆, 30 °C): major isomer δ 14.8 (PMe₂Ph); minor isomer δ 11.4 (PMe₂Ph).

[Ti(OC₆H₃Ph₂-2,6,2)(CHPhCH₂)(PMe₂Ph)] (17). ¹H NMR (C₆D₆, 30 °C): major isomer δ 6.50–7.40 (aromatics); 2.20 (ddd), 1.79 (ddd, 12 Hz, 7 Hz, CH₂); 1.10 (t, 12 Hz, CHPh); 0.63 (d), 0.03 (d, 7.2 Hz, PMe₂Ph); minor isomer δ 2.70 (dd, 12 Hz, CHPh); 0.98 (dd), 0.22 (dd, 12 Hz, 6 Hz, CH₂); 0.46 (d), –0.02 (d, 7.3 Hz, PMe₂Ph). ¹³C NMR (C₆D₆, 30 °C): major isomer δ 159.5 (Ti–O–C); 159.3 (Ti–O–C); 93.1 [d, ²J(¹³C–³¹P) = 2.9 Hz, CH₂]; 78.1 (CHPh); minor isomer δ 160.1 (Ti–O–C); 159.8 (Ti–O–C); 83.3 (CH₂); 80.1 [d, ²J(¹³C–³¹P) = 5.3 Hz, CHPh]. ³¹P NMR (C₆D₆, 30 °C): major isomer δ 15.2 [³J(³¹P–¹H) = 7.0 Hz, PMe₂Ph]; minor isomer δ 13.5 [³J(³¹P–¹H) = 6.0 Hz, PMe₂Ph].

[Ti(OC₆H₃Ph₂-2,6,2)(OCPh₂CH₂CH₂CH₂CH₂)] (18). Anal. Calcd for TiC₅₉H₅₀O₃: C, 82.90; H, 5.85. Found: C, 82.61; H, 6.14. ¹H NMR (C₆D₆, 30 °C): δ 6.88–7.42 (aromatics); 2.22 (m, CH₂CPh₂); 1.44 (m, TiCH₂CH₂); 1.21 (m, CH₂CH₂CPh₂); 1.04 (m, TiCH₂). ¹³C NMR (C₆D₆, 30 °C): δ 159.6 (Ti–O–C); 94.1 (OCPh₂); 87.5 (TiCH₂); 43.4 (TiCH₂CH₂); 32.4, 26.3 (CH₂CH₂CPh₂).

Preparation of [Ti(OC₆H₃Ph₂-2,6,2)(OCPh₂CH₂CH₂)] (19). Anal. Calcd for TiC₅₁H₄₀O₃: C, 81.81; H, 5.38. Found: C, 81.62; H, 5.20. ¹H NMR (C₆D₆, 30 °C): δ 6.85–7.38 (aromatics); 3.34 (t), 1.47 (t, 6.6 Hz, CH₂). Selected ¹³C NMR (C₆D₆, 30 °C): δ 160.0 (ipso OAr'); 148.8 (ipso OCPh₂); 92.4 (TiOCPh₂); 82.2 [¹J(¹³C–¹H) = 132.2 Hz, TiCH₂]; 53.0 [¹J(¹³C–¹H) = 131.0 Hz, TiCH₂CH₂].

[Ti(OC₆H₃Ph₂-2,6,2)(OCPh₂CH₂CH₂CPh₂O)] (20). Anal. Calcd for TiC₆₄H₅₀O₄: C, 82.57; H, 5.41. Found: C, 82.82; H, 5.52. ¹H NMR (C₆D₆, 30 °C): δ 6.78–7.41 (aromatics); 2.51 (s, CH₂). ¹³C NMR (C₆D₆, 30 °C): δ 160.0 (Ti–O–C); 95.1 (TiOCPh₂); 38.0 (CH₂).

[Ti(OC₆H₃Ph₂-2,6,2)(OCPh₂CH₂CH(C₄H₈)CHCH₂)] (21). Anal. Calcd for TiC₅₇H₅₀O₃: C, 82.39; H, 6.07. Found: C, 82.61; H, 6.16. ¹H NMR (C₆D₆, 30 °C): δ 6.83–7.71 (aromatics); 2.37 (dd, 16 Hz, 8 Hz), 2.18 (d, CH₂CPh₂); 0.63–1.38 (m, other aliphatic protons); 0.40 (bdd), 0.22 (bdd, TiCH₂). ¹³C NMR (C₆D₆, 30 °C): δ 160.0, 159.0 (Ti–O–C); 97.5 (OCPh₂); 93.1 (TiCH₂); 52.5 (OCPh₂CH₂); 49.3, 39.6 (CH); 37.8, 37.6, 26.8, 26.6 (CHCH₂CH₂).

Reaction of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{C}_4\text{H}_6(\text{Ph})_2\}]$ (7) with $\text{Ph}_2\text{C}=\text{O}$. A sample of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{C}_4\text{H}_6(\text{Ph})_2\}]$ (7) was dissolved in benzene- d_6 (1 mL) in a J. Young valve solvent sealed NMR tube. To this solution was added solid benzophenone. The ^1H NMR at 30 °C of the resulting dark red solution was immediately obtained and contains **22**, **23** and **24**: δ 6.50–7.80 (aromatics); 5.62 (broad d), 5.10 (broad d, $\text{PhCH}=\text{CH}_2$); 4.51 (broad, $\text{Ti}-\text{OCPh}_2\text{CHPh}$ of **24**); 3.71 (t, TiCHPh of **23**); 3.20–3.45 (m, TiCHPhCH_2 of **23**); 3.38 (dd), 3.01 (broad d, $\text{Ti}-\text{OCPh}_2\text{CH}_2$ of **24**). Over the course of one day, or if more benzophenone was added, the solution became light orange in color. The ^1H NMR of this solution showed that the signals for **23** had disappeared and the ^1H NMR signals for **24** had become more intense.

Reaction of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{C}_4\text{H}_6(\text{Ph})_2\}]$ (7) with $\text{Ph}_2^{13}\text{C}=\text{O}$. A sample of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{C}_4\text{H}_6(\text{Ph})_2\}]$ (7) was placed in a J. Young valve solvent sealed NMR tube along with benzene- d_6 (1 mL). To this was added one crystal of solid ^{13}C -labeled benzophenone. The ^{13}C NMR, at 30 °C, of the resulting red solution clearly shows the presence of **22*** (δ 112.8), **23*** (δ 90.3), and **24*** (δ 99.0 and 97.5). As more benzophenone was added (one crystal at a time) the signal at δ 90.3 disappeared and the signals at δ 112.8, 99.0, and 97.5 increased in intensity. This corresponds to the same trend seen in the ^1H NMR above using natural abundance benzophenone.

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{OCPh}_2)(\text{PMe}_3)]$ (25). Anal. Calcd for $\text{TiC}_{52}\text{H}_{45}\text{O}_3\text{P}$: C, 78.38; H, 5.69; P, 3.89. Found: C, 79.27; H, 5.90; P, 3.73. ^1H NMR (C_6D_6 , 30 °C): δ 6.79–7.49 (aromatics); –0.23 (d, PMe_3). ^{13}C NMR (C_6D_6 , 30 °C): δ 160.5 ($\text{Ti}-\text{O}-\text{C}$); 91.4 (CPh_2); 12.0 (d, PMe_3). Hydrolysis product Ph_2CHOH : ^1H NMR (C_6D_6 , 30 °C): δ 6.95–7.30 (aromatics) 5.48 (d, OH); 1.69 (d, CH).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{OCPh}_2\text{CH}_2\text{CHMe})]$ (26). The major isomer was $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{OCPh}_2\text{CH}_2\text{CHMe})]$ while the minor isomer was $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{OCPh}_2\text{CHMeCH}_2)]$. ^1H NMR (C_6D_6 , 30 °C): major isomer δ 6.7–7.5 (aromatics); 3.25 (dd, 13 Hz, 12 Hz), 2.94 (dd, 14 Hz, 6 Hz, CH_2); 1.88 (m, CHMe); 0.60 (d, 6 Hz, CHMe); minor isomer δ 3.69 (m, CHMe); 1.12 (dd), 0.99 (dd, CH_2); 0.93 (d, CHMe). ^{13}C NMR (C_6D_6 , 30 °C): major isomer δ 159.7 ($\text{Ti}-\text{O}-\text{C}$); 149.4, 148.6 (ipso OCPh_2); 95.4 (TiCHMe); 89.8 (TiOCPh_2); 59.0 (CH_2); 21.8 (CHMe).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{N}(\text{CH}_2\text{Ph})\text{CHPhCH}_2\text{CH}(\text{C}_4\text{H}_8)\text{CHCH}_2\}]$ (27). Anal. Calcd for $\text{TiC}_{58}\text{H}_{53}\text{NO}_2$: C, 82.54; H, 6.33; N, 1.66. Found: C, 82.58; H, 6.27; N, 1.66. ^1H NMR (C_6D_6 , 30 °C): δ 6.69–7.51 (aromatics); 4.30 (d), 3.01 (d, 15 Hz, CH_2Ph); 3.67 (d, 12 Hz, CHPh); 2.58 (m, CHPhCH_2); 0.31 (m), 0.02 (m, TiCH_2); 0.70–1.50 (other aliphatic protons). ^{13}C NMR (C_6D_6 , 30 °C): δ 160.5, 160.2 ($\text{Ti}-\text{O}-\text{C}$); 87.6 (TiCH_2); 63.3 (NCHPh); 58.1 (NCH_2Ph); 48.1 (CH); 43.4, 39.6, 27.5, 27.1 (CHCH_2CH_2). Hydrolysis product $\text{BzNHCHPhCH}_2\text{CH}(\text{C}_4\text{H}_8)\text{CH}_3$: ^1H NMR (C_6D_6 , 30 °C): δ 6.85–7.49 (aromatics); 3.63 (dd, CH_2Ph); 3.49 (q, NH); 0.77 (d, CH_3); 0.6–2.1 (other aliphatic protons). Mass spectroscopy: parent M^+ at 307 amu.

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{N}(\text{Ph})\text{CHPhCHPhN}(\text{Ph})\}]$ (28). ^1H NMR (C_6D_6 , 30 °C): δ 5.84–7.60 (aromatics); 5.73, 5.00 (m's, olefinic 1,7-octadiene); 5.41, 4.93 (*s,s,cis*- and *trans*- NPhCHPh); 2.00–1.00 (aliphatic 1,7-octadiene). ^{13}C NMR (C_6D_6 , 30 °C): δ 73.7, 70.8 (*cis*- and *trans*- NPhCHPh).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2(\text{NPhCHPhCH}_2\text{CHPh})]$ (29). Anal. Calcd for $\text{TiC}_{57}\text{H}_{45}\text{NO}_2$: C, 83.08; H, 5.51; N, 1.70. Found: C, 82.74; H, 5.63; N, 1.82. ^1H NMR (C_6D_6 , 30 °C): δ 6.50–7.50 (aromatics); 5.91 (d), 5.47 (d, 7.7 Hz, *ortho*- NPh); 4.37 (d, 3.7 Hz, TiNPhCHPh); 3.32 (ddd, 16.2 Hz, 14.5 Hz, 5.2 Hz), 3.04 (dd, 14.3 Hz, 3.5 Hz, TiCHPhCH_2); 2.42 (d, 15.8 Hz, TiCHPh). ^{13}C NMR (C_6D_6 , 30 °C): δ 160.7, 160.1 ($\text{Ti}-\text{O}-\text{C}$); 107.5 (TiCHPh); 59.0 (TiNPhCHPh); 30.7 (TiCHPhCH_2).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{N}(\text{CH}_2\text{Ph})\text{CHPhCH}_2\text{CH}_2\}]$ (30). Anal. Calcd for $\text{TiC}_{58}\text{H}_{49}\text{NO}_2$: $30\cdot\text{C}_6\text{H}_6$: C, 82.94; H, 5.88; N, 1.67. Found: C, 82.63; H, 5.95; N, 2.05. ^1H NMR (C_6D_6 , 30 °C): δ 6.63–7.43 (aromatics); 3.89 (d), 3.24 (d, 15.1 Hz, CH_2Ph); 3.75 (t, 6.9 Hz, CHPh); 2.21 (m, TiCH_2CH_2); 1.39 (m), 1.15 (m, TiCH_2). ^{13}C NMR (C_6D_6 , 30 °C): δ 160.4, 160.1 ($\text{Ti}-\text{O}-\text{C}$); 85.9 (TiCH_2); 64.2 (CHPh); 57.1; $\text{CH}_2\text{-Ph}$); 33.7 (TiCH_2CH_2). Hydrolysis product $[\text{PhCH}_2\text{NHCHPhCH}_2\text{CH}_3]$: ^1H NMR (C_6D_6 , 30 °C): δ 7.08–7.30 (aromatics); 3.66 (d), 3.44 (d, 13.6 Hz, CH_2Ph); 3.41 (t, CHPh); 2.7 (broad, NH); 1.63 (dq, CH_2); 0.70 (t, 7.4 Hz, CH_3). ^{13}C NMR (C_6D_6 , 30 °C): δ 64.6 (CHPh); 51.8 (CH_2Ph); 31.4 (CH_2); 10.9 (CH_3).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{N}(\text{CH}_2\text{Ph})\text{CHPhCHPhN}(\text{CH}_2\text{Ph})\}]$ (31). Anal. Calcd for $\text{TiC}_{64}\text{H}_{52}\text{N}_2\text{O}_2$: C, 82.74; H, 5.64; N, 3.02. Found: C, 82.83; H, 5.67; N, 3.02. ^1H NMR (C_6D_6 , 30 °C): δ 6.25–7.69 (aromatics); 4.68 (s, CHPh); 3.99 (d), 2.99 (d, 16 Hz, CH_2Ph). ^{13}C NMR (C_6D_6 , 30 °C): δ 160.6 (ipso OAr''); 76.3 (CHPh); 55.7 (CH_2Ph).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{N}(\text{CH}_2\text{Ph})\text{CHPhCEtCEt}\}]$ (32). Anal. Calcd for $\text{TiC}_{56}\text{H}_{49}\text{NO}_2$: C, 82.44; H, 6.05; N, 1.72. Found: C, 82.84; H, 6.27; N, 1.87. ^1H NMR (C_6D_6 , 30 °C): δ 6.76–7.61 (aromatics); 4.35 (s, CHPh); 3.14 (d), 2.90 (d, 15.7 Hz, CH_2Ph); 1.40 (q, 7.6 Hz), 1.26 (m, CH_2CH_3); 0.55 (t), 0.40 (t, 7.5 Hz, CH_2CH_3). ^{13}C NMR (C_6D_6 , 30 °C): δ 218.2 (TiCEt); 160.7, 160.4 ($\text{Ti}-\text{O}-\text{C}$); 139.8 (TiCEtCEt); 61.7 (CHPh); 55.5 (CH_2Ph); 27.2, 23.8 (CH_2CH_3); 14.8, 13.1 (CH_2CH_3). Hydrolysis product $\text{CHEt}=\text{CEtCHPhNHBz}$: ^1H NMR (C_6D_6 , 30 °C): δ 7.04–7.42 (aromatics); 5.62 (t, 7.3 Hz, CHEt); 4.14 (s, CHPh); 3.63 (dd, 16.9 Hz, 3.5 Hz, CH_2Ph); 1.97 (m, CH_2CH_3); 1.32 (broad, NH); 0.94 (t), 0.80 (t, 7.5 Hz, CH_2CH_3). ^{13}C NMR (C_6D_6 , 30 °C): δ 68.7 (CHPh); 52.2 (CH_2Ph); 21.6, 21.4 (CH_2CH_3); 14.9, 14.5 (CH_2CH_3). HRMS: calcd for 280.2065, found 280.2062.

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{N}(\text{CH}_2\text{Ph})\text{CHPhCHCPh}\}]$ (33). Anal. Calcd for $\text{TiC}_{58}\text{H}_{45}\text{NO}_2$: C, 83.34; H, 5.43; N, 1.68. Found: C, 83.00; H, 5.54; N, 1.71. ^1H NMR (C_6D_6 , 30 °C): δ 6.43–7.61 (aromatics); 6.74 (d, 3.1 Hz, CPhCH); 4.50 (d, 3.1 Hz, CHPh); 3.23 (d); 3.05 (d, 15.8 Hz, CH_2Ph). ^{13}C NMR (C_6D_6 , 30 °C): δ 207.5 (TiCPh); 160.9, 160.6 ($\text{Ti}-\text{O}-\text{C}$); 138.4 (CPhCH); 58.9 (CHPh); 56.6 (CH_2Ph). Hydrolysis product *trans*- CHPhCHCHPhNHBz : ^1H NMR (C_6D_6 , 30 °C): δ 7.0–7.45 (aromatics); 6.51 (d, 15.8 Hz, $\text{C}=\text{CHPh}$); 6.25 (dd, 15.8 Hz, 7.0 Hz, $\text{CH}=\text{C}$); 4.26 (d, 7.0 Hz, NCHPh); 3.65 (s, NCH_2Ph); 1.89 (broad, NH). ^{13}C NMR (C_6D_6 , 30 °C): δ 65.1 (CHPh); 51.8 (CH_2Ph). HRMS: calcd for 299.1674, found 299.1668.

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NPhCHPhCHC}(\text{SiMe}_3)\}]$ (34). Anal. Calcd for $\text{TiC}_{54}\text{H}_{47}\text{NO}_2\text{Si}$: C, 79.29; H, 5.80; N, 1.71. Found: C, 78.90; H, 6.06; N, 1.63. ^1H NMR (C_6D_6 , 30 °C): δ 7.57 (d, *ortho*- $\text{Ph}_2-2,6$); 7.39 (d, 4.0 Hz, TiNPhCHPhCH); 6.40–7.30 (aromatics); 5.45 (d, *ortho*- NPh); 4.62 (d, 4.0 Hz, TiNPhCHPh); –0.25 [s, $\text{Si}(\text{CH}_3)_3$]. ^{13}C NMR (C_6D_6 , 30 °C): δ 228.4 ($\text{TiNPhCHPhCHCSiMe}_3$); 160.7, 160.5 ($\text{Ti}-\text{O}-\text{C}$); 137.4 (TiNPhCHPhCH); 54.5 (TiNPhCHPh); –1.0 [$\text{Si}(\text{CH}_3)_3$]. Hydrolysis product *trans*- $\text{Me}_3\text{SiCH}=\text{CHCHPhNHPH}$: ^1H NMR (C_6D_6 , 30 °C): δ 6.60–7.30 (aromatics); 6.41 (d, *ortho*- NPh); 6.10 [dd, $^3J(\text{H}-\text{H}) = 18.5$ Hz and 5.0 Hz, Me_3SiCHCH]; 5.88 [d, $^3J(\text{H}-\text{H}) = 18.5$ Hz, Me_3SiCH]; 4.78 (b, $\text{Me}_3\text{SiCHCHCHPh}$); 3.73 (broad d, NHPH); –0.02 [s, $\text{Si}(\text{CH}_3)_3$].

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NCPHCHPhN}(\text{CH}_2\text{Ph})\}]$ (35). Anal. Calcd for $\text{TiC}_{63}\text{H}_{50}\text{N}_2\text{O}_2$: $35\cdot\text{C}_6\text{H}_6$: C, 82.70; H, 5.51; N, 3.06. Found: C, 83.32; H, 5.78; N, 2.99. ^1H NMR (C_6D_6 , 30 °C): δ 6.73–7.62 (aromatics); 5.81 (s, CHPh); 3.72 (d), 2.96 (d, 15.8 Hz, CH_2Ph). ^{13}C NMR (C_6D_6 , 30 °C): δ 186.4 (TiNCPH); 161.4, 160.7 (ipso OAr''); 90.6 (CHPh); 53.5 (CH_2Ph).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NHCPHCPH}(\text{CH}_2\text{Ph})\}]$ (36). A sample of $[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NCPHCHPhN}(\text{CH}_2\text{Ph})\}]$ (30, 0.50 g, 0.62 mmol) was dissolved in benzene (10 mL). A small amount of PhCN was added, and the solution was refluxed for 1 h. Excess solvent was removed under vacuum, and the product **36** was recovered as a dark red oil. Selected ^{13}C NMR (C_6D_6 , 30 °C): δ 159.2 ($\text{Ti}-\text{O}-\text{C}$); 117.0, 115.0 (CPhCPh); 58.4 (CH_2Ph).

$[\text{Ti}(\text{OC}_6\text{H}_3\text{Ph}_2-2,6)_2\{\text{NHCPHCPH}(\text{CH}_2\text{Ph})\}(\text{py})]$ (37). Anal. Calcd for $\text{TiC}_{62}\text{H}_{49}\text{N}_3\text{O}_2$: C, 81.30; H, 5.39; N, 4.59. Found: C, 81.67; H, 5.57; N, 4.49. ^1H NMR (C_6D_6 , 30 °C): δ 9.91 (broad, NH); 6.13–7.60 (aromatics); 4.62 (s, CH_2Ph). ^{13}C NMR (C_6D_6 , 30 °C): δ 160.6 (ipso OAr''); 56.4 (CH_2Ph).

Acknowledgment. We thank the National Science Foundation (Grant CHE-9700269) for financial support of this research.

Supporting Information Available: Experimental details of the synthesis of new compounds; description of the experimental procedures for X-ray diffraction studies; tables of thermal parameters, bond distances and angles, intensity data, torsion angles, and multiplicities for **4**, **7**, **10**, **12**, **14a**, **25** and **35** (172 pages). See any current masthead page for ordering information and Internet access instructions.