

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231448698>

Stereochemical analysis of the methyl transfer catalyzed by cobalamin-dependent methionine synthase from E. coli B

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MAY 1986

Impact Factor: 12.11 · DOI: 10.1021/ja00271a081

CITATIONS

45

READS

6

9 AUTHORS, INCLUDING:



Verna Frasca

Malvern Instruments

17 PUBLICATIONS 1,160 CITATIONS

SEE PROFILE



Rowena Green Matthews

University of Michigan

172 PUBLICATIONS 13,349 CITATIONS

SEE PROFILE

is prohibitive¹⁷ (theoretical calculations¹⁸ place barriers between 62 and ~80 kcal mol⁻¹ on this transformation), the strained nature of **1** suggests that it should be able to undergo ring opening to the triyne. Indeed, while **1b** is stable in the neat state to 400 °C (10 min, 50% recovery), flash pyrolysis (700 °C, 10⁻² mm) leads to the desired isomerization (40%; clearly identifiable by spectral data, particularly ¹³C NMR²⁰). This retrocyclization is the first which unravels a benzene ring to its three-component alkyne units in a purely thermal process without additional reagents which would be normally required to overcome thermodynamic obstacles.¹⁹ Our observations lend support to the contention that **1b** may have been an intermediate in the flash-thermolytical decomposition of a tricinoline to the same triyne.²⁰

With **1b** and two of its derivatives at hand, a detailed investigation of their physical, chemical, and physiological properties will be the subject of future efforts.

Acknowledgment. This paper is dedicated to the memory of Carol L. Goodman, who died after a heroic battle with life. This work was supported by the National Institutes of Health (CA-20713); K.P.C.V. is a Miller Research Professor in Residence (1985-1986). R.D. was the recipient of a NATO Science Fellowship (1984-1985). The X-ray structural analysis was carried out by Dr. F. J. Hollander, staff crystallographer. We thank J. C. Armstrong for carrying out the pyrolysis experiments on **1b**.

Supplementary Material Available: A listing of positional and thermal parameters and tables of bond lengths and angles and structure factors for **1a** (45 pages). Ordering information is given on any current masthead page.

(17) Dower, W. V.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1982**, *104*, 6878; *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 685; *Angew. Chem. Suppl.* **1982**, 1545; *Tetrahedron*, in press.

(18) Bach, R. D.; Wolber, G. J.; Schlegel, H. B. *J. Am. Chem. Soc.* **1985**, *107*, 2837. Houk, K. N.; Gandour, R. W.; Strozier, R. W.; Rondan, N. G.; Paquette, L. A. *Ibid.* **1979**, *101*, 6797.

(19) Parker, W. L.; Hexter, R. M.; Siedle, A. R. *J. Am. Chem. Soc.* **1985**, *107*, 4584.

(20) Barton, J. W.; Shepherd, M. K. *Tetrahedron Lett.* **1984**, 25, 4967.

Stereochemical Analysis of the Methyl Transfer Catalyzed by Cobalamin-Dependent Methionine Synthase from *Escherichia coli* B

T. M. Zydowsky,[†] L. F. Courtney,[‡] V. Frasca,[§]
K. Kobayashi,[†] H. Shimizu,[†] L.-D. Yuen,[†] R. G. Matthews,[§]
S. J. Benkovic,^{*,†} and H. G. Floss^{*,†}


Department of Chemistry, The Ohio State University
Columbus, Ohio 43210

Department of Chemistry, Pennsylvania State University
University Park, Pennsylvania 16802

Department of Biological Chemistry and Biophysics
Research Division, University of Michigan
Ann Arbor, Michigan 48109

Received January 31, 1986

The last step in the biosynthesis of methionine involves transfer of the methyl group of N⁵-methyltetrahydrofolate (5-CH₃-H₄-folate) or its polyglutamate analogue to the sulfur of homocysteine:




[†] The Ohio State University.


[‡] Pennsylvania State University.

[§] University of Michigan.

Scheme I



Scheme II



Two classes of methionine synthases are known to catalyze this reaction, one contains cobalamin as a cofactor and the other is cobalamin-independent.¹ The first class is found in mammalian tissues and the second class in plants, and microorganisms may have either one or both enzymes.

To further our understanding of the catalytic mechanism of methionine synthases, we studied the steric course of the methyl transfer from 5-CH₃-H₄folate to homocysteine catalyzed by the cobalamin-dependent enzyme from *E. coli*. The requisite substrate, 5-CH₃-H₄folate, carrying a chiral methyl group, was prepared by sequential reduction of 5,10-methylenetetrahydrofolate (5,10-CH⁺-H₄folate) with NaB²H₄ in the presence of diethylaniline, known to produce monodeuterated 5,10-methylenetetrahydrofolate (5,10-CH₂-H₄folate) with over 80% stereoselectivity,² followed by further reduction of the 5,10-CH₂-H₄folate with a 2-fold excess of tritiated sodium borohydride in 50 mM Tris buffer, pH 7.5³ (Scheme III). The diastereomer of opposite methyl configuration was generated from 5,10-CH₂-H₄folate produced by NaBH₄ reduction of 11-deuterio-5,10-CH⁺-H₄folate. The 5-CH₃-H₄folate was purified by passage over a column of DEAE Sephadex and elution with a linear gradient of 0.2-2.0 M triethylammonium bicarbonate. Fractions containing 5-CH₃-H₄folate were pooled and concentrated by lyophilization. The two samples of 5-CH₃-H₄folate were degraded by diazotization and KMnO₄ oxidation to give methylamine^{4,5} which by standard procedures⁶ was converted stereospecifically to acetic acid for chirality analysis⁷⁻⁹ of

(1) Matthews, R. G. In *Folates and Pterins: Volume 1 - Chemistry and Biochemistry of Folates*; Blakley, R. L., Benkovic, S. J., Eds.; Wiley: New York, 1984; pp 497-553.

(2) Sliker, L. J.; Benkovic, S. J. *J. Am. Chem. Soc.* **1984**, *106*, 1833-1838.

(3) Nixon, P. F.; Bertino, J. R. *Methods Enzymol.* **1980**, *66*, 547-553.

(4) Asano, Y.; Woodard, R. W.; Houck, D. R.; Floss, H. G. *Arch. Biochem. Biophys.* **1984**, *231*, 253-256.

(5) Gautier, A. E. Ph.D. Dissertation No. 6583, ETH Zürich, 1980.


(6) Woodard, R. W.; Mascaro, L.; Hörhammer, R.; Eisenstein, S.; Floss, H. G. *J. Am. Chem. Soc.* **1980**, *102*, 6314-6318.

(7) Cornforth, J. W.; Redmond, J. W.; Eggerer, H.; Buckel, W.; Gutschow, C. *Eur. J. Biochem.* **1970**, *14*, 1-13.

(8) Lüthy, J.; Rétey, J.; Arigoni, D.; *Nature (London)* **1969**, *221*, 1213-1215.

(9) Floss, H. G.; Tsai, M. D. *Adv. Enzymol.* **1979**, *50*, 243-302.

Scheme III



the methyl group (Scheme I). *F* values⁹ of 37.2 and 37.5 (duplicate analyses) for the acetate from the first sample and 59.8 and 61.4 from the second indicated predominant *S* and *R* configuration, respectively, of the two acetate specimens. Since the degradation sequence involves one inversion of configuration in the cyanide displacement step to give acetonitrile, it follows that the first and the second 5-CH₃-H₄folate sample contained $44 \pm 1\% \text{ ee R}$ and $37 \pm 3\% \text{ ee S}$ isomer, respectively (Scheme III). Hence the reduction of 5,10-CH₂-H₄folate had occurred stereoselectively, exhibiting a 2–3-fold preference for attack syn to C₆-H of the 5,10-CH₂-H₄folate molecule.

Samples of the chiral methyl-*R* and methyl-*S* 5-CH₃-H₄folate were then incubated with purified cobalamin-dependent methionine synthase from *E. coli*¹⁰ in 0.1 M potassium phosphate buffer, pH 7.2, in the presence of 25 mM dithiothreitol, 50 μM aquocobalamin, 0.5 mM homocysteine, and 19 μM *S*-adenosylmethionine to give methionine in about 80% yield. Methionine was separated from residual 5-CH₃-H₄folate by passage over a column of AG1X8 (Bio-Rad) and then purified by HPLC on an ODS column equilibrated with 0.1 M ammonium acetate, pH 3.55. Methionine eluted at 5–7 min,¹¹ and residual ammonium acetate was removed by lyophilization. The purified methionine samples were degraded by the procedure of Arigoni and co-workers^{12,13} (Scheme II) to recover the methyl group as acetate for chirality analysis. This degradation sequence involves two inversions of the methyl group configuration, one in the displacement of methyl from *S*-methylmethionine by *p*-nitrothiobenzoate anion and another in the reaction of methyl *p*-nitrothiobenzoate with cyanide to give acetonitrile. Thus, the configuration of the acetate from the degradation directly reflects that of the methionine methyl group. *F* values of 42.5 and 44.2 (duplicate experiments) for the material derived from methyl-*S*-5-CH₃-H₄folate and 56.3 and 55.7 for that from methyl-*R*-5-CH₃-H₄folate indicated that the product of the enzyme reaction contained methyl groups of predominantly the same configuration as the substrate (Scheme III).

The results demonstrate that the cobalamin-dependent methionine synthase from *E. coli* transfers the methyl group of 5-CH₃-H₄folate stereoselectively to the sulfur of homocysteine to generate methionine with net retention of configuration. The observed steric course is consistent with the postulated mechanism of the reaction, which invokes two sequential transfers of the methyl group, one from 5-CH₃-H₄folate to cobalt to generate enzyme-bound methylcobalamin and a second from cobalt to sulfur to produce methionine. There is a substantial decrease in the chiral purity of the methyl group during the overall process. Since the degradation procedures for both CH₃-H₄folate and methionine are highly stereospecific,^{4,5,12,13} it follows that the enzymatic methyl transfer is accompanied by nearly 50% racemization. This observation may provide a clue for the further mechanistic analysis of the reaction catalyzed by cobalamin-dependent methionine synthase.

Acknowledgment. We thank Kyungok Lee for the careful execution of the chirality analyses of acetate and the National Institutes of Health (Grants GM 32333 to H.G.F., GM 24908 to R.G.M., and GM 24729 to S.J.B. and postdoctoral fellowships ES 05306 to T.M.Z. and GM 08522 to L.F.C.) for financial support.

β-Eliminations in Isonitriles

Beverly A. Jones, Manju Varma, and Charles J. M. Stirling*

Department of Chemistry
University College of North Wales
Bangor, Gwynedd LL57 2UW, United Kingdom

Received January 6, 1986

The physical organic chemistry of the nitrile group has been the subject of intensive investigations¹ which have shown that is

(10) Frasca, V.; Matthews, R. G., unpublished method. The specific activity of the purified enzyme preparations used for these experiments was 0.86–3.72 IU (μmol min⁻¹ mg⁻¹), and the enzyme was about 25–80% pure as assessed by polyacrylamide gel electrophoresis in the presence of sodium dodecyl sulfate.

(11) Daubner, S. C.; Krapf, E.; Matthews, R. G. *Biochemistry of S-Adenosylmethionine and Related Compounds*; Usdin, E., Borchardt, R., Creveling, R., Eds.; McMillan Press: London, 1982; pp 617–620.

(12) Arigoni, D. *Ciba Found. Symp.* 1978, 60, 243–261.

(13) Virchaux, P. N. H. Ph.D. Dissertation No. 6565, ETH Zürich, 1981.

(1) "The Chemistry of the Cyano Group"; Rappoport, Z., Ed.; Wiley: New York, 1970.