

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/24344996>

FvVE1 Regulates Biosynthesis of the Mycotoxins Fumonisin and Fusarins in *Fusarium verticillioides*

ARTICLE in JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY · MAY 2009

Impact Factor: 2.91 · DOI: 10.1021/jf900783u · Source: PubMed

CITATIONS

52

READS

39

7 AUTHORS, INCLUDING:



[Robert Butchko](#)

28 PUBLICATIONS 1,243 CITATIONS

SEE PROFILE



[Hamed K Abbas](#)

United States Department of Agriculture

186 PUBLICATIONS 3,552 CITATIONS

SEE PROFILE

Published in final edited form as:

J Agric Food Chem. 2009 June 10; 57(11): 5089–5094. doi:10.1021/jf900783u.

FvVE1* Regulates Biosynthesis of Fumonisin and Fusarins in *Fusarium verticillioides

KYUNG MYUNG[†], SHAOJIE LI[†], ROBERT A.E. BUTCHKO[‡], MARK BUSMAN[‡], ROBERT H PROCTOR[‡], HAMED K. ABBAS[§], and ANA M. CALVO^{†,*}

[†]Department of Biological Sciences, Northern Illinois University, 1425 W. Lincoln Hwy., Dekalb, Illinois 60115, USA

[‡]Mycotoxin Research Unit, US Department of Agriculture-ARS, 1815 N. University Street, Peoria, Illinois 61604, USA

[§]Crop Genetics and Production Research Unit, US Department of Agriculture-ARS, CG&PRU, P.O. Box 345 Stoneville, Mississippi 38776, USA.

Abstract

The *veA* gene positively regulates sterigmatocystin production in *Aspergillus nidulans* and aflatoxin production in *A. parasiticus* and *A. flavus*. Whether *veA* homologs have a role in regulating secondary metabolism in other fungal genera is unknown. In this study, we examined the role of the *veA* homolog, *FvVE1*, on production of two mycotoxin families, fumonisins and fusarins, in the important corn pathogen *F. verticillioides*. We found that *FvVE1* deletion completely suppressed fumonisin production on two natural substrates, corn and rice. Furthermore, our results revealed that *FvVE1* is necessary for the expression of the pathway-specific regulatory gene *FUM21* and structural genes in the fumonisin biosynthetic gene (*FUM*) cluster. *FvVE1* deletion also blocked production of fusarins. The effects of *FvVE1* deletion on the production of these toxins were found to be the same in two separate mating types. Our results strongly suggest that *FvVE1* play an important role in regulating mycotoxin production in *F. verticillioides*.

Keywords

Fusarium verticillioides; fumonisin; fusarins; *FvVE1*; *veA*; secondary metabolism

INTRODUCTION

The filamentous fungus *Fusarium verticillioides* (syn. *F. moniliforme*, teleomorph *Gibberella moniliformis*) is one of the most common causes of corn ear rot worldwide and can produce multiple families of mycotoxins (1,2). Consequently, *F. verticillioides* mycotoxins are commonly detected in corn (maize, *Zea mays* L.) and often contaminate corn-based human food and animal feed (1,3). Fumonisin are currently considered the most agriculturally significant *F. verticillioides* mycotoxins because they can cause several animal diseases and are epidemiologically associated with some human diseases (4,5). Fumonisin are polyketide-derived metabolites that can inhibit ceramide synthase, a key enzyme in sphingolipid metabolism, and induce apoptosis (1,5). Fumonisin B₁ (FB₁) is typically the most abundant fumonisin in contaminated corn and accounts for approximately 70% of the total fumonisin

*Author to whom correspondence should be addressed [telephone: (815) 753-0451; fax (815) 753-0461; e-mail E-mail: amcalvo@niu.edu].

content. Fumonisin B₂ (FB₂) and fumonisin B₃ (FB₃) are also common in corn but typically comprise 10–20% of the total fumonisin content (1).

In *F. verticillioides*, fumonisin biosynthetic genes (*FUM*) are clustered. The cluster consists of seventeen genes, designated as *FUM1* through *FUM3* and *FUM6* through *FUM21* (6,7,8). Disruption of *FUM1*, *FUM6*, and *FUM8* has been shown to abolish fumonisin production (7, 8). In many cases, genes responsible for the synthesis of fungal secondary metabolites such as sterigmatocystin, gibberellins, aurofusarin, trichothecenes, and lovastatin, are also found clustered and specific regulatory genes are located within these gene clusters (9-11). *FUM21*, a predicted a Zn(II)2Cys6 DNA-binding transcription factor located in the *FUM* cluster, positively regulates *FUM* gene expression and is required for fumonisin synthesis (12). Nevertheless, the regulatory mechanism controlling fumonisin biosynthesis is poorly understood. Among the genes involved in fumonisin gene regulation are *FCCI*, *PAC1*, and *ZFR1* (13-15). *FCCI* encodes a cyclin-like protein (Fcc1) that positively regulates fumonisin biosynthesis and conidiation (13) and interacts with *FCK1*, a cyclin-dependent kinase (Fck1) (16). *PAC1* is required for growth at alkaline pH and may act as a repressor of fumonisin biosynthesis (14). *ZFR1* encodes a zinc binuclear cluster-type protein (Zfr1) which functions as a positive regulator of fumonisin biosynthesis (15). Studies by Flaherty and Woloshuk (15) indicated that Fcc1 is required for Zfr1 function. On the other hand, Pac1 and Fcc1 seem to act independently of each other in regulating fumonisin biosynthesis (15).

In addition to fumonisins, *F. verticillioides* produces other mycotoxins. Among them are the polyketide compounds fusarins (17-19). Fusarins have been reported to induce mutagenesis in mammalian cells *in vitro* (17) and to cause immunosuppression (18). Although a polyketide synthase gene required for fusarin biosynthesis has been identified in several *Fusarium* species (20,21), nothing is known about how fusarin biosynthesis is regulated.

In *Aspergillus* spp., the velvet gene (*veA*) regulates the biosynthesis of several secondary metabolites, including the polyketide toxins sterigmatocystin and aflatoxin (22-24). Whether *veA* homologs have a similar role in regulation of toxin production in other fungal genera has not been investigated. Previously, we identified *FvVE1*, a *veA* homolog in *F. verticillioides*, and demonstrated that it functions in regulation of morphogenesis (25). In this study, we investigate the role of *FvVE1* in secondary metabolism in *F. verticillioides*, specifically in the biosynthesis of fumonisin and fusarins. Our results suggest that *FvVE1* regulates biosynthesis of both fumonisin and fusarins in this important plant pathogenic fungus.

MATERIALS AND METHODS

Strains and media

The strains used in this study are: M-3125 (*MAT1-1*, *FvVE1*); M-3120 (*MAT1-2*, *FvVE1*); M312501 (*MAT1-1*, $\Delta Fvve1::HygB$); M31206 (*MAT1-2*, $\Delta Fvve1::HygB$); M312501C1 (*MAT1-1*, $\Delta Fvve1::HygB$, *FvVE1::GenR*); M31206C5 (*MAT1-2*, $\Delta Fvve1::HygB$, *FvVE1::GenR*). *MAT1-1* and *MAT1-2* are the two different mating type idiomorphs (alleles) in *F. verticillioides*. M-3120 and M-3125 are strain designations from the Fusarium Research Center culture collection (Pennsylvania State University, University Park, PA). The *FvVE1* deletion strains and complementation strains were generated in both mating types as described by Li *et al* (25). In brief, the $\Delta Fvve1$ mutant strains were generated by gene replacement via double homologous recombination events using the hygromycin B resistance gene (*HygB*) as selectable marker (25). Complementation strains were obtained by transformation of the $\Delta Fvve1$ mutants with wild-type *FvVE1* allele using the geneticin resistant gene, *GenR*, as a selectable marker (25).

V8 agar medium (10% V8 juice, 0.1% CaCO₃, and 1.5% agar) was used for production of conidia. Corn medium and rice medium were prepared as previously described (26,27) with some modifications. In this study we mixed 25 g of corn kernels and 40 mL distilled water in 250 mL flasks and 50 g of long-grain rice and 60 mL distilled water in 250 mL flasks. For RNA experiments, cracked corn kernel cultures were prepared by thoroughly mixing 250 g cracked corn kernels and 100 mL water and autoclaving. After cooling, the moistened kernels were combined with 25 mL of a suspension of *F. verticillioides* conidia (1×10^7 conidia per mL water) prepared from seven-day-old, V8 agar cultures of the fungus. This mixture was then distributed among eight 100-mm plastic Petri plates and incubated in the dark at 22 °C.

RNA Preparation and Northern Blots

At 36, 48 and 72 hours of incubation, 10 g of cracked corn culture was frozen in liquid nitrogen, placed at -80 °C until the nitrogen evaporated, and then lyophilized. Total RNA was isolated from the lyophilized material with TRIzol (Invitrogen Life Technologies, Carlsbad, California) using the protocol for samples with high polysaccharide content as described by the manufacturer. For Northern blot analysis, 5 µg of total RNA for each sample was subjected to electrophoresis in a 1.3% agarose gel containing 1.8% formaldehyde and then transferred to nylon membrane following standard protocols (28). ³²P-labeled hybridization probes were prepared with the Ready-to Go DNA labeling kit (Amersham Biosciences, Little Chalfont, Buckinghamshire), and the hybridization, wash, and autoradiography procedures followed standard protocols (28). Templates for hybridization probes corresponding to *FUM1*, *FUM8*, and *TEF1* were prepared by PCR that employed genomic DNA from wild-type *F. verticillioides* strain M-3120 and following primer pairs: The primers used to amplify DNA templates for Northern blot hybridization probes were: for *FUM1*, rp405 (5'-TGGGACACAGTTCTCAAGGAGA-3') and rp408 (5'-CAAGCTCCTGTGACAGAGATAC-3'); for *FUM8*, rp679 (5'-CGTAGTAGGAATGAGAAGGATG-3') and rp680 (5'-GCAAGCTTTGTGGCTGATTGTC-3'); and for *TEF1*, rp992 (5'-ATGGGTAAGGARGACAAGAC-3') and rp993 (5'-GGARGTACCAGTSATCATGTT-3'). *TEF1*, encoding the transcription elongation factor 1α, was used as loading control.

Reverse transcription PCR

Total RNA was treated with Turbo DNA-free DNase (Applied Biosystems, Carlsbad, California) following manufacturers recommended protocol. RNA from DNase treated samples was quantified on a Nanodrop spectrophotometer (Thermo Scientific, Delaware) and diluted to a concentration of 40 ng/µL. A total of 60 ng RNA was used per reverse transcription (RT) PCR reaction. RT-PCR was accomplished with the Easy-A One-tube RT-PCR System (Stratagene, La Jolla, California) following manufacturers recommended protocol. Primers used are: for *TEF1*, rb291 (5'-ATGGGTAAGGAGGACAAGAC-3') and rb292 (5'-GGAAGTACCAGTGATCATGTT-3'); for *FUM21*, rb373 (5'-TAAATGCGAGACAGCATTTGCGGG-3') and rb374 (5'-TGCATCTTGCCCTACTCAATCGGA-3'); for *FUM8*, rb379 (5'-TCCATGTTTACGGGCGCATTTGTC-3') and rb380 (5'-TCGTGAAACCTAGACGCTTGCTGA-3'); for *ZFRI*, rb384 (5'-ATCCACGAAGGAGGCATGTTGGTA-3') and rb385 (5'-AGGCGGATACAAAGAACGACAGGT-3'); and for *FCCI*, rb391 (5'-AATGTTTCCGCTTCCGCA-3') and rb394 (5'-TGCCGCTTCTCCTTAGGTTCT-3'). When possible, primers were designed to amplify different size fragments from genomic DNA and cDNA. *TEF1* was used as control reference to indicate amounts of total RNA. Primers for *TEF1* amplify a 771-bp fragment from genomic DNA and a 324-bp fragment from cDNA. *FUM21* primers amplify 920-bp and 707-bp fragments from genomic DNA and cDNA respectively. *FUM8* primers amplify 789-bp and 638-bp fragments from genomic DNA and

cDNA respectively. Primer pairs for *ZFRI* and *FCCI* amplify the same size fragments from genomic DNA and cDNA (566 bp and 725 bp, respectively due to the absence of introns in these genes).

Fumonisin analysis

A plug (1.6 cm diameter) containing mycelia and conidia from a 7-day old V8 agar culture was used as inoculum. In each case, the cultures were mixed twice during the first three days of incubation by shaking for 30 sec to ensure homogenization. Cultures were incubated for a total period of 2 weeks. After that time, the samples were lyophilized and ground to powder. Fumonisin analysis was performed as previously described by Abbas *et al.* (29). Samples were analyzed by liquid chromatography/electrospray ionization/mass spectrometry (LC/ESI/MS). Samples (10 g) from the ground corn and rice cultures were extracted with 50 mL of 70% methanol and filtered through No. 1 Whatman filter paper. An aliquot (10 mL) was applied to a SAX clean-up column (Varian, Harbor City, CA). The sample was reconstituted in 1 mL of acetonitrile:water (1:1) and diluted if necessary. The LC/ESI/MS analysis was performed on a Thermo Finnigan LCQ Advantage, coupled to a Thermo Finnigan Surveyor MS, and a Thermo Finnigan Surveyor MS Pump (Thermo Electron Corp., West Palm Beach, FL). A 10 μ L aliquot was injected and each sample was evaluated in full-scan mode, using the appropriate mass ranges, fumonisin B₁; 722 (M+H), fumonisin B₂ and B₃; 706 (M+H), fumonisin B₄; 690 (M+H), fumonisin (FA₁, FA₂, FA₃); 764, 748, 748 (M+H), and fumonisin C₁; 708 (M+H). MS/MS was performed on 722 (M+H) for further confirmation of FB₁. The column used for fumonisin analysis was a 3.0 mm \times 150 mm i.d, 5 μ m, Intersil ODS-3 column (MetaChem Technologies Inc., Torrance, CA). The mobile phase at initial elution starting condition consisted of water: 1% acetic acid in methanol (65:35) at 300 μ L per min, followed by water: 1% acetic acid in methanol:methanol (5:35:65) at 10 min. The gradient was held constant for 10 min, and returned to the initial starting conditions for 4 min for column equilibration. Quantitation of FB₁, FB₂, and FB₃ was carried out by the external standard method where FB₄ was calculated as a percentage of FB₁. Other derivatives of fumonisin were monitored for qualitative purposes only.

Fumonisin and fusarins analysis was also carried out on crack-corn cultures used for RNA analysis to further investigate the correlation between fumonisin production and gene expression levels. Ten grams of cracked corn kernel culture were extracted in 25 ml acetonitrile:water (1:1, v/v) on a rotary shaker set at 250 rpm. After 2.5 hours of shaking the mixture was centrifuged at 500 g for 5 minutes and the supernatant was recovered for analysis by reversed phase liquid chromatography-mass spectrometry (LC-MS) in electrospray mode as previously described (30). Briefly, 10 grams of cracked corn kernel culture were extracted in 25 ml acetonitrile:water (1:1, v/v) on a rotary shaker set at 250 rpm. After 2.5 hours of shaking the mixture was centrifuged at 500 g for 5 minutes and the supernatant was recovered for analysis by reversed phase liquid chromatography-mass spectrometry (LC-MS) in electrospray mode. The LC-MS system consisted of a ThermoFinnigan LCQ Deca mass spectrometer coupled to a ThermoSpectraPhysics HPLC with a C18 column. The column used for fumonisin analysis was the same as the one described above (Intersil ODS-3 column MetaChem Technologies Inc., Torrance, CA). Samples were run on a gradient of 35 to 95% (v/v) methanol over 35 minutes at a flow rate of 0.3 ml per minute. Fumonisin and fusarins were detected by monitoring masses of 240 through 1000. The identities of fumonisins and fusarins were confirmed by retention time and the presence of appropriate [M+H]⁺ ions. Quantification was accomplished by comparison of the integrated intensity of ions corresponding to fumonisin and fusarin standards (30).

RESULTS and DISCUSSION

Fumonisin production on natural substrates

Fumonisin is currently considered the most economically important mycotoxin produced by the corn pathogen *F. verticillioides* because of their widespread occurrence in corn and their potential health effects on humans and animals (1,2,4,5). Our current studies indicated that fumonisin production is affected in the $\Delta Fvve1$ mutants. In corn and rice cultures (Figure 1), FB₁ was produced by the wild-type strains as well as by the complemented strains (Figure 2). Trace amounts of FB₂ and FB₃ were also detected in corn and rice cultures of the wild-type and complemented strains. In contrast, neither FB₁, FB₂, nor FB₃ was detected in rice or corn cultures of the $\Delta Fvve1$ mutants (Figure 2). These results were consistent in both mating-type genetic backgrounds. Our findings revealed that the novel regulatory factor *FvVe1* encoded by the *FvVE1* gene is required for fumonisin production when the fungus grows on the natural substrates corn and rice (Figures 1 and 2).

Expression of fumonisin biosynthetic genes

Previous studies showed that VeA, the *FvVe1* homolog in *Aspergillus* species, is required for expression of sterigmatocystin/aflatoxin biosynthetic genes and for concomitant production of the toxins (22-24). However, the possible role of VeA homologs in activation of mycotoxin biosynthetic genes in other fungal genera was not known until now. To investigate whether the expression of fumonisin biosynthetic genes is regulated by the *FvVE1* gene, transcription levels of *FUM1* and *FUM8* (essential fumonisin biosynthetic genes) were examined in the wild-type, $\Delta Fvve1$ mutant and complemented strains grown on cracked-corn medium. The Northern analysis in Figure 3 shows that transcription of *FUM1* was first detected at 48 h after inoculation in both wild-type and complemented strains. *FUM8* expression followed a similar pattern where transcripts started to accumulate slightly earlier in the wild-type strain. However, *FUM1* and *FUM8* transcripts were not detected in the $\Delta Fvve1$ mutant (Figure 3). The lack of fumonisin biosynthetic gene expression in the *FvVE1* deletion mutant is most likely responsible for the lack of fumonisin production in this strain.

Expression of *FUM21*, *FCC1* and *ZFR1*

With the goal of further elucidating the mechanism through which *FvVE1* regulates the expression of the *FUM* gene cluster and the concomitant fumonisin production, we investigated whether the expression of *FUM21* was altered by the *FvVE1* deletion. *FUM21* gene encodes a putative Zn(II)2Cys6 DNA-binding transcriptional activator that is likely specific for *FUM* cluster genes (12). Furthermore, a functional *FUM21* is necessary for fumonisin production in *F. verticillioides*. Genes encoding cluster-specific regulatory proteins have been previously found in other secondary metabolism gene clusters. Well known examples of these regulators are *aflR* and *aflJ*, demonstrated to govern the expression of sterigmatocystin/aflatoxin gene clusters in *Aspergillus* spp. (31-33). We have previously shown that *veA* is necessary for the expression of *aflR* and *aflJ* in *Aspergillus* (22-24). In the present study, in order to investigate whether *FvVE1* plays a role in regulating the expression of *FUM21* we chose RT-PCR analysis because of its ability to detect transcripts that are present at low levels. *FUM21* transcripts were absent in the $\Delta Fvve1$ mutant cultures under conditions that allow expression of this gene in the wild type and complementation strains (Figure 4). *FUM8* was also included in the RT-PCR analysis as an internal control for comparison of the RT-PCR and Northern experiments, which yielded essentially the same results. The absence of *FUM21* transcripts in the $\Delta Fvve1$ mutant indicates that a functional *FvVE1* is necessary for *FUM21* expression.

Genes outside the *FUM* cluster can also regulate fumonisin production in *F. verticillioides*. For example, deletion of the C-type cyclin-like gene, *FCC1*, abolished fumonisin production on corn kernels and in a defined medium at high pH (14). To test whether *FvVE1* controls this

regulatory gene, the transcription levels of *FCC1* were examined in wild-type, $\Delta Fvve1$ mutant and complemented strains grown on cracked-corn medium (Figure 4). Our results showed that *FCC1* expression was very low and only detected in the *FvVE1* deletion mutant at 48 h after inoculation. Although this result differs from those previously reported (13), where *FCC1* expression was detected at higher levels in cracked-corn cultures, under the experimental conditions assayed in our study fumonisin production was detected as well as *FUM1* and *FUM8* expression, indicating that our culture system yielded reliable results. Fumonisin production in the *fcc1* mutant is not blocked in the defined medium at low pH (14). Changes in pH did not rescue fumonisin production in the $\Delta Fvve1$ mutant (data not shown) as in the case of the *FCC1* mutant (14). *FCC1* is required for function of the Zfr1, a putative Zn(II) 2Cys6 transcription factor postulated to control fumonisin production by regulating genes involved in the perception or uptake of carbohydrates (15, 34). In our study, *ZFR1* transcripts were detected in the wild type, $\Delta Fvve1$ mutant and complementation strain, particularly in the $\Delta Fvve1$ mutant where *ZFR1* transcription occurred earlier and was more abundant than in strains with a functional *FvVE1* (Figure 4). This suggests that *FvVE1* negatively influences *ZFR1* expression. Further studies will focus on elucidating possible interactions between *ZFR1* and *FvVE1*, and whether *FvVE1* has a role in carbohydrate metabolism in *F. verticillioides*.

Cultures utilized for RNA studies were also analyzed for the fumonisins. In agreement with previous experiments (Figures 1 and 2), wild-type and complementation strains produced fumonisins but the $\Delta Fvve1$ mutant did not (Table 1). Even when incubation time was increased to 144 hours after inoculation of the crack-corn medium, fumonisin were not detected in $\Delta Fvve1$ cultures.

Fusarin analysis

The VeA homologs in *Aspergillus* regulate not only production of sterigmatocystin and aflatoxin, but also production of other secondary metabolites such as aflatrem, cyclopiazonic acid, and penicillin (22-24). In addition to fumonisins, *F. verticillioides* produces an array of other secondary metabolites, including the mycotoxins fusarins. These toxins have been reported to be mutagenic as well as immunosuppressive (17-19). In order to evaluate the role of *FvVE1* in fusarin biosynthesis, we examined the production of this compound in the wild-type, *FvVE1* deletion mutant and complementation strains on cracked-corn medium. Fusarins were detected in extracts from cultures of wild-type and complementation strains, but were not detected in extracts of $\Delta Fvve1$ cultures (Table 1), indicating that *FvVE1* is also necessary for fusarin biosynthesis in *F. verticillioides*. To our knowledge, this is the first report of a gene described to regulate fusarin production in *F. verticillioides*. Interestingly, a recent report by Estrada and Avalos (35) showed that the white-collar gene *wcoA* modulates fusarin production in a light-dependent manner in *F. fujikuroi*. In *Aspergillus nidulans*, VeA forms a nuclear protein complex that includes light-sensing proteins, such as the red phytochrome-like FphA and the white-collar LreA and LreB proteins responsive to blue light (36). In future studies we will investigate if a similar protein complex that includes *FvVe1* also exists in *F. verticillioides*.

As in the case of *veA* regulation of secondary metabolism in *Aspergillus* (22-24), the effect of *FvVE1* on secondary metabolism in *F. verticillioides* could also be broad. The differences in pigmentation observed in the natural substrate cultures (Figure 1, and data not shown) indicate that the synthesis of other unknown metabolites is also regulated by *FvVE1*.

In conclusion, we demonstrated that *F. verticillioides FvVE1* is required for fumonisin and fusarin production on the natural substrates corn and rice. We also showed that *FvVE1* is necessary for the expression of the fumonisin biosynthetic enzyme-encoding genes *FUM1* and *FUM8* as well as the transcription factor gene *FUM21*. This study also revealed that *FvVE1*

affects production of a second family of *F. verticillioides* secondary metabolites, namely fusarins. The consistent blockage of fumonisin on natural substrates, particularly in corn, suggests that *FvVE1* could be a potential target to control fumonisin contamination of food commodities. We are currently exploring this possibility by investigating whether the *FvVE1* deletion affects the ability of *F. verticillioides* to produce fumonisins in living corn plants.

ACKNOWLEDGMENT

We thank Bobbie Johnson and Jennifer Tonos for their help in analyzing fumonisins and Marcie Moore for the Northern blot analysis. We also thank Dapeng Bao for his technical support. This work was funded by NIH GM074267-01A1.

Abbreviation Used

FB₁, fumonisin B₁; FB₂, fumonisin B₂; FB₃, fumonisin B₃; veA, Velvet gene; FvVE1, *Fusarium verticillioides* veA.

LITERATURE CITED

1. Nelson PE, Desjardins AE, Plattner RD. Fumonisins, mycotoxins produced by *Fusarium* species: biology, chemistry, and significance. *Annu. Rev. Phytopathol* 1993;31:233–252. [PubMed: 18643768]
2. Rheeder JP, Marasas WFO, Vismer HF. Production of fumonisin analogs by *Fusarium* species. *Appl. Environ. Microbiol* 2002;68:2101–2105. [PubMed: 11976077]
3. Bezuidenhout SC, Gelderblom WCA, Gorst-Allman CP, Marthinus Horak R, Marasas WFO, Spiteller G, Vleggaar R. Structure elucidation of the fumonisins, mycotoxins from *Fusarium moniliforme*. *J. Chem. Soc., Chem. Commun* 1988:743–745.
4. Gelderblom WCA, Jaskiewicz K, Marasas WFO, Thiel PG, Horak MJ, Vleggaar R, Kriek NPJ. Fumonisins—novel mycotoxins with cancer-promoting activity produced by *Fusarium moniliforme*. *Appl. Environ. Microbiol* 1988;54:1806–1811. [PubMed: 2901247]
5. Desai K, Sullards MC, Allegood J, Wang E, Schmelz EM, Hartl M, Humpf HU, Liotta DC, Peng Q, Merrill AH Jr. Fumonisins and fumonisin analogs as inhibitors of ceramide synthase and inducers of apoptosis. *Biochim. Biophys. Acta* 2002;1585:188–192. [PubMed: 12531553]
6. Proctor RH, Brown DW, Plattner RD, Desjardins AE. Co-expression of 15 contiguous genes delineates a fumonisin biosynthetic gene cluster in *Gibberella moniliformis*. *Fungal Genet. Biol* 2003;38:237–249. [PubMed: 12620260]
7. Bojja RS, Cerny RL, Proctor RH, Du L. Determining the biosynthetic sequence in the early steps of the fumonisin pathway by use of three gene-disruption mutants of *Fusarium verticillioides*. *J. Agric. Food Chem* 2004;52:2855–2860. [PubMed: 15137825]
8. Seo JA, Proctor RH, Plattner RD. Characterization of four clustered and coregulated genes associated with fumonisin biosynthesis in *Fusarium verticillioides*. *Fungal Genet. Biol* 2001;34:155–165. [PubMed: 11728154]
9. Hohn TM, Krishna R, Proctor RH. Characterization of a transcriptional activator controlling trichothecene toxin biosynthesis. *Fungal Genet. Biol* 1999;26:224–235. [PubMed: 10361036]
10. Kennedy J, Auclair K, Kendrew SG, Park C, Vederas JC, Hutchinson CR. Modulation of polyketide synthase activity by accessory proteins during lovastatin biosynthesis. *Science* 1999;284:1368–1372. [PubMed: 10334994]
11. Malz S, Grell MN, Thrane C, Maier FJ, Rosager P, Felk A, Albertsen KS, Salomon S, Bohn L, Schafer W, Giese H. Identification of a gene cluster responsible for the biosynthesis of aurofusarin in the *Fusarium graminearum* species complex. *Fungal Genet. Biol* 2005;42:420–433. [PubMed: 15809006]
12. Brown DW, Butchko RAE, Busman M, Proctor RH. The *Fusarium verticillioides* *FUM* Gene Cluster Encodes a Zn(II)2Cys6 Protein That Affects *FUM* Gene Expression and Fumonisins. *Euk. Cell* 2007;6:1210–1218.

13. Shim WB, Woloshuk CP. Regulation of fumonisin B₁ biosynthesis and conidiation in *Fusarium verticillioides* by a cyclin-like (C-type) gene, *FCC1*. Appl. Environ. Microbiol 2001;67:1607–1612. [PubMed: 11282612]
14. Flaherty JE, Pirttilä AM, Bluhm BH, Woloshuk CP. *PAC1*, a pH-regulatory gene from *Fusarium verticillioides*. Appl. Environ. Microbiol 2003;69:5222–5227. [PubMed: 12957906]
15. Flaherty JE, Woloshuk CP. Regulation of fumonisin biosynthesis in *Fusarium verticillioides* by a zinc binuclear cluster-type gene, *ZFRI*. Appl. Environ. Microbiol 2004;70:2653–2659. [PubMed: 15128515]
16. Bluhm BH, Woloshuk CP. Fck1, a C-type cyclin-dependent kinase, interacts with Fcc1 to regulate development and secondary metabolism in *Fusarium verticillioides*. Fungal Genet. Biol 2006;43:146–154. [PubMed: 16504555]
17. Cheng SJ, Jiang YZ, Li MH, Lo HZ. A mutagenic metabolite produced by *Fusarium moniliforme* isolates in Linxian County, China. Carcinogenesis 1985;6:903–905. [PubMed: 4006077]
18. Dong ZY, Zhan YH. Inhibitory effect of a mycotoxin, fusarin C, on macrophage activation and macrophage mediated cytotoxicity to tumor cells in mice. J. Exp. Clin. Cancer Res 1987;6:31–38.
19. Desjardins AE, Proctor RH. Molecular biology of *Fusarium* mycotoxins. Int. J. Food Microbiol 2007;119:47–50. [PubMed: 17707105]
20. Gaffoor I, Brown DW, Plattner R, Proctor RH, Qi W, Trail F. Functional analysis of the polyketide synthase genes in the filamentous fungus *Gibberella zeae* (anamorph *Fusarium graminearum*). Eukaryot. Cell 2005;4:1926–1933. [PubMed: 16278459]
21. Song ZS, Cox RJ, Lazarus CM, Simpson TJ. Fusarin C biosynthesis in *Fusarium moniliforme* and *Fusarium venenatum*. ChemBioChem 2004;5:1196–1203. [PubMed: 15368570]
22. Kato N, Brooks W, Calvo AM. The expression of sterigmatocystin and penicillin genes in *Aspergillus nidulans* is controlled by *veA*, a gene required for sexual development. Eukaryot. Cell 2003;2:1178–1186. [PubMed: 14665453]
23. Calvo AM, Bok J, Brooks W, Keller NP. *veA* is required for toxin and sclerotial production in *Aspergillus parasiticus*. Appl. Environ. Microbiol 2004;70:4733–4739. [PubMed: 15294809]
24. Duran RM, Cary JW, Calvo AM. Production of cyclopiazonic acid, aflatoxin, and aflatoxin by *Aspergillus flavus* is regulated by *veA*, a gene necessary for sclerotial formation. Appl. Microbiol. Biotechnol 2007;73:1158–1168. [PubMed: 16988822]
25. Li S, Myung K, Guse D, Donkin B, Proctor RH, Grayburn WS, Calvo AM. FvVE1 regulates filamentous growth, the ratio of microconidia to macroconidia and cell wall formation in *Fusarium verticillioides*. Mol Microbiol 2006;62:1418–32. [PubMed: 17054442]
26. Abbas HK, Mirocha CJ, Shier WT. Mycotoxins produced from fungi isolated from foodstuffs and soil: Comparison of toxicity in fiberblasts and rat feeding test. Appl. Environ. Microbiol 1984;48:654–661. [PubMed: 6541890]
27. Abbas HK, Cartwright RD, Xie W, Mirocha CJ, Richard JL, Dvorak TJ, Sciumbato GL, Shier WT. Mycotoxins production by *Fusarium proliferatum* isolates from rice with *Fusarium* sheath rot disease. Mycopathologia 1999;147:97–104. [PubMed: 10967968]
28. Sambrook, J.; Fritsch, EF.; Maniatis, T. Molecular Cloning: A Laboratory Manual. Vol. 2nd Ed.. Cold Spring Harbor Laboratory Press; Cold Spring Harbor, NY:
29. Abbas HK, Cartwright RD, Xie W, Shier WT. Aflatoxin and fumonisin contamination of corn (maize, *Zea mays*) hybrids in Arkansas. Crop Protection 2006;25:1–9.
30. Plattner RD, Weisleder D, Poling SM. Analytical determination of fumonisins and other metabolites produced by *Fusarium moniliforme* and related species on corn. Adv. Exp. Med. Biol 1996;392:57–64. [PubMed: 8850605]
31. Yu JH, Butchko RA, Fernandes M, Keller NP, Leonard TJ, Adams TH. Conservation of structure and function of the aflatoxin regulatory gene *aflR* from *Aspergillus nidulans* and *A. flavus*. Curr. Genet 1996;29:549–555. [PubMed: 8662194]
32. Payne GA, Nystrom GJ, Bhatnagar D, Cleveland TE, Woloshuk CP. Cloning of the *afl-2* gene involved in aflatoxin biosynthesis from *Aspergillus flavus*. Appl. Environ. Microbiol 1993;59:156–162. [PubMed: 8439147]
33. Chang PK. The *Aspergillus parasiticus* protein AFLJ interacts with the aflatoxin pathway-specific regulator AFLR. Mol. Genet. Genomics 2003;268:711–719. [PubMed: 12655397]

34. Bluhm BH, Kim H, Butchko RA, Woloshuk CP. Involvement of ZFR1 of *Fusarium verticillioides* in kernel colonization and the regulation of FST1, a putative sugar transporter gene required for fumonisin biosynthesis on maize kernels. *Mol. Plant Pathol* 2008;9:203–211. [PubMed: 18705852]
35. Estrada AF, Avalos J. The White Collar protein WcoA of *Fusarium fujikuroi* is not essential for photocarotenogenesis, but is involved in the regulation of secondary metabolism and conidiation. *Fungal Genet. Biol* 2008;45:705–718. [PubMed: 18203635]
36. Calvo AM. The VeA regulatory system and its role in morphological and chemical development in fungi. *Fungal Genet. Biol.* 2008;45:1053–1061. [PubMed: 18457967]




Figure 1. Photographs of wild-type (WT, M-3120 and M-3125), *FvVE1* deletion mutant ($\Delta Fvve1$, M31206 and M312501) and complemented (Com., M31206C5 and M312501C1) strains of *MAT1-1* and *MAT1-2* respectively, in corn and rice cultures.





Figure 2.

Production of fumonisins in wild-type (WT), *FvVE1* deletion mutant ($\Delta Fvve1$) and complemented (Com) strains of *MAT1-1* and *MAT1-2* in corn (A) and rice (B) cultures. Total fumonisins (B_1 , B_2 and B_3 combined) were analyzed at 2 weeks after inoculation. Bars indicate standard errors of 3 independent cultures. N.D. = not detected.

**Figure 3.**

Northern analysis of *FUM1*, and *FUM 8* gene expression from wild-type M-3120, *FvVE1* deletion mutant M31206 ($\Delta Fvve1$), and complementation (Com) strain M31206C5. Total RNA was isolated from mycelial tissue grown on cracked-corn medium at 36, 48 and 72 h after inoculation. *TEF1* was used as loading control.




Figure 4.

RT-PCR analysis of *FUM21*, *FUM 8*, *ZFR1* and *FCC1* gene expression from wild-type M-3120, *FvVE1* deletion mutant M31206 ($\Delta Fvve1$), and complementation (Com) strain M31206C5. Total RNA was isolated from mycelial tissue grown on cracked-corn medium at 36, 48 and 72 h after inoculation. *TEF1* was used as loading control. A no reverse transcriptase control reaction is shown for *TEF1* primers indicating no genomic DNA remained after DNase treatment.

Table 1
Fumonisin and Fusarin Analysis of Cracked-corn Cultures.

Time	Strain	Fumonisin ^a	Fusarins ^b
36 hours	Wild type	0	0
	$\Delta FvveI$	0	0
	Comp	0	196
48 hours	Wild type	1	0
	$\Delta FvveI$	0	0
	Comp	0	864
72 hours	Wild type	98	330
	$\Delta FvveI$	0	0
	Comp	127	1709
96 hours	Wild type	178	673
	$\Delta FvveI$	0	0
	Comp	316	2066
144 hours	Wild type	281	1945
	$\Delta FvveI$	0	0
	Comp	334	2246

^a μg fumonisins B₁, B₂ and B₃ combined per gram of cracked corn culture

^b μg fusarins C₁, C₂ and C₃ combined per gram of cracked corn culture