

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7801705>

# New Dammarane-Type Saponins from the Galls of *Sapindus mukorossi*

ARTICLE in JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY · JULY 2005

Impact Factor: 2.91 · DOI: 10.1021/jf047963s · Source: PubMed

CITATIONS

25

READS

83

7 AUTHORS, INCLUDING:



**Yao-Haur Kuo**

National Research Institute of Chinese Medic...

219 PUBLICATIONS 3,461 CITATIONS

SEE PROFILE



**Kuo-Hsiung Lee**

University of North Carolina at Chapel Hill

566 PUBLICATIONS 12,536 CITATIONS

SEE PROFILE



**Fang-Rong Chang**

Kaohsiung Medical University

325 PUBLICATIONS 5,236 CITATIONS

SEE PROFILE



**Yang-Chang Wu**

China Medical University (ROC)

482 PUBLICATIONS 7,487 CITATIONS

SEE PROFILE

New Dammarane-Type Saponins from the Galls of *Sapindus mukorossi*YAO-HAUR KUO,<sup>†</sup> HUI-CHI HUANG,<sup>‡,§</sup> LI-MING YANG KUO,<sup>†</sup> YA-WEN HSU,<sup>†</sup>  
KUO-HSIUNG LEE,<sup>#</sup> FANG-RONG CHANG,<sup>‡</sup> AND YANG-CHANG WU<sup>\*,‡</sup>

National Research Institute of Chinese Medicine, Taipei 112, Taiwan, Republic of China; Graduate Institute of Natural Products and Graduate Institute of Pharmaceutical Sciences, Kaohsiung Medical University, Kaohsiung 807, Taiwan, Republic of China; and Natural Products Laboratory, Division of Medicinal Chemistry and Natural Products, School of Pharmacy, University of North Carolina, Chapel Hill, North Carolina 27599

Five new dammarane-type saponins, 3 $\beta$ ,7 $\beta$ ,20(S),22-tetrahydroxydammar-24-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside, 3 $\beta$ ,7 $\beta$ ,20(S),22,23-pentahydroxydammar-24-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside, 3 $\beta$ ,7 $\beta$ ,20(S),22,25-pentahydroxydammar-23-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside, 25-methoxy-3 $\beta$ ,7 $\beta$ ,20(S),22-tetrahydroxydammar-23-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside, and 25-methoxy-3 $\beta$ ,7 $\beta$ ,20(R)-trihydroxydammar-23-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside, named sapinmusaponins A (1), B (2), C (3), D (4), and E (5), respectively, together with three known phenylpropanoid glycosides (6–8), were isolated from the galls of *Sapindus mukorossi*. The structures of these saponins were elucidated on the basis of spectroscopic analyses and chemical methods. Preliminary bioassay data revealed that saponins 1 and 3–5 showed moderate cytotoxic activity (ED<sub>50</sub> ~ 9–18  $\mu$ g/mL) against human tumor cell lines (Hepa59T/VGH, NCI, HeLa, and Med) and that 1–5 were inactive in vitro against HIV replication in H9 lymphocytes.

**KEYWORDS:** Dammarane; saponin; sapinmusaponins A–E; *Sapindus mukorossi*; Sapindaceae; cytotoxicity

## INTRODUCTION

*Sapindus mukorossi* (Sapindaceae), also known as soap-nut tree, is an important economic agricultural product in tropical and subtropical regions of Asia. The pericarp of *S. mukorossi* has been traditionally used as an expectorant, as well as a source of natural surfactants (1–3). Previous reports on *S. mukorossi* dealt with the isolation and identification of triterpenoids, saponins (2, 4, 5), fatty acids (6), and flavonoids (7), from the pericarp, stem, and fruit of the plant. Other literature reports discussed the biological activities, including molluscicidal (8), anti-inflammatory (2), and cytotoxic (9) effects, for this plant. We reported previously that extracts of *S. mukorossi* showed molluscicidal effects against the golden apple snail and isolated a series of hederagenin-based acetylated saponins from the plant pericarp. Recently, we found the ethanol extract of galls from the stems of *S. mukorossi* exhibited marginal cytotoxicity against

human liver carcinoma (Hepa59T/VGH), human large lung cell carcinoma (NCI), cervical epitheloid carcinoma (HeLa), and medulloblastoma (Med) tumor cells. Usually, galls grow on the stems or leaves of the tree and are caused by parasite organisms such as insects, mites, mistletoes, fungi, and bacteria (10). Different kinds of galls occur in many plant families, including Anacardiaceae, Fagaceae, Leguminosae, Myrtaceae, Lauraceae, and Fagaceae (11). Wu-Bei-Zi, a gall that grows on *Rhus chinensis* (Anacardiaceae), is a famous Chinese medicine, used as an astringent, insecticide, anti-inflammatory agent, and antibacterial drug (12, 13). However, the chemical constituents and pharmacological effects of the gall of *S. mukorossi* have not been reported. Thus, we were prompted to investigate the bioactive constituents from collected galls. We report herein on the isolation and structural elucidation of 1–5 using spectroscopic analyses, including 1D and 2D NMR techniques (COSY, HMQC, HMBC, and NOESY), and chemical methods. Biological evaluation of the newly isolated saponins in cytotoxicity and anti-HIV assays is also described.

## MATERIALS AND METHODS

**General Experimental Procedures.** Infrared (IR) spectra were measured on a Mattson Genesis II spectrophotometer (Thermo Nicolet, Madison, WI) using a KBr matrix. FABMS data were obtained on a

\* Author to whom correspondence should be addressed (telephone +886-2-28201999, ext. 7051, or +886-7-3121101, ext. 2197; fax +886-2-28236150 or +886-7-3114773; e-mail kuoyh@nricm.edu.tw or yachwu@kmu.edu.tw).

<sup>†</sup> National Research Institute of Chinese Medicine.

<sup>‡</sup> Graduate Institute of Natural Products, Kaohsiung Medical University.

<sup>§</sup> Graduate Institute of Pharmaceutical Sciences, Kaohsiung Medical University.

<sup>#</sup> University of North Carolina.

**Table 1.** Selected  $^1\text{H}$  NMR Data for the Aglycon Moieties of Compounds 1–5<sup>a</sup> ( $\delta$ , in  $\text{CD}_3\text{OD}$ )

| no.            | 1                           | 2                   | 3                                       | 4                   | 5                   |
|----------------|-----------------------------|---------------------|---|---------------------|---------------------|
| 1              | 1.66 m, 0.93 m              | 1.66 m, 0.94 m      | 1.63 m, 0.94 m                          | 1.61 m, 0.93 m      | 1.63 m, 0.96 m      |
| 2              | 1.98 m, 1.66 m              | 1.98 m, 1.66 m      | 1.98 m, 1.66 m                          | 1.98 m, 1.66m       | 1.90 m, 1.65 m      |
| 3              | 3.17 dd (11.2, 4.0)         | 3.17 dd (11.8, 4.0) | 3.17 dd (11.2, 4.0)                     | 3.16 m              | 3.17 dd (11.0, 4.0) |
| 5              | 0.81 m                      | 0.81 m              | 0.81 m                                  | 0.81 m              | 0.81 m              |
| 6              | 1.63 m, 1.55 m              | 1.63 m, 1.52m       | 1.63 m, 1.55 m                          | 1.63 m, 1.55 m      | 1.62 m, 1.55 m      |
| 7              | 3.70 dd (10.0, 4.8)         | 3.72 d (4.8)        | 3.70 m                                  | 3.70 m              | 3.69 dd (11.2, 4.4) |
| 9              | 1.24 m                      | 1.25 m              | 1.22 m                                  | 1.22 m              | 1.24 m              |
| 11             | 1.52 m, 1.32m               | 1.49 m, 1.31m       | 1.53 m, 1.32 m                          | 1.52 m, 1.31 m      | 1.52 m, 1.30 m      |
| 12             | 1.71 m, 1.35 m              | 1.68 m, 1.32 m      | 1.71 m, 1.33 m                          | 1.70 m, 1.33 m      | 1.68 m, 1.33 m      |
| 13             | 1.75 m                      | 1.73m               | 1.73 m                                  | 1.73 m              | 1.70 m              |
| 15             | 1.66 m, 1.35 m              | 1.66 m, 1.35 m      | 1.69 m, 1.36 m                          | 1.69 m, 1.34 m      | 1.67 m, 1.35 m      |
| 16             | 1.98 m, 1.27 m              | 1.98 m, 1.25 m      | 1.98 m, 1.26 m                          | 1.99 m, 1.26 m      | 1.96 m, 1.26 m      |
| 17             | 1.92 m                      | 2.14 dd (4.0, 10.0) | 2.01 m                                  | 2.01 m              | 1.98 m              |
| 18             | 0.97 s                      | 0.98 s              | 0.97 s                                  | 0.97 s              | 0.97 s              |
| 19             | 0.86 s                      | 0.86 s              | 0.85 s                                  | 0.85 s              | 0.85 s              |
| 21             | 1.05 s                      | 1.15 s              | 0.98 s                                  | 0.99 s              | 1.11 s              |
| 22             | 3.41 m                      | 3.15 d (2.4)        | 3.82 m/4.28 <sup>c</sup> d (7.2)        | 3.87 d (7.2)        | 2.23 m, 2.22 m      |
| 23             | 2.25 dd (14.8, 6.8), 2.05 m | 4.67 dd (8.8, 2.0)  | 5.57 m/6.48 <sup>c</sup> dd (15.6, 7.2) | 5.76 dd (16.0, 7.2) | 5.68 dt (16.0, 7.2) |
| 24             | 5.24 t (6.8)                | 5.43 dt (8.8, 1.2)  | 5.74 m/6.29 <sup>c</sup> d (15.6)       | 5.64 d (16.0)       | 5.45 d (16.0)       |
| 26             | 1.70 s                      | 1.73 s              | 1.28 s                                  | 1.27 s              | 1.25 s              |
| 27             | 1.61 s                      | 1.70 s              | 1.28 s                                  | 1.27 s              | 1.25 s              |
| 28             | 1.05 s                      | 1.04 s              | 1.04 s                                  | 1.04 s              | 1.04 s              |
| 29             | 0.85 s                      | 0.85 s              | 0.86 s                                  | 0.86 s              | 0.86 s              |
| 30             | 0.93 s                      | 0.94 s              | 0.94 s                                  | 0.94 s              | 0.91 s              |
| $\text{OCH}_3$ |                             |                     |   | 3.16 s              | 3.14 s              |

<sup>a</sup> Assignments confirmed by decoupling,  $^1\text{H}$ – $^1\text{H}$  COSY, TOCSY, NOESY, HMQC, and HMBC. <sup>b</sup> Overlapped signal. <sup>c</sup> Chemical shifts and coupling constants were measured in  $\text{C}_5\text{D}_5\text{N}$ .

JEOL SX-102A instrument (JEOL USA, Inc., Peabody, MA). High-resolution FABMS were measured on a Finnigan/Thermo Quest MAT mass spectrometer (Scientific Instrument Services, Inc., Ringoes, NJ). NMR spectra were performed on Bruker NMR spectrometers (Unity Plus 400 MHz) (Bruker BioSpin GmbH, Rheinstetten, Germany) using  $\text{CD}_3\text{OD}$  and  $\text{C}_5\text{D}_5\text{N}$  as solvent for measurement. Diaion HP-20 (Mitsubishi Chemical Co., Tokyo, Japan), Sephadex LH-20, and silica gel (Merck 70–230 and 230–400 mesh) (Merck & Co., Inc., Whitehouse Station, NJ) were used for column chromatography, and precoated silica gel (Merck 60 F-254) plates were used for TLC. The spots on TLC were detected by spraying with 50%  $\text{H}_2\text{SO}_4$  and then heating at 100 °C. HPLC separations were performed on a Shimadzu LC-6AD series apparatus with an RID-10A refractive index, equipped with a  $250 \times 20$  mm i.d. preparative Cosmosil 5SL-II column (TimTec, Inc., Newark, DE).

**Plant Materials.** The galls of *S. mukorossi* were collected in October 2001 in Taipei County, Taiwan, and identified by Professor Muh-Tsuen Kao of the National Institute of Chinese Medicine.

**Extraction and Isolation.** The dried galls of *S. mukorossi* (8.5 kg) were extracted three times with ethanol (40 L). Removal of solvent in a vacuum gave the ethanol extract, which was partitioned with  $\text{MeOH}/\text{CHCl}_3/\text{H}_2\text{O}$  (7:10:3, 1000 mL  $\times$  3) to give  $\text{CHCl}_3$  and  $\text{H}_2\text{O}$  layers. The  $\text{CHCl}_3$  layer (190 g) was partitioned with methanol/*n*-hexane (1:3, 800 mL  $\times$  3) to give methanol and *n*-hexane layers. After the evaporation of methanol layer in a vacuum, the residue (100 g) was purified by chromatography on a silica gel column (41  $\times$  10.5 cm) with a stepwise gradient of  $\text{CHCl}_3/\text{MeOH}$  (1:0, 40:1, 30:1, 20:1, 10:1, 8:1, 6:1, 4:1, 2:1, 1:1, 0:1, each 1 L) to afford 13 fractions (fr 1–13). Fraction 12 (18.8 g) was chromatographed on a Diaion HP-20 porous polymer resin column (23  $\times$  4.5 cm), eluting with 10, 40, 60, 80, and 100% MeOH (each 2 L), respectively, to yield 5 fractions (fr 12.1–12.5). Fraction 12.3 (3.0 g) was further separated by chromatography on a Sephadex LH-20 column (30  $\times$  2.8 cm) with MeOH (1.5 L) to yield 5 fractions (fr 12.3.1–12.3.5). Fraction 12.3.2 (2.0 g) was rechromatographed on a Sephadex LH-20 column (40  $\times$  1.5 cm) with MeOH (1.0 L), and 4 fractions (fr 12.3.2.1–12.3.2.4) were obtained. Fraction 12.3.2.2 (1.1 g) was further purified by HPLC on an ODS column (250  $\times$  20.0 mm i.d., flow rate = 5 min/mL, with 80% MeOH to afford **1** (17.0 mg), **4** (6.0 mg), and **5** (21.0 mg). Using the same column as that of fr 12.3.2.2 on HPLC with 75% MeOH, **6** (0.5 mg), **7** (1.3 mg), and **8** (1.0 mg) were isolated from fr 12.3.3 (0.3 g), and **2**

(23.0 mg) and **3** (4.0 mg) were yielded from fr 12.3.4 (0.7 g) eluting with 60% MeOH.

**3 $\beta$ ,7 $\beta$ ,20(S),22-Tetrahydroxydammar-24-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside (**1**)** was obtained as a white amorphous powder: mp 169–171 °C;  $[\alpha]_{\text{D}}^{24} +50.0^\circ$  (*c* 0.3, MeOH); IR  $\nu_{\text{max}}$  (KBr) 3387, 1629, 1047  $\text{cm}^{-1}$ ; FABMS,  $m/z$  807  $[\text{M} + \text{Na}]^+$ ; HRFABMS,  $m/z$  807.4864 (calcd, 807.4870,  $\text{C}_{42}\text{H}_{72}\text{O}_{13}\text{Na}$ ).  $^1\text{H}$  and  $^{13}\text{C}$  NMR data are shown in **Tables 1–3**.

**3 $\beta$ ,7 $\beta$ ,20(S),22,23-Pentahydroxydammar-24-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside (**2**)** was obtained as a white amorphous powder: mp 254–256 °C;  $[\alpha]_{\text{D}}^{24} +22.0^\circ$  (*c* 0.3, MeOH); IR  $\nu_{\text{max}}$  (KBr) 3389, 1640, 1047  $\text{cm}^{-1}$ ; HRFABMS,  $m/z$  823.4813 (calcd 823.4819,  $\text{C}_{42}\text{H}_{72}\text{O}_{14}\text{Na}$ ).  $^1\text{H}$  and  $^{13}\text{C}$  NMR data are shown in **Tables 1–3**.

**3 $\beta$ ,7 $\beta$ ,20(S),22,25-Pentahydroxydammar-23-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside (**3**)** was obtained as a white amorphous powder: mp 198–200 °C;  $[\alpha]_{\text{D}}^{24} +66.0^\circ$  (*c* 0.3, MeOH); IR  $\nu_{\text{max}}$  (KBr) 3382, 1641, 1045  $\text{cm}^{-1}$ ; HRFABMS,  $m/z$  823.4824 (calcd 823.4819,  $\text{C}_{42}\text{H}_{72}\text{O}_{14}\text{Na}$ ).  $^1\text{H}$  and  $^{13}\text{C}$  NMR data are shown in **Tables 1–3**.

**25-Methoxy-3 $\beta$ ,7 $\beta$ ,20(S),22-tetrahydroxydammar-23-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside (**4**)** was obtained as a white amorphous powder: mp 189–191 °C;  $[\alpha]_{\text{D}}^{24} +46.6^\circ$  (*c* 0.4, MeOH); IR  $\nu_{\text{max}}$  (KBr) 3394, 1654, 1048  $\text{cm}^{-1}$ ; HRFABMS,  $m/z$  837.4982 (calcd 837.4976,  $\text{C}_{43}\text{H}_{74}\text{O}_{14}\text{Na}$ ).  $^1\text{H}$  and  $^{13}\text{C}$  NMR data are shown in **Tables 1–3**.

**25-Methoxy-3 $\beta$ ,7 $\beta$ ,20(R)-trihydroxydammar-23-ene-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside (**5**)** was obtained as a white amorphous powder: mp 190–192 °C;  $[\alpha]_{\text{D}}^{24} +10.2^\circ$  (*c* 0.3, MeOH); IR  $\nu_{\text{max}}$  (KBr) 3409, 1597, 1049  $\text{cm}^{-1}$ ; HRFABMS,  $m/z$  821.5016 (calcd 821.5027,  $\text{C}_{43}\text{H}_{74}\text{O}_{13}\text{Na}$ ).  $^1\text{H}$  and  $^{13}\text{C}$  NMR data are shown in **Tables 1–3**.

**4-Allyl-2-methoxyphenyl-6-O- $\beta$ -D-apiosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucoside (**6**)** was obtained as an amorphous powder; IR  $\nu_{\text{max}}$  (KBr) 3370, 1640, 1599, 1500  $\text{cm}^{-1}$ ; FABMS,  $m/z$  481  $[\text{M} + \text{Na}]^+$  (**14**).

**4-Allyl-2-methoxyphenyl-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (**7**)** was obtained as a white amorphous powder: IR  $\nu_{\text{max}}$  (KBr) 3450, 1630, 1600, 1490  $\text{cm}^{-1}$ ; FABMS,  $m/z$  495  $[\text{M} + \text{Na}]^+$  (**15**).

**Table 2.**  $^{13}\text{C}$  and  $^1\text{H}$  NMR Data for the Sugar Moieties of Compounds 1–5 ( $\delta$ , in  $\text{CD}_3\text{OD}$ )

| no. | 1                   |                             | 2                   |                             | 3                   |                             | 4                   |                             | 5                   |                             |
|-----|---------------------|-----------------------------|---------------------|-----------------------------|---------------------|-----------------------------|---------------------|-----------------------------|---------------------|-----------------------------|
|     | $\delta_{\text{C}}$ | $\delta_{\text{H}}$ (J, Hz) | $\delta_{\text{C}}$ | $\delta_{\text{H}}$ (J, Hz) | $\delta_{\text{C}}$ | $\delta_{\text{H}}$ (J, Hz) | $\delta_{\text{C}}$ | $\delta_{\text{H}}$ (J, Hz) | $\delta_{\text{C}}$ | $\delta_{\text{H}}$ (J, Hz) |
| 1'  | 105.57              | 4.39 d (7.6)                | 105.56              | 4.39 d (7.6)                | 105.59              | 4.39 d (7.3)                | 105.56              | 4.40 d (7.6)                | 105.58              | 4.40 d (7.2)                |
| 2'  | 78.96               | 3.39 m                      | 78.97               | 3.39 dd (8.4, 7.6)          | 78.76               | 3.39 m                      | 78.94               | 3.39 m                      | 78.93               | 3.39 d (8.0, 7.2)           |
| 3'  | 79.50               | 3.45 t (8.8)                | 79.48               | 3.43 t (8.4)                | 79.53               | 3.44 t (8.4)                | 79.48               | 3.45 t (8.4)                | 79.51               | 3.45 t (8.0)                |
| 4'  | 72.17               | 3.27 t (8.8)                | 72.15               | 3.28 t (8.4)                | 72.15               | 3.26 t (8.4)                | 72.15               | 3.27 t (8.4)                | 72.04               | 3.27 t (8.0)                |
| 5'  | 77.61               | 3.23 m                      | 77.58               | 3.23 dd (5.6, 2.0)          | 77.62               | 3.23 m                      | 77.58               | 3.23 dd (5.2, 2.0)          | 77.61               | 3.23 dd (5.2, 2.0)          |
| 6'  | 62.79               | 3.83 dd (11.6, 1.8)         | 62.76               | 3.84 dd (12.0, 2.0)         | 62.78               | 3.83 dd (11.0, 1.2)         | 62.76               | 3.84 dd (12.0, 2.0)         | 62.78               | 3.84 dd (12.0, 2.0)         |
|     |                     | 3.65 dd (11.6, 5.2)         |                     | 3.65 dd (12.0, 5.6)         |                     | 3.64 dd (11.0, 5.6)         |                     | 3.65 dd (12.0, 5.2)         |                     | 3.65 dd (12.0, 5.2)         |
| 1'' | 101.85              | 5.35 br s                   | 101.85              | 5.35 d (1.2)                | 101.85              | 5.35 br s                   | 101.84              | 5.35 d (1.6)                | 101.84              | 5.36 d (1.6)                |
| 2'' | 72.08               | 3.95 m                      | 72.03               | 3.94 dd (3.2, 1.6)          | 72.06               | 3.94 m                      | 72.02               | 3.94 dd (2.8, 1.6)          | 72.13               | 3.94 dd (3.2, 1.6)          |
| 3'' | 72.04               | 3.74 dd (9.2, 3.6)          | 72.03               | 3.74 dd (9.6, 3.2)          | 72.06               | 3.72 dd (9.2, 3.2)          | 72.02               | 3.72 dd (9.2, 3.2)          | 72.08               | 3.74 dd (9.6, 3.2)          |
| 4'' | 73.97               | 3.35 t (9.2)                | 73.95               | 3.37 t (9.6)                | 73.97               | 3.35 t (9.2)                | 73.95               | 3.35 t (9.2)                | 73.96               | 3.35 t (9.6)                |
| 5'' | 69.98               | 3.98 m                      | 69.98               | 3.96 m                      | 69.98               | 3.97 m                      | 69.97               | 3.96 dd (6.0, 3.2)          | 69.96               | 3.97 dd (6.4, 3.2)          |
| 6'' | 17.98               | 1.19 d (6.4)                | 17.98               | 1.19 d (6.0)                | 17.98               | 1.20 d (6.4)                | 17.98               | 1.20 d (6.0)                | 17.97               | 1.21 d (6.4)                |

**Table 3.**  $^{13}\text{C}$  NMR Data for the Aglycon Moieties of Compounds 1–5 ( $\delta$ , in  $\text{CD}_3\text{OD}$ )

| no.            | 1      | 2      | 3      | 4      | 5      |
|----------------|--------|--------|--------|--------|--------|
| 1              | 40.40  | 40.40  | 40.41  | 40.39  | 40.39  |
| 2              | 27.45  | 27.45  | 27.46  | 27.44  | 27.45  |
| 3              | 90.03  | 90.04  | 90.03  | 90.03  | 90.00  |
| 4              | 40.26  | 40.25  | 40.26  | 40.25  | 40.26  |
| 5              | 55.47  | 55.45  | 55.46  | 55.45  | 55.45  |
| 6              | 29.10  | 29.08  | 29.10  | 29.08  | 29.11  |
| 7              | 75.94  | 75.93  | 75.95  | 75.94  | 75.89  |
| 8              | 47.37  | 47.38  | 47.35  | 47.34  | 47.33  |
| 9              | 51.89  | 51.89  | 51.89  | 51.87  | 51.84  |
| 10             | 37.88  | 37.89  | 37.89  | 37.88  | 37.87  |
| 11             | 22.63  | 22.57  | 22.59  | 22.58  | 22.63  |
| 12             | 26.34  | 26.64  | 26.44  | 26.41  | 26.31  |
| 13             | 43.88  | 44.07  | 44.35  | 44.35  | 44.28  |
| 14             | 50.62  | 50.45  | 50.45  | 50.43  | 50.97  |
| 15             | 35.54  | 35.42  | 35.42  | 35.39  | 35.57  |
| 16             | 28.83  | 28.50  | 28.70  | 28.67  | 28.77  |
| 17             | 46.49  | 46.12  | 46.22  | 46.23  | 50.03  |
| 18             | 10.45  | 10.05  | 10.49  | 10.50  | 10.43  |
| 19             | 16.90  | 16.89  | 16.69  | 16.90  | 16.91  |
| 20             | 78.81  | 80.14  | 78.76  | 78.59  | 76.09  |
| 21             | 19.15  | 19.03  | 18.73  | 18.86  | 26.02  |
| 22             | 77.61  | 77.58  | 78.49  | 78.52  | 45.29  |
| 23             | 31.05  | 69.33  | 126.97 | 130.85 | 128.12 |
| 24             | 123.21 | 126.99 | 142.01 | 138.83 | 138.73 |
| 25             | 133.59 | 135.54 | 71.17  | 76.43  | 76.54  |
| 26             | 26.03  | 26.02  | 29.78  | 26.10  | 26.31  |
| 27             | 18.12  | 18.33  | 29.84  | 26.19  | 26.41  |
| 28             | 28.39  | 28.39  | 28.38  | 28.39  | 28.36  |
| 29             | 17.10  | 17.11  | 17.11  | 17.11  | 17.07  |
| 30             | 16.69  | 16.67  | 16.64  | 16.62  | 16.65  |
| $\text{OCH}_3$ |        |        |        | 50.78  | 50.03  |

**4-Allyl-2-methoxyphenyl-6-O- $\alpha$ -L-arabinopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (8)** was obtained as a white amorphous powder:  $[\alpha]_{\text{D}}^{24} +30^\circ$  ( $c$  0.2, MeOH); IR  $\nu_{\text{max}}$  (KBr) 3470, 1720, 1600, 1430  $\text{cm}^{-1}$ ; FABMS,  $m/z$  481  $[\text{M} + \text{Na}]^+$  (16).

**Acid Hydrolysis of 1.** Compound 1 (6.0 mg) was treated with 2 N methanolic HCl (2 mL) under reflux at  $90^\circ\text{C}$  for 1 h. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ , and the aqueous layer was neutralized with  $\text{Na}_2\text{CO}_3$  and filtered. The dried filtrate was acetylated with pyridine– $\text{Ac}_2\text{O}$ . GC–MS analysis showed peracetylramnose and peracetylglucose (1:1), in comparison with reference compounds.

**Cytotoxicity Assay.** The assay using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) against Hepa59T/VGH (human liver carcinoma), NCI (human large lung cell carcinoma), HeLa (human cervical epitheloid carcinoma), and Med (human medulloblastoma) tumor cells was based on the reported methods (17). The tumor cells were purchased from the American Type Culture Collection (ATCC). In brief, the cells were cultured in RPMI-1640 medium supplemented with serum in 5%  $\text{CO}_2$  and incubated at  $37^\circ\text{C}$ . Test samples and control drug standard were prepared at concentrations of 1, 10, 20, and 40

$\mu\text{g/mL}$ . After seeding of 2880 cells/well in a 96-well microplate for 4 h, 20  $\mu\text{L}$  of sample or standard agent was placed in each well and incubated at  $37^\circ\text{C}$  for 3 days, and then 20  $\mu\text{L}$  of MTT was added for 5 h. After removal of the medium and addition of DMSO (200  $\mu\text{L}$ /well) into the microplate with shaking for 10 min, the formazan crystals (the product of MTT reacting with dehydrogenase existing in mitochondria) were redissolved and their absorbance was measured on a model MR 7000 microtiter plate reader (Dynatech International Corp., Edgewood, NY) at a wavelength of 550 nm. The  $\text{ED}_{50}$  was defined by comparison with the untreated cells as the concentration of test sample resulting in 50% reduction of absorbance.

**HIV Inhibition Assay.** HIV inhibition was measured as described previously (18).


## RESULTS AND DISCUSSION

Bioassay-directed fractionation of the EtOH extract of galls from *S. mukorossi* resulted in the isolation of dammarane-type saponins 1–5 and phenylpropanoid glycosides 6–8.

The HRFABMS of compound 1 showed a pseudomolecular ion at  $807.4864 [\text{M} + \text{Na}]^+$ , consistent with the formula of  $\text{C}_{42}\text{H}_{72}\text{O}_{13}$ . The IR spectrum showed absorption bands for hydroxyl groups at  $3387 \text{ cm}^{-1}$  and olefinic groups at 1629 and  $1047 \text{ cm}^{-1}$ . The  $^1\text{H}$  NMR spectrum (Tables 1 and 2) displayed signals for six tertiary methyls ( $\delta_{\text{H}}$  0.85, 0.86, 0.93, 0.97, and  $1.05 \times 2$ ), two allylic methyls ( $\delta_{\text{H}}$  1.61 and 1.70), three oxygenated methines [ $\delta_{\text{H}}$  3.17 (dd,  $J = 11.2, 4.0 \text{ Hz}$ , H-3), 3.70 (dd,  $J = 10.0, 4.8 \text{ Hz}$ , H-7), and 3.41 (m, H-22)], and one olefinic proton [ $\delta_{\text{H}}$  5.24 (t,  $J = 6.8 \text{ Hz}$ , H-24)], as well as two anomeric protons [ $\delta_{\text{H}}$  4.39 (d,  $J = 7.6 \text{ Hz}$ , H-1') and 5.35 (br s, H-1'')]. The signals for eight tertiary methyls, eight methylenes, eight methines (including an olefinic carbon signal at  $\delta$  133.59 and 123.21 and three oxygenated carbons at  $\delta$  75.94, 77.61, and 90.03), and one oxygenated quaternary carbon at  $\delta_{\text{C}}$  78.81 were found in the  $^{13}\text{C}$  NMR (Table 3) and DEPT spectra of 1. After subtraction of the 12 carbon resonances of the sugar moieties (Table 2), the remaining 30 signals were attributable to a dammarane-type triterpene aglycon (19–24). On the basis of the triplet coupling pattern of the olefinic proton in the  $^1\text{H}$  NMR spectrum and the long-range correlation in the HMBC spectrum, the double bond was likely substituted with one methylene and two geminal methyl groups, suggesting the partial structure of the side chain (25).

The planar structure of 1 was further clarified by  $^1\text{H}$ – $^1\text{H}$  COSY, TOCSY, and HMBC experiments. The TOCSY spectrum (Figure 1) indicated the four partial structures for a dammarane-type triterpene as drawn with bold lines ( $\text{C}_1$ – $\text{C}_3$ ,  $\text{C}_5$ – $\text{C}_7$ ,  $\text{C}_9$ – $\text{C}_{17}$ , and  $\text{C}_{22}$ – $\text{C}_{24}$ ). Detailed inspection of the HMBC spectrum showed correlations between the following






**Figure 1.** Structures of dammarane saponins and phenylpropanoid glycosides isolated from the galls of *S. mukorossi*.

proton and carbon pairs: H<sub>3</sub>-18/C-7, 8, 9; H<sub>3</sub>-19/C-1, 5, 9, 10; H<sub>3</sub>-21/C-17, 22; H<sub>3</sub>-26, 27/C-24, 25; H<sub>3</sub>-28, 29/C-3, 4, 5. These data confirmed the positions of the seven methyl groups of **1** as shown in **Figure 1**. Moreover, the secondary hydroxy groups were located at C-7 and C-22 on the basis of the correlations between H-7 and C-6, 14, 18 and between H-22 and C-21, 23, 24, respectively.

Acid hydrolysis (26) of **1** gave D-glucose and L-rhamnose as the component sugars, which was confirmed by GLC analysis. HMBC spectroscopic data were used to assign the linkage positions of these sugars in **1**. Thus, the C-1 of rhamnose was linked to the C-2 of glucose on the basis of correlations between C-1 ( $\delta_C$  101.85) of the terminal rhamnose and H-2' of glucose ( $\delta_H$  3.39). Also, the glucose was linked at C-3 of the triterpene, due to a cross-peak between C-1' ( $\delta_C$  105.57) of glucose and H-3 ( $\delta_H$  3.17) of the aglycon. Moreover, the coupling pattern (dd,  $J$  = 11.2, 4.0 Hz) of H-3 indicated that the glucose–aglycon linkage has the  $\beta$  orientation. Coupling constants of the H-1 protons for the glucose (d,  $J$  = 7.6 Hz) and rhamnose (br s) (22) fully established the sugars of **1** as 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside.

The relative stereochemistries of **1** were deduced by NOESY experiments. As shown in **Figure 2**, correlations between the proton pairs, H-3/H-5 with H<sub>3</sub>-28 and H-7/H-5 with H<sub>3</sub>-30, indicated that the configurations of the oxygenated geminal protons are H-3 $\alpha$  and H-7 $\alpha$ . Moreover, H-17 showed NOESY correlations with H<sub>3</sub>-30 and H-16 $\alpha$ ; therefore, the side chain at C-17 was  $\beta$ -oriented. These data together with 2D-NOE correlation between H<sub>3</sub>-21 and H-13 established **1** as 3 $\beta$ ,7 $\beta$ ,20-(*S*),22-tetrahydroxydammar-24-ene-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside. This compound has been named sapinmusaponin A (**1**).

The molecular formula of **2** was deduced as C<sub>42</sub>H<sub>72</sub>O<sub>14</sub> from a pseudomolecular ion at 823.4813 [M + Na]<sup>+</sup> in the HR-



**Figure 2.** Key TOCSY and HMBC correlations of **1** and **5**.

FABMS. The IR spectrum showed absorption bands for hydroxyl (3389 cm<sup>-1</sup>) and olefinic (1640, 1047 cm<sup>-1</sup>) groups. The <sup>1</sup>H and <sup>13</sup>C NMR spectra of **2** showed signal patterns similar to those of **1**, except for the appearance of signals for an oxymethine [ $\delta_H$  4.67 (dd,  $J$  = 8.8, 2.0 Hz),  $\delta_C$  69.33] in **2**. Further detailed analysis of **2** using a TOCSY experiment suggested the same four partial structures of **1** as shown in **Figure 1**. The side chains of **1** and **2** differ by an additional hydroxyl group found in **2**. Moreover, correlations between  $\delta_H$  4.67 (H-23) and  $\delta_C$  126.99 (C-24) and 135.54 (C-25) were found in the HMBC spectrum, suggesting that the hydroxyl group is attached to C-23. After acid hydrolysis, compound **2**, as well as the isolates **3**–**5** described subsequently, also gave D-glucose and L-rhamnose as component sugars.

The stereochemistry at C-20 was assigned as *S*, due to 2D-NOE correlations between H<sub>3</sub>-21 and H-13/H-16 $\beta$ , as well as between H-17 and H<sub>3</sub>-30 as shown in **Figure 2**. Thus, **2** was elucidated as 3 $\beta$ ,7 $\beta$ ,20-(*S*),22,23-pentahydroxydammar-24-ene-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside and named sapinmusaponin B.

Sapinmusaponin C (**3**) had a molecular formula of C<sub>42</sub>H<sub>72</sub>O<sub>14</sub> based on a pseudomolecular ion at 823.4824 [M + Na]<sup>+</sup> in its HRFABMS. The IR spectrum showed the absorption bands for hydroxyl (3382 cm<sup>-1</sup>) and olefinic (1641 and 1045 cm<sup>-1</sup>) groups. As found in **1** and **2**, compound **3** possessed two sugars and a dammarane triterpene skeleton based on their similar <sup>1</sup>H and <sup>13</sup>C NMR spectra. Comparison of the <sup>1</sup>H spectra of **1** and **3** showed that the H<sub>3</sub>-26 and H<sub>3</sub>-27 signals of **3** were found at higher field ( $\delta_H$  1.70 and 1.61 in **1**;  $\delta_H$  1.28  $\times$  2 in **3**) and that the vinyl quaternary carbon at C-25 in **1** was replaced by an oxygenated carbon in **3**. Also, two vinyl carbons signals at  $\delta_C$  126.97 and 142.01 were observed in **3** and assigned as C-23 and 24, respectively, due to the long-range correlations between H-24 and C-26/C-27 and between H-22 and C-23/C-24 in the HMBC spectrum. Furthermore, when the <sup>1</sup>H NMR spectrum of **3** was measured in C<sub>5</sub>D<sub>5</sub>N rather than CD<sub>3</sub>OD, the coupling constants ( $J$  = 15.6 Hz) for H-23 and H-24 were clearly observed and, therefore, *trans*-geometry was assigned to the double bond (27). On the basis of the HMBC spectrum (**Figure 1**), the five hydroxyl groups ( $\delta_C$  90.03, 75.95, 78.76, 78.49, and 71.17) in **3** were assigned at C-3, C-7, C-20, C-22, and C-25, respectively.

The stereochemistry of each of the hydroxyl groups was based on a NOESY experiment (**Figure 3**). H-3 was correlated with H-5 and H<sub>3</sub>-28 and H-7 with H-5 and H<sub>3</sub>-30, indicating their orientations as H-3 $\alpha$  and H-7 $\alpha$ , respectively. Moreover, correlations of H<sub>3</sub>-21 with H-13 and H<sub>3</sub>-18 determined that C-20 has an *S* configuration. Together with above data, **3** was tentatively elucidated as 3 $\beta$ ,7 $\beta$ ,20-(*S*),22,25-pentahydroxydammar-23-ene-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside.




Figure 3. Main NOESY correlations for sapinmusaponins A–E (1–5).

The molecular formula of sapinmusaponin D (**4**) was determined as  $C_{43}H_{74}O_{14}$  from the HRFABMS, which exhibited a pseudomolecular ion peak at 837.4982  $[M + Na]^+$ . The IR spectrum showed absorption bands for hydroxyl ( $3394\text{ cm}^{-1}$ ) and olefinic ( $1654$  and  $1048\text{ cm}^{-1}$ ) groups. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **4** contained similar patterns as found in **3**, except for the appearance of an additional methoxy signal ( $\delta_{\text{H}}$  3.16 and  $\delta_{\text{C}}$  50.78) in **4**. Consequently, C-25 ( $\delta_{\text{C}}$  76.43) in **4** was shifted to lower field (+5.26), and C-26 ( $\delta_{\text{C}}$  26.10) and C-27 ( $\delta_{\text{C}}$  26.19) were shifted to higher field (−3.68 and −3.65, respectively), compared with **3**. In detailed inspection of the HMBC spectrum, a cross-peak between  $\delta_{\text{H}}$  3.16 ( $\text{OCH}_3$ ) and  $\delta_{\text{C}}$  76.43 (C-25) placed the methoxyl group at the C-25 position. Like **3**, the configurations of **4** were based on NOE studies and, therefore, **4** was tentatively identified as 25-methoxy- $3\beta,7\beta,20$ -(*S*),22-tetrahydroxydammar-23-ene-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside.

Sapinmusaponin E (**5**) was obtained as a white powder. Its HRFABMS spectrum revealed the molecular formula of  $C_{43}H_{74}O_{13}$  from a pseudomolecular ion at 821.5016  $[M + Na]^+$ . As found in **4**, the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **5** indicated a dammarane-type triterpene with glucose and rhamnose moieties. The NMR spectra of **4** and **5** revealed similar structures, except that **5** had only a methylene group at C-22 rather than the hydroxymethine found in **4**. Other obvious carbon shifts included a downfield shift of the C-21 signal in **5** ( $\delta_{\text{C}}$  18.86 in **4** and  $\delta_{\text{C}}$  26.02 in **5**) and upfield shifts of C-20 ( $\delta_{\text{C}}$  76.09) and C-22 ( $\delta_{\text{C}}$  45.29), compared with **4**. On the basis of the HMBC spectrum (Figure 1), cross-peaks between  $\delta_{\text{H}}$  2.23, 2.22 (H-22) and  $\delta_{\text{C}}$  76.09 (C-20), 128.12 (C-23), and 138.73 (C-24) further demonstrated the presence of a methylene group at C-22. Together with the NOE correlations similar to those of **4**, **5** was thus determined to be 25-methoxy- $3\beta,7\beta,20$ -(*R*)-trihydroxydammar-23-ene-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside.

Compounds **6–8** were identified as known phenylpropanoid glycosides, 4-allyl-2-methoxyphenyl-6-*O*- $\beta$ -D-apiosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucoside (**6**) (**14**), 4-allyl-2-methoxyphenyl-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (**7**) (**15**), and 4-allyl-2-methoxyphenyl-6-*O*- $\alpha$ -L-arabinopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (**8**) (**16**), respectively, by analysis of their spectroscopic data (FAB-MS,  $^1\text{H}$  NMR, and  $^{13}\text{C}$  NMR spectra) and comparison with literature values.

The isolated dammarane-type saponins, **1–5**, were evaluated in a cytotoxicity assay (Hepa59T/VGH, NCI, HeLa, and Med). The quantities of **6–8** were insufficient for the bioassay. The cytotoxicity data are shown in Table 4 and show that **3** and **4** exhibited weak cytotoxic effects against all four tested cell lines ( $\text{ED}_{50} \sim 9\text{--}17\text{ }\mu\text{g/mL}$ ). Compound **1** had a weak response against Hepa59T/VGH and Med tumor cells, and **5** demonstrated weak cytotoxicity against HeLa and Med tumor cells. These results were consistent with literature data that some dammarane glycosides from processed ginseng had moderate cytotoxicity against SK-Hep-1 hepatoma cancer cells (28). Notably, com-

Table 4. Cytotoxic Activity of Compounds 1–5

| compd    | $\text{ED}_{50}$ ( $\mu\text{g/mL}$ ) for cell line <sup>a</sup> |                |      |      |
|----------|--|----------------|------|------|
|          | Hepa59T/VGH  | NCI            | HeLa | Med  |
| <b>1</b> | 15.2   | — <sup>b</sup> | —    | 14.9 |
| <b>2</b> | —  | —              | —    | —    |
| <b>3</b> | 17.3   | 12.7           | 17.1 | 9.9  |
| <b>4</b> | 12.1   | 9.1            | 10.3 | 10.8 |
| <b>5</b> | —  | —              | 16.9 | 18.5 |

<sup>a</sup> Hepa59T/VGH, human liver carcinoma; NCI, human large lung cell carcinoma; HeLa, human cervical epitheloid carcinoma; Med, human medulloblastoma. <sup>b</sup> —, inactive,  $\text{ED}_{50} > 20\text{ }\mu\text{g/mL}$ .

pound **2**, which has three rather than one or two hydroxyl groups in the dammarane side chain, had the lowest inhibitory effects ( $\text{ED}_{50} > 20\text{ }\mu\text{g/mL}$ ).

Compounds **1–5** were also evaluated for in vitro inhibitory effects against HIV replication in H9 lymphocytes. None of these dammarane-type saponins suppressed HIV replication.

## LITERATURE CITED

- (1) Nakayama, K.; Fujino, H.; Kasai, R.; Tanaka, O.; Zhou, J. Solubilizing properties of saponins from *Sapindus mukorossi* Gaertn. *Chem. Pharm. Bull.* **1986**, *34*, 3279–3283.
- (2) Takagi, K.; Park, E. H.; Kato, H. Anti-inflammatory activity of hederaganin and crude saponin isolated from *Sapindus mukorossi* Gaertn. *Chem. Pharm. Bull.* **1980**, *28*, 1183–1188.
- (3) Kasai, R.; Fujino, H.; Kuzuki, T.; Wong, W. H.; Goto, C.; Yata, N.; Tanaka, O.; Yasuhara, F.; Yamaguchi, S. Acyclic sesquiterpene oligoglycoside from pericarps of *Sapindus mukorossi*. *Phytochemistry* **1986**, *25*, 871–876.
- (4) Watanabe, K.; Fujino, H.; Morita, T.; Kasai, R.; Tanaka, O. Solubilization of saponins of *Bupleuri radix* with ginseng saponins: cooperative effect of dammarane saponins. *Planta Med.* **1988**, *54*, 405–409.
- (5) Dhar, J. D.; Bajpai, V. K.; Setty, B. S.; Kamboj, V. P. Morphological changes in human spermatozoa as examined under scanning electron microscope after in vitro exposure to saponins isolated from *Sapindus mukorossi*. *Contraception* **1989**, *39*, 563–568.
- (6) Sengupta, A.; Basu, S. P.; Saha, S. Triglyceride composition of *Sapindus mukorossi* seed oil. *Lipids* **1975**, *10*, 33–40.
- (7) Zikova, N. I.; Krivenchuk, P. E. Chemical study of flavonoids from the leaves of *Sapindus mukorossi* Gaertn. *Farm. Zh.* **1970**, *25*, 43–45.
- (8) Hunag, H. C.; Liao, S. C.; Chang, F. R.; Kuo, Y. H.; Wu, Y. C. Molluscicidal saponins from *Sapindus mukorossi*, inhibitory agents of Golden Apple snails, *Pomacea canaliculata*. *J. Agric. Food Chem.* **2003**, *51*, 4916–4919.
- (9) Quetin-Leclercq, J.; Elias, R.; Balansard, G.; Bassleer, R.; Angenot, L. Cytotoxicity activity of some triterpenoid saponins. *Planta Med.* **1992**, *58*, 279–281.
- (10) Graham, N. S.; Karsten, S. The adaptive significance of insect gall morphology. *Trends Ecol. Evolution* **2003**, *18*, 215–222.
- (11) Abrahamson, W. G.; Melika, G.; Scrafford, R.; Csoka, G. Gall-inducing insects provide insights into plant systematic relationships. *Am. J. Bot.* **1998**, *85*, 1159–1165.
- (12) Li, Thomas S. C. Chinese and related North American herbs. *Phyletic Relationships between Chinese and Western Medicinal Herbs*; CRC Press: Washington, DC, 2002; p 280.
- (13) Huang, K. C. Antidiarrheal herbs. *The Pharmacology of Chinese Herbs*; CRC Press: Boca Raton, FL, 1991; p 239.
- (14) Takeda, Y.; Oosio, Y.; Masuda, T.; Honda, G.; Otsuka, H.; Seziki, E.; Yesilada, E. Iridoid and eugenol glycosides from *Nepeta cadmea*. *Phytochemistry* **1998**, *49*, 787–791.
- (15) Orihara, Y.; Furuya, T.; Hashimoto, N.; Deguchi, Y.; Tokoro, K.; Kanisawa, T. Biotransformation of isoeugenol and eugenol by cultured cells of *Eucalyptus perriniana*. *Phytochemistry* **1992**, *31*, 827–831.

- (16) Machida, K.; Nakano, Y.; Kikuchi, M. Phenolic glycosides from *Viburnum dilatatum*. *Phytochemistry* **1991**, *30*, 2013–2014.
- (17) Kuo, Y. H.; Li, S. Y.; Shen, Y. C.; Huang, H. C.; Hsu, Y. W.; Tseng, R. J.; Ou, J. C.; Chen, C. F. Sesquiterpene polyol esters and triterpenes from *Celastrus punctatus*. *Chin. Pharm. J.* **2001**, *53*, 257–268.
- (18) Kuo, Y. H.; Kuo-Yang, L. M. Antitumour and anti-AIDS triterpenes from *Celastrus hindsii*. *Phytochemistry* **1997**, *44*, 1275–1281.
- (19) Kasai, R.; Besso, H.; Tanaka, O.; Saruwatari, Y. I.; Fuwa, T. Saponins of red ginseng. *Chem. Pharm. Bull.* **1983**, *31*, 2120–2125.
- (20) Morita, T.; Kasai, R.; Tanaka, O.; Zhou, J.; Yang, T. R.; Shoji, J. Saponins of Zu-Tziseng, rhizomes of *Panax japonicus* C.A. Meyer var. major (Burk.) C.Y. Wu et K.M. Feng, collected in Yunnan, China. *Chem. Pharm. Bull.* **1982**, *30*, 4341–4346.
- (21) Jiang, Z. H.; Fukuoka, R.; Aoki, F.; Tanaks, T.; Kouno, I. Dammarane-type triterpene glycosides from the leaves of *Rhoiptelea chiliantha*. *Chem. Pharm. Bull.* **1999**, *47*, 257–262.
- (22) Bedir, E.; Toyang, N. J.; Khan, I. A.; Walker, L. A.; Clark, A. M. A new dammarane-type triterpene glycoside from *Polyscias fulva*. *J. Nat. Prod.* **2001**, *64*, 95–97.
- (23) Takemoto, T.; Arihara, S.; Yoshokawa, K. Studies on the constituents of Cucurbitaceae plants. XIV. On the saponin constituents of *Gynostemma pentaphyllum* Makino. *Yakugaku Zasshi* **1986**, *106*, 664–670.
- (24) Tori, M.; Mastsuda, R.; Sono, M.; Asakawa, Y. <sup>13</sup>C NMR assignment of dammarane triterpenes and dendropanoxide: application of 2D long-range <sup>13</sup>C–<sup>1</sup>H correlation spectra. *Magn. Reson. Chem.* **1988**, *26*, 581–590.
- (25) Bianchini, J. P.; Gaydou, E. M.; Rafaralahitsimba, G.; Waegell, B.; Zahra, J. P. Dammarane derivatives in the fruit lipids of *Olea madagascariensis*. *Phytochemistry* **1988**, *27*, 2301–2304.
- (26) Seo, Y.; Berger, J. M.; Hoch, J.; Neddermann, K. M.; Bursuker, I.; Mamber, S. W.; Kingston, D. I. A new triterpene saponin from *Pittosporum viridiflorum* from the Madagascar rainforest. *J. Nat. Prod.* **2002**, *65*, 65–68.
- (27) Benosman, A.; Richomme, P.; Sevenet, T.; Perroms, G.; Hadi, A. H. A.; Bruneton, J. Tirucallane triterpenes from the stem bark of *Aglaia leucophylla*. *Phytochemistry* **1995**, *40*, 1485–1487.
- (28) Park, I. H.; Piao, L. Z.; Kwon, S. W.; Lee, Y. J.; Cho, S. Y.; Park, M. K.; Park, J. H. Cytotoxic dammarane glycosides from processed ginseng. *Chem. Pharm. Bull.* **2002**, *50*, 538–40.

Received for review December 3, 2004. Revised manuscript received March 17, 2005. Accepted March 17, 2005.

JF047963S