

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51485671>

Enhanced Stability by Three-Dimensional Aromaticity of Endohedrally Doped Clusters $X_{10}Mo/-$ with $X = Ge, Sn, Pb$ and $M = Cu, Ag, Au$

ARTICLE *in* THE JOURNAL OF PHYSICAL CHEMISTRY A · AUGUST 2011

Impact Factor: 2.69 · DOI: 10.1021/jp111324n · Source: PubMed

CITATIONS

11

READS

45

2 AUTHORS:



Truong Tai

University of Leuven

65 PUBLICATIONS 506 CITATIONS

SEE PROFILE



Minh Tho Nguyen

University of Leuven

748 PUBLICATIONS 10,846 CITATIONS

SEE PROFILE




Figure 1. Shapes, point groups, and relative energies (kcal/mol) of some low-lying isomers of the anionic clusters $X_{10}M^-$ ($X = \text{Ge, Sn, Pb}$ and $M = \text{Cu, Ag, Au}$) (B3LYP/LanL2DZ): (a) $X_{10}\text{Cu}^-$; (b) $X_{10}\text{Ag}^-$; (c) $X_{10}\text{Au}^-$.

Motivated by these reasons, we set out to carry out a systematic investigation on the clusters with impurity $X_{10}M^{0/-}$ in both neutral and anionic states, in which $X = \text{Ge, Sn, Pb}$ and $M = \text{Cu, Ag, Au}$. The location of the lower-lying isomers of the clusters considered is carried out using a stochastic search method.²⁰ As experimental studies on silver- or gold-doped group 14 clusters are not available, predictions of their properties and additional confirmation on the effects of impurities on the caged clusters of the higher homologues would stimulate subsequent experimental studies. In addition, the high stability of the anionic systems $X_{10}M^-$ is also of interest for future investigations.

COMPUTATIONAL METHODS

Quantum chemical computations are carried out using the Gaussian 03²¹ and Molpro 06²² suites of programs. Geometries and harmonic vibrational frequencies of the low-lying isomers are

determined using density functional theory (DFT) method with the hybrid B3LYP functional, which involves the Becke three-parameter exchange²³ and Lee–Yang–Parr correlation²⁴ functionals. The initial search for all possible low-lying isomers of each of the clusters $X_{10}M^{0/-}$ considered is performed using a stochastic search algorithm, which was implemented by us,²⁵ on the basis of the Saunderson's procedure.¹⁶ First, the possible structures of the cluster $X_{10}M$ are generated by random “kick” method, and then rapidly optimized at the B3LYP/CEP-31G level.²⁶ In this “kick” procedure, the minimum and maximum distances between atoms are limited to 2 and 7 Å, respectively. Geometries of the stationary points located are then reoptimized using the B3LYP functional in conjunction with the LanL2DZ basis set, in which ab initio effective core potentials (ECPs) have been generated to replace the Coulomb, exchange and core-orthogonality effects of the chemically inert core electrons in transition metals.²⁷

To obtain more accurate energetic predictions, calculations of single-point electronic energies of the global minima are further




Figure 2. Shapes and point groups of some low-lying isomers of the neutral clusters $X_{10}M$ ($X = \text{Ge, Sn, Pb}$ and $M = \text{Cu, Ag, Au}$) (B3LYP/LanL2DZ level).

performed using the coupled-cluster theory CCSD(T) method,²⁸ making use of the correlation consistent aug-cc-pVTZ-PP basis set, in which PP stands for relativistic pseudopotentials for metal atoms.²⁹ Densities of states are computed using the Pymolize program.³⁰ The electron localizability indicator (ELI-D) values are calculated by using DGrid-4.2 program suite,³¹ and the ELI isosurfaces are plotted using the Gopenmol software.³²

RESULTS AND DISCUSSION

Geometries of Lower-Lying Isomers. The structures given hereafter are labeled by $XY.n$ for the anions, where $X = \text{I, II, III}$ stands for Ge, Sn, Pb and $Y = \text{A, B, C}$ stands for Cu, Ag and Au, respectively. While the shapes, point groups, electronic states, and relative energies of the lower-lying isomers $X_{10}M^-$ are shown in Figure 1, the structures of the global minima $X_{10}M$ are given in Figure 2. The global minima of all copper doped group IVA clusters at the anionic state $X_{10}\text{Cu}^-$ ($X = \text{Ge, Sn, Pb}$) have high symmetry structure (D_{4d} , Figure 1a). It is interesting to see that the same structures D_{4d} are found to be the global minima for neutrals $X_{10}\text{Cu}$ (Figure 2). At the B3LYP/LanL2DZ level, the Ge–Cu bond length of the neutral Ge_{10}Cu is equal to 2.55 Å, which represents the shortest distance as compared to those of Sn–Cu (2.80 Å) and Pb–Cu (2.86 Å). Following attachment of one excess electron, the distances of X–Cu of the anion $X_{10}\text{Cu}^-$ are slightly shortened ($d(\text{Ge–Cu}) = 2.52$ Å, $d(\text{Sn–Cu}) = 2.75$ Å, and $d(\text{Pb–Cu}) = 2.82$ Å), due to the closed shell electronic configuration of the anionic systems.

Following replacement of Cu by heavier impurities Ag and Au, the D_{4d} structure becomes unfavorable for doped Ge clusters. The C_2 structure IB.1 (Figure 1b) in which the Ag atom is doped outside of the Ge_{10} cage is found to be the global minimum for $\text{Ge}_{10}\text{Ag}^-$ with 16.5 kcal/mol lower in energy as compared to the D_{4d} structure IB.6. The structures IB.2 and IB.3 are the next isomers with relative energies of 6.0 and 8.2 kcal/mol, respectively. At the neutral state, a C_s structure (Figure 2), which is a lower symmetry of IB.1, is the global minimum for the neutral Ge_{10}Ag , while the D_{4d} structure is the next isomer with 10.7 kcal/mol higher in energy.

For $\text{Ge}_{10}\text{Au}^-$ anionic clusters (Figure 1c), two structures, including a D_{5h} structure IC.1 in which Au is located at center of a D_{5h} cage, and a C_1 structure IC.2 in which one Ge atom of the Ge_{11} is replaced by one Au atom are almost degenerate in energy. The next isomer is a D_{4d} structure IC.3 that is 12.2 kcal/mol less

stable than the first one. Following detachment of one electron, a C_s structure (Figure 2) that is distorted from a high symmetry IB.1 (D_{5h}) under a Jahn–Teller effect, is found to be the most stable isomer for the neutral Ge_{10}Au . Two structures, including a C_1 structure and a D_{4d} structure (Figure 2) are also quite stable with relative energies of 2.7 and 5.2 kcal/mol, respectively.

Let us note that the size of D_{4d} cages is smaller compared to that of D_{5h} counterparts. At the B3LYP/LanL2DZ level, the Au–Ge bond length of IC.1 (D_{5h}) is equal to 2.72 Å and that is larger than that of 2.64 Å of IC.3 (D_{4d}). This difference is not quite large but may be reason for why gold can be located at the center of the D_{5h} cage, while similar structures of silver dopant, which has a smaller atom radius, are not stable. Moreover, even for the gold doped germanium clusters $\text{Ge}_{10}\text{Au}^{0/-}$, there is a competition between the endohedral IC.1 and exohedral IC.2 structures. Their difference in relative energies is quite small for the neutrals Ge_{10}Au , and nearly zero for the anions $\text{Ge}_{10}\text{Au}^-$. These clusters are less stable as compared to other endohedral forms shown in the following sections.

Due to the larger size of the Sn_{10} and Pb_{10} cages, as compared to the Ge_{10} cage, it is expected that Ag and Au impurities can be encapsulated to form compact structures. Our calculations point out that all clusters $X_{10}M^{0/-}$ with $X = \text{Sn, Pb}$ and $M = \text{Ag, Au}$ are bicapped square antiprism structures (D_{4d}) (Figures 1 and 2). The X–M bonding lengths are slightly increased when moving down from Cu to Au as shown in Table 1.

Binding and Adiabatic-Vertical Detachment Energies. To probe the inherent stability of the anionic clusters considered, the average binding energies (BEs), vertical (VDE), and adiabatic (ADE) detachment energies of the lowest-lying $X_{10}M^{0/-}$ isomers are evaluated and compared to those of the pure dianionic clusters X_{10}^{2-} (D_{4d}). The average binding energies (BE) can be defined as follows:

$$BE = [nE(X) + E(M) - E(X_{10}M)] / (n + 1) \quad \text{for } X_{10}M \quad (1)$$

$$BE = [nE(X) + E(M^-) - E(X_{10}M^-)] / (n + 1) \quad \text{for } X_{10}M^- \quad (2)$$

$$BE = [(n - 2)E(X) + 2E(X^-) - E(X_{10}^{2-})] / n \quad \text{for } X_{10}^{2-} \quad (3)$$

where $E(X)$, $E(M)$, $E(X^-)$, and $E(M^-)$ with $X = \text{Ge, Sn, Pb}$ and $M = \text{Cu, Ag, Au}$ are the total energies of the atoms at the neutral and anionic states, respectively.

While the BE, ADE, and VDE values of the lowest-lying isomers are given in Table 1, the plots of the BE values are illustrated in Figure 3a. Compared to the $X_{10}M$ systems containing 42 valence electrons reported earlier, namely $\text{BE}(\text{Ge}_{10}\text{Zn}) = 3.1$ eV, $\text{BE}(\text{Sn}_{10}\text{Zn}) = 2.8$ eV,¹³ $\text{BE}(\text{Pb}_{10}\text{Mg}) = 1.77$ eV, and $\text{BE}(\text{Pb}_{10}\text{Ba}) = 1.83$ eV,¹⁴ the isoelectronic species $X_{10}M^-$ are thus characterized by large BE values that ultimately indicate their higher stability. Although the neutrals $X_{10}M$ have somewhat lower stability than the corresponding anions due to their opened electronic shell, they also possess relatively high BE values. Additionally, it can be seen from Figure 3a that the doping by coinage metals (Cu, Ag and Au) results in increasing binding energies of the impure clusters $X_{10}M$ as compared to those of the pure X_{10}^{2-} .

Table 1. Selected Geometrical Parameters, HOMO–LUMO Gaps (HLG) (eV), Vertical (VDE) and Adiabatic (ADE) Detachment Energies (eV), Embedding Energy (EE) (eV), and Average Binding Energy (BE) (eV) of the Global Minima $X_{10}M^{0/-}$ and X_{10}^{2-}

structure	d_{X-M} (Å) ^a	VDE ^b	ADE	BE	EE	HLG ^c
$Ge_{10}Cu$ (D_{4d} , 2A_1)	2.552; 2.662	6.47		2.98	3.18	
$Ge_{10}Ag$ (C_{3v} , $^2A'$)	2.830	6.36		2.78	1.02	
$Ge_{10}Au$ (C_{3v} , $^2A''$)	2.720	6.76		2.82	1.46	
$Sn_{10}Cu$ (D_{4d} , 2A_1)	2.795; 2.977	6.14		2.68	3.76	
$Sn_{10}Ag$ (D_{4d} , 2A_1)	2.880; 3.015	6.03		2.54	2.20	
$Sn_{10}Au$ (D_{4d} , 2A_1)	2.883; 3.019	5.93		2.65	3.33	
$Pb_{10}Cu$ (D_{4d} , 2A_1)	2.864; 3.093	5.93		2.43	3.56	
$Pb_{10}Ag$ (D_{4d} , 2A_1)	2.931; 3.117	5.84		2.33	2.47	
$Pb_{10}Au$ (D_{4d} , 2A_1)	2.948; 3.131	5.72		2.43	3.53	
$Ge_{10}Cu^-$ (D_{4d} , 1A_1)	2.520; 2.849	3.06	2.79	3.15	5.10	2.48
$Ge_{10}Ag^-$ (C_{3v} , 1A)	2.711	3.31	3.05	2.97	2.52	2.21
$Ge_{10}Au^-$ (D_{3h} , 1A_1)	2.721	3.17	3.02	2.92	3.08	2.19
$Sn_{10}Cu^-$ (D_{4d} , 1A_1)	2.753; 3.204	3.11	2.91	2.87	5.80	2.39
$Sn_{10}Ag^-$ (D_{4d} , 1A_1)	2.845; 3.218	2.99	2.79	2.71	4.01	2.16
$Sn_{10}Au^-$ (D_{4d} , 1A_1)	2.848; 3.236	2.90	2.68	2.71	4.06	2.07
$Pb_{10}Cu^-$ (D_{4d} , 1A_1)	2.819; 3.314	2.97	2.84	2.61	5.53	2.58
$Pb_{10}Ag^-$ (D_{4d} , 1A_1)	2.892; 3.320	2.85	2.71	2.49	4.19	2.39
$Pb_{10}Au^-$ (D_{4d} , 1A_1)	2.908; 3.346	2.75	2.59	2.49	4.17	2.26
Ge_{10}^{2-} (D_{4d} , 1A_1)				2.92		2.86
Sn_{10}^{2-} (D_{4d} , 1A_1)				2.55		2.39
Pb_{10}^{2-} (D_{4d} , 1A_1)				1.97		0.51

^aThe first value is bond length between metal (M) and X atoms of cage X_8 . The second value is bond length between metal (M) and X atoms in the C_4 axis. ^bValues of VDE, ADE, EE, and BE calculated at the CCSD(T)/aug-cc-pVTZ-PP level. ^cHOMO–LUMO gaps are obtained at the B3LYP/LanL2DZ level.

Let us stress that the average binding energy (BE) is only one of the energetic properties used to evaluate the stability of clusters. It can be seen that the difference in BEs between $Ge_{10}X$ ($X = Ag, Au$) and $Ge_{10}Cu$ is relatively larger than those of the corresponding $Sn_{10}X$ or $Pb_{10}X$ clusters. For the neutral state, the difference in BEs between $Ge_{10}Cu$ and $Ge_{10}Au$ is 0.16 eV (0.2 eV for $Ge_{10}Ag$). This difference between $Sn_{10}Cu$ and $Sn_{10}Au$ amounts to only 0.03 eV (0.14 eV for $Sn_{10}Ag$), whereas the BE of $Pb_{10}Cu$ and $Pb_{10}Au$ is nearly the same. This indicates a lower stability of the opened structure $Ge_{10}Ag$ as compared to the packing structure $Ge_{10}Cu$. The lower stability of $Ge_{10}Au$ can be understood from the larger size of the D_{3h} cage. Similar observations can be found for the anionic clusters. For the VDE and ADE values, it can be observed that the presence of dopants forms similar trends in both the neutral and anionic states. The VDE (and ADE for anionic clusters) values are remarkably high, and only slightly decrease when moving down from Cu to Au.

Embedding Energy (EE). Embedding energy is the energy gain in incorporating a dopant $M = Cu, Ag,$ and Au into a X_{10} cage ($X = Ge, Sn, Pb$) and defined as follows:

$$EE(X_{10}M) = E(X_{10}) + E(M) - E(X_{10}M) \quad \text{for a neutral } X_{10}M$$

$$EE(X_{10}M^-) = E(X_{10}) + E(M^-) - E(X_{10}M^-) \quad \text{for an anion } X_{10}M^-$$




Figure 3. Plots of (a) binding energies (eV), (b) embedding energies (eV) and (c) HOMO-LUMO gaps of the global minima $X_{10}M^{0/-}$ and X_{10}^{2-} .

where $E(X_{10})$ is the total energy of the global minima X_{10} (C_{3v}).²⁵

Embedding energy can thus be used for evaluating the stability of endohedrally doped structures. The EE values of the global minima $X_{10}M$ at both neutral and cationic states are given in Table 1, while the relevant plots are depicted in Figure 3b. At the first glance, there exists a similar trend between the binding energy and embedding energy values. The EE values of the anions $X_{10}M^-$ with closed electronic shells are considerably larger than those of the open shell neutrals $X_{10}M$. The embedding energies of endohedral structures $Ge_{10}Cu^{0/-}$ are found to be remarkably large as compared to those of exohedral structures $Ge_{10}Ag^{0/-}$. In addition, the EE values of the clusters $Ge_{10}Au^{0/-}$




Figure 6. Isosurfaces of p-ELI plots for $\text{Ge}_{10}\text{Cu}^-$ at ELI values of (a) 0.6 and (b) 0.8.

consequence, a large energy gap between frontier orbitals emerges leading to a high stability of $\text{Ge}_{10}\text{Cu}^-$.

The plots of total DOS of the global minima X_{10}M^- (Figure 5) show that addition of a coinage metal dopant lowers the energy levels of the mixed clusters, relative to the pure X_{10}^{2-} dianionic cages. The shifting of energy levels of X_{10}M^- consistently increases in going from Cu to Au as dopant.

Electron Localization. An inspection of the electron localizability indicator (ELI),³³ which is proved to be an effective tool to probe the chemical bonding in organic compounds as well as in transition metal compounds,^{34,35} is performed. Let us show an analysis of the ELI of $\text{Ge}_{10}\text{Cu}^-$. The isosurfaces of partial-ELI distribution for this anion given in Figure 6 reveal that the green localization domains arising from σ -orbitals are distributed between Ge atoms. Consequently, they are responsible for bonding between Ge atoms of the cages. The red domains composed of π -orbitals are distributed globally over Ge atoms of the cage and is expected to be responsible for π -aromaticity.

Three-Dimensional Aromaticity. Aromaticity is well recognized as a measure of stability of chemical compounds. While aromaticity of planar structures can be qualitatively probed by using Hückel rule of $(4n + 2)$ valence electrons,³⁶ the $2(N + 1)^2$ electron count rule, which is recently proposed by Hirsch et al.,³⁷ is proven to be an effective criterion for spherical structures. In the latter counting framework, the π -electron system of fullerenes can approximately be considered as a spherical electron gas which surrounds the surface of a sphere (I_h). The wave functions of this electron gas can be characterized by the angular momentum quantum numbers ($l = 0, 1, 2, 3, \dots$) that are comparable to the atomic s, p, d, f, ... orbitals. According to the classical Pauli principle, if a spherical system bearing $2(N + 1)^2$ π -electrons fully fills all π -shells, it then is stabilized and exhibits an aromatic character.

Let us now consider the MO pictures of the global minimum $\text{Ge}_{10}\text{Cu}^-$ shown in Figure 4. The valence orbitals of $\text{Ge}_{10}\text{Cu}^-$ can be divided into two subsets, the first contains the MOs of the shells $1\text{S}(\text{A}_1, \sigma)$, $1\text{P}(\text{B}_2, \sigma)$, $1\text{P}(\text{E}_1, \sigma)$, $1\text{D}(\text{A}_1, \sigma)$, $1\text{D}(\text{E}_2, \sigma)$, $1\text{D}(\text{E}_3, \sigma)$ and $1\text{F}(\text{A}_1, \sigma)$, $1\text{F}(\text{E}_2, \sigma)$, $1\text{F}(\text{E}_3, \sigma)$, $1\text{F}(\text{E}_4, \sigma)$ and $1\text{G}(\text{A}_1, \sigma)$ that are occupied by 34 σ -electrons. The second subset includes the MOs of the shells $2\text{S}(\text{A}_1, \pi)$, $2\text{P}(\text{B}_2, \pi)$, and $2\text{P}(\text{E}_1, \pi)$, which are occupied by 8 valence π -electrons. As a consequence, this π -electron system makes $\text{Ge}_{10}\text{Cu}^-$ three-dimensionally π -aromatic satisfying the $2(N + 1)^2$ count rule. Similar systems of 8 valence π -electrons are also found as the global minima of the clusters X_{10}M^- , including $\text{Sn}_{10}\text{Cu}^-$, $\text{Sn}_{10}\text{Ag}^-$, $\text{Sn}_{10}\text{Au}^-$, $\text{Pb}_{10}\text{Cu}^-$, $\text{Pb}_{10}\text{Ag}^-$, and $\text{Pb}_{10}\text{Au}^-$, and accordingly, they also have a spherical π -aromatic character.

CONCLUDING REMARKS

In the present theoretical study, we perform a systematic investigation on the coinage metal encapsulated group 14

clusters $\text{X}_{10}\text{M}^{0/-}$ at the neutral and anionic states, including $\text{X} = \text{Ge}, \text{Sn}, \text{Pb}$ and $\text{M} = \text{Cu}, \text{Ag}, \text{Au}$. The global minima are located using a stochastic search method. Except for the Ge_{10}Ag , the global minima of most $\text{X}_{10}\text{M}^{0/-}$ systems considered are endohedral structure with impurities encapsulated at the center of X_{10} cages. The calculated results reveal a high stability of systems having large frontier orbital energy gaps, high vertical and adiabatic detachment energies, large embedding and binding energies. These predictions are in good agreement with available experimental results for enhanced stability of the X_{10}Cu clusters with $\text{X} = \text{Sn}$ and Pb . Electronic structure and chemical bonding of the global minima are also considered using density of states (DOS) of molecular orbitals, and topology analysis of the electron localizability indicator (ELI). The global minima with D_{4d} structure turn out to be magic clusters that can be rationalized in terms of their high spherical aromaticity. We would hope that appropriate experimental studies probing these enhanced stability systems could be carried out in a near future.

ASSOCIATED CONTENT

S Supporting Information. Shapes and relative energies of low-lying isomers of X_{10}Cu^- , X_{10}Ag^- , and X_{10}Au^- . Total density of states plots of global minima of X_{10}M^- and X_{10}^{2-} . This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: minh.nguyen@chem.kuleuven.be.

ACKNOWLEDGMENT

We are indebted to the KULeuven Research Council (GOA, IDO, and IUAP programs). T.B.T. thanks the Arenberg Doctoral School of for a scholarship. M.T.N. thanks the HoChiMinh City Institute for Computational Science and Technology (ICST) for supporting his stays in Vietnam.

REFERENCES

- (1) Chen, X.; Deng, K.; Liu, Y.; Tang, C.; Yuan, Y.; Tan, W.; Wang, X. *J. Chem. Phys.* **2008**, *129*, 094301.
- (2) Schafer, S.; Schafer, R. *ChemPhysChem* **2008**, *9*, 1925.
- (3) Spiekermann, A.; Hoffmann, S. D.; Fassler, T. F. *Angew. Chem., Int. Ed.* **2006**, *45*, 3459.
- (4) Scharfe, S.; Fassler, T. F.; Stegmaier, S.; Hoffmann, S. D.; Ruhland, K. *Chem.—Eur. J.* **2008**, *14*, 4479.
- (5) Kumar, V.; Kawazoe, Y. *Appl. Phys. Lett.* **2003**, *83*, 2677.
- (6) Matxain, J. M.; Piris, M.; Formoso, E.; Mercero, J. M.; Lopez, X.; Ugalde, J. M. *ChemPhysChem* **2007**, *8*, 2096.
- (7) Cui, L. F.; Huang, X.; Wang, L. M.; Li, J.; Wang, L. S. *Angew. Chem., Int. Ed.* **2007**, *46*, 742.
- (8) Lu, J.; Nagase, S. *Chem. Phys. Lett.* **2003**, *372*, 394.
- (9) Li, G.; Zhang, X.; Tang, Z.; Gao, Z. *Chem. Phys. Lett.* **2002**, *359*, 203.
- (10) Xing, X.; Tian, Z.; Liu, H.; Tang, Z. *Rapid. Commun. Mass Spectrom.* **2003**, *17*, 1411.
- (11) (a) Tai, T. B.; Nguyen, M. T. *Chem. Phys. Lett.* **2010**, *492*, 290. (b) Tai, T. B.; Nguyen, H. M. T.; Nguyen, M. T. *Chem. Phys. Lett.* **2011**, *502*, 187.
- (12) Chen, Z.; Neukermans, S.; Wang, X.; Janssens, E.; Zhou, Z.; Silverans, R. E.; King, R. B.; Schleyer, P. v. R.; Lievens, P. *J. Am. Chem. Soc.* **2006**, *128*, 12829.

- (13) Kumar, V.; Singh, A. K.; Kawazoe, Y. *Nano Lett.* **2004**, *4*, 677.
- (14) Rajesh, C.; Majumder, C. *J. Chem. Phys.* **2008**, *128*, 024308.
- (15) (a) Ngan, V. T.; Grune, P.; Claes, P.; Janssens, E.; Fielicke, A.; Nguyen, M. T.; Lievens, P. *J. Am. Chem. Soc.* **2010**, *132*, 15589. (b) Veldeman, N.; Gruene, P.; Fielicke, A.; Claes, P.; Ngan, V. T.; Nguyen, M. T.; Lievens, P. Endohedrally Doped Silicon Clusters. In *Handbook of Nanophysics. Clusters and Fullerenes*; Sattler, Klaus D., Ed.; CRC Press: Boca Raton, FL, 2010; Chapter 5.
- (16) (a) Gopakumar, G.; Wang, X.; Lin, L.; De Haeck, J.; Lievens, P.; Nguyen, M. T. *J. Phys. Chem. C* **2009**, *113*, 10858. (b) Hou, X. J.; Gopakumar, G.; Lievens, P.; Nguyen, M. T. *J. Phys. Chem. A* **2007**, *111*, 13544.
- (17) Neukermans, S.; Wang, X.; Veldeman, N.; Janssens, E.; Silverans, R. E.; Lievens, P. *Int. J. Mass. Spectrom.* **2006**, *252*, 145.
- (18) Wang, J.; Han, J. G. *J. Chem. Phys.* **2005**, *123*, 244303.
- (19) Li, X. J.; Su, K. H. *Theor. Chem. Acc.* **2009**, *124*, 345.
- (20) Saunders, M. J. *Comput. Chem.* **2003**, *25*, 621.
- (21) Frisch, M. J.; et al. *Gaussian 03*, Revision D.02; Gaussian, Inc.: Wallingford, CT, 2004.
- (22) Werner, H. J.; et al. *Molpro*, a package of ab initio programs, version 2008.1.
- (23) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648.
- (24) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B.* **1988**, *37*, 785.
- (25) Tai, T. B.; Nguyen, M. T. *J. Chem. Theor. Comput.* **2011**, *7*, 1119.
- (26) Stevens, W. J.; Krauss, M.; Basch, H.; Jasien, P. R. *Can. J. Chem.* **1992**, *70*, 612.
- (27) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 270.
- (28) Bartlett, R. J.; Musial, M. *Rev. Mod. Phys.* **2007**, *79*, 291 and references therein.
- (29) Peterson, K. A. *J. Chem. Phys.* **2003**, *119*, 11099.
- (30) Tenderholt, A. *Pymolyze: a program to analyze quantum chemical computations*. 2006
- (31) Kohout, M. *DGrid-4.2*; Max-Planck Institut für Chemische Physik und Fester Stoffe: Dresden, 2007.
- (32) Laaksonen, L. *J. Mol. Graph.* **1992**, *10*, 33.
- (33) Wagner, F. R.; Bezuly, V.; Kohout, M.; Grin, Y. *Chem.—Eur. J.* **2007**, *13*, 5724.
- (34) Kohout, M.; Wagner, F. R.; Grin, Y. *Theor. Chem. Acc.* **2002**, *108*, 150.
- (35) (a) Tai, T. B.; Nguyen, M. T. *Chem. Phys. Lett.* **2010**, *489*, 75. (b) Ngan, V. T.; Nguyen, M. T. *J. Phys. Chem. A* **2010**, *114*, 7609. (c) Holtz, T.; Veldeman, N.; De Haeck, J.; Veszpremi, T.; Lievens, P.; Nguyen, M. T. *Chem.—Eur. J.* **2009**, *15*, 3970. (d) Lin, L.; Holtz, T.; Gruene, P.; Claes, P.; Meijer, G.; Fielicke, A.; Lievens, P.; Nguyen, M. T. *ChemPhysChem* **2008**, *9*, 2471.
- (36) Boldyrev, A. I.; Wang, L. S. *Chem. Rev.* **2005**, *105*, 3716.
- (37) Hirsch, A.; Chen, Z.; Jiao, H. *Angew. Chem., Int. Ed.* **2000**, *39*, 3915.