

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231574549>

Occurrence of longicaudatine, A new type of bis-indole base and bisnor-C-alkaloid H in Strychnos Species

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · JUNE 1983

Impact Factor: 4.72 · DOI: 10.1021/jo00159a015

CITATIONS

7

READS

35

9 AUTHORS, INCLUDING:



[Georges Massiot](#)

Université de Reims Champagne-Ardenne

217 PUBLICATIONS 3,093 CITATIONS

SEE PROFILE

[Catherine Mirand](#)

Université de Reims Champagne-Ardenne

34 PUBLICATIONS 340 CITATIONS

SEE PROFILE



[K. Husnu Can Baser](#)

Near East University

1,074 PUBLICATIONS 10,098 CITATIONS

SEE PROFILE



[Peter J Hylands](#)

King's College London

94 PUBLICATIONS 2,266 CITATIONS

SEE PROFILE

Occurrence of Longicaudatine, a New Type of Bis-Indole Base and Bisnor-C-alkaloid H in *Strychnos* Species

G. Massiot, M. Zeches, C. Mirand, and L. Le Men-Olivier

Faculte de Pharmacie, ERA 319 CNRS, 51096 Reims Cedex, France

C. Delaude

Université de Liège, Sart Tilman, Liège, Belgium

K. H. C. Başer,* R. Bavovada, N. G. Bisset, and P. J. Hylands

*Pharmacognosy Research Laboratories, Department of Pharmacy, Chelsea College, University of London,
London SW3 6LX, England*

J. Strömbom

Department of Pharmacognosy, Faculty of Pharmacy, Biomedicum, S-751 23 Uppsala, Sweden

R. Verpoorte

*Department of Pharmacognosy, Subfaculty of Pharmacy, Gorlaeus Laboratories, University of Leiden,
2300 RA Leiden, The Netherlands*

Received July 16, 1982

The structure of longicaudatine, a novel bis-indole alkaloid isolated from several *Strychnos* species, has been elucidated chiefly by 400-MHz ¹H NMR and ¹³C NMR spectroscopy. In some species this alkaloid co-occurs with bisnor-C-alkaloid H, an isomeric base which has similar chromatographic and chromogenic properties.

During the screening of Asian *Strychnos* material, tertiary alkaloid extracts from the root bark of *S. axillaris* Colebr., *S. ignatii* Berg., and *S. nux-vomica* L. were observed to include several minor components which on thin-layer chromatograms immediately colored blue when sprayed with iron(III) chloride-perchloric acid reagent.¹ One of these blue-coloring bases was later isolated from the root bark of *S. nux-vomica*,² but the minute amount obtained at the time precluded any attempt to determine the structure. A further quantity of the compound has now been obtained from the root bark of *S. lucida* R.Br.³ and it is also present in *S. wallichiana* Steud. ex DC.⁴ The substance appears to be identical with an alkaloid occurring in greater amount in the stem bark of the African

species *S. dolichothyrsa* Gilg ex Onochie et Hepper⁵ and also detected in *S. urceolata* Leeuwenberg,⁶ *S. afzelii* Gilg,⁷ and *S. chrysophylla* Gilg.⁴

The various products have proved to be closely related to, if not identical with, each other. Initially, the spec-

(1) (a) Bisset, N. G.; Phillipson, J. D. *Lloydia* 1976, 39, 263. (b) Bisset, N. G.; Başer, K. H. C.; Phillipson, J. D.; Bohlin, L.; Sandberg, F. *Ibid.* 1977, 40, 546.

(2) Başer, K. H. C. Ph. D. Thesis, University of London, 1978, pp 82, 186.

(3) Bavovada, R., 1980, unpublished results.

(4) Strömbom, J., 1981, unpublished results.

(5) (a) Verpoorte, R. Ph.D. Thesis, University of Leiden, 1976, p 115. (b) Verpoorte, R.; Verzijl, M. J.; Baerheim Svendsen, A. *Planta Med.* 1982, 34, 21.

(6) Verpoorte, R.; Kodde, E. W.; Baerheim Svendsen, A. *Planta Med.* 1978, 34, 62.


(7) Verpoorte, R.; Kodde, E. W.; van Doorne, H.; Baerheim Svendsen, A. *Planta Med.* 1978, 33, 237.

* Present address: Faculty of Pharmacy, University of Anatolia, Eskişehir, Turkey.


trosopic data were interpreted as indicating that the compound had the structure of the dimeric base bisnor-C-alkaloid H, but acid hydrolysis experiments established that, in fact, two dimeric alkaloids were involved. Reevaluation of the data was in progress when the draft of a proposed paper by the French group was received. This independent study revealed that the root bark of *S. longicaudata* Gilg from Zaire is an important source of the two blue-coloring dimeric alkaloids: one was identified as a novel type of dimer, for which the name longicaudatine is proposed, and the other as bisnor-C-alkaloid H.⁸ Longicaudatine has also been isolated from the root bark of *S. ngouniensis* Pellegr.⁸ Subsequent comparison have shown that the various research groups have been working with one or the other or both of these compounds, and in order to avoid unnecessary duplication in the literature, this joint paper is being published.

Results and Discussion

Longicaudatine (1). In its mass spectrum this alkaloid shows a molecular ion at m/z 568 which analyses for C₃₈H₄₀N₄O (calcd m/z 568.320, found m/z 568.318). Major fragments are seen at m/z 249, 250, 251, and 319; the first three of these peaks are typical for indoloquinolizidines such as usambarensine.⁹ In the IR spectrum there are absorption bands due to the presence of NH (3420 and 3290 cm⁻¹) and double bonds (1630 and 1600 cm⁻¹).



A more detailed understanding of the structure of the molecule has been gained from its 400-MHz ¹H NMR spectrum in which most of the hydrogens give separated resonances. Signals for eight aromatic hydrogens, an exchangeable NH (at δ 9.6), and an ethylidene side chain (q, δ 5.60; d, δ 1.50) are observed. Other lower field signals are those assigned to H-17' (s, δ 6.06) and to the system of mutually coupled hydrogens CH(19)-CH₂(18)-O (brt, δ 6.08; 2 dd, δ 4.25 and 4.45) previously recognized in the spectra of strychnine (2)¹⁰ and related alkaloids. The assignments of the ¹H NMR spectrum presented in Table I are based on extensive decoupling experiments.




The structure **1** proposed for longicaudatine is based to a great extent on the ¹³C NMR spectra. At low field there are 20 signals (11 d and 9 s) and at higher field 18 signals (1 q, 9 t, 7 d, and 1 s). Together, these account

Table I. Longicaudatine (1) and Strychnine (2)
400-MHz ¹H NMR Spectral Data (Selected Values Only)

atoms	chemical shift, ^a δ	
	(2)	(1)
H-2	3.85	3.62 (d, 14)
H-3	3.92	3.83 (br t)
H-5a	3.18	3.15 (dd, 15, 5)
H-5b	2.86	2.64 (ddd, 15, 7, 1.5)
H-6a	1.87	1.67 (dd, 12, 5)
H-6b		1.60 (ddd, 12, 7, 1.5)
H-14a	2.34	2.30 (m)
H-14b	1.43	1.41 (br d)
H-15	3.13	3.17 (m)
H-16	1.25	1.48 (dt, 12, 2.5)
H-17	4.27	4.10 (d, 2.5)
H-18a	4.05	4.25 (dd, 14, 6)
H-18b	4.13	4.45 (dd, 14, 7)
H-19	5.88	6.08 (br t)
H-21a	3.69	3.75 (br d, 15)
H-21b	2.71	2.80 (br d, 15)
H-3'		4.20 (br)
H-15'		3.40 (t, 5)
H-17'		6.06 (s)
H-18'		1.50 (d, 7)
H-19'		5.60 (a, 7)
H-21'a		3.07 (br d, 13)
H-21'b		3.60 (br d, 13)

^a Multiplicities and coupling constants (in hertz) are given in parentheses.

carbons in the molecule. Apart from the slight discrepancies caused by the attachment between the two halves of the molecule, there is general agreement between the ¹³C NMR spectra of the pertinent parts of strychnine (2), geissoschizine (4), and longicaudatine (1) (Table II). The



signals due to the strychnan half also show much similarity with the spectrum of bisnordihydrotoxiferine (3a) (Table II).¹¹

The absence of low-field aminomethylene signals suggests a *cis*-quinolizidine arrangement for the corynanthean part of longicaudatine, as in geissoschizine,¹² geissospermine,¹³ and various other yohimbine- and heteroyohimbine-type alkaloids.¹⁴ That the relative configuration of the strychnan part is the same as in strychnine is clear from the NMR spectra: by the excellent agreement between the chemical shifts of C-2, C-3, C-7, and C-15 in longicaudatine and strychnine and by the observation of the appropriate coupling constants between H-2, H-16, and H-17.¹⁰

Bisnor-C-alkaloid H (3b). The second blue-coloring alkaloid isolated is isomeric with longicaudatine. Its mass spectrum shows a molecular ion at m/z 568 and fragments are observed at m/z 550 (M^+ - 18), 537 (M^+ - 31), and 430 (M^+ - 138); this fragmentation pattern is similar to

(8) Massiot, G.; Zeches, M.; Mirand, C.; Le Men-Olivier, L.; Delaude, C., unpublished manuscript.

(9) (a) Angenot, L.; Bisset, N. G. *J. Pharm. Belg.* **1971**, 26, 585. (b) Angenot, L.; Coune, C. A.; Tits, M. J. G.; Yamada, K. *Phytochemistry* **1978**, 17, 1687.

(10) (a) Carter, J. C.; Luther, G. W., III; Long, T. C. *J. Magn. Reson.* **1974**, 15, 122. (b) Galeffi, C.; Nicoletti, M.; Messina, I.; Marini-Bettolo, G. B. *Tetrahedron* **1979**, 35, 2545.

(11) Verpoorte, R.; Hylands, P. J.; Bisset, N. G. *Org. Magn. Reson.* **1977**, 9, 567.

(12) Damak, M.; Ahond, A.; Potier, P.; Janot, M.-M. *Tetrahedron Lett.* **1976**, 4731.

(13) Chiaroni, A.; Riche, C.; Pais, M.; Goutarel, R. *Tetrahedron Lett.* **1976**, 4729.

(14) Wenkert, E.; Chang, C. J.; Chawla, H. P. S.; Cochran, D. W.; Hagaman, E. W.; King, J. C.; Orita, K. *J. Am. Chem. Soc.* **1976**, 98, 3645.

Table II. ^{13}C NMR Spectral Data of Longicaudatine (1), Strychnine (2), Bisnordihydrotoxiferine (3a), and Geissoschizine (4)


atom	chemical shift, ^e δ			atom	chemical shift, ^e δ	
	1	2	3a		1	4
C-2	60.5	60.1	72.3	C-2'	134.3 ^a	132.8
C-3	59.9	60.1	68.0	C-3'	54.7	53.6
C-5	51.4 ^b	50.2	52.8 ^c	C-5'	51.4 ^b	50.5
C-6	39.9	42.8 ^d	42.6	C-6'	18.3	20.4
C-7	51.0	51.9	54.3	C-7'	108.1	108.1
C-8	136.9 ^a	132.6	137.2	C-8'	127.8	126.4
C-9	119.4	122.3	119.1	C-9'	117.9	118.2
C-10	122.6	124.3	122.6	C-10'	118.9	119.6
C-11	128.1 ^e	128.6	128.2	C-11'	121.8	121.9
C-12	108.5	116.3	107.0	C-12'	110.5	110.9
C-13	147.0	142.2	146.1	C-13'	135.6 ^a	136.5
C-14	27.1	26.8	24.5	C-14'	32.3	33.8
C-15	32.2	31.5	29.8	C-15'	38.6	27.7
C-16	42.7	48.2	117.8	C-16'	116.5	107.5
C-17	82.6	77.5	129.9	C-17'	127.0 ^e	161.5
C-18	64.9	64.6	12.9	C-18'	13.1	13.1
C-19	126.2	127.7	115.8	C-19'	120.7	121.9
C-20	142.1	140.3	141.1	C-20'	132.0 ^a	133.1
C-21	53.2 ^b	52.7	54.8 ^c	C-21'	55.4 ^b	59.1
C-22		42.3 ^d				

^{a-d} These values may be interchanged. ^e Chemical shifts are relative to Me_4Si .

Table III. Occurrence of Longicaudatine and Bisnor-C-alkaloid H in *Strychnos* Species

species	longicaudatine				bisnor-C-alkaloid H identified by				
	UV	IR	MS	NMR	TLC	UV	IR	MS	¹ H NMR
	identified by								
	TLC								
<i>S. longicaudata</i>	+	+	+	+					
<i>S. ngouniensis</i>	+	+	+	+					
<i>S. lucida</i>	+	+	+	+					
<i>S. ignatii</i>	+	+		+					
<i>S. nux-vomica</i>	+	+		+					
<i>S. wallichiana</i>	+	+		+					
<i>S. chrysophylla</i>	+								
<i>S. dolichothyrsa</i>	+	+		+					
<i>S. afzelii</i>	+	+		+					
<i>S. urceolata</i>	+	+		+					


that given by the toxiferine derivative alcuronium.¹⁵ The UV spectrum of bisnor-C-alkaloid H shows the presence of the typical toxiferine-type chromophore, and the IR spectrum is like that of longicaudatine in having absorption bands at 1640 and 1600 cm^{-1} (enamine system). The ^1H NMR spectrum has two singlets at δ 6.27 and 6.33 (H-17 and H-17'), a triplet at δ 5.57 (H-19), a quartet at δ 5.36 (H-19'), a singlet at δ 5.34 (H-2 and H-2'), a doublet at δ 4.33 (H-18), and a multiplet at δ 3.70 (H-15 and H-15'). These data accord well with those for bisnordihydrotoxiferine (3a), taking into account the effects of the hydroxyl group introduced at C-18.¹⁶ Further evidence for the structure 3b has been obtained by acid hydrolysis,



which yields two monomers, Wieland-Gumlich aldehyde and 18-deoxy Wieland-Gumlich aldehyde, both identified by co-TLC with reference samples and by using various solvent systems and the selective spray reagents iron(III)

chloride in perchloric acid and cerium(IV) sulfate in sulfuric acid. Similar treatment of longicaudatine does not result in any noticeable hydrolysis.

The occurrence of the two isomeric alkaloids is summarized in Table III. Longicaudatine (1) is found in species containing mainly toxiferine-type alkaloids but also in those having only strychnine-type alkaloids. It is the first bis-indole alkaloid similar to geissospermine to be isolated from the genus *Strychnos* and is probably formed by an enamine coupling between geissoschizal and Wieland-Gumlich aldehyde (see eq 1); both alkaloids are



known to be present in *S. nux-vomica*.¹¹ Bisnor-C-alkaloid H (3b) occurs in plants together with the other two tertiary toxiferine-type dimers bisnordihydrotoxiferine (3a) and caracurine V. Only longicaudatine has been detected in the Asian species of *Strychnos*, and there is no evidence as yet for the concomitant presence of bisnor-C-alkaloid H. The blue-coloring alkaloid noted on the chromatograms of *S. axillaris* extracts is more polar than either of the two compounds described here, and the elucidation of its structure awaits more detailed investigation.

(15) Angenot, L. Ph.D. Thesis, Université de Liège, 1973, p 111.

(16) Verpoorte, R.; Baerheim Svendsen, A. *Lloydia* 1976, 39, 357.

(17) Heimberger, S. I.; Scott, A. I. *J. Chem. Soc., Chem. Commun.* 1973, 217.

Longicaudatine has strong reserpine-like activity, which is different from the activities described by Sandberg et al.¹⁸ for fractionated extracts of *S. longicaudata*.

Experimental Section

Longicaudatine (1): mp 350 °C dec; $[\alpha]_D^{+141}$ (c 0.5, CHCl₃); UV λ_{\max} 223 nm (log *e* 4.66), 270 (sh), 284 (4.28), 290 (4.26), 307 (sh, 3.97); in perchloric acid a strong absorption maximum appears at 268 nm; IR ν_{\max} (KBr) 3420, 2900, 1630, 1600, 1480, 740 cm⁻¹. The 400-MHz ¹H NMR spectrum was recorded in CDCl₃ on material containing 0.25 molecule of acetone of crystallization (Table I). ¹³C NMR spectra were recorded at 62 and 25.15 MHz, in CDCl₃. Proton-noise-decoupled, off-resonance-decoupled, and selective decoupled (δ 9.6, 6.06, and 4.27 in the ¹H NMR spectrum) spectra were determined (see Table II). On TLC plates a blue color is obtained with the spray reagents 0.2 M iron(III) chloride in 35% aqueous perchloric acid and 1% cerium(IV) sulfate in 10% aqueous sulfuric acid.

(18) Sandberg, F.; Lunell, E.; Ryrberg, K. J. *Acta Pharm. Suec.* 1969, 6, 79.

Bisnor-C-alkaloid H (3b): UV λ_{\max} 293, 322 nm (sh); IR ν_{\max} (KBr) 2920, 1640, 1600, 1480, 1460, 740 cm⁻¹. The 300-MHz ¹H NMR spectrum was recorded in CDCl₃ (see text). On TLC plates a blue color was obtained with the spray reagent 0.2 M iron(III) chloride in 35% aqueous perchloric acid and a purple color with 1% cerium(IV) sulfate in 10% aqueous sulfuric acid.

Acknowledgment. The French group thanks Dr. S. K. Kan for 400-MHz ¹H NMR spectra and V. Michon for ¹³C NMR spectra. They are especially grateful to Prof. M. Aourousseau and Dr. O. Albert for pharmacological evaluation of longicaudatine and to Prof. J. Levy for stimulating discussions. C. D. acknowledges support by the Ministère de la Coopération au Développement, Belgium. The Chelsea College group is greatly indebted to Dr. Hawkes of the ULIRS WH-400 NMR Service at Queen Mary College, University of London, for detailed spectra of longicaudatine and for carrying out numerous decoupling experiments. R. V. thanks C. Erkelens for recording ¹H and ¹³C NMR spectra.

Registry No. 1, 85335-06-4; 3b, 67739-70-2.