

Flavonol Glycosides and Novel Iridoid Glycoside from the Leaves of *Morinda citrifolia*

Shengmin Sang,[†] Xiaofang Cheng,[‡] Nanqun Zhu,[†] Ruth E. Stark,[‡] Vladimir Badmaev,[§]
Geetha Ghai,[†] Robert T. Rosen,[†] and Chi-Tang Ho*,[†]

Department of Food Science and Center for Advanced Food Technology, Rutgers University,
65 Dudley Road, New Brunswick, New Jersey 08901-8520, Department of Chemistry,
College of Staten Island, City University of New York, New York 10314, and Sabinsa Corporation,
121 Ethel Road, Piscataway, New Jersey 08854

One new iridoid glycoside and five known flavonol glycosides have been isolated from the leaves of *Morinda citrifolia*. The new iridoid exists as an epimeric mixture in solution. Complete assignments of the proton and carbon chemical shifts for the individual epimers were accomplished on the basis of high-resolution 1D and 2D NMR data. Their antioxidative activities were measured. All of these compounds showed DPPH free radical scavenging activity at the concentration of 30 μ M.

Keywords: Iridoid; epimer; *Morinda citrifolia*; Rubiaceae; DPPH; antioxidant; flavonoid

INTRODUCTION

Morinda citrifolia (Rubiaceae), commonly known as noni, is native to certain regions bordering the Indian Ocean, such as Tahiti and Samoa, as well as Hawaii. The plant is a small evergreen tree growing in the open coastal regions and in forest areas up to about 1300 feet above sea level. This plant is identifiable by its straight trunk, large green leaves, and its distinctive, ovoid, "grenade-like" yellow fruit. The fruit can grow to a size of 12 cm and results from coalescence of the inferior ovaries of many closely packed flowers. It has a foul taste and a soapy smell when mature. The bark, stem, roots, leaves, and fruits have been used traditionally as a folk remedy for many diseases including diabetes, hypertension, and cancer (1, 2). In previous chemical studies, anthraquinones, including damnacanthal, 7-hydroxy-8-methoxy-2-methyl-anthraquinone, morenone 1, and morenone 2, have been found in the roots of noni (3, 4). In the seeds of noni, ricinoleic acid has been found (5). From the heartwood, two known anthraquinones (morindone and physcion) and one new anthraquinone glycoside have been isolated (6). Studies on the chemical components of the flowers of noni have resulted in the identification of one anthraquinone glycoside and two flavone glycosides (7, 8). The volatile compounds in the ripe fruits are characterized by a large amount of carboxylic acids, especially octanoic acid and hexanoic acid (9). Several nonvolatile compounds, including acetyl derivatives of asperuloside and glucose, have been identified in noni fruit (10). Recently, our research group reported five novel glycosides, including glycosides of octanoic and hexanoic acid, from the Hawaiian noni fruits (11, 12). However, only two compounds, β -sitosterol and ursolic acid, have been isolated previously from the leaves of this plant (13). We have examined the chemical components of the polar *n*-butanol-soluble

fraction of the ethanol extract of noni leaves (14). In this report, we describe the isolation and structure elucidation of a new iridoid glucoside (1), named citrifolinin B, together with those of five known flavonol glycosides from the *n*-butanol fraction of *Morinda citrifolia* extract. The DPPH free radical scavenging activities for these compounds were also determined.

MATERIALS AND METHODS

General Procedures. ¹H (400 and 600 Hz), ¹³C (100 and 150 Hz), and 2D NMR spectra were obtained on a Varian AM-600 and AM-400 NMR spectrometer with TMS as internal reference. APCI MS was obtained on a Fisons/VG Platform II mass spectrometer. Thin-layer chromatography was performed on Sigma-Aldrich TLC plates (250 μ m thickness, 2–25 μ m particle size), with compounds visualized by spraying with 5% (v/v) H₂SO₄ in ethanol solution.

Plant Material. The dried leaves of *Morinda citrifolia* were collected from Bengal, India in 1999 and were identified by Dr. Vladimir Badmaev at the Sarbinsa Corporation. A voucher specimen (HS16) was deposited in the Department of Food Science, Cook College, Rutgers University.

Extraction and Isolation Procedures. The dried noni leaves (5 kg) were extracted with 95% EtOH (4 L) at 50 °C for 1 day. The extract was concentrated to dryness under reduced pressure, and the residue was suspended in water (500 mL) and partitioned successively with hexane (3 \times 500 mL), ethyl acetate (3 \times 500 mL), and *n*-butanol (3 \times 500 mL). The butanol fraction was subjected to a Diaion HP-20 column, and eluted with a water–EtOH (water, 30% EtOH, 70% EtOH, 95% EtOH) solvent system. The fraction (5 g) eluted by 30% EtOH was subjected to silica gel column chromatography with an ethyl acetate–MeOH–H₂O–hexane solvent system (10:1:1:0.5 \rightarrow 5:1:1:0). In total, 10 fractions were collected. Fraction 1 eluted by ethyl acetate–MeOH–H₂O–hexane (10:1:1:0.5) was subjected to RP-18 silica gel column chromatography with 30% MeOH to give 3 fractions (I–III). Fraction II was rechromatographed on a Sephadex LH-20 eluted by 95% EtOH to give compound 2 (16 mg). Fractions 3 and 4 eluted by ethyl acetate–MeOH–H₂O (10:1:1) were subjected to Sephadex LH-20 eluted by 95% EtOH to afford 21 mg of compound 3 and 25 mg of compound 4, respectively. Fraction 6 eluted by ethyl acetate–MeOH–H₂O (5:1:1) was subjected to RP-18 silica gel column chromatography with 30% MeOH, and then Sephadex LH-20 eluted by 95% EtOH to give 100 mg of compound 1, 70 mg of compound 5, and 80 mg of compound 6.

* To whom correspondence should be addressed. Phone: 732-932-9611, ext. 215. Fax: 732-932-6776. E-mail: ho@aesop.rutgers.edu.

[†] Rutgers University.

[‡] City University of New York.

[§] Sabinsa Corporation.




Figure 1. Structures of compounds 1–6.

Spectral Identification of Known Compounds. Quercetin-3-*O*- β -D-glucopyranoside (**2**): yellow powder. APCI-MS, m/z : 463 [M-H]⁻. ¹H NMR (CD₃OD, 400 MHz): δ 7.73 (1H, d, J = 2.0 Hz), 7.60 (1H, dd, J = 8.8, 2.0 Hz), 6.88 (1H, d, J = 8.8 Hz), 6.39 (1H, d, J = 1.6 Hz), 6.21 (1H, d, J = 1.6 Hz), 5.24 (1H, d, J = 7.8 Hz); 3.20–3.80 (6H, m). ¹³C NMR (CD₃OD) δ : 179.5 (s), 166.6 (s), 163.1 (s), 158.6 (s), 150.0 (s), 146.0 (s), 135.7 (s), 123.3 (s), 123.2 (d), 117.7 (d), 116.1 (d), 108.7 (s), 105.6 (d), 104.5 (d), 95.0 (d), 78.5 (d), 78.2 (d), 75.8 (d), 71.3 (d), 62.7 (t).

Kaempferol-3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**3**): yellow powder. APCI-MS, m/z : 593 [M-H]⁻. ¹H NMR (CD₃OD, 400 MHz): δ 8.07 (2H, d, J = 8.8 Hz), 6.90 (2H, d, J = 8.8 Hz), 6.40 (1H, d, J = 1.8 Hz), 6.21 (1H, d, J = 1.8 Hz), 5.15 (1H, d, J = 7.6 Hz), 4.54 (1H, s), 3.20–3.90 (10H, m), 1.14 (3H, d, J = 5.8 Hz). ¹³C NMR (CD₃OD) δ : 179.5 (s), 166.4 (s), 163.0 (s), 161.6 (s), 159.5 (s), 158.6 (s), 135.6 (s), 132.5 (d), 122.9 (s), 116.2 (d), 105.7 (s), 104.8 (d), 102.5 (d), 100.2 (d), 95.1 (d), 78.3 (d), 77.3 (d), 75.9 (d), 74.0 (d), 72.4 (d), 72.2 (d), 71.6 (d), 69.8 (d), 68.7 (t), 18.0 (q).

Quercetin-3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**4**): yellow powder. APCI-MS, m/z : 609 [M-H]⁻. ¹H NMR (CD₃OD, 400 MHz): δ 7.68 (1H, d, J = 2.0 Hz), 7.64 (1H, dd, J = 8.8, 2.0 Hz), 6.85 (1H, d, J = 8.8 Hz), 6.39 (1H, d, J = 1.6 Hz), 6.21 (1H, d, J = 1.6 Hz), 5.10 (1H, d, J = 7.8 Hz), 4.56 (1H, s), 3.20–3.90 (10H, m), 1.14 (3H, d, J = 5.8 Hz). ¹³C NMR (CD₃OD) δ : 179.4 (s), 166.1 (s), 162.9 (s), 159.4 (s), 158.5 (s), 149.9 (s), 145.9 (s), 135.8 (s), 124.3 (s), 123.3 (d), 118.4 (d), 117.3 (d), 105.7 (s), 103.1 (d), 102.0 (d), 100.7 (d), 95.6 (d), 78.8 (d), 77.8 (d), 75.4 (d), 74.1 (d), 72.5 (d), 72.0 (d), 71.7 (d), 69.6 (d), 68.7 (t), 18.4 (q).

Quercetin-3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)-[α -L-rhamnopyranosyl-(1 \rightarrow 6)]- β -D-galactopyranoside (**5**): yellow powder. APCI-MS, m/z : 771 [M-H]⁻; ¹H NMR (C₅D₅N, 600 MHz) δ : 8.39 (1H, d, J = 2.0 Hz), 8.14 (1H, dd, J = 8.8, 2.0 Hz), 7.39 (1H, d, J = 8.8 Hz), 6.91 (1H, d, J = 1.6 Hz), 6.79 (1H, d, J = 1.6

Hz), 6.08 (1H, d, J = 7.8 Hz), 5.81 (1H, d, J = 7.8 Hz), 5.37 (1H, s), 3.80–4.60 (16H, m), 1.25 (3H, d, J = 5.8 Hz). ¹³C NMR (C₅D₅N) δ : 178.7 (s), 163.9 (s), 162.0 (s), 158.0 (s), 156.9 (s), 150.8 (s), 146.8 (s), 136.0 (s), 122.9 (s), 122.1 (d), 117.9 (d), 116.9 (d), 108.8 (s), 104.4 (d), 102.5 (d), 101.5 (d), 100.4 (d), 95.1 (d), 79.2 (d), 78.5 (d), 78.4 (d), 77.5 (d), 75.4 (d), 74.8 (d), 73.3 (d), 72.5 (d), 72.1 (d), 71.3 (d), 71.1 (d), 69.6 (d), 68.4 (t), 62.4 (t), 18.5 (q).

Kaempferol-3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)-[α -L-rhamnopyranosyl-(1 \rightarrow 6)]- β -D-galactopyranoside (**6**): yellow powder. APCI-MS, m/z : 755 [M-H]⁻. ¹H NMR (C₅D₅N, 600 MHz) δ : 8.43 (1H, d, J = 8.8 Hz), 7.31 (1H, d, J = 8.8 Hz), 7.05 (1H, s), 6.79 (1H, s), 5.83 (1H, d, J = 7.6 Hz), 5.30 (1H, s), 3.80–4.60 (16H, m), 1.25 (3H, d, J = 5.8 Hz). ¹³C NMR (C₅D₅N) δ : 178.4 (s), 164.2 (s), 162.4 (s), 162.2 (s), 158.9 (s), 157.3 (s), 136.0 (s), 132.2 (d), 121.1 (s), 116.5 (d), 107.1 (s), 104.5 (d), 102.8 (d), 101.8 (d), 100.7 (d), 95.5 (d), 79.5 (d), 78.8 (d), 78.7 (d), 77.8 (d), 76.2 (d), 75.1 (d), 74.2 (d), 72.9 (d), 72.4 (d), 71.7 (d), 71.5 (d), 69.9 (d), 68.9 (d), 62.8 (t), 18.5 (q).

Determination of the DPPH Radical Scavenging Capacity. This method was adapted from Chen and Ho (26). DPPH radicals were prepared in ethanol as a 1.0×10^{-4} M solution. This DPPH solution was mixed with different tested compounds (final concentration was 30 μ M) and kept in a dark area for 0.5 h. The absorbance of the samples was measured on a spectrophotometer (Milton Roy, model 301) at 517 nm against a blank of ethanol without DPPH. All tests were run in triplicate and averaged.

RESULTS AND DISCUSSION

The *n*-butanol fraction of *Morinda citrifolia* leaves extract was chromatographed successively on Diaion HP-20, silica gel, Sephadex LH-20, and RP-18 silica gel columns to afford one new compound and five known




Figure 2. Significant HMBC (H—C) and NOESY correlations of compound **1**.

Table 1. ^{13}C (150 MHz) and ^1H (600 MHz) NMR Spectral Data for Compounds **1a** and **1b** (CD_3OD) (δ in ppm, J in Hz)

	1a		1b	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}
1	94.7 d	5.95 s	94.1 d	5.93 s
3	154.8 s	7.57 s	154.5 s	7.56 s
4	107.5 s		107.4 s	
5	33.7 d	3.32 m	33.6 d	3.32 m
6	60.9 d	3.88 d 3.6	60.6 d	3.87 d 3.6
7	58.9 d	3.58 d 3.6	58.2 d	3.56 d 3.6
8	82.5 s		82.2 s	
9	46.9 d	2.34 d, 9.6	46.8 d	2.33 d, 9.6
10	98.4 d	4.45 s	98.4 d	4.34 s
11	168.8 s		168.7 s	
11-OCH ₃	52.0 s	3.79 s	52.0 s	3.79 s
1'	100.2 d	4.63 d, 7.8	100.0 d	4.62 d, 7.8
2'	74.6 d	3.19 m	74.6 d	3.19 m
3'	78.3 d	3.32 m	78.3 d	3.32 m
4'	71.6 d	3.31 m	71.6 d	3.31 m
5'	78.0 d	3.33 m	78.0 d	3.33 m
6'	62.8 t	3.92 d, 9.6	62.8 t	3.92 d, 9.6
		3.71 dd, 6.6, 11.4		3.71 dd, 6.6, 11.4

compounds. Their structures (Figure 1) were established by interpretation and full assignments of 1D and 2D NMR spectroscopic data and comparison with literature data.

Compound **1** was obtained as an amorphous solid. The molecular formula, $\text{C}_{17}\text{H}_{22}\text{O}_{12}$ was determined by negative-ion APCI-MS ($[\text{M} - \text{H}]^-$ at m/z 417), as well as from its ^{13}C NMR data, and indicated seven degrees of unsaturation. Of great interest was the doubling of the signals in both the ^1H and the ^{13}C NMR spectra of **1**,

indicating the presence of an epimeric mixture. The ratio of the epimers, as deduced from the integration in the ^1H NMR spectrum, was 5:4 (**1a**:**1b**). The ^1H NMR spectrum of **1a** showed a singlet for a carbomethoxy group at δ 3.79, a singlet for the C-1 proton at δ 5.95, a doublet ($J = 2.0$ Hz) for the characteristic C-3 proton of iridoids at δ 7.57, two doublets ($J = 3.6$ Hz, each) for the C-6, C-7-epoxy protons at δ 3.88 and 3.58, and a doublet ($J = 7.8$ Hz) for the C-1' proton at δ 4.63, suggesting that the cyclopentanopyran ring system and the sugar moiety of **1a** were identical to those of 6 β ,7 β -epoxysplendoside (**15**). The major difference was at position 10 in the ^1H NMR spectrum. The methene group at position 10 of 6 β ,7 β -epoxysplendoside was substituted by a methine group bearing two oxygen substituents (δ 4.45, s, 1H) in compound **1a**. This also was supported by the ^{13}C NMR data (δ 98.4, d). In addition, **1a** had seven degrees of unsaturation, whereas 6 β ,7 β -epoxysplendoside only had six degrees of unsaturation. These results indicated that an epoxyl group existed between C-8 and C-10.

The ^{13}C NMR spectrum of **1a** exhibited 17 carbon signals (Table 1), with 10 representing the aglycon, one methoxy group (δ 52.0), and six for the glucopyranose unit (δ 100.2, d, C-1'; δ 74.6, d, C-2'; δ 78.3, d, C-3'; δ 71.6, d, C-4'; δ 78.0, d, C-5'; and δ 62.8, t, C-6'). The β -anomeric configuration for the glucose was judged from its large $^3J_{\text{H}1,\text{H}2}$ coupling constants ($J = 7.8$ Hz) (**16**). HMBC and NOESY correlations between C-1/H-1', H-1/C-1', and H-1/H-1' (Figure 2) suggested that the β -glucopyranose unit was attached at the C-1 position of the aglycon.

The configuration of the C-6, C-7-epoxide group was confirmed from the NMR spectra of **1a** as β ($J_{\text{H}1,9} < 1$ Hz, and $\delta_{\text{C}1} = 94.7 < 99$ ppm) (**17**, **18**). To determine the stereochemistry at C-8 and C-10, NOESY measurements were carried out on **1**. In the NOESY spectrum, the presence of strong cross-peaks between H-1 and H-10a indicated that the linkage between C-8 and C-10 was α -oriented and H-10 was β in **1a**. Thus, compound **1a** was determined as citrifolinin **Ba** (Structure **1a** in Figure 1).

Compound **1b** was almost identical with compound **1a** in all respects except at position 10 (Table 1). In the ^1H NMR spectrum of compound **1b**, H-10 appeared at δ 4.34 ppm instead of δ 4.45 ppm as observed in compound **1a**. So **1b** must be the C-10 epimers of **1a**.




Figure 3. Proposed biosynthetic pathway of Citrifolinoside **B** (**1a**+**1b**).

H-10 was in α position instead of β as observed in **1a**. This was approved by the NOESY spectrum of compound **1**. H-10a showed strong cross-peak with H-1. However, H-10b did not show cross-peak with H-1. Therefore, compound **1b** was determined as citrifolinin **Bb** (Structure **1b** in Figure 1). Full assignments of the ^1H and ^{13}C NMR signals of **1a** and **1b** were accomplished using HMBC, HMQC, ^1H - ^1H COSY, TOCSY, and NOESY experiments (Table 1).

The different stereochemistry at C-10 in **1a** and **1b** could be explained by involvement of the intramolecular acetalation of compound **7** which was the $6\beta,7\beta$ -epoxy derivative of 10-dehydrogardenoside (**19**) in the proposed biogenetic pathway (Figure 3). According to this pathway, we presumed the existence of compound **7** as a low concentration in these leaves. Because of the instability, as well as the low content of this compound, we did not isolate it.

Actually, the R_f values of these two epimers are significantly different. It should be easy to separate these two compounds. However, when we tried to use the RP-C18 silica gel column, silica gel column, and the prepared TLC plate to separate these two compounds, all efforts failed because of the instability of these two compounds. There is a rapid equilibrium between them.

In addition to the new iridoid glycosides, five known flavonol glycosides, quercetin-3- O - β -D-glucopyranoside (**2**), kaempferol-3- O - α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**3**), quercetin-3- O - α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**4**), quercetin-3- O - β -D-glucopyranosyl-(1 \rightarrow 2)-[α -L-rhamnopyranosyl-(1 \rightarrow 6)]- β -D-galactopyranoside (**5**), and kaempferol-3- O - β -D-glucopyranosyl-(1 \rightarrow 2)-[α -L-rhamnopyranosyl-(1 \rightarrow 6)]- β -D-galactopyranoside (**6**), have also been isolated in this study. Their structures were identified by comparison of their NMR and MS data with those reported in the literature (11, 20, 21). It is easy to determine the kinds and the positions of sugar moieties by comparing their ^{13}C data with those in the literature (22), because the ^{13}C data are very different between glucose and galactose (22). All of these compounds are being reported from the leaves of this plant for the first time.

The antioxidative activities of isolated compounds were measured. All of these compounds showed DPPH free radical scavenging activity at the concentration of 30 μM with 7.7, 85.8, 4.5, 79.9, 81.3, and 28.6%. Flavonoids are phenolic substances isolated from a wide range of vascular plants, with over 8000 individual compounds known. It is suggested that their antioxidative activity is related to their conjugated rings and hydroxyl groups (23), so it is not a surprise that quercetin glycosides showed stronger antioxidant activity than kaempferol glycosides.

LITERATURE CITED

- (1) Hirazumi, A.; Furusawa, E.; Chou, S. C. Anticancer activity of *Morinda citrifolia* (noni) on intraperitoneally implanted lewis lung carcinoma in syngeneic mice. *Proc. West Pharmacol. Soc.* **1994**, *37*, 145–146.
- (2) Hirazumi, A.; Furusawa, E.; Chou, S. C. Immunomodulation contributes to the anticancer activity of *Morinda citrifolia* (noni) fruit juice. *Proc. West Pharmacol. Soc.* **1996**, *39*, 25–27.
- (3) Rusia, K.; Srivastava, S. K. A new anthraquinone from the roots of *Morinda citrifolia* Linn. *Curr. Sci.* **1989**, *58*, 249.
- (4) Jain, R. K.; Ravindra, K.; Srivastava, S. D. Two new anthraquinones in the roots of *Morinda citrifolia*. *Proc. Natl. Acad. Sci., India, Sect. A* **1992**, *62*, 11–13.
- (5) Daulatabad, C. D.; Mulla, G. M.; Mirajkar, A. M. Ricinoleic acid in *Morinda citrifolia* seed oil. *J. Oil Technol. Assoc. (India)* **1989**, *21*, 26–27.
- (6) Srivastava, M.; Singh, J. A new anthraquinone glycoside from *Morinda citrifolia*. *Int. J. Pharmacogn.* **1993**, 182–184.
- (7) Tiwari, R. D.; Singh, J. Structural study of the anthraquinone glycosides from the flowers of *Morinda citrifolia*. *J. Indian Chem. Soc.* **1977**, *54*, 429–430.
- (8) Singh, J.; Tiwari, R. D. Flavone glycosides from the flowers of *Morinda citrifolia*. *J. Indian Chem. Soc.* **1976**, *52*, 424.
- (9) Farine, J. P.; Legal, L.; Moreteau, B.; Le Quere, J. L. Volatile components of ripe fruits of *Morinda citrifolia* and their effects on *Drosophila*. *Phytochemistry* **1996**, *41*, 433–438.
- (10) Levand, O.; Larson, H. Some chemical constituents of *Morinda citrifolia*. *Planta Med.* **1979**, *36*, 186–187.
- (11) Wang, M. F.; Kikuzaki, H.; Csiszar, K.; Boyd, C. D.; Maunakea, A.; Fong, S. F. T.; Ghai, G.; Rosen, R. T.; Nakatani, N.; Ho, C.-T. Novel trisaccharide fatty acid ester identified from the fruits of *Morinda citrifolia* (Noni). *J. Agric. Food Chem.* **1999**, *47*, 4880–4882.
- (12) Wang, M. F.; Kikuzaki, H.; Jin, Y.; Nakatani, N.; Zhu, N.; Csiszar, K.; Boyd, C. D.; Rosen, R. T.; Ghai, G.; Ho, C.-T. Novel glycosides from noni (*Morinda citrifolia*). *J. Nat. Prod.* **2000**, *63*, 1182–1183.
- (13) Ahmad, V. U.; Bano, M. Isolation of β -sitosterol and ursolic acid from *Morinda citrifolia* Linn. *J. Chem. Soc. Pak.* **1980**, *2*, 71–72.
- (14) Sang, S. M.; He, K.; Liu, G. M.; Zhu, N.; Wang, M.; Jhoo, J. W.; Zheng, Q.; Dong, Z.; Ghai, G.; Rosen, R. T.; Ho, C.-T. Citrifolinin A, a new unusual iridoid with inhibition of Activator Protein-1 (AP-1) from the leaves of noni (*Morinda citrifolia* L.). *Tetrahedron Lett.* **2001**, *42*, 1823–1825.
- (15) Jensen, S. R.; Nielsen, B. J. Iridoid glucosides in Fouquieriaceae. *Phytochemistry* **1982**, *21*, 1623–1629.
- (16) Agrawal, P. K. NMR spectroscopy in the structural elucidation of oligosaccharides and glycosides. *Phytochemistry* **1992**, *31*, 3307–3321.
- (17) Damtoft, S.; Jensen, S. R.; Nielsen, B. J. ^{13}C and ^1H NMR spectroscopy as a tool in the configurational analysis of iridoid glucosides. *Phytochemistry* **1981**, *20*, 2717–2732.
- (18) Peng, J. N.; Feng, X. Z.; Liang, X. T. Two new iridoids from *Hedyotis chrysotricha*. *J. Nat. Prod.* **1999**, *62*, 611–612.
- (19) Uesato, S.; Ali, E.; Nishimura, H.; Kawamura, I.; Inouye, H. Four iridoids from *Randia canthioides*. *Phytochemistry* **1982**, *21*, 353–357.
- (20) Foo, L. Y.; Lu, Y.; Molan, A. L.; Woodfield, D. R.; McNabb, W. C. The phenols and prodelphinidins of white clover flowers. *Phytochemistry* **2000**, *54*, 539–548.
- (21) Agrawal, P. K. *Carbon-13 NMR of Flavonoids*; Studies in Organic Chemistry Series, 39. Elsevier: New York, 1989.
- (22) Agrawal, P. K. NMR spectroscopy in the structural elucidation of oligosaccharides and glycosides. *Phytochemistry* **1992**, *31*, 3307–3330.
- (23) Pietta, P. G. Flavonoids as antioxidants. *J. Nat. Prod.* **2000**, *63*, 1035–1042.

Received for review April 16, 2001. Revised manuscript received June 28, 2001. Accepted July 9, 2001. Funds to purchase the NMR spectrometer were provided by the New York State Higher Education Applied Technology Program, the New York State Graduate Research Initiative, and the City University of New York (CUNY). The NMR Facility is operated by the CUNY Center for Applied Biomedicine and Biotechnology (CABB) and by the College of Staten Island.