

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/15338633>

Highly κ receptor-selective dynorphin A analogues with modifications in position 3 of dynorphin A(1-11)-NH₂

ARTICLE in JOURNAL OF MEDICINAL CHEMISTRY · MARCH 1995

Impact Factor: 5.45 · DOI: 10.1021/jm00004a002 · Source: PubMed

CITATIONS

23

READS

11

9 AUTHORS, INCLUDING:



Feng-Di T Lung

Tunghai University

41 PUBLICATIONS 1,142 CITATIONS

SEE PROFILE



Jean-Philippe Meyer

Evolva Holding

21 PUBLICATIONS 466 CITATIONS

SEE PROFILE



Bih Show Lou

Chang Gung University

88 PUBLICATIONS 726 CITATIONS

SEE PROFILE



Victor J Hruby

The University of Arizona

555 PUBLICATIONS 17,650 CITATIONS

SEE PROFILE

Highly κ Receptor-Selective Dynorphin A Analogues with Modifications in Position 3 of Dynorphin A(1–11)-NH₂

Feng-Di T. Lung,[†] J.-Ph. Meyer,[†] Guigen Li,[†]
B.-S. Lou,[†] Dagmar Stropova,[‡] Peg Davis,[‡]
Henry I. Yamamura,[‡] Frank Porreca,[‡] and
Victor J. Hruby*,[†]

Departments of Chemistry and Pharmacology,
University of Arizona, Tucson, Arizona 85721

Received October 11, 1994

The heptadecapeptide dynorphin A (Dyn A(1–17); H-Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Ile-Arg-Pro-Lys-Leu-Lys-Trp-Asp-Asn-Gln-OH) was first isolated from porcine pituitary and recognized as a potent opioid agonist.^{2–4} It is now well established that there are at least three types of opioid receptors, namely, μ , δ , and κ .^{5–7} However, the physiological and pharmacological roles of these receptors still need clarification, thus requiring the design and synthesis of highly potent and selective ligands. Research in this area has expanded rapidly in the past decade, with considerable effort devoted to the development of δ and μ opioid receptor-selective peptide ligands primarily based on enkephalin.^{8–11} The potential of targeting the κ opioid receptor as an effector for analgesia has yet to be explored in detail.¹² Previous structure–activity studies have shown that the truncated derivative, dynorphin A(1–11)-NH₂, retains the high binding potency of dynorphin A at the κ receptor. Thus we and others have primarily used Dyn A(1–11)-NH₂ as a template to examine the structure–activity relationships of dynorphin.^{13,14} Since Tyr¹ and Phe⁴ are reported to be important for opioid agonist activity, the effects of the glycine residues in positions 2 and 3 of Dyn A on the relative orientations of the two aromatic rings may be biologically important.¹³ To assess these possible effects, we have substituted Gly³ by D- and L-alanine residues to form the linear peptides [D-Ala³]Dyn A(1–11)-NH₂ and [Ala³]Dyn A(1–11)-NH₂. We report here that both analogues are very potent for κ receptors and very selective for κ vs μ and κ vs δ receptors. These results suggest that substitution of lipophilic residues and/or certain D-amino acids at position 3 of Dyn A(1–11)-NH₂ may lead to novel Dyn A analogues which exhibit enhanced selectivities for κ receptors, while retaining strong affinities.

Peptide Synthesis and Purification. Dynorphin A analogues were synthesized by solid phase methods that were reported previously for the synthesis of other Dyn A analogues.^{14,15} Side chain-protected N^α-Boc amino acids were purchased from Bachem (Torrance, CA), whereas the other amino acids were synthesized by standard methods in our laboratory. The synthesized analogues were purified by RP-HPLC (linear gradient of 10–90% acetonitrile in 0.1% TFA in water over 40 min) and characterized by FAB-MS and amino acid analysis. The purity of the synthetic peptides was assessed by TLC (single spot in four different solvent

systems, ninhydrin detection) and HPLC (one single peak, UV detection at 280 and 225 nm, using two different linear gradients).

Opioid Receptor Binding Affinities and Selectivities in the Guinea Pig Brain (GPB). The peptides were evaluated for their receptor binding affinities at κ , δ , and μ receptors by measuring the inhibition of binding of [³H]U-69,593 (N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide) (κ), [³H]cyclo[D-Pen²,p-Cl-Phe⁴,D-Pen⁵]enkephalin (δ), and [³H]DAMGO ([D-Ala²,MePhe⁴,Glyol⁵]enkephalin) (μ) to opioid receptors in guinea pig brain (GPB) homogenates (Table 1).^{14,15} As shown in Table 1, substitution of Gly³ with D- and L-alanine lead to analogues 2 and 3 that displayed high affinities at the central κ opioid receptor (IC₅₀ = 0.76 and 1.1 nM, respectively) similar to that of Dyn A(1–11)-NH₂ (1) (IC₅₀ = 0.58 nM) and, rather surprisingly, greatly enhanced selectivities for κ vs μ and κ vs δ receptors (IC₅₀ ratios of 350 and 1300 for 2 and 190 and 660 for 3, respectively, compared to 17 and 44 for 1), due to poor affinities for μ and δ receptors (Table 1). The binding affinity obtained for analogue 3 differs somewhat from one previously published for [Ala³]Dyn A(1–13).¹⁶ No satisfactory explanation can be given for this discrepancy except that this latter study used a 1–13 analogue, a different radioligand, and a much shorter incubation time in the binding assay (30 min vs 180 min).

Analogue 2 is one of the most κ receptor-selective dynorphin-like peptides reported and can be compared to [N-benzyl-Tyr¹,D-Pro¹⁰]Dyn A(1–13)-NH₂, a peptide that exhibits so far the highest reported selectivity for the central κ vs μ and δ receptors ($\kappa/\mu/\delta$ K_i ratio = 1/1070/6080).¹⁷ The fact that a higher κ selectivity can be observed with both analogues 2 and 3, by incorporating the two alanine enantiomers, suggests that an important factor could be related more to the increase in lipophilicity than to a specific orientation of the methyl group of alanine. Another possibility is that replacing an α -helix-breaking residue like glycine could increase the α -helical content of the message segment of dynorphin A, which has been postulated to be important for κ site selection.¹⁸

Biological Activities in the Guinea Pig Ileum (GPI). The κ (and μ) opioid activities of these peptides were measured by their ability to inhibit the electrically evoked contraction of the guinea pig ileum (GPI) (Table 2)^{14,15} and the effect of the μ opioid receptor antagonist CTAP on the IC₅₀ value.¹⁹ The results obtained in GPI bioassay show that, as no shift in activity can be observed upon addition of the μ antagonist CTAP, all analogues tested interact only or specifically with the peripheral κ receptors. Though they are still potent at these opioid receptors, [D-Ala³]Dyn A(1–11)-NH₂ and [L-Ala³]Dyn A(1–11)-NH₂ display somewhat lower potencies than the standard 1 (IC₅₀ = 8.1 and 1.7 nM, respectively, for 2 and 3, vs 1.1 nM for 1).

Conclusion. It has been found that substitution of Gly³ with D- and L-alanine in Dyn A(1–11)-NH₂ (analogues 2 and 3) resulted in a dramatic increase in selectivity for κ receptors, when compared to Dyn A(1–11)-NH₂. Though it still is not possible to say whether these results are due to (1) an increase in lipophilicity of the peptide, (2) a possible enhancement of the α -hel-

* Author to whom reprint requests and correspondence should be addressed at the Department of Chemistry.

[†] Department of Chemistry.

[‡] Department of Pharmacology.

Table 1. Opioid Receptor Binding Affinities and Selectivities of Dyn A(1–11)-NH₂ Analogues in Guinea Pig Brain (GPB) Homogenate

analogues of Dyn A(1–11)-NH ₂	IC ₅₀ (nM) ^a			selectivity	
	κ	μ	δ	μ/κ	δ/κ
Dyn A(1–11)-NH ₂ (1)	0.58 ± 0.03	9.90 ± 2.02	25.5 ± 3.4	17	44
[D-Ala ³]Dyn A(1–11)-NH ₂ (2)	0.76 ± 0.28	260 ± 57	1000 ± 422	350	1300
[Ala ³]Dyn A(1–11)-NH ₂ (3)	1.10 ± 0.40	210 ± 40	730 ± 50	190	660

^a The radioligands used were [³H]U-69,593 (κ receptor), [³H]DAM-GO (μ receptor), and [³H]cyclo[D-Pen²,p-Cl-Phe⁴,D-Pen⁵]enkephalin (δ receptor). Results are given ± SEM.

Table 2. Bioassays with the Smooth Muscle Tissue of the Guinea Pig Ileum (GPI)

analogues of Dyn A(1–11)-NH ₂	IC ₅₀ (nM)	
	GPI	shift ^a
Dyn A(1–11)-NH ₂ (1)	1.1 ± 0.3	ns
[D-Ala ³]Dyn A(1–11)-NH ₂ (2)	8.1 ± 2.3	ns
[Ala ³]Dyn A(1–11)-NH ₂ (3)	1.7 ± 0.2	ns

^a ns: no significant shift observed with 1000 nM CTAP used as a μ antagonist. Results are given ± SEM.

ical content of the message segment of this peptide, (3) a more favorable spatial arrangement of the relative positions of the aromatic residues, or (4) a combination of these different effects, they are highly interesting, as they provide us with new lead compounds for further enhancement of potency and selectivity. Several other peptides, incorporating different modifications at position 3, are currently being synthesized in our laboratory, will be examined for their binding affinities and selectivities, and also will be studied by circular dichroism and nuclear magnetic resonance techniques, in order to determine if conformational changes take place in these analogues and if those changes can account for the increased selectivity of these analogues for κ receptors.

Acknowledgment. This work was supported by grants from the National Institute of Drug Abuse (DA04248 and DA06284) and the USPHS NS 19972.

References

- (1) Symbols and abbreviations are in accord with the recommendations of the IUPAC-IUB Commission on Nomenclature (*J. Biol. Chem.* **1972**, *247*, 977–983). All optically active amino acids are of the L variety unless otherwise stated. Other abbreviations are Dyn A, dynorphin A; GPB, guinea pig brain; GPI, guinea pig ileum; HPLC, high-performance liquid chromatography; TFA, trifluoroacetic acid; CTAP, cyclo-[D-Phe-Cys-Tyr-D-Trp-Arg-Thr-Pen-Thr-NH₂].
- (2) Goldstein, A.; Tachibana, S.; Lowney, L. I.; Hunkapiller, M.; Hood, L. Dynorphin-(1–13), an Extraordinary Potent Opioid Peptide. *Proc. Natl. Acad. Sci. U.S.A.* **1979**, *76*, 6666–6670.
- (3) Cox, B. M.; Opheim, K. E.; Teschenmacher, H.; Goldstein, A. A Peptide-Like Substance from Pituitary that Acts Like Morphine. 2. Purification and Properties. *Life Sci.* **1975**, *16*, 1777–1782.
- (4) Goldstein, A.; Fischli, W.; Lowney, L. I.; Hunkapiller, M.; Hood, L. Porcine Pituitary Dynorphin: Complete Amino Acid Sequence of the Biologically Active Heptadecapeptide. *Proc. Natl. Acad. Sci. U.S.A.* **1981**, *78*, 7219–7223.
- (5) Miller, R. J. Multiple Opiate Receptors for Multiple Opioid Peptides. *Med. Biol.* **1982**, *60*, 1–6.
- (6) Wood, P. L. Multiple Opiate Receptors: Support for Unique Mu-sites, Delta-sites, and Kappa-sites. *Neuropharmacology* **1982**, *21*, 487–497.
- (7) Paterson, S. J.; Roberson, L. E.; Kosterlitz, H. W. Classification of Opioid Receptors. *Br. Med. Bull.* **1983**, *39*, 31–36.
- (8) Hruby, V. J.; Gehrig, C. Recent Developments in the Design of Receptor Specific Opioid Peptides. *Med. Res. Rev.* **1989**, *9*, 343–401.
- (9) Schiller, P. W. Development of Receptor-Specific Opioid Peptide Analogues. In *Progress in Medicinal Chemistry*; Ellis, G. P., West, G. B., Eds.; Elsevier: Amsterdam, 1991; Vol. 38, pp 301–304.
- (10) Portoghesi, P. S. The Role of Concepts in Structure-Activity Relationship Studies of Opioid Ligands. *J. Med. Chem.* **1992**, *35*, 1927–1937.
- (11) Hruby, V. J. Design of Conformationally Constrained Cyclic Peptides with High Delta and Mu Potency. In *Opioid Peptides: Medicinal Chemistry*; Rapaka, R. S., Barnett, G., Hawks, R. L., Eds.; NIDA Research Monograph Series 69; NIDA: Rockville, MD, 1986; pp 128–147.
- (12) Milan, M. J. κ-Opioid Receptors and Analgesia. *Trends Pharmacol. Sci.* **1990**, *11*, 70–76.
- (13) Chavkin, C.; Goldstein, A. Specific Receptor for the Opioid Peptide Dynorphin: Structure-Activity Relationships. *Proc. Natl. Acad. Sci. U.S.A.* **1981**, *78*, 6543–6547.
- (14) Meyer, J.-P.; Collins, N.; Lung, F.-D.; Davis, P.; Zalewska, T.; Porecca, F.; Yamamura, H. I.; Hruby, V. J. Design, Synthesis and Biological Properties of Highly Potent Cyclic Dynorphin A Analogues. Analogues Cyclized Between Positions 5 and 11. *J. Med. Chem.* **1994**, *37*, 3910–3917.
- (15) Kawasaki, A. K.; Knapp, R. J.; Kramer, T. H.; Walton, A.; Wire, W. S.; Hashimoto, S.; Yamamura, H. I.; Porreca, F.; Burks, T. F.; Hruby, V. J. Design and Synthesis of Highly Potent and Selective Cyclic Dynorphin A Analogues. 2. New Analogues. *J. Med. Chem.* **1993**, *36*, 750–757.
- (16) Turcotte, A.; Lalonde, J.-M.; St-Pierre, S.; Lemaire, S. Dynorphin-(1–13). I. Structure-Function Relationships of Ala-Containing Analogs. *Int. J. Pept. Protein Res.* **1984**, *23*, 361–367.
- (17) Choi, H.; Murray, T. F.; DeLander, G. E.; Caldwell, V.; Aldrich, J. V. N-Terminal Alkylated Derivatives of [D-Pro¹⁰] Dynorphin A-(1–13) Are Highly Selective for Kappa-Opioid Receptors. *J. Med. Chem.* **1992**, *35*, 4638–4639.
- (18) Schwyzler, R. Estimated Conformation, Orientation, and Accumulation of Dynorphin A(1–13)-tridecapeptide on the Surface of Neutral Lipid Membranes. *Biochemistry* **1986**, *25*, 4281–4286.
- (19) Pelton, J. T.; Kazmierski, W. M.; Gulya, K.; Yamamura, H. I.; Hruby, V. J. Design and Synthesis of Conformationally Constrained Somatostatin Analogs with High Potency and Specificity for Mu Opioid Receptors. *J. Med. Chem.* **1986**, *29*, 2370–2375.

JM940665+