

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/248738387>

Germacranolides and a New Type of Guaianolide from *Acanthospermum hispidum*

ARTICLE in JOURNAL OF NATURAL PRODUCTS · OCTOBER 2000

Impact Factor: 3.8 · DOI: 10.1021/np9905057

CITATIONS

25

READS

40

6 AUTHORS, INCLUDING:



[Alicia Bardon](#)

National University of Tucuman

68 PUBLICATIONS 549 CITATIONS

[SEE PROFILE](#)



[Cesar Catalan](#)

National University of Tucuman

215 PUBLICATIONS 2,475 CITATIONS

[SEE PROFILE](#)



[Luis Ricardo Hernandez](#)

Universidad de las Americas Puebla

35 PUBLICATIONS 198 CITATIONS

[SEE PROFILE](#)



[Pedro Joseph-Nathan](#)

Center for Research and Advanced Studies of...

253 PUBLICATIONS 2,242 CITATIONS

[SEE PROFILE](#)

JOURNAL OF NATURAL PRODUCTS

© Copyright 2000 by the American Chemical Society and the American Society of Pharmacognosy

Volume 63, Number 10

October 2000

Full Papers

Germacranolides and a New Type of Guaianolide from *Acanthospermum hispidum*

Elena Cartagena,[†] Alicia Bardón,[‡] César A. N. Catalán,[‡] Zaida N. J. de Hernández,[§] Luis R. Hernández,[§] and Pedro Joseph-Nathan^{*,§}

Instituto de Química Orgánica, Facultad de Bioquímica, Química y Farmacia, Universidad Nacional de Tucumán, Ayacucho 491, S. M. de Tucumán 4000, Argentina, and Departamento de Química, Centro de Investigación y de Estudios Avanzados, Instituto Politécnico Nacional, Apartado 14-740, México, D. F. 07000, México

Received October 11, 1999

The aerial parts of an Argentinian collection of *Acanthospermum hispidum* afforded 26 sesquiterpene lactones, including the two guaianolides (**1** and **2**) having a novel oxygen bridge between C-4 and C-14, three new *cis,cis*-germacranolides (**4**, **7**, and **8**), and two new melampolides (**25** and **26**). Guaianolides **1** and **2** seem to derive biosynthetically from the germacranolide **27** having the ${}_1D^{14,15}D_5$ conformation. The structures were elucidated using extensive spectroscopic analysis.

Previous investigations of *Acanthospermum* species^{1–4} have led to the isolation of *cis,cis*-germacranolides and melampolides,² in agreement with the fact that many species of the genera *Acanthospermum*, *Melampodium*, and *Lecocarpus*, belonging to the tribe Heliantheae, subtribe Melampodiinae, contain melampolides.⁵ In view of the fact that these compounds show cytotoxic and in vivo anticancer activity,³ and as a continuation of our work on sesquiterpene lactones of the Argentinian species of Asteraceae,^{6–8} we have carried out an exhaustive examination of the minor constituents of *A. hispidum* (Asteraceae), a shrub indigenous to northern Argentina. The aerial parts afforded the new guaianolides hispidunolides A (**1**) and B (**2**) with an unprecedented oxygen bridge between C-4 and C-14; the new *cis,cis*-germacranolides **4**, **7**, and **8**; the new melampolides **25** and **26**; and the known sesquiterpene lactones **3**, **5**, **6**, **9–18**, previously isolated from American^{1,2} and African^{3,4} collections of *A. hispidum*; compounds **19–23**, previously found in species of *Lecocarpus* from Ecu-

ador;^{5,9} compound **24**, previously isolated from an Australian collection of *Siegesbeckia orientalis*;¹⁰ and lolilide.¹¹

Results and Discussion


Hispidunolide A (**1**) showed IR bands for alcohol, γ -lactone, and ester groups at 3450, 1760, and 1735 cm^{-1} , respectively. It has the molecular formula $\text{C}_{22}\text{H}_{28}\text{O}_8$ as followed from its mass spectrum, which showed a $[\text{M}]^+$ at m/z 420, accounting for nine degrees of unsaturation. Mass spectral peaks at m/z 360 $[\text{M} - \text{CH}_3\text{COOH}]^+$, 335 $[\text{M} - \text{C}_5\text{H}_9\text{O}]^+$ and 85 $[\text{C}_5\text{H}_9\text{O}]^+$ indicated the presence of an acetate and a saturated five-carbon atom ester. The ^1H NMR spectrum showed the typical signals of a 2-methylbutyrate residue at δ 2.37 (qt, $J = 7.0, 7.0$ Hz), 1.62 (ddq, $J = 13.5, 7.0, 7.0$ Hz), 1.44 (ddq, $J = 13.5, 7.0, 7.0$ Hz), 1.08 (d, $J = 7.0$ Hz), and 0.89 (t, $J = 7.0$ Hz). An α -methylene- γ -lactone moiety was evident by the two doublets at δ 6.28 ($J = 3.5$ Hz) and 5.56 ($J = 3.0$ Hz) in the ^1H NMR spectrum. The ^{13}C NMR spectrum showed 22 signals corresponding to three CH_3 , five CH_2 , eight CH , and six quaternary carbons, as deduced from a DEPT experiment, in agreement with the molecular formula obtained from the mass spectrum. The ^{13}C NMR spectrum also indicated the presence of a lactone moiety, which showed signals at δ 168.8 (C-12), 134.6 (C-11), and 122.1

* To whom correspondence should be addressed. Tel.: (52)5747-7112. Fax: (52)5747-7113. E-mail: pjoseph@nathan.chem.cinvestav.mx.


[†] Universidad Nacional de Tucumán.

[‡] Universidad Nacional de Tucumán, Research Member of the National Research Council of Argentina (CONICET).

[§] Centro de Investigación y de Estudios Avanzados.




- 1 R = 2-MeBu
2 R = Ang



	R	R'	R''
3	Ac	H	<i>i</i> -Bu
4	Ac	H	Ang
5	Ac	H	2-MeBu
6	Ac	H	<i>i</i> -Val
7	Ang	H	Ac
8	H	OAc	2-MeBu

(C-13). The guaianolide skeleton was easily deduced from the ^1H NMR spectrum and sequential spin decoupling involving H-1, H-5, H-6, and H-7. The vinyl proton signal at C-14 appeared at δ 6.30 in agreement with the chemical-shift range for protons attached to the α -carbon in enol-ethers.¹² The COSY experiment showed that the signal at δ 6.30 (H-14) was long-range coupled with the signals at δ 5.06 and 2.91, corresponding to H-9 and H-1, respectively. The signals at δ 114.2 and 140.5 were assigned to the enol-ether carbons C-10 and C-14, respectively. To establish the relative configuration of the fragment C-1–C-5–C-6–C-7 and that of C-4, the minimum energy conformations of **1**, having either a C4 β –O–C14 or a C4 α –O–C14 bridge, was calculated using the PCMODEL program.¹³ These calculations showed that the dihedral angles and coupling constant values for the fragment CH(1)–CH(5)–CH(6)–CH(7) in the C4 α –O–C14 isomer are: $\text{H}\beta\text{C}(1)\text{--H}\beta\text{C}(5) = 56^\circ$ ($J = 3.6$ Hz), $\text{H}\beta\text{C}(5)\text{--H}\beta\text{C}(6) = -49^\circ$ ($J = 5.1$ Hz), and $\text{H}\beta\text{C}(6)\text{--H}\alpha\text{C}(7) = -174^\circ$ ($J = 11.2$ Hz); while for the C4 β –O–C14 isomer the values are: $\text{H}\alpha\text{C}(1)\text{--H}\alpha\text{C}(5) = -48^\circ$ ($J = 5.0$ Hz), $\text{H}\alpha\text{C}(5)\text{--H}\beta\text{C}(6) = -142^\circ$ ($J = 6.6$ Hz), and $\text{H}\beta\text{C}(6)\text{--H}\alpha\text{C}(7) = -165^\circ$ ($J = 10.8$ Hz). The latter set of values was in good agreement with the observed coupling constants, as can be seen in Table 1. To confirm the β -orientation of the vinyl oxygen at C-4, an NOE experiment irradiating the H-9 signal showed enhancement of the signal at δ 6.30 (6%) corresponding to H-14. The minimum energy conformation of **1** is shown in Scheme 1. The individual assignment of the protons attached to C-3 was deduced from the minimum energy conformation of hispidunolide A (**1**), in which the dihedral angles and calculated coupling constants are: $\text{H}\alpha\text{C}(2)\text{--H}\alpha\text{C}(3) = 14^\circ$ ($J = 11.5$ Hz), $\text{H}\alpha\text{C}(2)\text{--H}\beta\text{C}(3) = -106^\circ$ ($J = 1.5$ Hz), $\text{H}\beta\text{C}(2)\text{--H}\alpha\text{C}(3) = 134^\circ$ ($J = 6.5$ Hz), and $\text{H}\beta\text{C}(2)\text{--H}\beta\text{C}(3) = 13^\circ$ ($J = 11.5$ Hz), in good agreement with the experimental coupling



	R	R'	R''	R'''
9	CHO	Me	OAc	2-MeBu
10	CHO	CH ₂ OH	H	<i>i</i> -Bu
11	CHO	CH ₂ OH	OAc	2-MeBu
12	CHO	CH ₂ OH	H	2-MeBu
13	CHO	CH ₂ OH	H	<i>i</i> -Val
14	CHO	CH ₂ OH	OMe	Ang
15	CH ₂ OH	CH ₂ OH	OAc	<i>i</i> -Bu
16	CH ₂ OH	CH ₂ OH	OAc	Ang
17	CH ₂ OH	CH ₂ OH	OAc	2-MeBu
18	CHO	CH ₂ OH	OH	2-MeBu
19	CHO	CH ₂ OH	OAc	Ang
20	CHO	CH ₂ OH	H	Ang
21	CHO	CH ₂ OH	OMe	2-MeBu
22	CHO	CH ₂ OH	OMe	<i>i</i> -Bu
23	CHO	CH ₂ OH	OH	Ang
24	CHO	CH ₂ OH	OAc	<i>i</i> -Bu
25	CH ₂ OH	Me	OAc	Ang
26	CH ₂ OH	Me	OAc	2-MeBu

constants given in Table 1. The small coupling constant between H-7 and H-8 indicated that the ester residue at C-8 is β -oriented. Therefore, with the small coupling constant between H-8 and H-9 also taken into account, the acetate group at C-9 is α -oriented, as occurs in many known sesquiterpene lactones isolated from this species^{1,2} and also found in the present investigation.

The mass spectrum of hispidunolide B (**2**) showed $[\text{M}]^+$ at m/z 418, corresponding to the molecular formula $\text{C}_{22}\text{H}_{26}\text{O}_8$ in agreement with the ^{13}C NMR spectrum and DEPT experiment, which showed 22 signals, three of them corresponding to CH_3 , four to CH_2 , eight to CH , and seven to quaternary carbons. Relevant mass peaks at m/z 358 $[\text{M} - \text{CH}_3\text{COOH}]^+$, 335 $[\text{M} - \text{C}_5\text{H}_7\text{O}]^+$, and 83 $[\text{C}_5\text{H}_7\text{O}]^+$ were indicative of an acetate and an unsaturated five-carbon atom ester. Both the ^1H and ^{13}C NMR signals of hispidunolide B (**2**) indicated the presence of the same skeleton as in hispidunolide A (**1**), with the only difference being the ester moiety at C-8, since for **2** the ^1H NMR spectrum shows signals at δ 1.96 (dq, $J = 7.0, 1.5$ Hz), 1.81 (dq, $J = 1.5, 1.5$ Hz), and 6.15 (qq, $J = 7.0, 1.5$ Hz), which indicated the presence of an angelate residue. The ^{13}C NMR spectrum further confirmed the angelate moiety due to the signals at δ 165.6, 141.1, 126.4, 20.5, and 16.0.¹⁴ It is interesting to note that, from the biosynthetic point of view, hispidunolides A (**1**) and B (**2**) might be biosynthesized through a hetero Diels–Alder transformation¹⁵ from germacranolide **27**, which has the ${}_1\text{D}^{14}$, ${}_{15}\text{D}_5$ conformation,¹⁶ as shown in Scheme 1.

Compound **3** was previously reported by Kraus et al.,⁴ but no spectral data were provided to support the structure. Therefore, the ^1H and ^{13}C NMR data of **3** are included in Tables 2 and 4, respectively.

Compound **4** was isolated as a gum. The ^1H NMR data indicated that we were dealing with a germacranolide-type

Table 1. ^1H and ^{13}C NMR^b Data (CDCl_3 , TMS) for Hispidunolide A (**1**) and B (**2**)^c

	1		2	
	δH	δC	δH	δC
1	2.91 br t (5.5)	32.8	2.94 br t (5.5)	32.8
2a	1.99 ^d	29.3	1.99 ^d	29.4
2b	1.99 ^d		1.99 ^d	
3 α	1.77 ddd (10.5, 10.0, 3.5)	36.9	1.78 ddd (10.5, 10.0, 3.5)	36.9
3 β	2.23 br t (10.5)		2.23 br t (10.5)	
4		88.6		88.6
5	2.52 t (5.5)	45.8	2.53 t (5.5)	45.9
6	4.46 dd (9.5, 5.5)	75.6	4.48 dd (9.5, 5.5)	75.8
7	3.48 dddd (9.5, 3.5, 3.0, 1.5)	46.0	3.51 dddd (9.5, 3.5, 3.0, 1.5)	46.0
8	5.48 t (1.5)	71.9	5.56 t (1.5)	72.1
9	5.06 d (1.5)	73.7	5.15 d (1.5)	73.6
10		114.2		114.3
11		134.6		134.6
12		168.8 ^e		168.8 ^e
13a	6.28 d (3.5)	122.1	6.29 d (3.5)	122.3
13b	5.56 d (3.0)		5.60 d (3.0)	
14	6.30 br s	140.5	6.31 br s	140.6
15a	3.91 br s	64.3	3.96 d (17.0)	64.3
15b	3.91 br s		3.90 d (17.0)	
OAc	2.15 s	168.9, ^e 21.0	2.16 s	169.0 ^e , 21.0

^a 300 MHz. *J* values are given in Hz in parentheses. ^b 75.4 MHz. ^c Other signals for **1**: 2-MeBu: δH : 2.37 (qt, 7.0, 7.0, H-2'); 1.62 (ddq, 13.5, 7.0, 7.0, H-3'a); 1.44 (ddq, 13.5, 7.0, 7.0, H-3'b); 1.08 (d, 7.0, H-5'); 0.89 (t, 7.0, H-4'); δC : 174.9 (C-1'); 41.1 (C-2'); 26.6 (C-3'); 16.8 (C-5'); 11.6 (C-4'). For **2**: Ang: δH : 6.15 (qq, 7.0, 1.5, H-3'); 1.96 (dq, 7.0, 1.5, H-4'); 1.81 (dq, 1.5, 1.5, H-5'); δC : 165.6 (C-1'); 141.1 (C-3'); 126.4 (C-2'); 20.5 (C-5'); 16.0 (C-4'). ^d Overlapping signals. ^e Interchangeable.

Scheme 1. Proposed Biosynthetic Path and Minimum Energy Conformation of **1** and **27**




Table 2. ^1H NMR Data (δ , CDCl_3 , TMS) for Germacranolides **3**, **4**, **7**, and **8**^a

	3	4	7	8
H-1	6.62 ddd (8.0, 7.0, 1.5)	6.63 ddd (8.5, 6.5, 1.5)	6.67 br t (6.5)	6.78 dd (9.0, 6.0)
H-2 α	2.70 m ^b	2.70 m ^b	2.70 m ^b	3.25 dddd (15.0, 8.0, 6.0, 2.0)
H-2 β	2.85 dddd (15.0, 7.0, 4.0, 1.0)	2.85 dddd (15.0, 6.5, 4.0, 1.0)	2.85 dddd (15.0, 6.5, 4.0, 1.0)	2.80 m ^b
H-3 α	2.69 m ^b	2.69 m ^b	2.69 m ^b	2.58 ddd (14.0, 8.0, 2.0)
H-3 β	2.40 ddd (14.5, 7.5, 4.0)	2.40 ddd (14.5, 7.5, 4.0)	2.40 ddd (14.5, 7.5, 4.0)	2.99 ddd (14.0, 11.0, 8.0)
H-5	5.55 br d (9.5)	5.57 br d (9.5)	5.55 br d (9.5)	5.58 br d (9.5)
H-6	5.46 dd (9.5, 4.0)	5.48 dd (9.5, 4.0)	5.41 dd (9.5, 4.0)	5.43 dd (9.5, 4.5)
H-7	2.64 dddd (4.0, 3.0, 2.5, 2.5)	2.67 dddd (4.0, 3.0, 2.5, 2.5)	2.75 dq (4.0, 3.0, 2.5, 1.5)	2.75 m ^b
H-8	5.93 ddd (10.0, 7.0, 2.5)	5.98 ddd (10.0, 7.0, 2.5)	6.13 ddd (10.0, 7.0, 1.5)	6.53 dd (9.0, 2.5)
H-9 α	3.07 br ddd (14.0, 7.0, 1.5)	3.07 br ddd (14.0, 7.0, 1.5)	3.07 br dd (14.0, 7.0)	
H-9 β	2.40 ddd (14.0, 10.0, 1.5)	2.57 ddd (14.0, 10.0, 1.5)	2.57 ddd (14.0, 10.0, 2.0)	5.80 dd (9.0, 2.0)
H-13a	6.35 d (3.0)	6.36 d (3.0)	6.40 d (3.0)	6.41 d (3.0)
H-13b	5.71 d (2.5)	5.72 d (2.5)	5.79 d (2.5)	5.86 d (2.5)
H-14	9.41 d (1.5)	9.43 d (1.5)	9.45 d (2.0)	9.40 d (2.0)
H-15a	4.49 s	4.53 dd (13.5, 1.5)	4.49 s	4.08 s
H-15b	4.49 s	4.46 dd (13.5, 1.0)	4.49 s	4.08 s
OAc	2.12 s	2.12 s	2.12 s	2.00 s
H-2'	2.49 sept (7.0)			2.31 sext (7.0)
H-3'a	1.12 d (7.0)	6.09 qq (7.0, 1.5)	6.15 ^b	1.60 m ^b
H-3'b				1.39 m
H-4'	1.09 d (7.0)	1.96 dq (7.0, 1.5)	1.96 dq (7.0, 1.5)	0.85 t (7.0)
H-5'		1.81 dq (1.5, 1.5)	1.85 dq (1.5, 1.5)	1.05 d (7.0)

^a 300 MHz. *J* values are given in Hz in parentheses. ^b Overlapping signals.

sesquiterpene lactone containing acetate and angelate esters. These data are similar to those of related *cis,cis*-germacranolides with 2-methylbutyrate or isovalerate ester

residues attached to C-8.^{2,4} The presence of a 1,10-*cis*-double bond with an aldehyde group at C-10 followed from the chemical shifts of H-1 (δ 6.63, ddd) and H-14 (δ 9.43,

Table 3. ¹H NMR Data (δ, CDCl₃, TMS) for Melampolides **10**, **14**, **16**, **25**, and **26**^{a,b}

	10	14	16	25	26
H-1	6.63 ddd (9.5, 7.0, 2.0)	6.82 dd (10.0, 7.5)	5.80 dd (9.0, 8.0)	5.77 dd (9, 7.5)	5.75 br dd (8.5, 7.5)
H-2a	2.47 ^c	2.74 m	2.42 m ^c	2.44 m ^c	2.44 m ^c
H-2b	2.42 ^c	2.62 ^c	2.34 m ^c	2.3 ^c	2.3 ^c
H-3α	2.04 ddd (12.5, 12.5, 2.0)	2.04 ddd (12.0, 12.0, 2.0)	1.90 ddd (12.5, 12.5, 2.5)	2.0 ^c	2.0 ^c
H-3β	2.83 ddd (12.5, 6.0, 2.5)	2.85 ddd (12.0, 5.5, 2.5)	2.69 ddd (12.5, 5.5, 2.5)	2.3 ^c	2.3 ^c
H-5	5.16 br d (10.5)	5.03 br d (10.0)	5.15 br d (10.0)	5.03 br d (10.5)	5.01 br d (10.5)
H-6	5.23 t (10.5)	5.20 t (10.0)	5.35 t (10.0)	5.10 dd (10.5, 9)	5.10 dd (10.5, 9)
H-7	2.48 ^c	2.63 ^c	3.35 dddd (10.0, 3.5, 3.0, 2.0)	3.31 dddd (9.0, 3.5, 3.0, 2.0)	3.29 dddd (9.0, 3.5, 3.0, 2.0)
H-8	6.36 ddd (10.0, 8.0, 2.0)	6.65 dd (8.5, 1.5)	6.16 dd (9.5, 2.0)	6.16 dd (9.5, 2.0)	6.06 dd (9.5, 2.0)
H-9α	2.75 ddd (14.0, 8.0, 2.0)				
H-9β	2.06 ddd (14.0, 10.0, 1.5)	3.88 dd (8.5, 2.0)	5.53 d (9.5)	5.42 d (9.5)	5.35 d (9.5)
H-13a	6.24 d (3.5)	6.30 d (3.5)	6.24 d (3.5)	6.24 d (3.5)	6.24 d (3.5)
H-13b	5.58 d (3.0)	5.87 d (3.0)	5.68 d (3.0)	5.68 d (3.0)	5.63 d (3.0)
H-14a	9.46 d (1.5)	9.52 d (2.0)	4.40 br d (12.5)	4.38 d (12.5)	4.37 br d (12.5)
H-14b			4.23 br d (12.5)	4.21 br d (12.5)	4.19 d (12.5)
H-15a	4.53 d (12.5)	4.47 br d (13.0)	4.49 br s	1.97 br s	1.98 br s
H-15b	4.32 br d (12.5)	4.37 br d (13.0)	4.49 br s		

^a 300 MHz. *J* values are given in Hz in parentheses. ^b Other signals (δ), for **10**: *i*-Bu: 2.53 (sept, 7.0, H-2'); 1.14 (d, 7.0, H-4'); 1.12 (d, 7.0, H-3'a). For **14**: Ang: 6.05 (qq, 7.0, 1.5, H-3'); 1.96 (dq, 7.0, 1.5, H-4'); 1.88 (dq, 1.5, 1.5, H-5'); OMe: 3.10 s. For **16**: Ang: 6.11 (qq, 7.0, 1.5, H-3'); 1.95 (dq, 7.0, 1.5, H-4'); 1.82 (dq, 1.5, 1.5, H-5'); Ac: 1.95 s. For **25**: Ang: 6.10 (qq, 6.0, 1.5, H-3'); 1.95 (overlapped, H-4'); 1.82 (dq, 1.5, 1.5, H-5'); Ac: 1.94 s. For **26**: 2-MeBu: 2.30 (m, H-2'); 1.60 (m, H-3'a); 1.39 (m, H-3'b); 1.06 (d, 7.0, H-5'); 0.86 (t, 7.5, H-4'). Ac: 1.97 s. ^cOverlapped signals.

Table 4. ¹³C NMR Data (δ, CDCl₃, TMS) for Compounds **3**, **8**, **13**, **16**, **17**, **24**, and **26**^a

	3	8	13 ^b	16	17	24	26
1	153.2	158.9	153.8	134.5 ^d	134.5 ^d	158.5	134.3 ^d
2	25.0	24.9	27.0	26.3	26.5 ^e	27.6	26.5
3	26.1	27.3 ^c	32.6	33.1	33.1	32.4	37.7
4	135.0 ^d	139.3 ^c	140.4 ^d	139.9	141.8	140.9 ^d	139.1
5	129.2	127.0	128.4	127.9	128.0	128.5	125.9
6	73.2	73.5	73.8	72.3	72.3	73.4	72.8
7	46.8	46.4	49.4	50.9	51.0	51.1	50.8
8	71.9	72.2	65.6	68.8	68.8	69.8	68.9
9	28.6	69.6	28.8	74.2	74.0	67.9	75.6
10	142.0	140.3 ^c	142.7 ^d	136.0	136.0	141.2 ^d	136.2
11	134.4 ^d	133.4	134.8	134.1 ^d	134.0 ^d	133.7	134.8 ^d
12	169.3	169.0	169.3	169.4	169.3	169.0	164.6
13	124.7	126.2	121.2	121.3	121.3	122.3	121.1
14	194.8	193.3	195.5	64.0	64.0	193.8	64.0
15	66.6	65.8	60.5	60.9	60.9	60.6	16.7
OAc	170.4	170.4		170.0	169.9	170.5	
	20.9	20.7		20.8	21.0	20.8	
1'	175.7	175.0	171.7	166.5	175.4	175.4	175.4
2'	34.0	41.3	43.3	142.0	41.4	34.1	41.4
3'	19.1	26.5 ^c	25.8	126.8	26.3 ^e	19.0	25.4
4'	18.6	11.6	22.4	15.8	11.6	19.0	11.7
5'		17.1	22.3	20.4	16.9		16.7

^a 75.4 MHz. ^b Distinction of C-2 and C-3' followed from APT measurements. ^c Distinction of C-4 from C-10 and of C-3 from C-3' followed from HMBC measurements. ^{d,e} Interchangeable signals.

d).¹ The *cis*-configuration of the 4,5-double bond was deduced from the typical chemical shifts of H-5 (δ 5.57, br d) and H-6 (δ 5.48, dd) and the value of *J*_{6,7} = 4.0 Hz.^{2,5} The angelate residue at C-8 is β-oriented because *J*_{7,8} = 2.5 Hz, in agreement with the well-known *syn*-periplanar orientation of H-8 and H-7.⁵ In addition, β-oriented ester residues at this position are frequent in germacranolides of the subtribe Melampodiinae. The large coupling constant between H-8 and H-9β (10.0 Hz) showed that H-9β is *trans* to H-8. This stereochemistry places H-9β and H-14 into a *W* relationship if the aldehyde carbonyl is oriented such that there is maximal overlap between the π orbitals of the 1(10)-double bond and the carbonyl group, an arrangement that accounts for the observed long-range coupling between H-9β and H-14.¹ An allylic coupling between H-1 and H-9α was also observed.

The ¹H NMR spectrum of **7** (Table 2) was very similar to that of **4**. It only differed in the H-8 and H-15 chemical

shifts, and therefore in **7** the angelate ester is located at C-15, while the acetate group is located at C-8.

The spectral features of **8** were similar to those described for *cis,cis*-germacranolides **4** and **7**, but no signals at δ 2.57 and 3.07 were found for the H-9 protons. Instead, a doublet at δ 5.80 (*J* = 9.0, 2.0 Hz) and a singlet at δ 2.00 revealed the presence of an α-oriented acetate at C-9, as further supported by the signals at δ 69.6 (C-9), 170.4, and 20.7 (OAc) in the ¹³C NMR spectrum. A singlet at δ 4.08 (2H) indicated that a hydroxyl group was bonded to a methylene group, and it was assigned to the protons at C-15. The 500-MHz HMBC contour plot showed, among others, correlations between C-5 and H-3, H-6, and H-6; between C-6 and H-5 and H-8; between C-7 and H-6, H-8, and the two hydrogens at C-13, between C-8, and H-6 and H-9, between C-9 and H-1, H-8, and H-14; between the acetate carbonyl and H-9, and between the 2-methylbutyrate carbonyl and H-8, the two H-3' signals, and the H-5' methyl, which further supported the structure of **8**. The experiment also allowed distinction of the C-4 and C-10 signals; the former had correlations with one H-2 and one H-3, H-5, and H-15, while the latter had correlations with one H-2, H-9, and H-14. There was also distinction of the C-3 and C-3' signals, as much as the former had correlations with H-5 and H-15, while the latter correlated with H-2', H-4', and H-5'.

The IR spectrum of **10** showed strong absorptions at 3450, 1760, 1735, and 1685 cm⁻¹, indicating the presence of a hydroxyl, γ-lactone, ester, and conjugated carbonyl with a double bond, respectively. The ¹H NMR spectrum exhibited signals ascribable to a germacranolide-type compound bearing an isobutyrate ester, because it was similar to the spectra of related melampolides.^{1,2,5,9} The presence of a 1,10-*cis*-double bond with an aldehyde group at C-10 followed from the chemical shifts of H-1 (δ 6.63, ddd) and H-14 (δ 9.46, d).¹ The *trans*-configuration of the 4,5-double bond was deduced from the typical chemical shifts of H-5 (δ 5.16, br d) and H-6 (δ 5.23, t), as well as a large *J*_{6,7} of 10.5 Hz.² The signals at δ 4.53 (d) and 4.32 (br d) were assigned to H-15a and H-15b, respectively. The signal at δ 6.36 (ddd) is typical for a proton attached to a carbon supporting an ester group and was assigned to H-8. The small coupling constant between H-7 and H-8 (2.0 Hz) indicated that the ester residue at C-8 is β-oriented. An

allylic coupling between H-1 and H-9 α ($J = 2.0$ Hz) and a W -type coupling between H-9 β and H-14 ($J = 1.5$ Hz) were observed. The typical signals for an α -methylene- γ -lactone moiety appeared at δ 6.24 (d, $J = 3.5$ Hz, H-13a) and 5.58 (d, $J = 3.0$ Hz, H-13b). A previous report on this compound by Kraus et al.⁴ provided no spectral data to support the structure. The ¹H NMR data are given in Table 3.

Compounds **25** and **26** were melampolides having an acetate and an angelate in the case of **25**, and an acetate and a 2-methylbutyrate in the case of **26**. They differed in the ester group attached to C-8, as can be seen from the ¹H NMR data given in Table 3. Neither the IR nor NMR spectra showed bands or signals for an aldehyde group. However, in the ¹H NMR spectrum of **25** an AB system at δ 4.38 (d, $J = 12.5$ Hz) and 4.21 (br d, $J = 12.5$ Hz), assigned to the H-14 protons, was present. The chemical shift of these protons and those of H-1 (δ 5.77, dd) were similar to those observed for related melampolides bearing a CH₂OH at C-10.¹ Similar signals were observed for **26**, as can be seen in Table 3. Noteworthy for melampolides with carbonyl groups attached to C-10 is the chemical shift of H-1, found 1 ppm downfield. The total signal assignment was achieved by comparison with a melampolide obtained by Herz and Kalyanaraman¹ by reduction of a precursor with a carbonyl attached to C-10.

The ¹H NMR spectrum of melampolide **14** differed from that of leocarpinolide J (**21**), previously isolated from *Lecocarpus leocarpoides*,⁵ only in the signals corresponding to the ester residue at C-8. Because **14** has an allylic methoxyl group and we used methanol for the HPLC separation, one could suspect it to be an artifact. However, leocarpinolide J and leocarpinolide M, which also contain an allylic methoxyl group at C-9, were found in the genus *Lecocarpus*,⁵ belonging to the same subtribe of *Acanthospermum*, for which no methanol was used during the isolation procedures. Compound **16** displayed similar spectral data to those of **25**. However, the ¹H and ¹³C NMR spectra of **16** accounted for an additional CH₂OH group at C-4. Compounds **14** and **16** were previously reported by Kraus et al.,⁴ but no spectral data were provided to support the structures. Herein we report the ¹H NMR data of **14** and **16** in Table 3 and the ¹³C NMR data of **13**,² **16**,⁴ **17**,¹ and **24**,¹⁰ which were not published previously, in Table 4.

Experimental Section

General Experimental Procedures. IR spectra were recorded on a Perkin-Elmer 16F PC FT-IR spectrophotometer. Optical rotations were performed on a Perkin-Elmer 241 polarimeter. NMR spectra were recorded on Varian XL-300GS or Unity-500 spectrometers. The EIMS were obtained on a Hewlett-Packard 5989-A spectrometer at 20 eV. For separation of mixtures, a Gilson HPLC instrument with a 305 pump and a 112-differential refractometer detector was used. Columns: A (Beckman ultrasphere C₁₈, 10 \times 250 mm) and B (Beckman ultrasphere C₈, 10 \times 250 mm) were employed. Retention times (t_R) were measured from the solvent peak. For column chromatography, Si gel Merck 70–230 or 230–400 mesh ASTM were used.

Plant Material. The aerial parts (flowers and leaves) of *A. hispidum* DC. were collected in Vipos, Tucumán Province, Argentina, in April 1995. A voucher specimen (LIL 604458) is on deposit at the Herbarium of Fundación Miguel Lillo, Tucumán, Argentina.

Extraction and Isolation. The plant material (330 g) was extracted with CHCl₃ (2 \times 3 L) at room temperature for 14 days to give 15.9 g (yield 4.8%) of a crude extract, which was suspended in EtOH (130 mL) at 55 °C, diluted with H₂O (100 mL), and extracted successively with hexane (3 \times 150 mL) and CHCl₃ (3 \times 150 mL). The chloroform extract on evaporation

at reduced pressure furnished a residue (3.38 g), which was column chromatographed over Si gel using CHCl₃ with increasing amounts of EtOAc (0–100%) and finally MeOH, to give nine fractions. Fractions containing sesquiterpene lactones, as evidenced by IR, were further processed.

A portion (200 mg) of fraction 2 (463 mg) was chromatographed by HPLC (Column A, MeOH–H₂O, 3:2, 1.5 mL min^{−1}) to give 5 mg of **3**, t_R 27 min, 4 mg of **1**, t_R 32 min; 2.3 mg of **4**, t_R 43 min; 33.3 mg of **6**, t_R 52 min; 2 mg of **9**, t_R 84 min; and mixtures further purified by HPLC (Column B, MeOH–H₂O, 3:2, 1.3 mL min^{−1}) to give 1.5 mg of **2**, t_R 30 min, and 0.9 mg of **5**, t_R 45 min.

Fraction 3 (168 mg) was chromatographed by HPLC (Column A, MeOH–H₂O, 4:3, 1.5 mL min^{−1}) to give 2.1 mg of loliolide,¹¹ t_R 6 min; 5.5 mg of **2**, t_R 23 min; 2.4 mg of **1**, t_R 25 min; 1.3 mg of **25**, t_R 76 min; 6.4 mg of **26**, t_R 88 min; and mixtures further purified by HPLC (Column B, MeOH–H₂O, 1:1, 1.3 mL min^{−1}) to give 1 mg of **10**, t_R 24 min; 0.7 mg of **19**, t_R 35 min; 7.4 mg of **11**, t_R 28 min; 4.4 mg of **20**, t_R 40 min; 8.7 mg of **12**, t_R 44 min; and 11.4 mg of **13**, t_R 48 min.

A portion (110 mg) of fraction 4 (193 mg) was processed by HPLC (Column A, MeOH–H₂O, 1:1, 2 mL min^{−1}) to give 1.7 mg of **24**, t_R 14 min; 0.6 mg of **7**, t_R 19 min; 5.8 mg of **13**, t_R 31 min; and mixtures further purified by HPLC (Column B, MeOH–H₂O, 1:1, 2 mL min^{−1}) to give 1 mg of **14**, t_R 12 min; 1.9 mg of **10**, t_R 14 min; 3.3 mg of **21**, t_R 16 min; 9.6 mg of **19**, t_R 20 min; 30.8 mg of **11**, t_R 25 min; and 1.6 mg of **12**, t_R 16 min.

A portion (200 mg) of fraction 5 (423 mg) was processed by HPLC (Column B, MeOH–H₂O, 6:5, 2 mL min^{−1}) to give 96 mg of **11**, t_R 45 min; 4.5 mg of a mixture of **13** and **24**, t_R 54 min; and mixtures further purified by HPLC (Column A, MeOH–H₂O, 1:1, 2 mL min^{−1}) to give 0.5 mg of **22**, t_R 13 min; 3.1 mg of **24**, t_R 19 min; and 4.7 mg, of **19**, t_R 30 min.

A portion (200 mg) of fraction 6 (487 mg) was chromatographed by HPLC (Column A, MeOH–H₂O, 1:1, 1.8 mL min^{−1}) to give 10.3 mg of **24**, t_R 15 min, and 35.8 mg of **11**, t_R 28 min.

A portion (200 mg) of fraction 7 (549 mg) was processed by HPLC (Column A, MeOH–H₂O, 1:1, 2 mL min^{−1}) to give 1.1 mg of **15**, t_R 18 min; 18.8 mg of **17**, t_R 30 min; 8.5 mg of **8**, t_R 61 min; and mixtures further purified by HPLC (Column B, MeOH–H₂O, 1:1, 2 mL min^{−1}) to give 1.5 mg of **15**, t_R 21 min; 4.7 mg of **16**, t_R 27 min; and 0.7 mg of **17**, t_R 32 min.

Fraction 8 (60 mg) was chromatographed by HPLC (Column B, MeOH–H₂O, 1:1, 2 mL min^{−1}) to give 2.3 mg of **23**, t_R 9 min, and 2.8 mg of **18**, t_R 10 min.

9-Acetyloxy-15-hydroxy-8-(2-methylbutanoyloxy)-10(14),11(13)-guaidiene-6,12-olide-4,14-oxide (hispidunolide A) (1): gum; $[\alpha]_D^{25}{}_{589} -46^\circ$, $[\alpha]_D^{25}{}_{578} -48^\circ$, $[\alpha]_D^{25}{}_{546} -56^\circ$, $[\alpha]_D^{25}{}_{436} -102^\circ$, $[\alpha]_D^{25}{}_{365} -174^\circ$ (c 5.0, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 210 (4.1) nm; IR (CHCl₃) ν_{max} 3450, 1760, 1735, 1635 cm^{−1}; ¹H and ¹³C NMR, see Table 1; EIMS (direct inlet) m/z 420 [M]⁺ (3), 402 (9) [M – H₂O]⁺, 360 [M – CH₃COOH]⁺ (1), 335 [M – CH₃CH₂CH(CH₃)CO]⁺ (11), 318 [M – CH₃CH₂CH(CH₃)COOH]⁺ (1), 317 [M – H₂O – CH₃CH₂CH(CH₃)CO]⁺ (4), 293 [M – CH₂CO – CH₃CH₂CH(CH₃)CO]⁺ (66), 275 [M – CH₃COOH – CH₃CH₂CH(CH₃)CO]⁺ (74), 258 [M – CH₃COOH – CH₃CH₂CH(CH₃)COOH]⁺ (16), 240 [M – H₂O – CH₃COOH – CH₃CH₂CH(CH₃)COOH]⁺ (34), 85 [CH₃CH₂CH(CH₃)CO]⁺ (39), 57 [C₄H₉]⁺ (100).

9-Acetyloxy-15-hydroxy-8-angeloyloxy-10(14),11(13)-guaidiene-6,12-olide-4,14-oxide (hispidunolide B) (2): gum; $[\alpha]_D^{25}{}_{589} -40^\circ$, $[\alpha]_D^{25}{}_{578} -42^\circ$, $[\alpha]_D^{25}{}_{546} -50^\circ$, $[\alpha]_D^{25}{}_{436} -87^\circ$, $[\alpha]_D^{25}{}_{365} -147^\circ$ (c 6.2, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 209 (4.5) nm; IR (CHCl₃) ν_{max} 3500, 1760, 1730, 1640 cm^{−1}; ¹H and ¹³C NMR, see Table 1; EIMS (direct inlet) m/z 418 [M]⁺ (1), 400 [M – H₂O]⁺ (4), 358 [M – CH₃COOH]⁺ (0.2), 335 [M – CH₃CHC(CH₃)CO]⁺ (12), 317 (2), 293 [M – CH₂CO – CH₃CHC(CH₃)CO]⁺ (35), 275 [M – CH₃COOH – CH₃CHC(CH₃)CO]⁺ (41), 258 [M – CH₃COOH – CH₃CHC(CH₃)COOH]⁺ (5), 240 [M – CH₃COOH – CH₃CHC(CH₃)COOH – H₂O]⁺ (12), 83 [CH₃CHC(CH₃)CO]⁺ (100), 55 [C₄H₇]⁺ (28).

15-Acetyloxy-8 β -angeloyloxy-14-oxo-(4Z)-acanthospermolide (4): gum; UV (EtOH) λ_{max} (log ϵ) 211 (5.3) nm; IR (CHCl₃) ν_{max} 2720, 1760, 1735, 1630 cm^{−1}; ¹H NMR, see Table

2; EIMS (direct inlet) m/z 402 $[M]^+$ (1), 342 (4), 302 (15), 242 (20), 213 (11), 82 (44), 54 (100).

8 β -Acetyloxy-15-angeloyloxy-14-oxo-(4Z)-acanthospermolide (7): gum; UV (EtOH) λ_{\max} (log ϵ) 212 (5.1) nm; IR (CHCl₃) ν_{\max} 2720, 1760, 1735, 1630 cm⁻¹; ¹H NMR, see Table 2; EIMS (direct inlet) m/z 402 $[M]^+$ (2), 342 (4), 302 (10), 242 (19), 213 (11), 82 (50), 54 (100).

9 α -Acetyloxy-8 β -(2-methylbutanoyloxy)-14-oxo-(4Z)-acanthospermolide (8): gum; $[\alpha]^{25}_{589} -73^\circ$, $[\alpha]^{25}_{578} -78^\circ$, $[\alpha]^{25}_{546} -90^\circ$, $[\alpha]^{25}_{436} -168^\circ$, $[\alpha]^{25}_{365} -278^\circ$ (c 6.3, CHCl₃); UV (EtOH) λ_{\max} (log ϵ) 200 (3.6) nm; IR (CHCl₃) ν_{\max} 2725, 1765, 1730, 1640 cm⁻¹; ¹H and ¹³C NMR, see Table 2 and Table 4, respectively; EIMS (direct inlet) m/z 420 $[M]^+$ (1), 402 (15), 342 (20), 240 (26), 212 (10), 85 (32), 57 (100).

9 α -Acetyloxy-8 β -angeloyloxy-14-hydroxyacanthospermolide (25): gum; UV (EtOH) λ_{\max} (log ϵ) 204 (4.1) nm; IR (CHCl₃) ν_{\max} 3455, 1760, 1735, 1630 cm⁻¹; ¹H NMR, see Table 3; EIMS (direct inlet) m/z 404 $[M]^+$ (0.2), 386 (9), 326 (15), 226 (22), 83 (100).

9 α -Acetyloxy-14-hydroxy-8 β -(2-methylbutanoyloxy)-acanthospermolide (26): gum; $[\alpha]^{25}_{589} -20^\circ$, $[\alpha]^{25}_{578} -22^\circ$, $[\alpha]^{25}_{546} -24^\circ$, $[\alpha]^{25}_{436} -38^\circ$, $[\alpha]^{25}_{365} -67^\circ$ (c 4.5, CHCl₃); UV (EtOH) λ_{\max} (log ϵ) 207 (3.8) nm; IR (CHCl₃) ν_{\max} 3450, 1760, 1735, 1635 cm⁻¹; ¹H and ¹³C NMR, see Table 3 and Table 4, respectively; EIMS (direct inlet) m/z 406 $[M]^+$ (1), 388 (9), 328 (17), 226 (11), 85 (32), 57 (100), 43 (56).

Acknowledgment. Financial support from CoNaCyT (México) and CYTED (Spain) is acknowledged. Work in Tucumán was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (CONICET) and Consejo de Investigaciones de la Universidad Nacional de Tucumán (CIUNT). We thank M. C. María Isabel

Chávez (Instituto de Química, University of Mexico) for the HMBC measurements. L.R.H. thanks Ministerio de Cultura y Educación de la Nación Argentina (Dirección Nacional de Cooperación e Integración Educativa Internacional) for a postdoctoral fellowship.

References and Notes

- (1) Herz, W.; Kalyanaraman, P. S. *J. Org. Chem.* **1975**, *40*, 3486-3491.
- (2) Bohlmann, F.; Jakupovic, J.; Zdero, C.; King, R. M.; Robinson, H. *Phytochemistry* **1979**, *18*, 625-630.
- (3) Jakupovic, J.; Baruah, R. N.; Bohlmann, F.; Msonthi, J. D. *Planta Med.* **1986**, *52*, 154-155.
- (4) Kraus, W.; Köll-Weber, M.; Maile, R.; Wunder, T.; Vogler, B. *Pure Appl. Chem.* **1994**, *66*, 2347-2352.
- (5) Macias, F. A.; Molinillo, J. M. G.; Fischer, N. H. *Phytochemistry* **1993**, *32*, 127-131.
- (6) de Hernández, Z. N. J.; Hernández, L. R.; Catalán, C. A. N.; Gedris, T. E.; Herz, W. *Phytochemistry* **1997**, *46*, 721-727.
- (7) Kotowicz, C.; Bardón, A.; Catalán, C. A. N.; Cerda-García-Rojas, C. M.; Joseph-Nathan, P. *Phytochemistry* **1998**, *47*, 425-428.
- (8) de Hernández, Z. N. J.; Catalán, C. A. N.; Hernández, L. R.; Guerra-Ramírez, D.; Joseph-Nathan, P. *Phytochemistry* **1999**, *51*, 79-82.
- (9) Macias, F. A.; Fischer, N. H. *Phytochemistry* **1992**, *31*, 2747-2754.
- (10) Zdero, C.; Bohlmann, F.; King, R. M.; Robinson, H. *Phytochemistry* **1991**, *30*, 1579-1584.
- (11) Hodges, R.; Porte, A. L. *Tetrahedron* **1964**, *20*, 1463-1467.
- (12) Pretsch, E.; Clerc, T.; Seibl, J.; Simon, W. *Spectral Data for Structure Determination of Organic Compounds*, 2nd ed.; Springer-Verlag: New York, 1989.
- (13) Burket, U.; Allinger, N. L. *Molecular Mechanics*; ACS Monograph 177; American Chemical Society: Washington, DC, 1982.
- (14) Joseph-Nathan, P.; Wesener, J. R.; Günther, H. A. *Org. Magn. Reson.* **1984**, *22*, 190-191.
- (15) Martin, S. F.; Chen, K. X.; Eary, C. T. *Org. Lett.* **1999**, *1*, 79-81.
- (16) Samek, Z.; Harmatha, J. *Collect. Czech. Chem. Commun.* **1978**, *43*, 2779-2799.

NP9905057