

Reviews

An Overview of Chemical Processes That Damage Cellular DNA: Spontaneous Hydrolysis, Alkylation, and Reactions with Radicals

Kent S. Gates*

Departments of Chemistry and Biochemistry, University of Missouri—Columbia, 125 Chemistry Building, Columbia, Missouri 65211

Received July 17, 2009

The sequence of heterocyclic bases on the interior of the DNA double helix constitutes the genetic code that drives the operation of all living organisms. With this said, it is not surprising that chemical modification of cellular DNA can have profound biological consequences. Therefore, the organic chemistry of DNA damage is fundamentally important to diverse fields including medicinal chemistry, toxicology, and biotechnology. This review is designed to provide a brief overview of the common types of chemical reactions that lead to DNA damage under physiological conditions.

Contents

1. Introduction	1747	5.7. Tandem Lesions	1754
2. Hydrolysis of DNA	1747	6. Conclusion	1756
2.1. Spontaneous Hydrolysis of the Phosphodiester Backbone Is Very Slow	1747		
2.2. Hydrolytic Deamination of DNA Bases	1748		
2.3. Spontaneous Hydrolysis of the Glycosidic Bonds Connecting the Nucleobases to the DNA Backbone	1749		
2.4. Properties of Abasic Sites Arising from Depurination	1749		
3. Overview: Common Reactions by Which Small Organic Molecules Damage DNA	1750		
4. DNA Alkylation	1750		
4.1. Preferred Alkylation Sites in DNA	1750		
4.2. The Chemical Fate of Alkylated DNA	1750		
4.2.1. Alkylated Phosphates Are Chemically Stable	1750		
4.2.2. Alkylation at Some Positions on the DNA Nucleobases Yields Chemically Stable Adducts	1751		
4.2.3. Alkylation of Some Endocyclic Nitrogens on the Nucleobases Yields Labile Lesions That Undergo Deglycosylation or Ring Opening	1751		
4.2.4. Alkylation at Some Endocyclic Nitrogens Accelerates Deamination	1751		
5. Reactions of Radicals with DNA	1752		
5.1. Abstraction of Hydrogen Atoms from the 2'-Deoxyribose Sugars of DNA: General Features	1752		
5.2. Alkali-Labile Strand Damage Stemming from Abstraction of the C1'-Hydrogen Atom	1752		
5.3. Direct Strand Cleavage Initiated by Abstraction of the 4'-Hydrogen Atom	1753		
5.4. Reactions of Radicals with the Nucleobases	1753		
5.5. Addition of Radicals to C5 of Thymine Residues: Generation of Thymine Glycol	1753		
5.6. Addition of Radicals to Guanine Residues: Generation of FAPy-G and 8-Oxo-G Residues	1754		

1. Introduction

The sequence of heterocyclic bases on the interior of the DNA double helix constitutes the genetic code that drives the operation of all living organisms (Figure 1) (1–4). Accurate readout of DNA sequence is necessary for the expression of functional mRNA and, ultimately, proteins in cells (5). In addition, during cell division, faithful replication of DNA is required to produce daughter cells containing exact copies of the genome (6). In light of these facts, it is not surprising that chemical modification of cellular DNA can have profound biological consequences. DNA damage can trigger changes in gene expression, inhibit cell division, or trigger cell death (7–9). In addition, attempts to replicate damaged DNA can introduce errors (mutations) into the genetic code (10). Thus, the organic chemistry of DNA damage is fundamentally important to diverse fields including medicinal chemistry, toxicology, and biotechnology (11–13). This review is designed to provide a brief overview of the most common types of chemical reactions that lead to DNA damage under physiological conditions.

2. Hydrolysis of DNA

2.1. Spontaneous Hydrolysis of the Phosphodiester Backbone Is Very Slow. Hydrolysis of the phosphodiester groups in the backbone of DNA is thermodynamically favored ($\Delta G^\circ = -5.3$ kcal/mol) (14) but extremely slow (Scheme 1) (15).¹ Work with carefully designed model compounds such as **1** indicate that the half-life of phosphodiester hydrolysis is

* To whom correspondence should be addressed. Tel: 573-882-6763. Fax: 573-882-2754. E-mail: gatesk@missouri.edu.

¹ In the interest of brevity, many of the reaction mechanisms shown in this review are schematic in nature. For example, protonation, deprotonation, and proton transfer steps may not be explicitly depicted or may not distinguish specific acid–base catalysis from general acid–base catalysis. In some cases, arrows indicating electron movement are used simply to






Figure 1. Structure of DNA.



approximately 30 000 000 years under physiologically relevant conditions (15–18). In short, this means that spontaneous hydrolysis of the phosphodiester linkages in DNA does not occur to a significant extent under biological conditions, although the reaction can be vastly accelerated by various catalysts including phosphodiesterases, lanthanide ions, and transition metal ions (19–23).


2.2. Hydrolytic Deamination of DNA Bases. Cytosine residues **2** in DNA can undergo hydrolytic deamination to yield uracil residues **3** (Scheme 2) (24–28). The reaction may proceed via either attack of hydroxide on the neutral nucleobase or attack of water on the N3-protonated base (Scheme 2) (24, 25, 29). Sensitive genetic reversion assays revealed that cytosine deamination occurs in duplex DNA with a half-life of 30 000–85 000 years at pH 7.4 and 37 °C (27, 28). In single-stranded DNA and at base mismatches in duplex DNA, deamination proceeds much faster ($t_{1/2} \sim 200$ years) presumably due to increased solvent accessibility of the base (26, 30). Deamination of guanine and adenine residues in DNA is much slower, occurring at only 2–3% of the rate of cytosine deamination (31).




Methylcytosine residues **4** are mutation hotspots in bacterial and eukaryotic genomes (28). The deamination of 5-methylcytosine residues occurs approximately 2–3 times faster than at unmodified cytosine residues (28); however, the increased mutation frequencies observed at methylcytosine positions are believed to stem not from increased deamination at these sites but from the fact that the resulting G–T mismatches are poorly

direct the reader's attention to sites where the "action" is occurring rather than to illustrate complete sequences of bond-making and bond-breaking.


Scheme 3



Scheme 4



Scheme 5



repaired and produce G-C→A-T transitions in one of the daughter cells (28, 32, 33). Mutagenesis resulting from deamination at 5-methylcytosine residues teaches us an important, general lesson: That is, DNA damage reactions that are terribly slow and low yielding can have profound biological consequences if the resulting lesion is not efficiently repaired and is mutagenic or cytotoxic.

Alkylation of the N3-position of cytosine and reactions that lead to saturation of the 5,6-double bond in cytosine and 5-methylcytosine accelerate deamination (34–41). Activation-induced cytidine deaminases catalyze the conversion of cytosine residues to uracil residues in single-stranded regions of DNA (42, 43). In addition, deamination is an important reaction that is associated with exposure of DNA to nitrosating agents and nitric oxide (44–47).

2.3. Spontaneous Hydrolysis of the Glycosidic Bonds Connecting the Nucleobases to the DNA Backbone. With regard to hydrolytic stability, the glycosidic bonds that hold the nucleobases to the sugar–phosphate backbone are weak points in the structure of DNA (Scheme 3). The pyrimidine bases cytosine and thymine are lost with rate constants of $1.5 \times 10^{-12} \text{ s}^{-1}$ ($t_{1/2} = 14\,700$ years), while the reaction is faster at the purine bases guanine and adenine, occurring with rate constants of $3.0 \times 10^{-11} \text{ s}^{-1}$ ($t_{1/2} = 730$ years) (48, 49). Accordingly, hydrolytic cleavage of the glycosidic bonds in DNA is often referred to as “depurination” because the reaction is much more facile at purines than at pyrimidines. Hydrolysis of the glycosidic bond in 2'-deoxypurines proceeds via a specific acid-catalyzed S_N1 reaction mechanism (50, 51). Equilibrium protonation increases the leaving group ability of the base and facilitates unimolecular, rate-limiting C–N bond cleavage that generates the free base 5 and an oxocarbenium ion 6 (Scheme 3). The oxocarbenium ion undergoes subsequent hydrolysis to yield an abasic site 7 (often referred to as an apurinic site or AP site). It is calculated that spontaneous depurination generates about 10 000 abasic sites

per cell per day (49). Indeed, steady state levels of 10 000–50 000 abasic sites have been detected in cells (52, 53). Depurination occurs about four times faster in single-stranded DNA than it does in duplex DNA (49).

2.4. Properties of Abasic Sites Arising from Depurination. Abasic sites (7, Scheme 3) generated by depurination are cytotoxic and mutagenic (54, 55). This lesion exists as an equilibrium mixture of the ring-closed acetal (7, 99%) and the ring-opened aldehyde (8, 1%) (56). Abstraction of the acidic α -proton adjacent to the aldehyde group in 8 leads to β -elimination of the phosphate residue on the 3'-side of the abasic site. This strand-cleavage reaction occurs with a half-life of 200 h under physiological conditions (pH 7.4 and 37 °C) (57, 58). Thermal workup, sodium hydroxide, or treatment with various amines such as piperidine, dimethylethylene diamine, or putrescine facilitate strand cleavage at abasic sites (59–62). Subsequent γ,δ -elimination of the phosphate on the 5'-side of the abasic site 9 is relatively slow under physiological conditions (57) but occurs readily under basic conditions (e.g., 0.05 M NaOH, 37 °C, and 2 h) to generate a strand break with 3'- and 5'-phosphate termini (63, 64).

In addition to generating single strand breaks, the aldehyde residue of the ring-opened form of the abasic site can react with the exocyclic N^2 -amino group of a guanine residue on the




Figure 2. Reactions of bifunctional electrophiles with DNA.

opposing strand to generate an interstrand cross-link **10** at 5'-d(CAp) sites (where Ap = the abasic site; Scheme 4) (65). This is especially intriguing because interstrand DNA cross-links are highly cytotoxic lesions (66–68).

3. Overview: Common Reactions by Which Small Organic Molecules Damage DNA

The vast majority of DNA damage caused by bioactive molecules such as drugs, toxins, and mutagens falls into just two general categories: (1) alkylation of a DNA nucleophile by an electrophile or (2) the reaction of a π -bond or C–H bond in DNA with a radical intermediate (Scheme 5). In the following sections, we will consider the reactions of electrophiles and radicals with DNA and examine the fate of the chemically modified DNA that is generated in these processes.

4. DNA Alkylation

4.1. Preferred Alkylation Sites in DNA. Almost all of the heteroatoms in the double helix have the potential to be alkylated. The preferred sites of alkylation in duplex DNA depend strongly on the nature of the alkylating agent. For example, in the case of diethylsulfate, the preferred sites of reaction follow the order: N7G \gg P–O $>$ N3A \gg N1A \sim N7A \sim N3G \sim N3C \gg O⁶G (69, 70). In contrast, the preferred sites for alkylation by ethyldiazonium ion follow the order: P–O \gg N7G $>$ O²T $>$ O⁶G $>$ N3A \sim O⁴C $>$ O⁴T $>$ N3G \sim N3T \sim N3C \sim N7A (69, 70). The N7-position of the guanine residue is the most nucleophilic site on the DNA bases and is a favored site of reaction for almost all small, freely diffusible alkylating agents. The preferences observed at other sites commonly have been rationalized in terms of hard–soft reactivity principles (71). Hard alkylating agents (defined by small size, positive charge, and low polarizability) such as diazonium ions display increased reactivity with hard oxygen nucleophiles in DNA (69, 70, 72–74). On the other hand, soft (large, uncharged, and polarizable) alkylating agents like dialkylsulfates favor reactions at the softer nitrogen centers in DNA.²

Typically, small diffusible alkylating agents react with DNA at multiple sites; however, connection of a nonselective alky-




Figure 3. Chemically stable lesions resulting from DNA alkylation.

lating agent to noncovalent DNA-binding units can confer sequence and atom site selectivity to the reaction (75–83). Bifunctional alkylating agents (molecules containing two electrophilic centers) can cross-link two nucleophilic centers in the DNA duplex. Examples of bifunctional DNA alkylating agents shown in Figure 2 include the endogenous lipid peroxidation products malondialdehyde (84) and acrolein (85) and the anticancer drugs mitomycin C (86) and mechlorethamine (87). Cross-links may present special challenges to DNA repair systems, and agents that generate cross-links can be exceptionally bioactive (66–68). Accordingly, a significant number of clinically used anticancer drugs are DNA cross-linking agents (66).

4.2. The Chemical Fate of Alkylated DNA. 4.2.1. Alkylated Phosphates Are Chemically Stable. Alkylation of a DNA phosphodiester residue yields a phosphotriester (**11**, Figure 3). Phosphotriesters undergo hydrolysis about 10^9 times more

² Fishbein and co-workers provided an excellent discussion of the mechanistic origin of hard–soft acid base effects in DNA alkylation reactions. On page 1466 of ref 74, these authors note that “the low degree of covalency in the transition state for S_N2 substitution on primary diazonium ions means that electrostatic stabilization of the transition state by the nucleophile will become a dominant interaction and this gives rise to enhanced oxygen atom alkylation by primary diazonium ions”.




Figure 4. Chemically labile lesions resulting from DNA alkylation.

rapidly than native DNA phosphodiester (15). Nonetheless, hydrolysis of phosphotriesters is slow under physiological conditions, and alkylated DNA phosphates are chemically stable both *in vitro* and *in vivo* (88, 89).

4.2.2. Alkylation at Some Positions on the DNA Nucleobases Yields Chemically Stable Adducts. Alkylation at the exocyclic nitrogen atoms N²G, N⁶A, and N⁴C, the amidic nitrogens at N1-G and N1-T, and the oxygens at O⁶-G and O⁴-T yield chemically stable adducts (Figure 3) (90–96). Special attention has been paid to the O⁶-G and O⁴-T adducts because these reactions alter the Watson–Crick hydrogen-bonding faces of these bases and cause miscoding and mutagenesis during DNA replication (97–99).

4.2.3. Alkylation of Some Endocyclic Nitrogens on the Nucleobases Yields Labile Lesions That Undergo Deglycosylation or Ring Opening. Alkylation of the endocyclic nitrogens N7G, N7A, N3G, N3A, N1A, and N3C can destabilize the nucleobases, facilitating deglycosylation and ring-opening reactions (Figure 4). The half-lives for deglycosylation of bases modified with simple alkyl groups in DNA generally follow the trend: N7dA (3 h) > N3dA (24 h) > N7dG (150 h) > N3dG (greater than 150 h) > O²dC (750 h) > O²dT (6300 h) > N3dC (>7700 h) (36, 96, 100–105). The mechanisms for these processes (Scheme 6) mirror that described above for acid-catalyzed deglycosylation reactions (Scheme 3) (106). Studies on the depurination of rates of N7-alkylguanine derivatives


indicate that electron-withdrawing groups on the alkyl substituent increase the rate of deglycosylation (103, 107, 108).

Alternatively, attack of water at the C8-position of an N7-alkylguanine residue initiates a ring-opening reaction that yields the corresponding 5-(alkyl)formamidopyrimidine (FAPy) derivative **12** (Scheme 7) (103, 109–116). In general, the ring-opening reaction that yields alkyl-FAPy derivatives is slow compared to the competing depurination reaction under physiological conditions (103, 117), although basic conditions (e.g., pH 10, 37 °C, and 4 h) can be used to generate alkyl-FAPy lesions from N7-alkylguanine lesions in oligonucleotides (116, 118). Nonetheless, FAPy lesions are formed *in vivo* and are persistent lesions found in animals following exposure to alkylating agents (119–121). In this regard, it may be important that the glycosidic bond in alkyl-FAPy residues is stable at neutral pH (122). Spontaneous deglycosylation of FAPy residues occurs only slowly, with an estimated (103, 123) half-life of about 1500 h in single-stranded DNA at pH 7.5 and 37 °C. The deglycosylation reaction is pH-dependent, however, and at pH 6.5, complete deglycosylation of 5-methyl-FAPy in an oligonucleotide was observed over 48 h (122). The stereochemistry of the attachment between the FAPy base and the deoxyribose sugar residue undergoes anomerization via an imine intermediate **13** (Scheme 7) (122, 124, 125). The half-life of this process is 3.5 h for the FAPy derivative of deoxyguanosine in a nucleoside derivative (123, 126) and 16 h for the FAPy derivative of aflatoxin B1 in a single-stranded DNA oligomer (124).


Adenine residues alkylated at the N1-position are also prone to hydrolytic ring opening. In this case, however, ring opening initiates a Dimroth rearrangement reaction that leads to an apparent migration of the alkyl group from the N1-position to the exocyclic N⁶-position of the nucleobase (Scheme 8) (127). The Dimroth rearrangement occurs with a half-life of approximately 150 h at pH 7 in the context of the nucleotide, 1-methyl-2'-deoxyadenosine-5'-phosphate (101). Importantly, this rearrangement can occur even when the adducted base is stacked within the DNA double helix (128).

4.2.4. Alkylation at Some Endocyclic Nitrogens Accelerates Deamination. Alkylation of the N3-position of cytosine (Scheme 9) vastly accelerates deamination of the nucleobase. For example, deamination of 3-methyl-2'-deoxycytidine, occurring with a half-life of 406 h, proceeds 4000 times faster than the same reaction in the native nucleoside (36). Alkylation of the endocyclic nitrogen in deoxycytidine also facilitates deglycosylation ($t_{1/2} = 7700$ h, lower pathway in Scheme 9), but in the context of the nucleoside, the deamination reaction is approximately 20 times faster and predominates (36). Alkylation at the N1-position of 2'-deoxyadenosine nucleosides typically leads to a Dimroth rearrangement rather than deamination (129); however, the hydroxyl group of N1-styrene adducts


Scheme 6




Scheme 7



Scheme 8



Scheme 9



facilitates deamination via intramolecular attack on C6 of the adducted base (Scheme 10) (130–132).

5. Reactions of Radicals with DNA

5.1. Abstraction of Hydrogen Atoms from the 2'-Deoxyribose Sugars of DNA: General Features. Abstraction of hydrogen atoms from the sugar–phosphate backbone of DNA generates 2-deoxyribose radicals that lead to final strand damage products via complex reaction cascades (133–137). In the case of highly reactive species such as the hydroxyl radical (HO^\bullet), the relative amounts of direct strand cleavage stemming from hydrogen atom abstraction at each position of the deoxyribose sugars are dictated by steric accessibility of the deoxyribose hydrogens rather than their C–H bond strengths (133–138). The steric accessibility of the deoxyribose hydrogens in DNA follows the trend: $\text{H5}' > \text{H4}' \gg \text{H3}' \sim \text{H2}' \sim \text{H1}'$ (138). Here, I will summarize DNA strand damage arising from hydrogen atom abstraction at the C1'- and C4'-positions of deoxyribose.


These examples were selected because they are perhaps the best-characterized pathways and are illustrative of the general types of reactions that stem from radical-mediated damage to the deoxyribose units of DNA. A number of excellent reviews provide more comprehensive coverage of radical-mediated strand damage reactions (133–137).

5.2. Alkali-Labile Strand Damage Stemming from Abstraction of the C1'-Hydrogen Atom. Although sequestered deep in the minor groove of duplex DNA (138), the C1'-hydrogen atom is an important reaction site for a variety of agents including $\text{Cu}(\text{o-phenanthroline})_2$, neocarzinostatin, espermicin, dynemicin, tirapazamine, and radiation (134, 139). In addition, some base radical adducts efficiently abstract the C1'-hydrogen atom (140).


In cells, 2-deoxyribose radicals generally are expected to react with either molecular oxygen, present at 60–100 μM , or thiols such as glutathione, which are present at 1–10 mM. The C1' radical abstracts a hydrogen atom from thiols with a rate constant of $1.8 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ in duplex DNA (141). This reaction has the potential to generate either the natural β -anomer (a “chemical repair” reaction) or the mutagenic α -anomer. Greenberg's group showed that, in duplex DNA, the C1' radical reacts with 2-mercaptoethanol to yield a 6:1 ratio of the natural β -anomer over the α -anomer (141).

Molecular oxygen reacts with the C1' radical **14** with a rate constant of $1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ to yield the peroxy radical **15** (Scheme 11) (142). The peroxy radical ejects superoxide radical anion (the pK_a of the relevant conjugate acid HOO^\bullet is 5.1) with a rate constant of $2 \times 10^4 \text{ s}^{-1}$ to generate the carbocation **16** (142) that, in turn, is attacked by water to give the presumed C1'-alcohol intermediate **17**. Loss of the DNA base yields the


Scheme 10



Scheme 11



Scheme 12




2-deoxyribose-2-phosphate abasic site **18**. The ratio of rate constants $k_{O_2}/k_{RSH} = 1100$ (143). This means that at normal cellular concentrations of oxygen ($\sim 70 \mu\text{M}$) and thiol (5 mM) more than 90% of the radical will be trapped by molecular oxygen.

The 2-deoxyribose-2-phosphate lesion **18** does not rapidly yield direct strand cleavage ("frank strand breaks") under physiological conditions (144–146). Rather, **18** is a base-labile lesion that is efficiently converted into a strand break upon treatment with piperidine (0.1 M, 20 min, and 90 °C) or hydroxide (at pH 10 and 37 °C, the $t_{1/2}$ for the cleavage reaction is 15 min in a single-stranded oligomer) (144, 145). Under physiologically relevant conditions, α,β -elimination of the 3'-phosphate residue slowly yields strand cleavage with a half-life of between 38 and 54 h in double-stranded DNA, depending on the identity of the base that opposes the lactone abasic site (Scheme 12) (144–146). The resulting 3'-ene-lactone intermediate **19** subsequently undergoes rapid γ,δ -elimination to give the methylene lactone byproduct **20** and an oligonucleotide with a 3'-phosphate residue (145, 146). The transient ene-lactone **19** has been trapped by a variety of chemical reagents (144). In addition, the 2-ribose-lactone and the ene-lactone form covalent cross-links with DNA-binding proteins (147, 148).

5.3. Direct Strand Cleavage Initiated by Abstraction of the 4'-Hydrogen Atom. Abstraction of the 4'-hydrogen atom from the sugar–phosphate backbone of duplex DNA is a major reaction for hydroxyl radical (HO^\bullet), tirapazamine, bleomycin, and several enediynes (136, 139, 149). The C4' radical **21** abstracts hydrogen atoms from biological thiols with a rate constant of about $2 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ in single-stranded DNA (150). In duplex DNA, this chemical repair reaction yields a

Scheme 14




10:1 ratio of natural:unnatural stereochemistry at the C4' center (151). Reaction of the C4' radical **21** with molecular oxygen yields the peroxy radical **22**, with a presumed rate constant of about $2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ (Scheme 13) (150). Reaction of the peroxy radical with thiol, occurring with a rate constant of $k \leq 200 \text{ M}^{-1} \text{ s}^{-1}$ (152), yields the hydroperoxide **23**. This intermediate has been observed directly by MALDI mass spectroscopy in a single-stranded 2'-deoxyribonucleotide in which a single C4' radical was generated at a defined site (153). In the case of agents such as bleomycin, peroxydinitrite, and the enediynes, the hydroperoxide **23** is thought to yield direct strand cleavage via a Criegee rearrangement, followed by hydrolysis and fragmentation reactions that generate the 5'-phosphate, 3'-phosphoglycolate, and base propenal end products (Scheme 13) (153). The base propenals generated in this reaction act as "malondialdehyde equivalents" that can go forward to cause further damage to DNA via covalent modification of guanine residues (Scheme 14) (154, 155).

5.4. Reactions of Radicals with the Nucleobases. Radicals react readily with the heterocyclic bases of DNA. For example, over 90% of the hydroxyl radicals generated by γ -radiolysis react at the nucleobases in polyU (see page 340 of ref 156). A large number of base damage products arising from the reaction of hydroxyl radical with DNA have been characterized (Figure 5) (157–162). These products arise via hydrogen atom abstraction or, more commonly, addition of hydroxyl radical to the π -bonds of the bases (Scheme 5). Here, we consider the formation of several common DNA base damage products arising from reactions at thymidine and deoxyguanosine. The pathways described here produce some of the most prevalent oxidative base damage products and also illustrate the general types of reactions that commonly occur following the attack of radicals on the nucleobases.

5.5. Addition of Radicals to C5 of Thymine Residues: Generation of Thymine Glycol. Radicals such as HO^\bullet readily add to the carbon–carbon π -bond of thymine residues (163). The addition of hydroxyl radical to the C5-position thymine residues generates the C6-thymidine radical (**24**, Scheme 15) (133). (To a lesser extent, hydroxyl radical also adds to the C6-position of thymine residues (133), but reactions stemming from this pathway will not be discussed here.) The base radical adduct **24** resulting from the addition of HO^\bullet at C5 may be reduced by reagents such as thiols or superoxide radical to yield the 5-hydroxy-6-hydrothymine residue (**25**, Scheme 15) (133, 164).

Scheme 13







Figure 5. Oxidatively damaged nucleobases.



Alternatively, the reaction with molecular oxygen produces the peroxyl radical **26**. On the basis of model reactions involving the uracil free base (165), it has been proposed that ejection of superoxide radical anion from **26**, followed by subsequent attack of water, generates thymine glycol **28** (133). However, recent studies did not detect production of superoxide from the structurally related peroxyl radical derived from 5,6-dihydro-2'-deoxyuridin-6-yl **30**. It was concluded that the rate for superoxide elimination is less than 10^{-3} s^{-1} (166). This suggests that, in biological systems, the peroxyl radical **26** may instead abstract a hydrogen atom from thiols to give the hydroperoxide **27**, which, in turn, is expected to undergo further thiol-mediated reduction to yield thymine glycol **28**. Thymine glycol is one of the major products stemming from oxidative damage of DNA (157–161). In the absence of thiol, the hydroperoxide **27** undergoes slow decomposition ($t_{1/2} = 1.5\text{--}10 \text{ h}$ for the nucleoside in water, with the trans isomer decomposing more


rapidly than the cis) to generate the ring-opened lesion **29** as the major product and thymine glycol **28** as a minor product (Scheme 15) (167).

5.6. Addition of Radicals to Guanine Residues: Generation of FAPy-G and 8-Oxo-G Residues. Guanine is a major target for oxidative damage by radicals (157, 168, 169). A primary pathway for the reaction of radicals with guanine residues involves addition to the C8-position (Scheme 16) (157, 170, 171). This process yields a “redox ambivalent” nucleobase radical **31** that can undergo either one-electron reduction or oxidation (157, 164, 171). Reduction leads to a ring-opened formamidopyrimidine (FAPy) lesion (**32**, Scheme 16) that is chemically stable in DNA under physiological conditions and is mutagenic (172–175). On the other hand, oxidation of the base radical adduct **31** yields 8-oxo-7,8-dihydroguanosine (**33**, 8-oxo-G). The 8-oxo-G lesion has been incorporated into synthetic 2'-deoxyoligonucleotides (176) but is prone to oxidative decomposition under biologically relevant conditions (157, 177). For example, two-electron oxidation of 8-oxo-G (**33**) yields 5-hydroxy-8-oxo-7,8-dihydroguanosine **34** (178). This compound can decompose to yield the spiroamino-dihydantoin **35** via a 1,2-shift (178). Alternatively, hydrolysis of **34** generates the ureidoimidazoline **36**, which undergoes decarboxylation to the guanidinohydantoin **37** (178). In nucleosides, guanidinohydantoin formation is favored at pH values <7 , while formation of the spiroaminohydantoin **35** is favored at pH values ≥ 7 (178–180). In the context of duplex DNA, oxidative degradation of 8-oxo-G residues yields the guanidinohydantoin residue **37** as the major product (177), although this lesion may be prone to further decomposition to products such as imidazolone and oxazolone (181, 182). Guanidinohydantoin **37**, spiroaminodihydantoin **35**, and 8-oxo-G **33** are mutagenic lesions in duplex DNA (182–186).


5.7. Tandem Lesions. In some cases, initially generated DNA radicals can undergo secondary reactions within the duplex. The resulting “tandem lesions”, comprised of cross-links or multiple adjacent damage sites, may present special challenges to DNA replication and repair systems (187, 188). For example, 5,6-dihydrothymidine radical (**38**, Scheme 17), in the presence of molecular oxygen, generates the corresponding peroxyl radical **39** that abstracts the C1'-hydrogen atom from the sugar residue on the 5'-side to produce a tandem lesion **40**, consisting of a damaged pyrimidine base adjacent to a 2-deoxy-ribofuranolactone residue (189, 190).

The 5-(2'-deoxyuridinyl)methyl radical (**41**, Scheme 18) generates intrastrand cross-links with adjacent purine residues (191, 192). This reaction is favored at 5'-GT sites (191). The 5-(2'-deoxycytidinyl)methyl radical can cross-link with guanine residues in a similar manner (193). The 5-(2'-deoxyuridinyl)-


Scheme 16



Scheme 17



Scheme 18




methyl radical **41** has also been shown to forge an interstrand cross-link with the opposing deoxyadenosine (194, 195).

Under low oxygen conditions, C5'-sugar radicals can react with the base residue on the same nucleotide. In purine nucleotides, the carbon-centered radical **42** can add to the C8-position of the nucleobase (Scheme 19). Oxidation of the intermediate nucleobase radical **43** yields the 8,5'-cyclo-2'-deoxypurine lesion **44** (161, 164, 187, 188, 196–199). Similarly, in pyrimidine nucleotides, the C5' radical can add to the C6-

position of nucleobase. Reduction of the resulting radical intermediate yields the 5',6-cyclo-5,6-dihydro-2'-deoxypyrimidine lesion **45** (200–202). These cyclolesions are chemically stable in DNA and may present special challenges to cellular repair systems (161, 164, 187, 188, 196–202).

Scheme 19



6. Conclusion

Francis Crick wrote, "DNA is such an important molecule that is almost impossible to learn too much about it" (203). Indeed, today, more than 50 years after the primary chemical structure of DNA was established (1), there is a continuing need to develop our understanding of biologically relevant DNA chemistry. Characterization of the chemical reactions of endogenous cellular chemicals, anticancer drugs, and mutagens with cellular DNA is important because the biological responses engendered by any given DNA-damaging agent are ultimately determined by the chemical structure of the damaged DNA. Improvements in methodologies for the detection and characterization of DNA damage such as LC/MS, high-field NMR, and the synthesis of non-natural 2'-deoxyoligonucleotides have continued to drive the field forward (143, 204–207). DNA-damaging agents can elicit a wide variety of cellular responses including cell cycle arrest, up-regulation of DNA repair systems, apoptosis, or mutagenesis (7–13); however, the relationships between the structure of damaged DNA and the biological responses that ensue are not yet well understood. Advances in this area have the potential to yield both fundamental and practical advances in cell biology, predictive toxicology, and anticancer drug development.

Acknowledgment. I am grateful to current and past research group members for helpful discussions on the topic of nucleic acid chemistry. I also acknowledge support from the NIH (CA 83925 and CA 119131) during the writing of this review.

References

- Todd, A. R. (1954) Chemical structure of the nucleic acids. *Proc. Natl. Acad. Sci. U.S.A.* 40, 748–755.
- Watson, J. D., and Crick, F. H. C. (1953) A structure for deoxyribose nucleic acid. *Nature* 171, 737–738.
- Venter, J. C. (2001) The sequence of the human genome. *Science* 291, 1304–1351.
- Pennisi, E. (2003) DNA's cast of thousands. *Science* 300, 282–285.
- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., and Walter, P. (2002) *Molecular Biology of the Cell*, 4th ed., Garland Science, New York.
- Alberts, B. (2003) DNA replication and recombination. *Nature* 421, 431–435.
- Zhou, B.-B. S., and Elledge, S. J. (2000) The DNA damage response: Putting checkpoints in perspective. *Nature* 408, 433–439.
- Norbury, C. J., and Hickson, I. D. (2001) Cellular responses to DNA damage. *Annu. Rev. Pharmacol. Toxicol.* 41, 367–401.
- Rouse, J., and Jackson, S. P. (2002) Interfaces between the detection, signaling, and repair of DNA damage. *Science* 297, 547–551.
- Guengerich, F. P. (2006) Interactions of carcinogen-bound DNA with individual DNA polymerases. *Chem. Rev.* 106, 420–452.
- Gates, K. S. (1999) Covalent modification of DNA by natural products. In *Comprehensive Natural Products Chemistry* (Kool, E. T., Ed.) pp 491–552, Pergamon, New York.
- Wolkenberg, S. E., and Boger, D. L. (2002) Mechanisms of in situ activation for DNA-targeting antitumor agents. *Chem. Rev.* 102, 2477–2495.
- Hurley, L. H. (2002) DNA and its associated processes as targets for cancer therapy. *Nat. Rev. Cancer* 2, 188–200.
- Dickson, K. S., Burns, C. M., and Richardson, J. P. (2000) Determination of the free-energy change for repair of a DNA phosphodiester bond. *J. Biol. Chem.* 275, 15828–15831.
- Schroeder, G. K., Lad, C., Wyman, P., Williams, N. H., and Wolfenden, R. (2006) The time required for water attack at the phosphorus atom of simple phosphodiester and of DNA. *Proc. Natl. Acad. Sci. U.S.A.* 103, 4052–4055.
- Williams, N. H., and Wyman, P. (2001) Base catalysed phosphate diester hydrolysis. *Chem. Commun.* 1268–1269.
- Cassano, A. G., Anderson, V. E., and Harris, M. E. (2002) Evidence for direct attack by hydroxide in phosphodiester hydrolysis. *J. Am. Chem. Soc.* 124, 10964–10965.
- Iché-Tarrat, N., Barthelat, J.-C., Rinaldi, D., and Vigroux, A. (2005) Theoretical studies of hydroxide-catalyzed P–O cleavage reactions of neutral phosphate triesters and diesters in aqueous solution: Examination of the change induced by H/Me substitution. *J. Phys. Chem. B* 109, 22570–22580.
- Williams, N. H., Takasaki, B., Wall, M., and Chin, J. (1999) Structure and nuclease activity of simple dinuclear metal complexes: Quantitative dissection of the role of metal ions. *Acc. Chem. Res.* 32, 485–493.
- Chen, W., Kitamura, Y., Zhou, J.-M., Sumaoka, J., and Komiyama, M. (2004) Site-specific DNA hydrolysis by combining Ce(IV)/EDTA with monophosphate-bearing oligonucleotides and enzymatic ligation of the scission fragments. *J. Am. Chem. Soc.* 126, 10285–10291.
- Galbur, E. A., and Stoddard, B. L. (2002) Catalytic mechanisms of restriction and homing endonucleases. *Biochemistry* 41, 13851–13860.
- Nomura, A., and Sugiura, Y. (2004) Sequence-specific and hydrolytic cleavage of DNA by zinc finger mutants. *J. Am. Chem. Soc.* 126, 15374–15375.
- Sreedhara, A., Freed, J. D., and Cowan, J. A. (2000) Efficient inorganic deoxyribonucleases. Greater than 50-million-fold rate enhancement in enzyme-like DNA cleavage. *J. Am. Chem. Soc.* 122, 8814–8824.
- Shapiro, R., and Danzig, M. (1972) Acidic hydrolysis of deoxycytidine and deoxyuridine derivatives. The general mechanism of deoxyribonucleoside hydrolysis. *Biochemistry* 11, 23–29.
- Shapiro, R., and Klein, R. S. (1966) The deamination of cytidine and cytosine by acidic buffer solutions. Mutagenic implications. *Biochemistry* 5, 2358–2362.
- Lindahl, T., and Nyberg, B. (1974) Heat-induced deamination of cytosine residues in deoxyribonucleic acid. *Biochemistry* 13, 3405–3410.
- Frederico, L. A., Kunkel, T. A., and Shaw, B. R. (1990) A sensitive genetic assay for detection of cytosine deamination: Determination of rate constants and the activation energy. *Biochemistry* 29, 2532–2537.
- Shen, J.-C., Rideout, W. M., and Jones, P. A. (1994) The rate of hydrolytic deamination of 5-methylcytosine in double-stranded DNA. *Nucleic Acids Res.* 22, 972–976.
- Garrett, E. R., and Tsau, J. (1972) Solvolyses of cytosine and cytidine. *J. Pharm. Sci.* 61, 1052–1061.
- Frederico, L. A., Kunkel, T. A., and Shaw, B. R. (1993) Cytosine deamination in mismatched base pairs. *Biochemistry* 32, 6523–6530.
- Karran, P., and Lindahl, T. (1980) Hypoxanthine in deoxyribonucleic acid: Generation of heat-induced hydrolysis of adenine residues and release in free form by a deoxyribonucleic acid glycosylase in calf thymus. *Biochemistry* 19, 6005–6011.
- Lutsenko, E., and Bhagwat, A. S. (1999) Principle causes of hot spots for cytosine to thymine mutations at sites of cytosine methylation in growing cells. A model, its experimental support and implications. *Mutat. Res.* 437, 11–20.
- Duncan, B. K., and Miller, J. H. (1980) Mutagenic deamination of cytosine residues in DNA. *Nature* 287, 560–561.
- Dong, M., Wang, C., Deen, W. M., and Dedon, P. C. (2003) Absence of 2'-deoxyoxanosine and presence of abasic sites in DNA exposed to nitric oxide at controlled physiological concentrations. *Chem. Res. Toxicol.* 16, 1044–1055.
- Shapiro, R., and Yamaguchi, H. (1972) Nucleic acid reactivity and conformation. I. Deamination of cytosine by nitrous acid. *Biochim. Biophys. Acta* 281, 501–506.
- Sowers, L. C., Sedwick, W. D., and Ramsay Shaw, B. (1989) Hydrolysis of N3-methyl-2'-deoxycytidine: Model compound for reactivity of protonated cytosine residues in DNA. *Mutat. Res.* 215, 131–138.

- (37) Peng, W., and Shaw, B. R. (1996) Accelerated deamination of cytosine residues in UV-induced cyclobutane pyrimidine dimers leads to CC-to-TT transitions. *Biochemistry* 35, 10172–10181.
- (38) Yebra, M. J., and Bhagwat, A. S. (1995) A cytosine methyltransferase converts 5-methylcytosine in DNA to thymine. *Biochemistry* 34, 14752–14757.
- (39) Zingg, J.-M., Shen, J.-C., Yang, A. S., Rapoport, H., and Jones, P. A. (1996) Methylation inhibitors can increase the rate of cytosine deamination by (cytosine-5)-DNA methyltransferase. *Nucleic Acids Res.* 24, 3267–3275.
- (40) Chen, H., and Shaw, B. R. (1993) Kinetics of bisulfite-induced cytosine deamination in single-stranded DNA. *Biochemistry* 32, 3535–3539.
- (41) Shapiro, R., DiFate, V., and Welcher, M. (1974) Deamination of cytosine derivatives by bisulfite. Mechanism of the reaction. *J. Am. Chem. Soc.* 96, 906–912.
- (42) Pham, P., Bransteitter, R., and Goodman, M. F. (2005) Reward versus risk: DNA cytidine deaminases triggering immunity and disease. *Biochemistry* 44, 2703–2715.
- (43) Samaranayake, M., Bujnicki, J. M., Carpenter, M., and Bhagwat, A. S. (2006) Evaluation of molecular models for the affinity maturation of antibodies: roles of cytosine deamination by AID and DNA repair. *Chem. Rev.* 106, 700–719.
- (44) Kosaka, H., Wishnok, J. S., Miwa, M., Leaf, C. D., and Tannenbaum, S. R. (1989) Nitrosation by stimulated macrophages, inhibitors, enhancers, and substrates. *Carcinogenesis* 10, 563–566.
- (45) Wink, D. A., Kasprzak, K. S., Maragos, C. M., Elespuru, R. K., Misra, M., Dunams, T. M., Cebula, T. A., Koch, W. H., Andrews, A. W., and Allen, J. S. (1991) DNA deaminating ability and genotoxicity of nitric oxide and its progenitors. *Science* 254, 1001–1003.
- (46) Wuenschell, G. E., O'Connor, T. R., and Termini, J. (2003) Stability, miscoding potential, and repair of 2'-deoxyxanthosine in DNA: Implications for nitric oxide-induced mutagenesis. *Biochemistry* 42, 3608–3616.
- (47) Lucas, L. T., Gatehouse, D., and Shuker, D. E. G. (1999) Efficient nitroso group transfer from N-nitrosoindoles to nucleotides and 2'-deoxyguanosine at physiological pH. *J. Biol. Chem.* 274, 18319–18326.
- (48) Lindahl, T., and Karlstrom, O. (1973) Heat-induced depyrimidination of deoxyribonucleic acid in solution. *Biochemistry* 12, 5151–5154.
- (49) Lindahl, T., and Nyberg, B. (1972) Rate of depurination of native deoxyribonucleic acid. *Biochemistry* 11, 3610–3618.
- (50) Zoltewicz, J. A., Clark, D. F., Sharpless, T. W., and Grahe, G. (1970) Kinetics and mechanism of acid-catalyzed hydrolysis of some purine nucleosides. *J. Am. Chem. Soc.* 92, 1741–1750.
- (51) Stivers, J. T., and Jiang, Y. J. (2003) A mechanistic perspective on the chemistry of DNA repair glycosylases. *Chem. Rev.* 103, 2729–2759.
- (52) Nakamura, J., Walker, V. E., Upton, P. B., Chiang, S. Y., Kow, Y. W., and Swenberg, J. A. (1998) Highly sensitive apurinic/apyrimidinic site assay can detect spontaneous and chemically induced depurination under physiological conditions. *Cancer Res.* 58, 222–225.
- (53) Nakamura, J., and Swenberg, J. A. (1999) Endogenous apurinic/apyrimidinic sites in genomic DNA of mammalian tissues. *Cancer Res.* 59, 2522–2526.
- (54) Auerbach, P., Bennett, R. A. O., Bailey, E. A., Krokan, H. E., and Demple, B. (2005) Mutagenic specificity of endogenously generated abasic sites in *Saccharomyces cerevisiae* chromosomal DNA. *Proc. Natl. Acad. Sci. U.S.A.* 102, 17711–17716.
- (55) Boiteux, S., and Guillet, M. (2004) Abasic sites in DNA: repair and biological consequences in *Saccharomyces cerevisiae*. *DNA Repair* 3, 1–12.
- (56) Wilde, J. A., Bolton, P. H., Mazumdar, A., Manoharan, M., and Gerlt, J. A. (1989) Characterization of the equilibrating forms of the abasic site in duplex DNA using 17O-NMR. *J. Am. Chem. Soc.* 111, 1894–1896.
- (57) Lindahl, T., and Andersson, A. (1972) Rate of chain breakage at apurinic sites in double-stranded deoxyribonucleic acid. *Biochemistry* 11, 3618–3623.
- (58) Crine, P., and Verly, W. G. (1976) A study of DNA spontaneous degradation. *Biochim. Biophys. Acta* 442, 50–57.
- (59) McHugh, P. J., and Knowland, J. (1995) Novel reagents for chemical cleavage at abasic sites and UV photoproducts in DNA. *Nucleic Acids Res.* 23, 1664–1670.
- (60) Sugiyama, H., Fujiwara, T., Ura, A., Tashiro, T., Yamamoto, K., Kawanishi, S., and Saito, I. (1994) Chemistry of thermal degradation of abasic sites in DNA. mechanistic investigation on thermal DNA strand cleavage of alkylated DNA. *Chem. Res. Toxicol.* 7, 673–683.
- (61) Maxam, A. M., and Gilbert, W. (1980) Sequencing end-labeled DNA with base-specific chemical cleavages. *Methods Enzymol.* 65, 499.
- (62) Mattes, W. B., Hartley, J. A., and Kohn, K. W. (1986) Mechanism of DNA strand breakage by piperidine at sites of N7 alkylation. *Biochim. Biophys. Acta* 868, 71–76.
- (63) Bailly, V., Derydt, M., and Verly, W. G. (1989) δ -Elimination in the repair of AP (apurinic/apyrimidinic) sites in DNA. *Biochem. J.* 261.
- (64) Bailly, V., and Verly, W. G. (1988) Possible roles of β -elimination and δ -elimination reactions in the repair of DNA containing AP (apurinic/apyrimidinic) sites in mammalian cells. *Biochem. J.* 253, 553–9.
- (65) Dutta, S., Chowdhury, G., and Gates, K. S. (2007) Interstrand crosslinks generated by abasic sites in duplex DNA. *J. Am. Chem. Soc.* 129, 1852–1853.
- (66) Rajski, S. R., and Williams, R. M. (1998) DNA cross-linking agents as antitumor drugs. *Chem. Rev.* 98, 2723–2795.
- (67) Noll, D. M., Mason, T. M., and Miller, P. S. (2006) Formation and repair of interstrand crosslinks in DNA. *Chem. Rev.* 106, 277–301.
- (68) Schärer, O. D. (2005) DNA interstrand crosslinks: Natural and drug-induced DNA adducts that induce unique cellular responses. *Chem-BioChem* 6, 27–32.
- (69) Beranek, D. T. (1990) Distribution of methyl and ethyl adducts following alkylation with monofunctional alkylating agents. *Mutat. Res.* 231, 11–30.
- (70) Singer, B., and Grunberger, D. (1983) *Molecular Biology of Mutagens and Carcinogens*, Plenum, New York.
- (71) Lo, T.-L. (1975) Hard soft acids bases (HSAB) principle in organic chemistry. *Chem. Rev.* 75, 1–20.
- (72) Moschel, R. C., Hudgins, W. R., and Dipple, A. (1979) Selectivity in nucleoside alkylation and aralkylation in relation to chemical carcinogenesis. *J. Org. Chem.* 44, 3324–3328.
- (73) Loechler, E. L. (1994) A violation of the Swain–Scott principle, and NOT S_N1 versus S_N2 reaction mechanisms, explains why carcinogenic alkylating agents can form different proportions of adducts at oxygen versus nitrogen in DNA. *Chem. Res. Toxicol.* 7, 277–280.
- (74) Lu, X., Heilman, J. M., Blans, P., and Fishbein, J. C. (2005) The structure of DNA dictates purine atom site selectivity in alkylation by primary diazonium ions. *Chem. Res. Toxicol.* 18, 1462–1470.
- (75) Yang, X.-L., and Wang, A. H.-J. (1999) Structural studies of atom-specific anticancer drugs acting on DNA. *Pharm. Ther.* 83, 181–215.
- (76) Fidler, A., Moes, G. W. H., Scheffer, A. G., ver der Schans, G. P., Baan, R. A., de Jong, L. P. A., and Benschop, H. P. (1994) Synthesis, characterization, and quantitation of the major adducts formed between sulfur mustard and DNA of calf thymus and human blood. *Chem. Res. Toxicol.* 7, 199–204.
- (77) Zang, H., and Gates, K. S. (2003) Sequence specificity of DNA alkylation by the antitumor natural product leinamycin. *Chem. Res. Toxicol.* 16, 1539–1546.
- (78) Asai, A., Hara, M., Kakita, S., Kanda, Y., Yoshida, M., Saito, H., and Saitoh, Y. (1996) Thiol-mediated DNA alkylation by the novel antitumor antibiotic leinamycin. *J. Am. Chem. Soc.* 118, 6802–6803.
- (79) Mehta, P., Church, K., Williams, J., Chen, F.-X., Encell, L., Shuker, D. E. G., and Gold, B. (1996) The design of agents to control DNA methylation adducts. Enhanced major groove methylation of DNA by an N-methyl-N-nitrosourea functionalized phenyl neutral red intercalator. *Chem. Res. Toxicol.* 9, 939–948.
- (80) Vasquez, K. M., Narayanan, L., and Glazer, P. M. (2000) Specific mutations induced by triplex-forming oligonucleotides in mice. *Science* 290, 530–533.
- (81) Wang, Y.-D., Dziegielewska, J., Wurtz, N. R., Dziegielewska, B., Dervan, P. B., and Beerman, T. A. (2003) DNA crosslinking and biological activity of a hairpin polyamide-chlorambucil conjugate. *Nucleic Acids Res.* 31, 1208–1215.
- (82) Dickinson, L. A., Burnett, R., Melander, C., Edelson, B. S., Arora, P. S., Dervan, P. B., and Gottesfeld, J. M. (2004) Arresting cancer proliferation by small-molecule gene regulation. *Chem. Biol.* 11, 1583–1594.
- (83) Povsic, T. J., and Dervan, P. B. (1990) Sequence-specific alkylation of double-helical DNA by oligonucleotide-directed triple-helix formation. *J. Am. Chem. Soc.* 112, 9428–9430.
- (84) Chaudhary, A. K., Nokubo, M., Reddy, G. R., Yeola, S. N., Morrow, J. D., Blair, I. A., and Marnett, L. J. (1994) Detection of endogenous malondialdehyde-deoxyguanine adducts in human liver. *Science* 265, 1580–1582.
- (85) Cho, Y.-J., Kim, H.-Y., Huang, H., Slutsky, A., Minko, I. G., Wang, H., Nechev, L. V., Kozekov, I. D., Kozekova, A., Tamura, P., Jacob, J., Voehler, M., Harris, T. M., Lloyd, R. S., Rizzo, C. J., and Stone, M. P. (2005) Spectroscopic characterization of interstrand carbino-lamine cross-links formed in the 5'-CpG-3' sequence by the acrolein-derived γ -OH-1,N²-propano-2'-deoxyguanosine DNA adduct. *J. Am. Chem. Soc.* 127, 17686–17696.
- (86) Tomasz, M. (1995) Mitomycin C: Small, fast and deadly (but very selective). *Chem. Biol.* 2, 575–579.

- (87) Millard, J. T., Raucher, S., and Hopkins, P. B. (1990) Mechlorethamine cross-links deoxyguanosine residues at 5'-GNC sequences in duplex DNA fragments. *J. Am. Chem. Soc.* 112, 2459–2460.
- (88) Le Pla, R. C., Guichard, Y., Bowman, K. J., Gaskell, M., Farmer, P. B., and Jones, G. D. D. (2004) Further development of ³²P-postlabeling for the detection of alkylphosphotriesters: Evidence for the long-term nonrandom persistence of ethyl-phosphotriester adducts in vivo. *Chem. Res. Toxicol.* 17, 1491–1500.
- (89) Beranek, D. T., Weiss, C. C., and Swenson, D. H. (1980) A comprehensive quantitative analysis of methylated and ethylated DNA using high-pressure liquid chromatography. *Carcinogenesis* 1, 595–606.
- (90) Bernadou, J., Blandin, M., and Meunier, B. (1983) A one step synthesis of O⁶-methyl-2'-deoxyguanosine. *Nucleosides Nucleotides* 2, 459–464.
- (91) Guy, A., Molko, D., Wagrez, L., and Teoule, R. (1986) Chemical synthesis of oligonucleotides containing N⁶-methyladenine residues in the GATC site. *Helv. Chim. Acta* 69, 1034–40.
- (92) Perrino, F. W., Blans, P., Harvey, S., Gelhaus, S. L., McGrath, C., Akman, S. A., Jenkins, S., LaCourse, W. R., and Fishbein, J. C. (2003) The N²-ethylguanine and the O⁶-ethyl- and O⁶-methylguanine lesions in DNA: Contrasting responses from the "bypass" DNA polymerase η and the replicative DNA polymerase α . *Chem. Res. Toxicol.* 16, 1616–1623.
- (93) Wilds, C. J., Noronha, A. M., Robidoux, S., and Miller, P. S. (2004) Mismatch-aligned N³T-alkyl-N³T interstrand cross-linked DNA: Synthesis and characterization of duplexes with interstrand cross-links of variable lengths. *J. Am. Chem. Soc.* 126, 9257–9265.
- (94) Spratt, T. E., and Campbell, C. R. (1994) Synthesis of oligonucleotides containing analogs of O⁶-methylguanine and reaction with O⁶-alkylguanine-DNA alkyltransferase. *Biochemistry* 33, 11364–11371.
- (95) Fischhaber, P. L., Gall, A. S., Duncan, J. A., and Hopkins, P. B. (1999) Direct demonstration in synthetic oligonucleotides that N,N'-bis(2-chloroethyl)-nitrosourea cross-links N1 of deoxyguanosine to N3 of deoxycytidine on opposite strands of duplex DNA. *Cancer Res.* 59, 4363–4368.
- (96) Singer, B., Kroger, M., and Carrano, M. (1978) O²- and O⁴-Alkyl pyrimidine nucleosides: Stability of the glycosyl bond and of the alkyl group as a function of pH. *Biochemistry* 17, 1246–1250.
- (97) Bignami, M., O'Driscoll, M., Aquilina, G., and Karran, P. (2000) Unmasking a killer: DNA O⁶-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462, 71–82.
- (98) Ellison, K. S., Dogliotti, E., Connors, E., Basu, A. K., and Essigmann, J. M. (1989) Site-specific mutagenesis by O⁶-alkylguanines located in the chromosomes of mammalian cells: Influence of the mammalian O⁶-alkylguanine-DNA alkyltransferase. *Proc. Natl. Acad. Sci. U.S.A.* 86, 8620–8624.
- (99) Pauly, G. T., and Moschel, R. C. (2001) Mutagenesis by O⁶-methyl-, O⁶-ethyl, and O⁶-benzylguanine and O⁴-methylthymine in human cells: Effects of O⁶-alkylguanine-DNA alkyltransferase and mismatch repair. *Chem. Res. Toxicol.* 14, 894–900.
- (100) Loeb, L. A., and Preston, B. D. (1986) Mutagenesis by apurinic/apyrimidinic sites. *Ann. Rev. Genet.* 20, 201–230.
- (101) Lawley, P. D., and Brookes, P. (1963) Further studies on the alkylation of nucleic acids and their constituent nucleotides. *Biochem. J.* 89, 127–138.
- (102) Lawley, P. D., and Warren, W. (1976) Removal of minor methylation products 7-methyladenine and 3-methylguanine from DNA of E. coli treated with dimethyl sulfate. *Chem.-Biol. Interact.* 12, 211–220.
- (103) Gates, K. S., Nooner, T., and Dutta, S. (2004) Biologically relevant chemical reactions of N⁷-alkyl-2'-deoxyguanosine adducts in DNA. *Chem. Res. Toxicol.* 17, 839–856.
- (104) Shipova, K., and Gates, K. S. (2005) A fluorimetric assay for the spontaneous release of an N⁷-alkylguanine residue from duplex DNA. *Bioorg. Med. Chem. Lett.* 15, 2111–2113.
- (105) Nooner, T., Dutta, S., and Gates, K. S. (2004) Chemical properties of the leinamycin 2'-deoxyguanosine adduct. *Chem. Res. Toxicol.* 17, 942–949.
- (106) Guthrie, R. D., and Jencks, W. P. (1989) IUPAC recommendations for the representation of reaction mechanisms. *Acc. Chem. Res.* 22, 343–349.
- (107) Moschel, R. C., Hudgins, W. R., and Dipple, A. (1984) Substituent-induced effects on the stability of benzylated guanines: Model systems for the factors influencing the stability of carcinogen-modified nucleic acids. *J. Org. Chem.* 49, 363–372.
- (108) Muller, N., and Eisenbrand, G. (1985) The influence of N⁷-substituents on the stability of N⁷-alkylated guanines. *Chem.-Biol. Interact.* 53, 173–181.
- (109) Lawley, P. D., and Brookes, P. (1961) Acidic dissociation of 7,9-dialkylguanines and its possible relation to mutagenic properties of alkylating agents. *Nature* 192, 1081–2.
- (110) Haines, J. A., Reese, C. B., and Todd, L. (1962) Methylation of guanosine and related compounds with diazomethane. *J. Chem. Soc.* 5281.
- (111) Townsend, L. B., and Robins, R. K. (1963) Ring cleavage of purine nucleosides to yield possible biogenetic precursors of pteridines and riboflavin. *J. Am. Chem. Soc.* 85, 242–243.
- (112) Box, H. C., Lilgam, K. T., French, J. B., Potienko, G., and Alderfer, J. L. (1981) ¹³C-NMR characterization of alkylated derivatives of guanosine, adenosine, and cytidine. *Carbohydrates, Nucleosides, Nucleotides* 8, 189.
- (113) Chetsanga, C. J., Bearie, B., and Makaroff, C. (1982) Alkaline opening of imidazole ring of 7-methylguanosine 1. Analysis of the resulting pyrimidine derivatives. *Chem.-Biol. Interact.* 41, 217–233.
- (114) Chetsanga, C. J., Bearie, B., and Makaroff, C. (1982) Alkaline opening of imidazole ring of 7-methylguanosine. 1. Analysis of the resulting pyrimidine derivatives. *Chem.-Biol. Interact.* 41, 217–233.
- (115) Hecht, S. M., Adams, B. L., and Kozarich, J. W. (1976) Chemical transformations of 7,9-disubstituted purines and related heterocycles. Selective reduction of imines and immonium salts. *J. Org. Chem.* 41, 2303–2311.
- (116) Humphreys, W. G., and Guengerich, F. P. (1991) Structure of formamidopyrimidine adducts as determined by NMR using specifically ¹⁵N-labeled guanosine. *Chem. Res. Toxicol.* 4, 632–636.
- (117) Hendler, S., Furer, E., and Srinivasan, P. R. (1970) Synthesis and chemical properties of monomers and polymer containing 7-methylguanine and an investigation of their substrate or template properties form bacterial deoxyribonucleic acid or ribonucleic acid polymerases. *Biochemistry* 9, 4141–4153.
- (118) Mao, H., Deng, Z., Wang, F., Harris, T. M., and Stone, M. P. (1998) An intercalated and thermally stable FAPY adduct of aflatoxin B1 in a DNA duplex: Structural refinement from ¹H NMR. *Biochemistry* 37, 4374–4387.
- (119) Croy, R. G., and Wogan, G. N. (1981) Temporal patterns of covalent DNA adducts in rat liver after single and multiple doses of aflatoxin B1. *Cancer Res.* 41, 197–203.
- (120) Beranek, D. T., Weiss, C. C., Evans, F. E., Chetsanga, C. J., and Kadlubar, F. F. (1983) Identification of N⁵-methyl-N⁵-formyl-2,5,6-triamino-4-hydroxypyrimidine as a major adduct in rat liver DNA after treatment with the carcinogens, N,N-dimethylnitrosamine or 1,2-dimethylhydrazine. *Biochem. Biophys. Res. Commun.* 110, 625–631.
- (121) Kadlubar, F. F., Beranek, D. T., Weiss, C. C., Evans, F. E., Cox, R., and Irving, C. C. (1984) Characterization of the purine ring-opened 7-methylguanine and its persistence in rat bladder epithelial DNA after treatment with the carcinogen N-methylnitrosourea. *Carcinogenesis* 5, 587–592.
- (122) Kristov, P. P., Brown, K. L., Kozekov, I. D., Stone, M. P., Harris, T. M., and Rizzo, C. J. (2008) Site-specific synthesis and characterization of oligonucleotides containing an N⁶-(2-deoxy-D-erythro-pentofuranosyl)-2,6-diamino-3,4-dihydro-4-oxo-5-N-methylformamidopyrimidine lesion, the ring-opened product from N⁷-methylation of deoxyguanosine. *Chem. Res. Toxicol.* 21, 2324–2333.
- (123) Greenberg, M. M., Hantosi, Z., Wiederholt, C. J., and Rithner, C. D. (2001) Studies on N⁴-(2-deoxy-D-pentofuranosyl)-4,6-diamino-5-formamidopyrimidine (Fapy-dA) and N⁶-(2-deoxy-D-pentofuranosyl)-6-diamino-5-formamido-4-hydroxypyrimidine (Fapy-dG). *Biochemistry* 40, 15856–15861.
- (124) Brown, K. L., Deng, J. Z., Iyer, R. S., Voehler, M. W., Stone, M. P., Harris, C. M., and Harris, T. M. (2006) Unraveling the aflatoxin-FAPY conundrum: Structural basis for differential replicative processing of isomeric forms of the formamidopyrimidine-type DNA adduct of aflatoxin B1. *J. Am. Chem. Soc.* 128, 15188–15199.
- (125) Patro, J. N., Haraguchi, K., Delaney, M. O., and Greenberg, M. M. (2004) Probing the configurations of formamidopyrimidine lesions Fapy-dA and Fapy-dG in DNA using endonuclease IV. *Biochemistry* 43, 13397–13403.
- (126) Bergdorf, L. T., and Carrel, T. (2002) Synthesis, stability, and conformation of the formamidopyrimidine G DNA lesion. *Chem.-Eur. J.* 8, 293–301.
- (127) Fujii, T., and Itaya, T. (1998) The Dimroth rearrangement in the adenine series: A review updated. *Heterocycles* 48, 359–390.
- (128) Barlow, T., Takeshita, J., and Dipple, A. (1998) Deamination and Dimroth rearrangement of deoxyadenosine-styrene oxide adducts in DNA. *Chem. Res. Toxicol.* 11, 838–845.
- (129) Macon, J. B., and Wolfenden, R. (1968) 1-Methyladenosine Dimroth rearrangement and reversible reduction. *Biochemistry* 7, 3453–3458.
- (130) Barlow, T., Ding, J., Vouros, P., and Dipple, A. (1997) Investigation of hydrolytic deamination of 1-(2-hydroxy-1-phenylethyl)adenosine. *Chem. Res. Toxicol.* 10, 1247–1249.
- (131) Koskinen, M., and Hemminki, K. (1999) Separate deamination mechanisms for isomeric styrene oxide induced N¹-adenine adducts. *Org. Lett.* 1, 1233–1235.

- (132) Barlow, T., and Dipple, A. (1999) Formation of deminated products in styrene oxide reactions with deoxycytidine. *Chem. Res. Toxicol.* 12, 883–886.
- (133) Breen, A. P., and Murphy, J. A. (1995) Reactions of oxyl radicals with DNA. *Free Radical Biol. Med.* 18, 1033–1077.
- (134) Greenberg, M. M. (1998) Investigating nucleic acid damage processes via independent generation of reactive intermediates. *Chem. Res. Toxicol.* 11, 1235–1248.
- (135) Pogozelski, W. K., and Tullius, T. D. (1998) Oxidative strand scission of nucleic acids: Routes initiated by hydrogen atom abstraction from the sugar moiety. *Chem. Rev.* 98, 1089–1107.
- (136) Pratviel, G., Bernadou, J., and Meunier, B. (1995) Carbon-hydrogen bonds of DNA sugar units as targets for chemical nucleases and drugs. *Angew. Chem., Int. Ed. Engl.* 34, 746–769.
- (137) von Sonntag, C., Hagen, U., Schon-Bopp, A., and Schulte-Frohlinde, D. (1981) Radiation-induced strand breaks in DNA: Chemical and enzymatic analysis of end groups and mechanistic aspects. *Adv. Radiat. Biol.* 9, 109–142.
- (138) Balasubramanian, B., Pogozelski, W. K., and Tullius, T. D. (1998) DNA strand breaking by the hydroxyl radical is governed by the accessible surface areas of the hydrogen atoms of the DNA backbone. *Proc. Natl. Acad. Sci. U.S.A.* 95, 9738–9743.
- (139) Chowdhury, G., Junnuthula, V., Daniels, J. S., Greenberg, M. M., and Gates, K. S. (2007) DNA strand damage analysis provides evidence that the tumor cell-specific cytotoxin tirapazamine produces hydroxyl radical and acts as a surrogate for O_2 . *J. Am. Chem. Soc.* 129, 12870–12877.
- (140) Carter, K. N., and Greenberg, M. M. (2003) Tandem lesions are the major products resulting from a pyrimidine nucleobase radical. *J. Am. Chem. Soc.* 125, 13376–13378.
- (141) Hwang, J.-T., and Greenberg, M. M. (1999) Kinetics and stereoselectivity of thiol trapping of deoxyuridin-1'-yl in biopolymers and their relationship to the formation of premutagenic α -deoxynucleotides. *J. Am. Chem. Soc.* 121, 4311–4315.
- (142) Emmanuel, C. J., Newcomb, M., Ferreri, C., and Chatgililoglu, C. (1999) Kinetics of 2'-deoxyuridin-1'-yl radical reactions. *J. Am. Chem. Soc.* 121, 2927–2928.
- (143) Greenberg, M. M. (2007) Elucidating DNA damage and repair processes by independently generating reactive and metastable intermediates. *Org. Biomol. Chem.* 5, 18–30.
- (144) Hwang, J.-T., Tallman, K. A., and Greenberg, M. M. (1999) The reactivity of the 2-deoxyribonolactone lesion in single-stranded DNA and its implication in reaction mechanisms of DNA damage and repair. *Nucleic Acids Res.* 27, 3805–3810.
- (145) Roupiez, Y., Lhomme, J., and Kotera, M. (2002) Chemistry of the 2-deoxyribonolactone lesion in oligonucleotides: Cleavage kinetics and products analysis. *J. Am. Chem. Soc.* 124, 9129–9135.
- (146) Zheng, Y., and Sheppard, T. L. (2004) Half-life and DNA strand scission products of 2-deoxyribonolactone oxidative DNA damage lesions. *Chem. Res. Toxicol.* 17, 197–207.
- (147) Kroeger, K. M., Hashimoto, M., Kow, Y. W., and Greenberg, M. M. (2003) Cross-linking of 2-deoxyribonolactone and its β -elimination product by base excision repair enzymes. *Biochemistry* 42, 2449–2455.
- (148) Hashimoto, M., Greenberg, M. M., Kow, Y. W., Hwang, J.-T., and Cunningham, R. P. (2001) The 2-deoxyribonolactone lesion produced in DNA by neocarzinostatin and other damaging agents forms cross-links with the base-excision repair enzyme endonuclease III. *J. Am. Chem. Soc.* 123, 3161–3162.
- (149) Junnuthula, V., Sarkar, U., Sinha, S., and Gates, K. S. (2009) Initiation of DNA strand cleavage by 1,2,4-benzotriazine 1,4-dioxides: Mechanistic insight from studies of 3-methyl-1,2,4-benzotriazine 1,4-dioxide. *J. Am. Chem. Soc.* 131, 1015–1024.
- (150) Dussy, A., Meggers, E., and Giese, B. (1998) Spontaneous cleavage of 4'-DNA radicals under aerobic conditions: Apparent discrepancy between trapping rates and cleavage products. *J. Am. Chem. Soc.* 120, 7399–7403.
- (151) Giese, B., Dussy, A., Meggers, E., Petretta, M., and Schwitter, U. (1997) Conformation, lifetime, and repair of 4'-DNA radicals. *J. Am. Chem. Soc.* 119, 11130–11131.
- (152) Hildebrand, K., and Schulte-Frohlinde, D. (1997) Time-resolved EPR studies on the reaction rates of peroxyl radicals of poly(acrylic acid) and of calf thymus DNA with glutathione. Re-examination of a rate constant for DNA. *Int. J. Radiat. Biol.* 71, 377–385.
- (153) Giese, B., Beyrich-Graf, X., Erdmann, P., Petretta, M., and Schwitter, U. (1995) The chemistry of single-stranded 4'-DNA radicals: Influence of the radical precursor on anaerobic and aerobic strand cleavage. *Chem. Biol.* 2, 367–375.
- (154) Dedon, P. C., Plastaras, J. P., Rouzer, C. A., and Marnett, L. J. (1998) Indirect mutagenesis by oxidative DNA damage: formation of the pyrimidopurine adduct of deoxyguanosine by base propenal. *Proc. Natl. Acad. Sci. U.S.A.* 95, 11113–11116.
- (155) Plastaras, J. P., Riggins, J. N., Otteneider, M., and Marnett, L. J. (2000) Reactivity and mutagenicity of endogenous DNA oxopropenylating agents: Base propenals, malondialdehyde, and N^{ϵ} -oxopropenyllysine. *Chem. Res. Toxicol.* 13, 1235–1242.
- (156) von Sonntag, C. (2006) *Free-Radical-Induced DNA Damage and Its Repair*, Springer-Verlag, Berlin.
- (157) Burrows, C. J., and Muller, J. G. (1998) Oxidative nucleobase modifications leading to strand scission. *Chem. Rev.* 98, 1109–1151.
- (158) Gajewski, E., Rao, G., Nackerdien, Z., and Dizdaroglu, M. (1990) Modification of DNA bases in mammalian chromatin by radiation-generated free radicals. *Biochemistry* 29, 7876–7882.
- (159) Wallace, S. S. (2002) Biological consequences of free radical-damaged DNA bases. *Free Radical Biol. Med.* 33, 1–14.
- (160) Dizdaroglu, M. (1994) Chemical determination of oxidative DNA damage by gas chromatography-mass spectrometry. *Methods Enzymol.* 234, 3–16.
- (161) Dizdaroglu, M., Jaruga, P., Birincioglu, M., and Rodriguez, H. (2002) Free radical-induced damage to DNA: Mechanisms and measurement. *Free Radical Biol. Med.* 32, 1102–1115.
- (162) Evans, M. D., Dizdaroglu, M., and Cooke, M. S. (2004) Oxidative DNA damage and disease: Induction, repair, and significance. *Mutat. Res.* 567, 1–61.
- (163) Jovanovic, S. V., and Simic, M. G. (1986) Mechanism of OH radical reactions with thymine and uracil derivatives. *J. Am. Chem. Soc.* 108, 5968–5972.
- (164) Birincioglu, M., Jaruga, P., Chowdhury, G., Rodriguez, H., Dizdaroglu, M., and Gates, K. S. (2003) DNA base damage by the antitumor agent 3-amino-1,2,4-benzotriazine 1,4-dioxide (Tirapazamine). *J. Am. Chem. Soc.* 125, 11607–11615.
- (165) Schuchmann, M. N., and von Sonntag, C. (1983) The radiolysis of uracil in oxygenated aqueous solutions. A study by product analysis and pulse radiolysis. *J. Chem. Soc. Perkin 2*, 1525–1531.
- (166) Carter, K. N., and Greenberg, M. M. (2003) Independent generation and study of 5,6-dihydro-2'-deoxyuridin-6-yl, a member of the major family of reactive intermediates formed in DNA from the effects of γ -radiolysis. *J. Org. Chem.* 68, 4275–4280.
- (167) Wagner, J. R., van Lier, J. E., Berger, M., and Cadet, J. (1994) Thymidine hydroperoxides: Structural assignment, conformational features, and thermal decomposition in water. *J. Am. Chem. Soc.* 116, 2235–2242.
- (168) Steenken, S., and Jovanovic, S. V. (1997) How easily oxidizable is DNA? One-electron reduction potentials of adenosine and guanosine radicals in aqueous solution. *J. Am. Chem. Soc.* 119, 617–618.
- (169) Lewis, F. D., Letsinger, R. L., and Wasielewski, M. R. (2001) Dynamics of photoinduced charge transfer and hole transport in synthetic DNA hairpins. *Acc. Chem. Res.* 34, 159–170.
- (170) Candeias, L. P., and Steenken, S. (2000) Reaction of HO \cdot with guanine derivatives in aqueous solution: Formation of two different redox-active OH-adduct radicals and their unimolecular transformation reactions. Properties of G(-H). *Chem.-Eur. J.* 6, 475–484.
- (171) Douki, T., Spinelli, S., Ravanat, J.-L., and Cadet, J. (1999) Hydroxyl radical-induced degradation of 2'-deoxyguanosine under reducing conditions. *J. Chem. Soc. Perkin 2*, 1875–1880.
- (172) Wiederholt, C. J., and Greenberg, M. M. (2002) Fapy-dG instructs Klenow Exo $^{-}$ to misincorporate deoxyadenosine. *J. Am. Chem. Soc.* 124, 7278–7279.
- (173) Haraguchi, K., Delaney, M. O., Wiederholt, C. J., Sambandam, A., Hantosi, Z., and Greenberg, M. M. (2002) Synthesis and characterization of oligodeoxynucleotides containing formamidopyrimidine lesions and nonhydrolyzable analogues. *J. Am. Chem. Soc.* 124, 3263–3269.
- (174) Haraguchi, K., and Greenberg, M. M. (2001) Synthesis of oligonucleotides containing FapydG (N6-(2-deoxy- α,β -D-erythro-pentofuranosyl)-2,6-diamino-4-hydroxy-5-formamidopyrimidine). *J. Am. Chem. Soc.* 123, 8636–8637.
- (175) Delaney, M. O., Wiederholt, C. J., and Greenberg, M. M. (2002) Fapy-dA induces nucleotide misincorporation translesionally by a DNA polymerase. *Angew. Chem.* 41, 771–3.
- (176) Bodepudi, V., Shibutani, S., and Johnson, F. (1992) Synthesis of 2'-deoxy-7,8-dihydro-8-oxo-guanosine and 2'-deoxy-7,8-dihydro-8-oxoadenosine and their incorporation into oligomeric DNA. *Chem. Res. Toxicol.* 5, 608–617.
- (177) Burrows, C. J., Muller, J. G., Kornysushyna, O., Luo, W., Duarte, V., Leipold, M. D., and David, S. S. (2002) Structure and potential mutagenicity of new hydantoin products from guanosine and 8-oxo-7,8-dihydroguanine oxidation by transition metals. *Environ. Health Perspect. Suppl.* 110, 713–717.
- (178) Luo, W., Muller, J. G., Rachlin, E. M., and Burrows, C. J. (2001) Characterization of hydantoin products from one-electron oxidation of 8-oxo-7,8-dihydroguanosine in a nucleoside model. *Chem. Res. Toxicol.* 14, 927–938.

- (179) Luo, W., Muller, E. M., and Burrows, C. J. (2000) Characterization of spiroiminodihydantoin as a product of one-electron oxidation of 8-oxo-7,8-dihydroguanosine. *Org. Lett.* 2, 613–616.
- (180) Ye, Y., Muller, J. G., Luo, W., Mayne, C. L., Shallop, A. J., Jones, R. A., and Burrows, C. J. (2003) Formation of ^{13}C -, ^{15}N -, and ^{18}O -labeled guanidinohydantoin from guanosine oxidation with singlet oxygen. Implications for structure and mechanism. *J. Am. Chem. Soc.* 125, 13926–13927.
- (181) Matter, B., Malejka-Giganti, D., Cxallany, A. S., and Tretyakova, N. (2006) Quantitative analysis of the oxidative DNA lesion, 2,2-diamino-4-(2-deoxy- β -D-erythro-pentofuranosyl)amino-5-(2H)-oxazolone (oxazolone), in vitro and in vivo by isotope dilution-capillary HPLC-ESI-MS/MS. *Nucleic Acids Res.* 34, 5449–5460.
- (182) Neeley, W. L., and Essigmann, J. M. (2006) Mechanisms of formation, genotoxicity, and mutation of guanine oxidation products. *Chem. Res. Toxicol.* 19, 491–505.
- (183) Kuchino, Y., Mori, F., Kasai, H., Inoue, H., Iwai, S., Miura, K., Ohtsuka, E., and Nishimura, S. (1987) Misreading of DNA templates containing 8-hydroxyguanosine at the modified base and at adjacent residues. *Nature* 327, 77–79.
- (184) Kornysheva, O., Berges, A. M., Muller, J. G., and Burrows, C. J. (2002) In vitro nucleotide misinsertion opposite the oxidized guanosine lesions spiroiminohydantoin and guanidinohydantoin and DNA synthesis past the lesions using *Escherichia coli* DNA polymerase I (Klenow fragment). *Biochemistry* 41, 15304–15314.
- (185) Henderson, P. T., Delaney, J. C., Muller, J. G., Neeley, W. L., Tannenbaum, S. R., Burrows, C. J., and Essigmann, J. M. (2003) The hydantoin lesions formed from oxidation of 7,8-dihydro-8-oxoguanine are potent sources of replication errors in vivo. *Biochemistry* 42, 9257–9262.
- (186) Shibutani, S., Takeshita, M., and Grollman, A. P. (1991) Insertion of specific bases during DNA synthesis past the oxidation damaged base 8-oxodG. *Nature* 349, 431–434.
- (187) Brooks, P. J., Wise, D. S., Berry, D. A., Kosmoski, J. V., Smerdon, M. J., Somers, R. L., Mackie, H., Spoonde, A. Y., Ackerman, E. J., Coleman, K., Tarone, R. E., and Robbins, J. H. (2000) The oxidative lesion 8,5'-cyclo-2'-deoxyadenosine is repaired by the nucleotide excision repair pathway and blocks gene expression in mammalian cells. *J. Biol. Chem.* 275, 22355–22362.
- (188) Kuraoka, I., Bender, C., Romieu, A., Cadet, J., Wood, R., and Lindahl, T. (2000) Removal of oxygen free-radical-induced 5',8-purine cyclodeoxynucleosides from DNA by the nucleotide excision repair pathway in human cells. *Proc. Natl. Acad. Sci. U.S.A.* 97, 3832–3837.
- (189) Greenberg, M. M., Barvian, M. R., Cook, G. P., Goodman, B. K., Matray, T. J., Tronche, C., and Venkatesan, H. (1997) DNA damage induced via 5,6-dihydrothymid-5-yl in single-stranded oligonucleotides. *J. Am. Chem. Soc.* 119, 1828–1839.
- (190) Tallman, K. A., and Greenberg, M. M. (2001) Oxygen-dependent DNA damage amplification involving 5,6-dihydrothymidin-5-yl in a structurally minimal system. *J. Am. Chem. Soc.* 123, 5181–5187.
- (191) Bellon, S., Ravanat, J.-L., Gasparutto, D., and Cadet, J. (2002) Cross-linked thymine-purine base tandem lesions: Synthesis, characterization, and measurement in γ -irradiated DNA. *Chem. Res. Toxicol.* 15, 598–606.
- (192) Box, H. C., Budzinski, E. E., Dawdzik, J. B., Wallace, J. C., and Iijima, H. (1998) Tandem lesions and other products in X-irradiated DNA oligomers. *Radiat. Res.* 149, 433–439.
- (193) Zhang, Q., and Wang, Y. (2005) Generation of 5-(2'-deoxycytidyl) methyl radical and the formation of intrastrand cross-link lesions in oligodeoxyribonucleotides. *Nucleic Acids Res.* 33, 1593–1603.
- (194) Ding, H., and Greenberg, M. M. (2007) γ -Radiolysis and hydroxyl radical produce interstrand cross-links in DNA involving thymidine. *Chem. Res. Toxicol.* 20, 1623–1628.
- (195) Ding, H., Majumdar, A., Tolman, J. R., and Greenberg, M. M. (2008) Multinuclear NMR and kinetic analysis of DNA interstrand cross-link formation. *J. Am. Chem. Soc.* 130, 17981–17987.
- (196) Dizdaroglu, M., Jaruga, P., and Rodriguez, H. (2001) Identification and quantification of 8,5'-cyclo-2'-deoxyadenosine in DNA by LC/MS. *Free Radical Biol. Med.* 30, 774–784.
- (197) Romieu, A., Gasparutto, D., Molko, D., and Cadet, J. (1998) Site-specific introduction of (5'S)-5',8-cyclo-2'-deoxyadenosine into oligodeoxyribonucleotides. *J. Org. Chem.* 63, 5245–5249.
- (198) Jaruga, P., Birincioglu, M., Rodriguez, H., and Dizdaroglu, M. (2002) Mass spectrometric assays for the tandem lesion 8,5'-cyclo-2'-deoxyguanosine in mammalian DNA. *Biochemistry* 41, 3703–3711.
- (199) Kuraoka, I., Robins, P., Masutani, C., Hanaoka, F., Gasparutto, D., Cadet, J., Wood, R. D., and Lindahl, T. (2001) Oxygen free radical damage to DNA: Translesion synthesis by human DNA polymerase η and resistance to exonuclease action at cyclopurine deoxynucleoside residues. *J. Biol. Chem.* 276, 49283–49288.
- (200) Muller, E., Gasparutto, D., Jaquinod, M., Romieu, A., and Cadet, J. (2000) Chemical and biochemical properties of oligonucleotides that contain 5'S,6S-cyclo-5,6-dihydro-2'-deoxyuridine and 5'S,6S-cyclo-5,6-dihydrothymidine, two main radiation-induced degradation products of pyrimidine 2'-deoxyribonucleosides. *Tetrahedron* 56, 8689–8701.
- (201) Muller, E., Gasparutto, D., and Cadet, J. (2002) Chemical synthesis and biochemical properties of oligonucleotides that contain the (5'S,5S,6S)-5',6-cyclo-5-hydroxy-5,6-dihydro-2'-deoxyuridine DNA lesion. *ChemBioChem* 3, 534–542.
- (202) Romieu, A., Gasparutto, D., and Cadet, J. (1999) Synthesis and characterization of oligodeoxyribonucleotides containing the two 5R and 5S diastereomers of 5'S,6S-5',6-cyclo-5,6-dihydrothymidine; radiation-induced tandem lesions of thymidine. *J. Chem. Soc. Perkin I*, 1257–1263.
- (203) Crick, F. H. C., Wang, J. C., and Bauer, W. R. (1979) Is DNA really a double helix? *J. Mol. Biol.* 129, 449–457.
- (204) Sturla, S. J. (2007) DNA adduct profiles: chemical approaches to addressing the biological impact of DNA damage from small molecules. *Curr. Opin. Chem. Biol.* 11, 293–299.
- (205) Cadet, J., Douki, T., Gasparutto, D., and Ravanat, J.-L. (2003) Oxidative damage to DNA: Formation measurement and biochemical features. *Mutat. Res.* 531, 5–23.
- (206) Jaruga, P., Gueldal, K., and Dizdaroglu, M. (2008) Measurement of formamidopyrimidines in DNA. *Free Radical Biol. Med.* 45, 1601–1609.
- (207) Butenandt, J., Burgdorf, L. T., and Carell, T. (1999) Synthesis of DNA lesions and DNA-lesion-containing oligonucleotides for DNA-repair studies. *Synthesis* 1085–1105.

TX900242K