

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/50302899>

Cytotoxic Clerodane Diterpenes from *Casearia rupestris*

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · MARCH 2011

Impact Factor: 3.8 · DOI: 10.1021/np100840w · Source: PubMed

CITATIONS

15

READS

72

10 AUTHORS, INCLUDING:



Gerardo Magela Vieira Júnior

Universidade Federal de Mato Grosso (UFMT)

22 PUBLICATIONS 350 CITATIONS

SEE PROFILE



Paulo Michel Pinheiro Ferreira

Universidade Federal do Piauí

58 PUBLICATIONS 559 CITATIONS

SEE PROFILE



Vanderlan S Bolzani

São Paulo State University

331 PUBLICATIONS 3,807 CITATIONS

SEE PROFILE



Alberto José Cavaleiro

São Paulo State University

111 PUBLICATIONS 1,208 CITATIONS

SEE PROFILE

Cytotoxic Clerodane Diterpenes from *Casearia rupestris*

Gerardo M. Vieira-Júnior,[†] Luiz A. Dutra,[†] Paulo M. P. Ferreira,[‡] Manoel O. de Moraes,[§] Leticia V. Costa Lotufo,[§] Cláudia do Ó Pessoa,[§] Roseli Buzanelli Torres,[‡] Nivaldo Borallo,[†] Vanderlan da S. Bolzani,[†] and Alberto J. Cavalheiro^{*,†}

[†]Institute of Chemistry, Nucleus of Bioassay, Biosynthesis and Ecophysiology of Natural Products (NuBBE), São Paulo State University, UNESP, C.P. 14801-970, Araraquara, SP, Brazil


[‡]Departamento de Ciências da Saúde, Campus Senador Helvídio Nunes de Barros, Universidade Federal do Piauí, UFPI, C.P. 64600-000, Picos, PI, Brazil

[§]Laboratório de Oncologia Experimental (LOE), Departamento de Fisiologia e Farmacologia, Universidade Federal do Ceará, UFC, C.P. 60430-270, Fortaleza, CE, Brazil

[†]Núcleo do Jardim Botânico, Instituto Agrônomo de Campinas, C.P. 13020-902, Campinas, SP, Brazil

Supporting Information

ABSTRACT: Four new clerodane diterpenes, casearupestrins A–D (**1–4**), were isolated from the leaves of *Casearia rupestris*. Compounds **1** and **4** were acetylated to yield 2,7-di-*O*-acetylcasearupestrin A (**5**) and 2,6-di-*O*-acetylcasearupestrin D (**6**). All compounds were evaluated for cytotoxicity against a small panel of human cancer cell lines. Casearupestrin A (**1**) exhibited the most potent activity against MDA/MB-435 (human melanoma) and SF-295 (human glioblastoma) cells, superior to that of the standard drug doxorubicin.



	R ₁	R ₂	R ₃	R ₄
1	OH	OAc	2 <i>Z</i> ,4 <i>E</i> -decadienoate	OH
2	OH	OAc	OH	2 <i>Z</i> ,4 <i>E</i> -decadienoate
3	OH	OCH ₃	OH	2 <i>Z</i> ,4 <i>E</i> -decadienoate
4	OH	OAc	OAc	2 <i>Z</i> ,4 <i>E</i> -decadienoate
5	OAc	OAc	2 <i>Z</i> ,4 <i>E</i> -decadienoate	OAc
6	OAc	OAc	OAc	2 <i>Z</i> ,4 <i>E</i> -decadienoate

The tropical genus *Casearia* (*Salicaceae sensu lato*) contains ca. 180 species,¹ and it is a rich source of clerodane-type diterpenes.^{2–13} The genus occurs widely in Brazil, where *C. sylvestris* is the most frequent species, occurring from the Amazon basin throughout the Cerrado (savanna) to the Atlantic Forest and Southern Brazil. The clerodanes are a group of diterpenoids with interesting biological effects such as antifungal,¹⁴ antileishmanial and antiparasitoid,¹⁵ antimalarial and antimycobacterial,¹⁶ antitumor,^{17,18} cytotoxic,^{1,14,16,19–31} immunomodulatory,³² and trypanocidal activities.³³ In our continuing investigation of the bioactive clerodane diterpenes from *Casearia* species, we undertook the first phytochemical investigation of *Casearia rupestris* Eichler. This shrub or small tree grows up to 6.0 m in height and is found in Brazil (Mato Grosso and Minas Gerais States) and Bolivia (Sta. Cruz State) in the Cerrado, grasslands, and sometimes pastures, at altitudes of 350–500 m.³⁴ Investigation of the cytotoxic Et₂O fraction yielded four new clerodane diterpenes (**1–4**) (Figure 1). Compounds **1** and **4** were acetylated, resulting in two new derivatives (**5**, **6**) (Figure 1). The

cytotoxic activity of the diterpenes (**1–6**) against a panel of human cancer cell lines (HL-60, MDA/MB-435, HCT-8, and SF-295) has been evaluated.

RESULTS AND DISCUSSION


Investigation of the Et₂O phase of the EtOH extract of the leaves of *C. rupestris* led to the isolation of four new clerodane diterpenoids, casearupestrins A–D (**1–4**). 2,7-Di-*O*-acetylcasearupestrin A (**5**) and 2,6-di-*O*-acetylcasearupestrin D (**6**) were obtained by acetylation of compounds **1** and **4**, respectively. The structures (Figure 1) of the compounds were elucidated on the basis of spectroscopic and spectrometric data (IR, UV, NMR, and MS) and by comparison with NMR data of reported compounds.⁶

Casearupestrin A (**1**) was obtained as a white, amorphous solid ($[\alpha]_D^{24} +4.3$ (c 1.0, MeOH)). The molecular formula

Received: November 17, 2010

Published: March 07, 2011





The figure shows the chemical structures of two clerodane diterpenes, 1 and 2, and a 2Z,4E-decadienoate moiety. Structure 1 is a complex polycyclic molecule with a decadienoate side chain. Structure 2 is a similar molecule with a different substitution pattern. The 2Z,4E-decadienoate moiety is shown as a separate structure with its numbering (1' to 9').

	R ₁	R ₂	R ₃	R ₄
1	OH	OAc	2Z,4E-decadienoate	OH
2	OH	OAc	OH	2Z,4E-decadienoate
3	OH	OCH ₃	OH	2Z,4E-decadienoate
4	OH	OAc	OAc	2Z,4E-decadienoate
5	OAc	OAc	2Z,4E-decadienoate	OAc
6	OAc	OAc	OAc	2Z,4E-decadienoate

Figure 1. Structure of the clerodane diterpenes from *Casearia rupestris*.

$C_{34}H_{48}O_9$ was determined by HRESIMS, based on the sodiated ion peak at m/z 623.3190 $[M + Na]^+$ (calcd 623.3190), which implies 11 degrees of unsaturation. The IR spectrum showed the presence of carbonyl esters (1746 and 1712 cm^{-1}). The UV spectrum displayed absorptions at λ_{max} 223 nm ($\log \epsilon$ 1.40) and 268 nm ($\log \epsilon$ 1.50), corresponding to diene and dienone chromophores, respectively. The ^{13}C and DEPT 135° NMR data showed the presence of 34 carbons divided into seven quaternary (including three carbonyl carbons), 13 methine, nine methylene, and five methyl carbons. The olefinic region of the ^1H NMR spectrum of **1** (Table 1) revealed proton signals with characteristic *cis/trans* coupling at δ_{H} 5.06 (1H, d, $J = 10.5$ Hz, H-15), 5.30 (1H, d, $J = 17.5$ Hz, H-15), and 6.47 (1H, dd, $J = 10.5$, 17.5 Hz, H-14), indicating a terminal monosubstituted olefinic moiety. Another terminal methylene, identified by two singlets at δ_{H} 4.94 and 5.03 (2H, H-16), was connected to a quaternary carbon (C-13). An additional olefinic doublet was observed at δ_{H} 6.04 (1H, d, $J = 4.0$ Hz, H-3) and correlated to a methine carbon at δ_{C} 128.6 (C-3) in the gHMBC spectrum. Two deshielded acetal protons were observed at δ_{H} 6.35 (1H, t, $J = 1.5$ Hz, H-18) and 6.49 (1H, s, H-19), which correlated to methine signals at δ_{C} 96.3 and 99.8, respectively. The ^1H NMR spectrum also revealed the presence of two methyl groups at δ_{H} 1.06 (1H, brd, $J = 4.0$ Hz, H-17) and 1.07 (1H, s, H-20) and three oxymethines at δ_{H} 4.45 (1H, brt, $J = 4.0$ Hz, H-2), 5.18 (1H, d, $J = 10.0$ Hz, H-6), and 3.66 (1H, t, $J = 10.0$ Hz, H-7). The spectroscopic data of **1** (Tables 1 and 2) are in accord with the basic skeleton of clerodane diterpenoids previously isolated from *Casearia* species.^{10,18,35} The acyclic moiety at C-9, including the diene group, was deduced from the correlations of H-12 with C-11, H-14 with C-13 and C-16, H-15 with C-13 and C-14, and H-16 with C-14 (Figure 2). The position of the hydroxy group at C-7 was deduced by the long-range correlation of the oxymethine signal at δ_{H} 3.66 (H-7) with δ_{C} 77.5 (C-6), 44.8 (C-8), and 11.7 (C-17) (Figure 2). A decadienoate moiety, identified by gHMBC and gCOSY correlations (Figure 2), was characterized by a *Z*- (δ_{H} 5.68, $J = 11.5$ Hz; 6.69, $J = 11.5$ Hz) and an *E*-double bond (δ_{H} 7.37, $J = 14.0$, 11.5 Hz; 6.16, $J = 14.0$,

7.0 Hz) and a triplet methyl (δ_{H} 0.98, $J = 7.0$ Hz). This same 2Z,4E-decadienoate unit has been verified in other clerodane diterpenes isolated from *C. pitumba* and *C. arguta*.^{6,36} Its position was determined at C-6 (δ_{C} 77.5), on the basis of the long-range correlation observed in the gHMBC spectrum between the oxymethine proton (H-6) and the carbonyl carbon (δ_{C} 167.3, C-1'). Finally, since H-19 (δ_{C} 6.49) was correlated with the carbonyl group (δ_{C} 171.4) of one acetate unit, the second acetate unit (δ_{C} 171.8 and 21.0) must be located at C-18 (Figure 2).

The relative configuration of **1** was determined by using coupling constants, 1D-NOE data, and ^{13}C NMR chemical shifts (Table 1 and Figure 3). The *trans* configuration between the C-17 and C-20 methyl groups was shown by the chemical shift at δ_{C} 26.4 for C-20, which also supported a *cis*-A/B ring junction.^{3,5,10} The J values of 3.5 and 13.5 Hz for the coupling of H-10 and H-1 revealed that H-10 (*rel*- α) is in the axial position. The NOE enhancement between H-19 and H-10, and H-19 and H-7, indicated the same α -orientation to H-19 and H-7 (Figure 2). The large coupling constant between H-6 (δ_{H} 5.18 – 1H, d, $J = 10.0$ Hz) and H-7 (δ_{H} 3.66 – 1H, t, $J = 10.0$ Hz) supported their 1,2-diaxial relationship. The NOE interaction between H-6 (*rel*- β) and H-8 implied that H-8 and Me-17 were at axial and equatorial positions, respectively. A small coupling constant of 4.0 Hz for H-2/H-1 suggested an equatorial orientation for H-2 (*rel*- β), which was confirmed by the broad triplet at δ_{H} 4.45 and the chemical shift of C-2 at δ_{C} 64.2. From the allylic coupling between H-3 and H-18 ($J = 1.5$ Hz) and NOE enhancement observed between H-18 and H-6 it can be deduced that H-18 also has a β -orientation. Therefore, casearupestrin A (**1**), a new clerodane diterpenoid, was assigned as *rel*-(2R,5S,6S,8R,9R,10S,18R,19R)-18,19-diacetoxy-18,19-epoxy-6-[(2'Z,4'E)-decadienoyloxy]-2,7-dihydroxyclo-3,13(16),14-triene.

The MS, UV, and IR data and the specific rotation value of casearupestrin B (**2**) were comparable with those of **1**. The ^1H and ^{13}C NMR spectroscopic data (Tables 1 and 2) revealed the same substituents and substitution patterns. However, changes in the chemical shift values related to H-6 (δ_{H} 3.75 – 1H, d, $J = 10.5$ Hz) and H-7 (δ_{H} 5.15 – 1H, t, $J = 10.5$ Hz) required placement of the decadienoate ester at C-7. The structure **2** was corroborated by the same correlations observed for **1** in the gHMBC, gHMBC, and gCOSY (Figure 2). The observed NOESY interactions and coupling constants for casearupestrin B (**2**) indicate the same relative configuration as **1** (Figure 3 and Table 1). Hence, casearupestrin B (**2**) was assigned as *rel*-(2R,5S,6S,8R,9R,10S,18R,19R)-18,19-diacetoxy-18,19-epoxy-7-[(2'Z,4'E)-decadienoyloxy]-2,6-dihydroxyclo-3,13(16),14-triene.

The structures of the remaining two compounds can be rationalized by comparison with the previous ones. Analysis of the NMR data of compound **3** revealed a strong similarity to compounds **1** and **2**. However, the molecular formula $C_{33}H_{48}O_8$, determined by the HRESIMS, revealed a sodiated ion peak at m/z 595.3241 $[M + Na]^+$, which implied 10 degrees of unsaturation. The main difference observed in the ^1H and ^{13}C NMR spectra of compound **3** in relation to compounds **1** and **2** (Tables 1 and 2) was the presence of signals at δ_{H} 3.36 (1H, s) and δ_{C} 55.4. These signals indicated that an acetoxy group at C-18 was replaced by a methoxy group, which was confirmed by the gHMBC correlation between H-18 and the methoxy group. The relative configuration of **3** was identical to **1** on the basis of the same NOESY interactions (Figure 3) and coupling constants. Thus, casearupestrin C (**3**) was assigned as *rel*-(2R,5S,6S,8R,9R,10S,18R,19R)-19-acetoxy-18,

Table 1. ¹H NMR Data for Compounds 1–6 (methanol-*d*₄, 500 and 300 MHz, *J* in Hz)

position	1	2	3	4	5	6
1	2.00 m 2.06 m	1.34 m 1.91 m	1.96 m	2.10 m	2.19 m	2.08 m 2.17 m
2	4.45 brt (4.0)	4.39 brt (4.0)	4.37 brt (4.0)	4.35 brt (4.0)	5.55 m	5.52 m
3	6.04 brd (4.0)	6.02 brd (4.0)	6.05 brd (4.0)	6.07 brd (4.0)	6.07 brd (5.5)	6.07 brd (4.0)
6	5.18 d (10.0)	3.75 d (10.5)	3.74 d (10.0)	5.28 brs	5.31 m	5.29 brd
7	3.66 t (10.0)	5.15 t (10.5)	5.16 t (10.0)	5.30 d (12.0)	5.37 m	5.32 brd
8	1.84 m	1.91 m	1.92 m	2.05 m	2.16 m	2.10 m
10	2.47 dd (3.5, 13.5)	2.46 dd (4.0, 13.5)	2.44 dd (4.5, 13.0)	2.51 dd (3.5, 13.5)	2.40 dd (5.0, 22.5)	2.40 dd (5.0, 22.0)
11	1.47 m	1.46 m	1.32 m	1.97 m	1.35 m	1.38 m
12	2.16 m 2.23 m	2.25 m	2.24 m	2.24 m	2.21 m	2.21 m
14	6.47 dd (10.5, 17.5)	6.46 m	6.51 brdd (8.5, 17.5)	6.47 dd (7.0, 11.0)	6.49 brdd (11.0, 18.5)	6.50 brdd (11.0, 18.0)
15	5.06 d (10.5) 5.30 d (17.5)	5.04 d (10.5) 5.29 d (17.5)	5.06 d (8.5) 5.30 d (17.5)	5.03 d (7.0) 5.30 d (11.0)	5.10 m 5.33 m	5.10 m 5.38 m
16	4.94 s 5.03 s	5.04 s 5.00 s	5.01 s 5.05 brs	5.06 s 5.40 brs	5.07 s	5.06 s
17	1.06 brd (4.0)	0.89 d (7.0)	0.91 brd (6.5)	0.93 brd (7.0)	0.92 brd (6.5)	0.95 brd (7.0)
18	6.35 t (1.5)	6.65 brt (2.0)	5.48 t (1.5)	6.33 t (1.5)	6.33 t (2.5)	6.36 t (3.0)
19	6.49 s	6.51 s	6.48 s	6.53 s	6.58 s	6.58 s
20	1.07 s	1.04 s	1.04 brs	0.98 s	1.08 s	1.08 s
2'	5.68 d (11.5)	5.63 d (11.5)	5.64 d (11.0)	5.40 d (11.0)	5.55 brd (19.0)	5.54 brd (19.0)
3'	6.69 t (11.5)	6.65 t (11.5)	6.67 t (11.0)	6.70 t (11.0)	6.74 t (19.0)	6.71 t (19.0)
4'	7.37 brdd (11.5, 14.0)	7.38 brdd (11.5, 14.0)	7.39 brdd (11.0, 15.5)	7.32 brdd (11.0, 15.5)	7.31 brdd (4.5, 19.0)	7.34 brdd (4.0, 19.0)
5'	6.16 dd (7.0, 14.0)	6.14 dd (7.0, 14.0)	6.15 dd (7.0, 15.5)	6.18 dd (7.5, 15.5)	6.24 m	6.20 m
6'	2.20 m	2.17 m	2.22 m	2.22 m	2.24 m	2.24 m
7'	1.27 m 1.69 m	1.91 m 1.34 m	1.48 m	1.46 m	1.49 m	1.49 m
8'	1.34 m	1.34 m	1.35 m	1.35 m	1.35 m	1.36 m
9'	1.34 m	1.34 m	1.35 m	1.35 m	1.35 m	1.36 m
10'	0.98 t (7.0)	0.92 t (7.0)	0.93 m	0.92 brt	0.92 m	0.93 brt
2-O ₂ CCH ₃					2.12 s	2.12 s
6-O ₂ CCH ₃				1.95 s	1.95 s	1.97 s
18-OCH ₃			3.36 s			
18-O ₂ CCH ₃	2.02 s	2.05 s		2.04 s	2.04 s	2.07 s
19-O ₂ CCH ₃	1.86 s	1.87 s	1.87 s	1.87 s	1.89 s	1.90 s

19-epoxy-7-[(2',4'E)-decadienoyloxy]-2,6-dihydroxy-18-methoxycyclo-3,13(16),14-triene.

Compound 4 had the molecular formula C₃₆H₅₀O₁₀, determined by HRESIMS (observed *m/z* 665.3297 [M + Na]⁺). The ¹H and ¹³C NMR data of 4 (casearupestrin D) (Tables 1 and 2) were similar to those of 2 except for replacement of a hydroxy group at C-6 by an acetoxy group [δ_{H} 1.95 (3H, s), δ_{C} 171.5, δ_{C} 20.9]. The ¹H and ¹³C NMR assignments of compound 4 were confirmed by 2D NMR correlations, and its configuration was established by the sign of the specific rotation, NOESY interactions, and coupling constants (Figure 3). Thus, casearupestrin D (4) was assigned as *rel*-(2*R*,5*S*,6*S*,8*R*,9*R*,10*S*,18*R*,19*R*)-6,18,19-triacetoxy-18,19-epoxy-7-[(2',4'E)-decadienoyloxy]-2-hydroxycyclo-3,13(16),14-triene.

Acetylation of 1 and 4 with Ac₂O–pyridine yielded two new clerodane diterpenes derivatives, 5 and 6, respectively (Tables 1 and 2, NMR data). The products were named 2,7-di-*O*-acetylcasearupestrin A (5) and 2,6-di-*O*-acetylcasearupestrin D (6).

The results of the biological tests of compounds 1–6 are shown in Table 3. Casearupestrins A (1), B (2), and D (4) showed significant cytotoxicity against four cell lines (HL-60, HCT-8, MDA/MB-435, and SF-295), with IC₅₀ values ranging from 0.10 to 1.3 μ M. Compound 1 had an IC₅₀ value of 0.36 μ M against MDA/MB-435 and SF-295 cancer cell lines, superior to that of the standard drug doxorubicin (Table 3). Acetylation of compounds 1 and 4 resulted in decreased cytotoxicity (Table 3), indicating that the C-2 OH group is essential for the cytotoxicity of these compounds.

EXPERIMENTAL SECTION

General Experimental Procedures. Optical rotations were measured on a Perkin-Elmer 341 LC polarimeter. UV spectra were measured on a Hewlett-Packard 8453 A spectrometer. The circular dichroism spectra were obtained on a JASCO PU-2089 HPLC system configured with a UV–vis (MD-2010) and a circular dichroism (CD-2095) detector. IR spectra were measured on a Nicolet Impact 400 spectrometer, using KBr disks. The 1D (¹H, ¹³C, DEPT, and NOE)

Table 2. ^{13}C NMR Data for Compounds 1–6 (methanol- d_4 , 125 and 75 MHz, δ)^a

position	1	2	3	4	5	6
1	30.6, CH ₂	30.6, CH ₂	30.6, CH ₂	30.5, CH ₂	27.8, CH ₂	27.8, CH ₂
2	64.2, CH	64.2, CH	64.3, CH	64.0, CH	67.5, CH	67.4, CH
3	128.6, CH	127.0, CH	126.5, CH	129.3, CH	125.2, CH	125.3, CH
4	141.9, C	143.5, C	144.0, C	141.1, C	144.2, C	144.1, C
5	54.3, C	55.2, C	55.1, C	54.3, C	54.3, C	54.2, C
6	77.5, CH	75.2, CH	75.3, CH	75.8, CH	74.2, CH	75.6, CH
7	72.2, CH	76.1, CH	76.1, CH	73.5, CH	74.7, CH	73.4, CH
8	44.8, CH	43.1, CH	43.0, CH	42.7, CH	42.5, CH	42.5, CH
9	39.8, C	40.2, C	40.2, C	40.2, C	40.2, C	40.2, C
10	37.2, CH	36.5, CH	36.3, CH	37.3, CH	38.3, CH	38.3, CH
11	29.4, CH ₂	29.5, CH ₂	30.5, CH ₂	30.5, CH ₂	30.6, CH ₂	30.6, CH ₂
12	25.0, CH ₂	25.1, CH ₂	25.0, CH ₂	25.0, CH ₂	25.1, CH ₂	25.1, CH ₂
13	147.0, C	146.9, C	147.0, C	146.8, C	146.6, C	146.6, C
14	141.5, CH	141.6, CH	141.6, CH	141.5, CH	141.4, CH	141.4, CH
15	113.2, CH ₂	113.1, CH ₂	113.1, CH ₂	113.2, CH ₂	113.2, CH ₂	113.2, CH ₂
16	115.5, CH ₂	115.7, CH ₂	115.6, CH ₂	115.1, CH ₂	115.9, CH ₂	115.9, CH ₂
17	11.7, CH ₃	11.5, CH ₃	11.5, CH ₃	11.3, CH ₃	11.3, CH ₃	11.3, CH ₃
18	96.3, CH	97.2, CH	105.7, CH	96.0, CH	96.0, CH	95.9, CH
19	99.8, CH	100.1, CH	99.7, CH	99.5, CH	99.5, CH	99.5, CH
20	26.4, CH ₃	26.2, CH ₃	26.3, CH ₃	26.0, CH ₃	25.8, CH ₃	25.8, CH ₃
1'	167.3, C	168.0, C	168.0, C	167.4, C	166.8, C	167.4, C
2'	115.9, CH	116.3, CH	116.3, CH	115.7, CH	114.6, CH	115.1, CH
3'	147.7, CH	147.1, CH	147.1, CH	148.1, CH	148.9, CH	148.2, CH
4'	128.3, CH	128.3, CH	128.2, CH	128.0, CH	128.1, CH	128.0, CH
5'	147.2, CH	146.9, CH	146.8, CH	148.0, CH	148.5, CH	148.0, CH
6'	33.9, CH ₂	33.9, CH ₂	33.9, CH ₂	33.9, CH ₂	33.9, CH ₂	33.9, CH ₂
7'	30.5, CH ₂	30.6, CH ₂	29.5, CH ₂	29.4, CH ₂	29.3, CH ₂	29.4, CH ₂
8'	32.4, CH ₂	32.5, CH ₂	32.5, CH ₂	32.5, CH ₂	32.5, CH ₂	32.5, CH ₂
9'	23.4, CH ₂	23.5, CH ₂	23.5, CH ₂	23.5, CH ₂	23.4, CH ₂	23.5, CH ₂
10'	14.3, CH ₃	14.3, CH ₃	14.3, CH ₃	14.3, CH ₃	14.3, CH ₃	14.3, CH ₃
2-O ₂ CCH ₃					172.1, C 21.0, CH ₃	172.1, C 21.0, CH ₃
6-O ₂ CCH ₃				171.5, C 20.9, CH ₃	171.6, C 20.6, CH ₃	171.5, C 20.9, CH ₃
18-OCH ₃			55.4, CH ₃			
18-O ₂ CCH ₃	171.8, C 21.0, CH ₃	172.0, C 21.0, CH ₃		171.9, C 20.9, CH ₃	172.0, C 20.9, CH ₃	171.8, C 20.9, CH ₃
19-O ₂ CCH ₃	171.4, C 21.8, CH ₃	171.5, C 21.8, CH ₃	171.9, C 21.9, CH ₃	171.9, C 21.9, CH ₃	170.9, C 21.7, CH ₃	171.0, C 21.7, CH ₃

^a ^{13}C NMR data were assigned by DEPT-135° and gHMQC experiments; multiplicities were determined with the assistance of DEPT-135° and 90° experiments.

and 2D (^1H – ^1H gCOSY, ^1H – ^{13}C gHMQC, gHMBC, and gNOESY) NMR experiments were recorded on a Varian INOVA 500 spectrometer (11.7 T) at 500 MHz (^1H) and 125 MHz (^{13}C) with pulse field gradient and a Varian INOVA 300 spectrometer (7.4 T) at 300 MHz (^1H) and 75 MHz (^{13}C), using methanol- d_4 as the internal standard for ^{13}C NMR chemical shifts and residual undeuterated solvent (methanol) as the internal standard for ^1H NMR, with the δ scale reported relative to TMS. Positive-ion HRMS spectra were recorded on an UltratOFq (Bruker Daltonics) ESI-qTOF mass spectrometer, using TFANa as the internal standard. Analytical HPLC was performed on a Varian Pro Star 230 with UV–vis detector (model 330), using a Phenomenex C₁₈ column (250 mm \times 4.6 mm, 5 μm). Open column chromatography was performed over silica gel (40–63 μm , Merck) or silica C18 (40 μm , J. T. Baker) or on Sephadex LH-20 (Pharmacia Biotech). TLC was performed using Merck silica gel 60 (>230 mesh) and precoated silica gel 60 PF₂₅₄ plates. Spots on TLC plates were observed under UV light and by spraying the plates with anisaldehyde–H₂SO₄ reagent, followed by heating at 120 °C.

Plant Material. *Casearia rupestris* leaves were collected in May 2007 in Campinas municipality, São Paulo State, Brazil. The plant material was identified by one of the authors (R.B.T.). A voucher specimen (IAC41542) has been deposited in the herbarium of the Instituto Agrônomico de Campinas (São Paulo State).

Extraction, Isolation, and Chemical Transformation. The dried and ground leaves (0.46 kg) were exhaustively extracted by successive maceration with hexanes, EtOH, and H₂O (3.0 L \times 3) at room temperature. The EtOH extract was concentrated under reduced pressure at 40 °C to yield 30.0 g of a crude extract. This extract was then diluted with MeOH–H₂O (3:1) and successively partitioned with hexanes (1.0 L \times 3), Et₂O (1.0 L \times 3), and EtOAc (1.0 L \times 3). After removal of the solvent, each extract yielded 10.8, 3.2, and 1.2 g, respectively. Part of the cytotoxic Et₂O (3.0 g) fraction was subjected to gel permeation chromatography on Sephadex LH-20 (78.0 \times 3.0 cm) eluted with MeOH, to afford 38 fractions. These fractions were combined on the basis of their TLC profiles. Fraction 12 (1.7 g) was subjected to column chromatography (silica gel–C₁₈, 78.0 \times 3.0 cm,




Figure 2. Important gHMBC and gCOSY correlations observed for 1.




Figure 3. Important NOE enhancements observed for compounds 1–4.

Table 3. Cytotoxic Activity of Compounds 1–4 from the Leaves of *Casearia rupestris* and Derivatives 5 and 6 on Human Tumor Cancer Cell Lines after 72 h Exposure

compound	cell lines (IC ₅₀ , μ M) ^a			
	HL-60	MDA/MB-435	HCT-8	SF-295
1	0.10	0.36	0.13	0.36
2	0.85	1.3	0.28	0.80
3	>5	>5	>5	>5
4	0.25	0.93	0.25	0.43
5	4.3	>5	>5	>5
6	>5	>5	>5	>5
doxorubicin	0.04	0.83	0.02	0.40

^aData are presented as IC₅₀ values and 95% confidence intervals obtained by nonlinear regression for leukemia (HL-60), melanoma (MDA-MB-453), colon (HCT-8), and glioblastoma (SF-295) cells from two independent experiments. Doxorubicin was used as positive control.

40 μ m) using a step gradient from H₂O–MeOH 23:77 to 00:100, yielding 55 fractions. Fraction 14 (175.0 mg) was subjected to silica gel column chromatography (21.8 \times 2 cm, 230–400 mesh) eluted with hexanes–EtOAc (6:4), to yield compound 1 (79.0 mg). Fraction 30 (88.0 mg) was subjected to silica gel column chromatography (21.8 \times 2 cm, 40–63 μ m) eluted with hexanes–EtOAc (4:6) to yield compound 2 (13.0 mg). Fraction 34 (150.5 mg) was subjected to silica gel column chromatography (21.8 \times 2 cm, 40–63 μ m) eluted with hexanes–EtOAc (1:1) to yield compound 3 (10.0 mg). Finally, fraction 36 (356.0 mg) was subjected to

silica gel column chromatography (21.8 \times 2 cm, 40–63 μ m) eluted with hexanes–EtOAc (1:1) to yield compound 4 (212.0 mg). Compounds 1 and 4 (20 mg each) were dissolved in pyridine (4.0 mL) and treated with Ac₂O (3.0 mL) at room temperature for 24 h. The solvents were removed by evaporation under reduced pressure, resulting in compounds 5 (18.0 mg) and 6 (18.0 mg).

Casearupestrin A (1): white, amorphous solid; $[\alpha]_D^{24} +4.3$ (c 0.1, CHCl₃); UV λ_{max}^{MeOH} nm (log ϵ) 223 (1.40), 268 (1.50); CD (c 0.02, MeOH) $\Delta\epsilon$ (nm) +6.5 (223), +6.6 (268); IR (KBr) ν_{max} cm^{−1} 3434, 2959, 2932, 1746, 1712, 1378, 1230, 1028; ¹H and ¹³C NMR (methanol-d₄), see Tables 1 and 2, respectively; HRTOF-ESIMS m/z 623.3190 [M + Na]⁺ (calcd for C₃₄H₄₈O₉Na, 623.3190).

Casearupestrin B (2): white, amorphous solid; $[\alpha]_D^{24} +76$ (c 0.1, CHCl₃); UV λ_{max}^{MeOH} nm (log ϵ) 223 (1.48), 268 (1.55); CD (c 0.03, MeOH) $\Delta\epsilon$ (nm) +5.4 (223), +6.0 (268); IR ν_{max} cm^{−1} 3436, 1746, 1712, 1636, 1177, 1000, 900; ¹H and ¹³C NMR (methanol-d₄), see Tables 1 and 2, respectively; HRTOF-ESIMS m/z 623.3206 [M + Na]⁺ (calcd for C₃₄H₄₈O₉Na, 623.3190).

Casearupestrin C (3): white, amorphous solid; $[\alpha]_D^{20} +72$ (c 0.1, CHCl₃); UV λ_{max}^{MeOH} nm (log ϵ) 223 (1.43), 268 (1.48); CD (c 0.03, MeOH) $\Delta\epsilon$ (nm) +5.2 (223), +5.7 (268); IR ν_{max} cm^{−1} 3436, 1714, 1636, 1178, 1001, 897; ¹H and ¹³C NMR (methanol-d₄), see Tables 1 and 2, respectively; HRTOF-ESIMS m/z 595.3241 [M + Na]⁺ (calcd for C₃₃H₄₈O₈Na, 595.3241).

Casearupestrin D (4): white, amorphous solid; $[\alpha]_D^{25} +81$ (c 0.1, CHCl₃); UV λ_{max}^{MeOH} nm (log ϵ) 223 (1.70), 268 (1.80); CD (c 0.04, MeOH) $\Delta\epsilon$ (nm) +5.2 (223), +5.0 (268); IR ν_{max} cm^{−1} 3446, 1746, 1225, 1029, 905; ¹H and ¹³C NMR (methanol-d₄), see Tables 1 and 2, respectively; HRTOF-ESIMS m/z 665.3297 [M + Na]⁺ (calcd for C₃₆H₅₀O₁₀Na, 665.3296).

2,7-Di-O-acetylcasearupestrin A (5): colorless oil; $[\alpha]_D^{25} +56$ (c 0.1, MeOH); UV λ_{max}^{MeOH} nm (log ϵ) 223 (1.43), 268 (1.60); IR ν_{max} cm^{−1} 3454, 2953, 1752, 1374, 1227, 1026; ¹H and ¹³C NMR (methanol-d₄), see Tables 1 and 2, respectively; HRTOF-ESIMS m/z 707.3401 [M + Na]⁺ (calcd for C₃₈H₅₂O₁₁Na, 707.3401).

2,6-Di-O-acetylcasearupestrin D (6): colorless oil; $[\alpha]_D^{25} +99$ (c 0.1, MeOH); UV λ_{max}^{MeOH} nm (log ϵ) 223 (1.40), 268 (1.55); IR ν_{max} cm^{−1} 3455, 2953, 1750, 1374, 1227, 1026; ¹H and ¹³C NMR (methanol-d₄), see Tables 1 and 2, respectively; HRTOF-ESIMS m/z 707.3401 [M + Na]⁺ (calcd for C₃₈H₅₂O₁₁Na, 707.3401).

Cytotoxicity Assay. The cytotoxicity of the clerodane diterpenes (1–6) was tested against four tumor cell lines: SF-295 (human glioblastoma), HCT-8 (human colon carcinoma), HL-60 (human leukemia), and MDA/MB-435 (human melanoma). Cells were maintained in RPMI 1640 medium, supplemented with 10% fetal bovine serum, 2 mM glutamine, 100 mg/mL streptomycin, and 100 U/mL penicillin at 37 °C with 5% CO₂. Cells were seeded in 96-well plates (105 cells/well for adherent cells or 0.3 \times 10⁶ cells/well for suspended cells in 100 μ L of medium). After 24 h, the compounds (0.39–25 μ g/mL) dissolved in DMSO were added to each well. Doxorubicin (0.01–0.58 μ g/mL) was used as the positive control. The proliferation of tumor cells was quantified by the ability of living cells to reduce the yellow dye 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) to a blue formazan product. At the end of 69 h incubation, the medium was replaced by fresh medium containing 0.5 mg/mL of MTT. Three hours later, the formazan product was dissolved in DMSO, and absorbance was measured using a multiplate reader at 595 nm (DTX-880, Beckman Coulter). The IC₅₀ values were obtained by nonlinear regression using the GRAPHPAD program (Intuitive Software for Science, San Diego, CA).³⁷

■ ASSOCIATED CONTENT

Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Tel: + 55 16 33016667. Fax: 55 16 33227932. E-mail: cavaleiro_alberto@yahoo.com.br.

■ ACKNOWLEDGMENT

The authors are grateful to the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and the Biota Instituto Virtual da Biodiversidade for financial support of Project No. 03/02176-7. We are grateful to Prof. Dr. W. Vilegas for the use of the polarimeter, UV, IR, and HPLC with circular dichroism detector. Finally, we thank Prof. Dr. N. Peoporine Lopes for the use of the LC-MS.

■ REFERENCES

- (1) Shen, Y.-C.; Wang, C.-H.; Cheng, Y.-B.; Wang, L.-T.; Guh, J.-H.; Chien, C.-T.; Khalil, A. T. *J. Nat. Prod.* **2004**, *67*, 316–321.
- (2) Carvalho, P. R. F.; Furlan, M.; Young, M. C. M.; Kingston, D. G. I.; Bolzani, V. D. S. *Phytochemistry* **1998**, *49*, 1659–1662.
- (3) Khan, M. R.; Gray, A. I.; Reed, D. R.; Sadler, I. H.; Waterman, P. G. *Phytochemistry* **1990**, *29*, 3591–3595.
- (4) Chen, T. B.; Wiemer, D. F. *J. Nat. Prod.* **1991**, *54*, 1612–1618.
- (5) Gibbons, S.; Gray, A.; Waterman, P. G. *Phytochemistry* **1996**, *41*, 565–570.
- (6) Guittet, E.; Stoven, V.; Lallemand, J.; Ramiandrasoa, F.; Kunesch, G. *Tetrahedron* **1988**, *44*, 2893–2901.
- (7) Mosaddik, M. A.; Waterman, P. G. *A. Nat. Prod. Commun.* **2006**, *1*, 601–607.
- (8) Mosaddik, M. A.; Forster, P. I.; Booth, R.; Waterman, P. G. *A. Nat. Prod. Commun.* **2006**, *1*, 441–448.
- (9) Mosaddik, M. A.; Forster, P. I.; Booth, R.; Waterman, P. G. *A. Biochem. Syst. Ecol.* **2007**, *35*, 631–633.
- (10) Santos, A. G.; Perez, C. C.; Tininins, A. G.; Bolzani, V. S.; Cavaleiro, A. J. *Quím. Nova* **2007**, *30*, 1100–1103.
- (11) Santos, A. G.; Ferreira, P. M. P.; Vieira-Júnior, G. M.; Perez, C. C.; Tininins, A. G.; Silva, G. H.; Bolzani, V. S.; Costa-Lotufo, L. V.; Pessoa, C. Ó.; Cavaleiro, A. J. *Chem. Biodiversity* **2010**, *7*, 205–215.
- (12) Shen, Y.-C.; Cheng, Y.-B.; Chen, Y.-H.; Khalil, A. T.; Ko, C.-L. *J. Chin. Chem. Soc.* **2005**, *52*, 1263–1268.
- (13) Vijayakumar, E. K. S.; Bal-Tembe, S.; Joshi, K. S.; Deore, V. B. *Indian J. Chem., Sec. B: Org. Chem. Incl. Med. Chem.* **2002**, *41*, 2706–2708.
- (14) Oberlies, N. H.; Burgess, J. P.; Navarro, H. A.; Pinos, R. E.; Fairchild, C. R.; Peterson, R. W.; Soejarto, D. D.; Farnsworth, N. R.; Kinghorn, A. D.; Wani, M. C.; Wall, M. E. *J. Nat. Prod.* **2002**, *65*, 95–99.
- (15) Jullian, V.; Bonduelle, C.; Valentin, A.; Acebey, L.; Duigou, A.-G.; Prévost, M.-F.; Sauvain, M. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5065–5070.
- (16) Kanokmedhakul, S.; Kanokmedhakul, K.; Kanarsa, T.; Buayairaksa, M. *J. Nat. Prod.* **2005**, *68*, 183–188.
- (17) Itokawa, H.; Totsuka, N.; Morita, H.; Takeya, K.; Watanabe, K.; Obata, E. *Chem. Pharm. Bull.* **1988**, *36*, 1585–1588.
- (18) Itokawa, H.; Totsuka, N.; Morita, H.; Takeya, K.; Itaka, Y.; Schenkel, E. P.; Motidome, M. *Chem. Pharm. Bull.* **1990**, *38*, 3384–3388.
- (19) Beutler, J. A.; MacCall, K. L.; Herbert, K.; Herald, D. L.; Pettit, G. R.; Johnson, T.; Shoemaker, R. H.; Boyd, M. R. *J. Nat. Prod.* **2000**, *63*, 657–661.
- (20) Beutler, J. A.; MacCall, K. L.; Herbert, K.; Johnson, T.; Shoemaker, R. H.; Boyd, M. R. *Phytochemistry* **2000**, *55*, 233–236.
- (21) Chen, C.-Y.; Cheng, Y.-B.; Chen, S.-Y.; Chien, C.-T.; Kuo, Y.-H.; Guh, J.-H.; Khalil, A. T.; Shen, Y.-C. *Chem. Biodiversity* **2008**, *5*, 162–167.
- (22) Kanokmedhakul, S.; Kanokmedhakul, K.; Buayairaksa, M. *J. Nat. Prod.* **2007**, *70*, 1122–1126.
- (23) Hayashi, K.-I.; Nakanishi, Y.; Bastow, K. F.; Cragg, G.; Nozaki, H.; Lee, K.-H. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 345–348.
- (24) Oberlies, N. H.; Burgess, J. P.; Navarro, H. A.; Pinos, R. E.; Soejarto, D. D.; Farnsworth, N. R.; Kinghorn, A. D.; Wani, M. C.; Wall, M. E. *J. Nat. Prod.* **2001**, *64*, 497–501.
- (25) Prakash, C. V. S.; Hoch, J. M.; Kingston, D. G. I. *J. Nat. Prod.* **2002**, *65*, 100–107.
- (26) Shen, Y.-C.; Wang, L.-T.; Wang, C.-H.; Khalil, A. T.; Guh, J.-H. *Chem. Pharm. Bull.* **2004**, *52*, 108–110.
- (27) Shen, Y.-C.; Cheng, Y.-B.; Ahmed, A. F.; Lee, C. L.; Chen, S.-Y.; Chien, C.-T.; Kuo, Y.-H.; Tzeng, G.-L. *J. Nat. Prod.* **2005**, *68*, 1665–1668.
- (28) Shen, Y.-C.; Lee, C. L.; Khalil, A. T.; Cheng, Y.-B.; Chien, C.-T.; Kuo, Y.-H. *Helv. Chim. Acta* **2005**, *88*, 68–77.
- (29) Vieira-Júnior, G. M.; Gonçalves, T. O.; Regasini, L. O.; Ferreira, P. M. P.; Pessoa, C. Ó.; Costa-Lotufo, L. V.; Torres, R. B.; Boralle, N. B.; Bolzani, V. S.; Cavaleiro, A. J. *J. Nat. Prod.* **2009**, *72*, 1847–1850.
- (30) Williams, R. B.; Norris, A.; Miller, J. S.; Birkinshaw, C.; Ratovoson, F.; Andriantsiferana, R.; Rasamison, V. E.; Kingston, D. G. I. *J. Nat. Prod.* **2007**, *70*, 206–209.
- (31) Ferreira, P. M. P.; Santos, A. G.; Tininins, A. G.; Costa, P. M.; Cavaleiro, A. J.; Bolzani, V. S.; Moraes, M. O.; Costa-Lotufo, L. V.; Montenegro, R. C.; Pessoa, C. Ó. *Chem.-Biol. Interact.* **2010**, *188*, 497–504.
- (32) Hunter, M. S.; Corley, D. G.; Carron, C. P.; Rowold, E.; Kilpatrick, B. F.; Durley, R. C. *J. Nat. Prod.* **1997**, *60*, 894–899.
- (33) Espindola, L. S.; Vasconcelos Júnior, J. R.; Mesquita, M. L.; Marquié, P.; Paula, J. E.; Mambu, L.; Santana, J. M. *Planta Med.* **2004**, *70*, 1093–1095.
- (34) Sleumer, H. O. *Flora Neotropica*; The New York Botanical Garden: New York, 1980; pp 1–499.
- (35) Morita, H.; Nakayama, M.; Kojima, H.; Takeya, K.; Itokawa, H.; Schenkel, E. P.; Motidome, M. *Chem. Pharm. Bull.* **1991**, *39*, 693–697.
- (36) Whitson, E. L.; Thomas, C. L.; Henrich, C. J.; Sayers, T. J.; McMahon, J. B.; McKee, T. C. *J. Nat. Prod.* **2010**, *73*, 2013–2018.
- (37) Mosmann, T. *J. Immunol. Methods* **1983**, *16*, 55–63.