

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231717394>

Bonducellpins A–D, New Cassane Furanoditerpenes of *Caesalpinia bonduc*

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · DECEMBER 1997

Impact Factor: 3.8 · DOI: 10.1021/np970308c

CITATIONS

32

READS

57

2 AUTHORS:



Sonia Peter

Barbados Community College

13 PUBLICATIONS 118 CITATIONS

SEE PROFILE



Winston Tinto

University of the West Indies at Cave Hill, Bar...

76 PUBLICATIONS 970 CITATIONS

SEE PROFILE

JOURNAL OF NATURAL PRODUCTS

© Copyright 1997 by the American Chemical Society and the American Society of Pharmacognosy

Volume 60, Number 12

December 1997

Full Papers

Bonducellpins A–D, New Cassane Furanoditerpenes of *Caesalpinia bonduc*

Sonia R. Peter and Winston F. Tinto*

Laboratory of Bioorganic Chemistry, Department of Biological and Chemical Sciences, University of the West Indies, Cave Hill Campus, Barbados

Stewart McLean, William F. Reynolds,* and Margaret Yu

Department of Chemistry, University of Toronto, Toronto, Ontario M5S 1A1, Canada

Received June 19, 1997[®]

Four cassane furanoditerpenes, designated bonducellpins A (**1**), B (**2**), C (**3**), and D (**4**), were isolated from the roots of *Caesalpinia bonduc*. The ¹H- and ¹³C-NMR spectra of all four compounds were completely assigned by using a combination of 2D NMR experiments, which included COSY, HMQC, HMBC, and NOESY sequences.

Plants belonging to the genus *Caesalpinia* have proven to be a rich source of cassane furanoditerpenes, some of which display interesting biological activity.^{1–4} *Caesalpinia bonduc* (L.) Roxb. (Fabaceae, subfamily Caesalpinioideae, tribe Caesalpinieae) is also known as *C. bonducella* and is widely distributed throughout the tropics and subtropics.^{5–7} The taxonomy of the family Fabaceae (previously Leguminosae) has been the subject of much debate, and plants of the genus *Caesalpinia* are sometimes referred to in the literature as belonging to the family Caesalpinaceae.⁷ *C. bonduc* has been the subject of several chemical investigations, wherein a number of cassane furanoditerpenes have been isolated.^{4,8–17} We have investigated the roots of *C. bonduc*, collected in Barbados, and report here the isolation of four new cassane furanoditerpenes, bonducellpins A (**1**), B (**2**), C (**3**), and D (**4**). The proton and carbon assignments as well as the relative stereochemistry of all four compounds were determined by 2D NMR spectroscopy.

Results and Discussion

Bonducellpin A (**1**) was isolated as white crystals, mp 118–119 °C, and had the molecular formula C₂₅H₃₄O₉.

The IR spectrum exhibited absorptions typical of hydroxyl (3447 cm⁻¹) and ester (1735 cm⁻¹) functionalities. The ¹H-NMR spectrum had oxymethine resonances associated with acetoxyl groups at δ 4.87 (t, J = 3.0 Hz, H-1) and δ 5.36 (dd, J = 9.8, 2.3 Hz, H-6) and a secondary hydroxyl at δ 3.92 (d, J = 9.8 Hz, H-7). A 1,2-disubstituted furan was evident from lowfield doublets at δ 7.25 (J = 2.5 Hz, H- α) and δ 6.17 (J = 2.5 Hz, H- β), while a methoxycarbonyl group had a sharp singlet at δ 3.73. The carbomethoxyl group was located at C-17 because an HMBC correlation was observed between its carbonyl at δ 173.4 and a proton at δ 3.49, attributable to H-14. HMBC correlations were also observed between H-14 and C-7, C-8, C-12, and C-13. The COSY spectrum established the spin system involving H-6, H-7, H-8, H-9, H₂-11, and H-14. The relative stereochemistry of **1** was determined by interpretation of the results of a NOESY experiment (Figure 1). In particular, H-14 had cross peaks with H-7 and H-9, which indicated that they were α -oriented, while the stereochemistry of H-1 followed from its cross peak with H₃-10 and from its vicinal couplings to the C-2 protons. These results are summarized in Tables 1 and 2 and led to the structural assignment of bonducellpin A (**1**).

Bonducellpin B (**2**), C₂₃H₃₀O₈, had IR absorbances due

[®] Abstract published in *Advance ACS Abstracts*, October 15, 1997.




Figure 1. Major NOESY correlations for compound **1**.

to hydroxyl (3392 cm^{-1}), ester (1740 cm^{-1}), and ketone (1720 cm^{-1}) functionalities. The ^1H - and ^{13}C -NMR spectra of **2** were similar to those of **1** except for the disappearance of the oxymethine proton at C-1 and its replacement with a ketone having a ^{13}C resonance at δ 211.8 (Tables 1 and 2). The location of the ketone at C-1 was confirmed because it showed HMBC correlations to H₂-2, H₂-3, H-9, and H₃-20. Bonducellpin B (**2**) is, therefore, the 1-keto analogue of **1**.

Bonducellpin C (**3**), $\text{C}_{23}\text{H}_{32}\text{O}_7$, was isolated as white crystals, mp $110\text{--}113^\circ\text{C}$. The IR spectrum had absorbances characteristic of hydroxyl (3401 cm^{-1}) and ester (1736 cm^{-1}) groups. The ^1H -NMR spectrum had resonances assignable to an acetoxy methine at δ 4.90 (br s), a secondary hydroxyl at δ 4.00 (m), and a methoxycarbonyl at δ 3.74. The acetoxy group was located at C-1 as the proton at δ 4.90 showed HMBC cross peaks to C-2, C-3, C-5, C-10, C-20, and the acetate carbonyl at δ 169.0. A signal at δ 3.47 (d, $J = 8.8\text{ Hz}$) was assigned to H-14, and it showed HMBC correlations to the methoxycarbonyl at δ 176.0, in addition to C-7, C-8, C-12, and C-13. When **3** was acetylated, the oxymethine resonance at δ 4.00 shifted downfield to δ 5.22 (ddd, $J = 10.2, 10.2, 5.7$). The foregoing evidence indicated that **3** was similar to **1** except that the acetoxy group at C-6 was replaced by a methylene group. The stereochemistry of all the chiral centers in **3** were identical to those in **1** for they both had similar coupling constants at these positions (Tables 1 and 2), and this was confirmed by interpretation of a NOESY spectrum of **3**.

Bonducellpin D (**4**), mp $215\text{--}217^\circ\text{C}$, had the molecular formula, $\text{C}_{22}\text{H}_{28}\text{O}_7$. The IR spectrum had absorbances characteristic of hydroxyl (3392 cm^{-1}), γ -lactone (1797 cm^{-1}), and ester (1735 cm^{-1}) functionalities. The ^1H -NMR spectrum had resonances due to oxymethine protons at δ 5.58 (d, $J = 9.8$, H-6), δ 4.74 (dd, $J = 13.1, 9.8$, H-7), and δ 3.70 (m, H-1). The HMQC spectrum revealed that H-7 was directly attached to a carbon at δ 82.7. The downfield nature of C-7 and the lack of a resonance due to a methoxycarbonyl group indicated that the γ -lactone was formed between the oxygen at C-7 and the C-17 carbonyl. Bonducellpins A–D (**1–4**) represent the first examples of cassane furanoditerpenes bearing a C-17 ester from the genus *Caesalpinia*; however, cassane furanoditerpenes with a C-17 carboxylic acid or ester were previously isolated from plants of the genus *Pterodon*, which is also a member of the Fabaceae (subfamily Papilionoideae, tribe Dipterygeae).^{18,19}

Experimental Section

General Experimental Procedures. Melting points were determined using a Koffler hotstage and are

uncorrected. The IR spectra were recorded on a Perkin-Elmer 1725X FT-IR spectrometer. UV spectra were obtained on a Hewlett-Packard 8452A spectrophotometer in MeOH. Optical rotations were recorded on a Perkin-Elmer 341 polarimeter in CHCl_3 solutions. All NMR spectra were obtained on a Varian UNITY 500 MHz spectrometer, in CDCl_3 using TMS as an internal standard.

Plant Material. The roots of *Caesalpinia bonduc* were collected in St. Andrew, Barbados, in February 1995. The plant was identified by Dr. Sean Carrington, Biological and Chemical Sciences Department, University of the West Indies, where a voucher specimen (No. SC1785) is kept.

Extraction and Isolation. The dried, ground roots (1.7 kg) were extracted with 95% EtOH (9.6 L) and the solvent evaporated *in vacuo* to give a brown syrup (180 g). The extract was dissolved in 10% aqueous MeOH (500 mL) and extracted with light petroleum ($6 \times 300\text{ mL}$). The aqueous MeOH layer was diluted with H_2O (200 mL) and extracted with CH_2Cl_2 ($6 \times 300\text{ mL}$), dried over anhydrous Na_2SO_4 , and the solvent evaporated to give a brown gum (24 g).

The CH_2Cl_2 extract was flash chromatographed over Si gel with light petroleum– Me_2CO (3:1) followed by reversed-phase preparative HPLC using MeOH– H_2O (75:25), to give compounds **1** (1.7 mg), **2** (12.8 mg), **3** (29.9 mg), and **4** (4.2 mg).

Bonducellpin A (1): colorless crystals; mp $118\text{--}119^\circ\text{C}$; $[\alpha]_D +4.6^\circ$ (c 0.33, CHCl_3); IR (CHCl_3) ν_{max} 3447, 1735 cm^{-1} ; UV (MeOH) λ_{max} (log ϵ) 226 (3.76) nm; EIMS m/z [M]⁺ 478 (12), 446 (2), 418 (10), 400 (9), 386 (23), 358 (18), 340 (56), 281 (100), 263 (36), 243 (44), 145 (21), 107 (25); HREIMS 478.2206 calcd for $\text{C}_{25}\text{H}_{34}\text{O}_9$ 478.2203; ^1H - and ^{13}C -NMR data, see Tables 1 and 2, respectively.

Bonducellpin B (2): colorless gum; $[\alpha]_D +18.7^\circ$ (c 0.14, CHCl_3); IR (CHCl_3) ν_{max} 3392, 1735, 1720 cm^{-1} ; UV (MeOH) λ_{max} (log ϵ) 218, 254 (3.74, 3.29) nm; EIMS m/z [M]⁺ 434 (29), 416 (2), 402 (7), 374 (35), 356 (26), 314 (52), 297 (39), 257 (28), 199 (45), 149 (100), 109 (41); HREIMS 434.1953 calcd for $\text{C}_{23}\text{H}_{30}\text{O}_8$ 434.1941; ^1H - and ^{13}C -NMR data, see Tables 1 and 2, respectively.

Bonducellpin C (3): colorless crystals; mp $112\text{--}113^\circ\text{C}$; $[\alpha]_D +12.6^\circ$ (c 0.78, CHCl_3); IR (CHCl_3) ν_{max} 3401, 1736 cm^{-1} ; UV (MeOH) λ_{max} (log ϵ) 216 (3.81) nm; EIMS m/z [M]⁺ 420 (15), 402 (3), 360 (9), 342 (53), 283 (27), 265 (100), 227 (22), 209 (31), 195 (51), 177 (42), 121 (42); HREIMS 420.2155 calcd for $\text{C}_{23}\text{H}_{32}\text{O}_7$ 420.2148; ^1H - and ^{13}C -NMR data, see Tables 1 and 2, respectively.


Acetate of 3. Acetylation of **3** (5.6 mg) with Ac_2O and pyridine (1:1) gave the acetate **3a** as a colorless gum (4.2 mg): $[\alpha]_D +26.5^\circ$ (c 0.05, CHCl_3); IR (CHCl_3) ν_{max} 3401, 1736 cm^{-1} ; UV (MeOH) λ_{max} (log ϵ) 216, 256 (3.80, 3.14) nm; ^1H NMR (CDCl_3 , 500 MHz) δ 7.24 (d, $J = 2.6\text{ Hz}$, H-16), 6.12 (d, $J = 2.6\text{ Hz}$, H-15), 5.22 (ddd, $J = 10.2, 10.2, 5.7\text{ Hz}$, H-7), 4.92 (br t, $J = 3.4\text{ Hz}$, H-1), 3.73 (s, OCH_3), 3.39 (br d, $J = 8.2\text{ Hz}$, H-14), 2.69 (ddd, $J = 12.1, 12.1, 5.6\text{ Hz}$, H-9), 2.52 (dd, $J = 16.2, 12.1\text{ Hz}$, H-11 β), 2.30 (dd, $J = 16.2, 5.6\text{ Hz}$, H-11 α), 2.17 (dd, $J = 12.7, 5.7\text{ Hz}$, H-6), 2.09 (s, 1- CH_3CO), 2.00 (s, 7- CH_3CO), 1.98 (m, H-2), 1.78 (m, H-2), 1.75 (m, H-3), 1.65 (ddd, $J = 12.7, 11.0, 3.0\text{ Hz}$, H-6), 1.22 (s, H-20), 1.15 (m, H-3), 1.08 (s, H-19), 1.04 (s, H-18); ^{13}C NMR (CDCl_3 , 125 MHz) δ 174.6 (C-17), 170.1 (7- CH_3CO), 169.0 (1- CH_3CO), 149.8 (C-12), 141.5 (C-16), 113.3 (C-

Table 1. ^1H -NMR Assignments for Bonducellpins A–D (1–4)^a

H	1	2	3	4
1	4.87 (t, 3)		4.90 (br s)	3.70 (m)
2	1.78 (m)	2.44 (m)	1.78 (m)	1.66 (m)
	1.95 (m)	2.65 (m)	1.99 (m)	2.06 (m)
3	1.12 (m)	1.78 (m)	1.15 (m)	1.11 (m)
	1.75 (m)	1.92 (m)	1.74 (m)	2.07 (m)
6	5.36 (dd, 9.8, 2.3)	5.31 (d, 10)	1.65 (ddd, 13.6, 11.3, 2.8)	5.58 (d, 9.8)
			2.02 (m)	
7	3.92 (dd, 14.7, 9.8)	3.75 (m)	4.00 (m)	4.74 (dd, 13.1, 9.8)
8	2.40 (m)	2.35 (m)	2.23 (m)	2.15 (m)
9	2.65 (ddd, 12, 12, 5)	2.67 (ddd, 11, 11, 5.3)	2.61 (ddd, 12.8, 12.8, 6.4)	3.19 (ddd, 13.7, 8.9, 7)
11 α	2.30 (dd, 15.5, 6)	3.38 (dd, 16, 5.3)	2.50 (ddd, 16, 6.4, 3.2)	2.79 (dd, 16.3, 7)
11 β	2.48 (dd, 15.5, 12)	2.45 (dd, 16, 11)	2.26 (dd, 16, 12.8)	2.57 (dd, 16.3, 8.9)
14	3.49 (br d, 8.5)	3.51 (br d 8)	3.47 (d, 8.8)	3.31 (ddd, 12.8, 1.4, 1.4)
15	6.17 (d, 2.5)	6.16 (d, 2)	6.17 (d, 2.5)	6.60 (d, 2.5)
16	7.25 (d, 2.5)	7.23 (d, 2)	7.24 (d, 2.5)	7.31 (d, 2.5)
18	1.14 (s)	1.18 (s)	1.05 (s)	1.12 (s)
19	1.15 (s)	1.27 (s)	1.09 (s)	1.16 (s)
20	1.26 (s)	1.50 (s)	1.19 (s)	1.13 (s)
1-Ac	2.10 (s)		2.10 (s)	
6-Ac	2.16 (s)	2.16 (s)		2.15 (s)
OMe	3.73 (s)	3.73 (s)	3.74 (s)	

^a Chemical shifts (δ) in ppm (mult., J in Hz).**Table 2.** ^{13}C NMR Assignments for Bonducellpins A–D (1–4)^a

C	1	2	3	4
1	75.6	211.8	75.5	72.6
2	22.2	35.2	22.5	25.7
3	32.2	38.6	30.0	32.1
4	38.6	38.1	38.4	39.5
5	79.5	82.1	78.5	84.4
6	76.1	76.6	36.2	72.9
7	76.5	76.4	73.4	82.7
8	41.6	41.8	42.4	44.5
9	35.9	37.4	36.5	32.3
10	45.1	55.7	43.6	47.6
11	21.6	24.1	21.5	21.2
12	149.6	151.2	150.0	151.9
13	113.6	112.2	113.6	113.8
14	46.2	45.8	46.4	41.6
15	108.4	108.3	108.5	107.8
16	141.6	141.2	141.4	141.7
17	175.3	175.6	176.0	173.4
18	30.6	28.6	28.0	30.4
19	24.6	26.6	25.0	24.2
20	17.1	15.5	17.7	16.9
1-Ac	168.9		169.0	
	21.4		21.5	
6-Ac	171.6	170.9		169.6
	21.8	21.6		21.7

^a Chemical shift (δ) in ppm.(1) X = H, α -OAc, R¹ = OAc, R² = H(2) X = O, R¹ = OAc, R² = H(3) X = H, α -OAc, R¹ = H, R² = H(3a) X = H, α -OAc, R¹ = H, R² = Ac

13), 108.3 (C-15), 78.3 (C-5), 76.0 (C-7), 75.4 (C-1), 51.9 (OCH₃), 45.9 (C-14), 43.5 (C-10), 38.9 (C-8), 38.4 (C-4), 36.6 (C-9), 32.0 (C-6), 29.9 (C-3), 27.9 (C-18), 24.9 (C-19), 22.4 (C-2), 21.4 (1-CH₃CO), 21.3 (C-11), 21.1 (7-CH₃CO), 17.6 (C-10); EIMS m/z [M]⁺ 462 (5), 402 (34), 370 (72), 342 (41), 309 (69), 286 (38), 265 (100), 209 (37), 195 (56), 145 (57), 109 (23); HREIMS 462.2251 calcd for C₂₅H₃₄O₈ 462.2254.

Bonducellpin D (4): colorless crystals; mp 216–217

°C, [α]_D +8.4° (c 0.35 CHCl₃) IR (CHCl₃) ν_{max} 3392, 1797, 1735 cm⁻¹; UV (MeOH) λ_{max} (log ϵ) 218, 260 (3.84, 3.21) nm; EIMS m/z [M]⁺ 404 (25), 386 (64), 371 (3), 335 (46), 307 (100), 262 (11), 249 (17), 217 (41), 188 (50), 145 (30), 121 (53); HREIMS 404.1850 calcd for C₂₂H₂₈O₇ 404.1835; ^1H - and ^{13}C -NMR data, see Tables 1 and 2, respectively.

Acknowledgment. Research in Toronto was funded by grants from the Natural Sciences and Engineering Research Council of Canada. One of us (S. P.) acknowledges the award of a University of the West Indies Postgraduate Scholarship.

References and Notes

- Ogawa, K.; Aoki, I.; Sashida, Y. *Phytochemistry* **1992**, *31*, 2897–2898.
- Kitagawa, I.; Simanjuntak, P.; Watano, T.; Shibuya, H.; Fujii, S.; Yamagata, Y.; Kobayashi, M. *Chem. Pharm. Bull.* **1994**, *42*, 1798–1802.
- Kitagawa, I.; Simanjuntak, P.; Mahmud, T.; Kobayashi, M.; Fujii, S.; Uji, T.; Shibuya, H. *Chem. Pharm. Bull.* **1996**, *44*, 1157–1161.
- Purushothaman, K. K.; Kalyani, K.; Subramanian, K. *Indian J. Chem.* **1981**, *20*, 625–626.
- Morton, J. F. *Atlas of Medicinal Plants of Middle America*; Bannerstone House: 1981; p 279.
- Adams, C. D. *Flowering Plants of Jamaica*; The University Press: Glasgow, 1972; p 320.
- Carrington, S. *Wild Plants of Barbados*; Macmillan Press: London, 1993; p 37.
- Ali, M. E.; Khuda, M. Q. *Chem. Ind. (London)* **1960**, 463–464.
- Khuda, M. Q.; Ali, M. E. *Pakistan J. Sci. Ind. Research* **1963**, 65–66.
- Canonica, L.; Jommi, G.; Manitto, P.; Pagoni, U. M.; Pelizzoni, F. *Gazz. Chim. Ital.* **1966**, 662–686.
- Canonica, L.; Jommi, G.; Manitto, P.; Pagoni, U. M.; Pelizzoni, F. *Gazz. Chim. Ital.* **1966**, 687–697.
- Canonica, L.; Jommi, G.; Manitto, P.; Pagoni, U. M.; Pelizzoni, F. *Gazz. Chim. Ital.* **1966**, 698–715.
- Balmain, A.; Connolly, J. D.; Ferrari, M.; Ghisalberti, E. L.; Pagnoni, U. M.; Pelizzoni, F. *Chem. Commun.* **1970**, 1244–1245.
- Connolly, J. D.; Orsini, F.; Pelizzoni, F.; Ricca, G. *Magn. Reson. Chem.* **1981**, *17*, 163–165.
- Balmain, A.; Bjämer, K.; Connolly, J. D.; Ferguson, G. *Tetrahedron Lett.* **1967**, 5027–5031.
- Pascoe, K. O. "Diterpenes from Jamaican Caesalpinaceae and Euphorbiaceae," Ph.D. Thesis, University of the West Indies, Mona, Jamaica, 1980; pp 9–10.
- Pascoe, K. O.; Burke, B. A.; Chan, W. R. *J. Nat. Prod.* **1986**, *49*, 913–915.
- Campos, A. M.; Silveira, E. R.; Braz-Filho, R.; Teixeira, T. C. *Phytochemistry* **1994**, *36*, 403–406.
- Demuner, A. J.; Barbosa, L. C. de A.; Veloso, D. P.; Alves, D. L. F.; Howarth, O. W. *J. Nat. Prod.* **1996**, *59*, 770–772.

NP970308C