

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231714429>

Synthesis of (+)-Cyclozonarone and the Absolute Configuration of Naturally Occurring (–)-Cyclozonarone

ARTICLE in JOURNAL OF NATURAL PRODUCTS · FEBRUARY 2001

Impact Factor: 3.8 · DOI: 10.1021/np0004146

CITATIONS

17

READS

3

5 AUTHORS, INCLUDING:



Jaime A Valderrama

Pontifical Catholic University of Chile

143 PUBLICATIONS 1,236 CITATIONS

SEE PROFILE



Mauricio Cuellar

Universidad de Valparaíso (Chile)

40 PUBLICATIONS 179 CITATIONS

SEE PROFILE



Veronica Armstrong

Pontifical Catholic University of Chile

18 PUBLICATIONS 210 CITATIONS

SEE PROFILE



Marcelo Preite

Pontifical Catholic University of Chile

33 PUBLICATIONS 168 CITATIONS

SEE PROFILE

Synthesis of (+)-Cyclozonarone and the Absolute Configuration of Naturally Occurring (–)-Cyclozonarone

Manuel Cortés,* Jaime A. Valderrama, Mauricio Cuellar, Verónica Armstrong, and Marcelo Preite

Facultad de Química, Pontificia Universidad Católica de Chile, Casilla 306, Correo 22, Santiago, Chile

Received August 23, 2000

Synthesis of (+)-cyclozonarone (**1**) has been achieved using (–)-polygodial (**3**) as chiral starting material. The absolute configuration of naturally occurring (–)-cyclozonarone was established as 5*R*,10*R* by comparison of spectral data and optical rotation with those of (+)-cyclozonarone.


Marine flora and fauna have proven to be a source of compounds showing a great variety of chemical structures and a wide range of biological activities.¹ Substances of mixed biogenesis that are based on farnesyl hydroquinone are found in brown algae of the genus *Dictyopteris*.²

Cyclozonarone (**1**) is a sesquiterpene benzoquinone derivative, which was isolated from the brown algae *Dictyopteris undulata* (family Dictyotaceae) and reported as a potent feeding-deterrent toward young abalones.³ The structure of the natural product was established by spectroscopic evidence. However, the absolute configuration remained unknown, since it was only suggested, based on biogenetic reasons.³ To establish the absolute configuration of (–)-cyclozonarone and as part of a program aimed at syntheses of drimane–quinone derivatives,⁴ in this work we describe the first enantiospecific synthesis of compound **1**.

Our synthetic strategy for the preparation of **1** was based on the Diels–Alder reaction of diene **2** with *p*-benzoquinone (Scheme 1). Since the previously reported synthesis of **2** via manool⁵ was not possible, because manool is commercially unavailable, alternative methods were used to prepare **2** starting from natural (–)-polygodial (**3**). Compound **3** was converted to the known bicyclic intermediates **4** and **5** according to our previously published procedure.⁶

Reaction of enone **5** with Tebbe reagent⁷ gave **2** in poor yield (16%; 23.4% based on recovered starting material). Alternatively, diene **2** was prepared from unsaturated diol **4** (Scheme 1). Catalytic hydrogenation of **4** afforded diol **6** (82% yield). Saturated diol **6** had been previously prepared from (–)-sclareol.⁸ Compound **6** was dimesylated to give **7** (93%). Double elimination of mesylic acid with potassium *tert*-butoxide in DMF afforded diene **2** in good yield (45%). Diene **2** was submitted to Diels–Alder reaction with *p*-benzoquinone in refluxing benzene. The initial adduct expected (**8**) could not be detected due to enolization–oxidation⁹ to quinone **9** (49% yield), which occurred during the workup and purification process. Together with **9**, minor amounts of **1** and **10** were detected in the mixture by ¹H NMR. The mixture of **1**, **9**, and **10** was subjected to DDQ oxidation in benzene under reflux conditions to afford (+)-cyclozonarone (**1**) (94% yield).

Compound **1** displays spectroscopic data identical to those of the natural (–)-cyclozonarone except that the opposite sign for optical rotation was observed ($[\alpha]_D^{16} +93.2$ (*c* 1.4, CHCl₃); lit.³ $[\alpha]_D^{19} -89.1$ (*c* 0.330, CHCl₃)). Thus, the absolute configuration of natural cyclozonarone is 5*R*,10*R*.




Experimental Section

General Experimental Procedures. Melting points were measured on a Stuart-Scientific SMP3 apparatus and are uncorrected. Optical rotations were obtained for CHCl₃ solutions on a AA-5 automatic polarimeter, and their concentrations are expressed in g/100 mL. NMR spectra were recorded on a Bruker AM-200 and a Bruker Avance DRX-300 spectrometer. Chemical shifts are expressed in ppm downfield relative to TMS (δ scale) in CDCl₃. Carbon multiplicity was established by a DEPT pulse sequence. Elemental analyses were obtained on a Fisons-Carlo-Erba FA-1108 Automost microanalyzer. For analytical TLC, Merck silica gel 60 in a 0.25 mm layer was used. Chromatographic separations were carried out by a conventional column on Merck silica gel 60 (230–400 mesh) using hexane–EtOAc gradients of increasing polarity. All organic extracts were dried over magnesium sulfate and evaporated under reduced pressure, below 65 °C. Compounds **4** and **5** were obtained according to ref 6.

Diene 2. Method A. Methylenation of enone **5** (240 mg, 1.17 mmol) with Tebbe reagent (0.5 M, 0.5 mL, 2.5 mmol) following the described experimental procedure⁷ afforded **2** (38 mg, 16%) as an oil. $[\alpha]_D^{16} -189.35$ (*c* 5.07, CHCl₃) (not described before). The ¹H NMR spectrum data were in good agreement with the previously reported data.⁵ ¹³C NMR (CDCl₃, 50 MHz) (not described before): δ 161.8 (C-9), 150.1

* To whom correspondence should be addressed. Phone: 56 2 68 64 435. Fax: 56 2 68 64 744. E-mail: mcortesm@puc.cl.

Scheme 1^a

^a Reagent (i) H₂, PtO₂; (ii) CH₃SO₂Cl, Py; (iii) *t*-BuOK, DMF; (iv) *p*-benzoquinone.

(C-8), 108.9 (C-12), 103.1 (C-11), 52.5 (C-5), 42.3 (C-3), 40.2 (C-10), 37.6 (C-1), 35.9 (C-7), 33.8 (C-4), 33.5 (C-13), 22.7 (C-6), 22.1 (C-14), 20.7 (C-15), 19.2 (C-2).

Method B. Catalytic hydrogenation of diol 4 (1.0 g, 4.2 mmol) with PtO₂ (100 mg) gave the known saturated diol 6⁸ (830 mg, 82%). Diol 6 (400 mg, 1.66 mmol) was esterified by CH₃SO₂Cl (0.8 mL, 9.96 mmol) and dry pyridine at -15 °C. After the usual workup and column chromatography, dimethylsilylate 7 was obtained (610 mg, 93%) as white crystals (hexane): mp 115–116 °C; [α]_D²⁵ +42 (c 2.38, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 4.47–4.41 (1H, dd, *J* = 10.0, 5.0, H-11), 4.34–4.20 (3H, m, *J* = 10.0, 2H-12 and 1H-11), 3.04 (3H, s, H₃CSO₂), 3.03 (3H, s, H₃CSO₂), 2.38–2.35 (1H, m, H-8), 2.05–2.0 (1H, m, Ha-7), 1.8–1.9 (1H, ddd, *J* = 5.0 Hz, H-9), 0.91–0.97 (1H, dd, *J* = 12.0, 2.0 Hz, H-5), 0.89 (3H, s, H-13), 0.80 (6H, s, H-15 and H-14); ¹³C NMR (CDCl₃, 75 MHz) δ 68.9 (C-12), 67.2 (C-11), 55.9 (C-5), 51.5 (C-9), 41.5 (C-3), 39.0 (C-1), 37.1 (C-10), 34.5 (C-8), 33.3 (C-13), 33.1 (C-4), 28.3 (C-7), 21.4 (C-14), 18.3 (C-2), 17.3 (C-6), 16.5 (C-15); *anal.* C 51.59%, H 8.39%, S 15.95%, calcd for C₁₇H₃₂O₆S₂ (396.56), C 51.49%, H 8.14%, S 16.14%. Compound 7 (470 mg, 1.206 mmol) in dry DMF (5.0 mL) was treated, at -15 °C under N₂ atmosphere, with *t*-BuOK (1.326 g, 11.23 mmol). The mixture was stirred at -15 °C for 8 h and then allowed to warm to room temperature. Usual workup and column chromatography gave diene 2 (108 mg, 45%), with physical properties and spectroscopic data identical to those obtained by method A.

Diels–Alder Reaction of 2 with *p*-Benzoquinone. *p*-Benzoquinone (60 mg, 0.33 mmol) was added to a solution of diene 2 (40 mg, 0.20 mmol) in dry benzene (6 mL). The reaction mixture was warmed to reflux temperature, under N₂ atmosphere, for 28 h. After evaporation of the solvent and column chromatography, quinone 9 (30 mg, 49%) was obtained as a yellow oil: [α]_D²⁵ +70 (c 3.0, CHCl₃); ¹H NMR (CDCl₃, 200 MHz) δ 6.7 (2H, s, H-3' and H-4'), 3.03–2.91 (4H, m, H-14

and H-15), 1.00 (3H, s, CH₃), 0.89 (3H, s, CH₃), 0.85 (3H, s, CH₃); ¹³C NMR (CDCl₃, 50 MHz) δ 187.2 (C-2')[#], 187.0 (C-5')[#], 140.3 (C-6'), 139.2 (C-1'), 136.4 (C-4'), 136.1 (C-3'), 134.2 (C-9), 122.7 (C-8), 51.1 (C-5), 41.6 (C-3), 37.4 (C-10), 36.8 (C-1), 33.3 (C-4), 33.2 (C-12), 31.5 (C-15), 29.9 (C-7), 22.9 (C-14), 21.6 (C-11), 19.4 (C-13), 18.9 (C-2), 18.6 (C-6) (# interchangeable signals).

Oxidation of 9. DDQ (70 mg, 0.31 mmol) was added to a solution of 9 (30 mg, 0.097 mmol) in benzene (6 mL). The mixture was warmed to reflux temperature for 4 h. After evaporation of the solvent and column chromatography, (+)-cyclozonarone (28 mg, 94%) was obtained as an oil: [α]_D²⁵ +93.18 (c 1.395, CHCl₃). ¹H NMR and ¹³C NMR spectra are in good agreement with those of natural (-)-cyclozonarone, except for opposite sign of optical rotation.

Acknowledgment. The authors thank FONDECYT (Research Grant 8980003) for financial support. M.C. thanks Dirección de Investigación, Pontificia Universidad Católica de Chile (DIPUC), for a fellowship.

References and Notes

- (1) Faulkner, D. J. *Nat. Prod. Rep.* **1998**, 113–158.
- (2) Ochi, M.; Kotsuki, H.; Murakoa, K.; Tokoroyama, T. *Bull. Chem. Soc. Jap.* **1979**, 52, 629–630.
- (3) Kurata, K.; Taniguchi, K.; Suzuki, M. *Phytochemistry* **1996**, 41, 749–752.
- (4) Valderrama, J. A.; Cortés, M.; Pessoa-Mahana, D.; Preite, M.; Benites, J. *Tetrahedron Lett.* **2000**, 41, 3563–3566.
- (5) Nakano, T.; Maillou, M. A. *Synth. Commun.* **1981**, 11, 463–473.
- (6) Peña, W.; López, J. T.; Cortés, M. *Synth. Commun.* **1989**, 19, 2841–2850.
- (7) Pine, S. H.; Shen, G. S.; Hoang, H. *Synthesis* **1991**, 165–167.
- (8) Barrero, A. F.; Cortés, M.; Manzaneda, E. A.; Cabrera, E.; Chahboun, R.; Lara, M.; Rivas, R. *J. Nat. Prod.* **1999**, 62, 1488–1491.
- (9) Araya-Maturana, R.; Cassels, B. K.; Delgado-Castro, T.; Valderrama, J. A.; Weiss-López, B. E. *Tetrahedron* **1999**, 55, 637–648.

NP0004146