

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/256075348>

Asteriscane-Type Sesquiterpenoids from the Soft Coral *Sinularia capillosa*

ARTICLE in JOURNAL OF NATURAL PRODUCTS · AUGUST 2013

Impact Factor: 3.8 · DOI: 10.1021/np400480p · Source: PubMed

CITATIONS

8

READS

56

6 AUTHORS, INCLUDING:



Dawei Chen

China National Center for Food Safety Risk Ass...

21 PUBLICATIONS 52 CITATIONS

SEE PROFILE



Dong Liu

Peking University

37 PUBLICATIONS 201 CITATIONS

SEE PROFILE



Leen van Ofwegen

Naturalis Biodiversity Center

102 PUBLICATIONS 1,020 CITATIONS

SEE PROFILE



Wenhan Lin

Beijing Medical University


167 PUBLICATIONS 2,205 CITATIONS

SEE PROFILE

Asteriscane-Type Sesquiterpenoids from the Soft Coral *Sinularia capillosa*Dawei Chen,[†] Wei Chen,[†] Dong Liu,[†] Leen van Ofwegen,[‡] Peter Proksch,[§] and Wenhan Lin^{*,†}[†]State Key Laboratory of Natural and Biomimetic Drugs, Peking University, Beijing 100191, People's Republic of China[‡]China National Center for Biotechnology Development, Beijing 100036, People's Republic of China[‡]National Museum of Natural History Naturalis, 2300 RA, Leiden, The Netherlands[§]Institute of Pharmaceutical Biology and Biotechnology, Heinrich-Heine University, 40225 Duesseldorf, Germany

S Supporting Information


ABSTRACT: Chemical examination of the soft coral *Sinularia capillosa* collected from the South China Sea resulted in the isolation of 14 new asteriscane-type sesquiterpenoids, namely, capillosananes A–N (1–14), four new *seco*-asteriscanes, capillosananes O–R (15–18), and (–)-sinularone A and sinularone A. Their structures were determined on the basis of extensive spectroscopic analyses, while the absolute configurations were determined by CD and ECD calculation, Mosher's method, and chemical conversion. This is the first report of asteriscane-type sesquiterpenoids from soft corals, and capillosananes Q (17) and R (18) represent new *seco*-asteriscane skeletons. Capillosanane A exhibited potent antifouling activity against *Balanus amphitrite*, with an IC₅₀ value of 9.70 μM, while capillosananes B and I and (–)-sinularone A inhibited inflammation-related TNF-α in vitro.



Asteriscane-type sesquiterpenoids feature a bicyclo[6.3.0]-undecane skeleton and are rarely found in nature.^{1–5} Asteriscanolide from the plant *Asteriscus aquaticus*¹ was the first asteriscane-type sesquiterpenoid, while an asteriscane from the aeolid nudibranch *Phyllodesmium magnum* is the only one isolated from marine organisms and was proposed to originate from a *Sinularia* sp. soft coral, a prey of this nudibranch.⁵ The soft coral genus *Sinularia* is recognized as a rich source of terpenoids, of which cembranoids are commonly regarded as the typical type of secondary metabolites used for chemical taxonomy. Sesquiterpenoids often co-occur with cembranoids in *Sinularia* species, but the number of sesquiterpenoids from the *Sinularia* genus is much less in comparison with cembranoid metabolites. The structural patterns of sesquiterpenoids derived from *Sinularia* species consist of cadinane-type,⁶ oppositane-type,^{7,8} africane-type,^{9–12} and others.^{13–15} In order to examine the chemical diversity and the structural variation of soft coral species inhabiting different geographic locations, we collected the specimen *Sinularia capillosa* from the coral reef near Hainan Island. Chromatographic (HPLC) and ¹H NMR spectroscopic analyses revealed that the EtOAc extract of this specimen featured numerous terpenoids, differing from those reported in the literature.^{16–19} This paper reports the structure elucidation of the new asteriscane-type sesquiterpenoids 1–18. The antifouling and anti-TNF-α activities of selected compounds were evaluated.

RESULTS AND DISCUSSION

The EtOAc-soluble fraction of the soft coral *S. capillosa* was examined by ¹H NMR spectroscopy and HPLC, which indicated the presence of diverse terpenoid analogues. Repeated column chromatography including reversed-phase



HPLC separation of the EtOAc fraction led to the isolation of 18 new asteriscane-based sesquiterpenoids, named capillosananes A–R (1–18).

Capillosanane A (1) had a molecular formula of C₁₅H₂₄O₂, as determined on the basis of the HRESIMS data, requiring four degrees of unsaturation. The IR absorptions suggested the presence of hydroxy (3488 cm⁻¹) and carbonyl (1700 cm⁻¹) functionalities. The APT spectrum displayed a total of 15 resonances involving four methyl, four methylene, three methine, and four quaternary carbons, of which a ketone and a vinyl group were characterized. Thus, a bicyclic nucleus is required from the molecular unsaturation. The COSY cross-

Received: June 18, 2013

Published: August 21, 2013

peaks established an alkyl spin system from C-1 to C-5 and extended to H₃-15 (δ_{H} 0.85, d, J = 6.9 Hz), whereas the HMBC correlations observed from H₃-12 (δ_{H} 1.03, s) and H₃-13 (δ_{H} 0.73, s) to C-1 (δ_{C} 47.1), C-10 (δ_{C} 53.1), and C-11 (δ_{C} 36.9) in addition to the correlations from the geminal protons H₂-10 (δ_{H} 2.12, 1.75) to C-1, C-2 (δ_{C} 40.1), C-9 (δ_{C} 142.5), and C-8 (δ_{C} 128.8) revealed a dimethylcyclopentane ring. Further HMBC correlations from the olefinic proton H-8 (δ_{H} 5.42, s) to C-2, C-7 (δ_{C} 76.5) and C-6 (δ_{C} 216.6) and from H₂-4 (δ_{H} 1.89, m) and H₂-5 (δ_{H} 3.14, 1.96) to C-6 resulted in a cyclooctenone substructure, which was fused to the cyclopentane ring across C-2 and C-9. The positions C-3 and C-7 were methylated. A D₂O exchangeable proton at δ_{H} 5.57 (s) showed HMBC correlations to C-6, C-7, and C-8, indicating that C-7 is hydroxylated. Accordingly, the gross structure of **1** was determined as 7-hydroxyasterisca-8-en-6-one. The NOE correlations from H₃-15 to H-10b (δ_{H} 2.12) and H-1b (δ_{H} 1.61) in addition to the correlation between H-2 (δ_{H} 2.43) and H-1a (δ_{H} 1.29) indicated the opposite orientation of H-2 and H₃-15. Additional cross-peaks from H-5b (δ_{H} 3.16) to H₃-15 and OH-7 (δ_{H} 5.57) assigned OH-7 the same face as H₃-15 (Figure 1). The absolute configuration of **1** was determined using the modified Mosher's method.²⁰ First, reduction of **1** by NaBH₄ yielded **1a** (Figure 2). The NOE interaction between H-6 (δ_{H} 3.49, dd, J = 2.5, 10.5 Hz) and H₃-15 (δ_{H} 0.83, d, J = 6.8 Hz) in association with the absence of NOE interaction

between H₃-14 (δ_{H} 1.36, s) and H-6 indicated H-6 of **1a** had the same orientation as that of H₃-15 and was on the opposite face compared to H₃-14. By a standard procedure, the (R)- and (S)-MPA (methoxyphenylacetic acid) esters of **1a** were synthesized. Analysis of the $\Delta\delta^{\text{RS}}$ (δ_{R} – δ_{S}) values (Figure 3) resulted in a 6R configuration. Thus, the remaining stereogenic centers in **1** were defined as 2R, 3R, and 7R.

Capillosanane B (**2**) exhibited a pseudomolecular ion peak at m/z 259.1666 $[\text{M} + \text{Na}]^+$ by HRESIMS, consistent with a molecular formula of C₁₅H₂₄O₂. Analyses of 1D and 2D NMR data revealed an asteriscane skeleton, structurally related to **1**. The APT spectrum displayed four olefinic carbons for two double bonds and two oxygenated methines. The HMBC correlations observed from H₃-12 (δ_{H} 1.00, s) and H₃-13 (δ_{H} 0.84, s) to C-10 (δ_{C} 85.4), H₃-14 (δ_{H} 1.67, s) to C-6 (δ_{C} 69.6), C-7 (δ_{C} 140.0), and C-8 (δ_{C} 120.9), and H₃-15 (δ_{H} 0.91, d, J = 6.9 Hz) to C-2 (δ_{C} 141.5) established conjugated double bonds at $\Delta^{2,9}$ and $\Delta^{7,8}$. The COSY correlations of the D₂O exchangeable protons at δ_{H} 4.63 and 5.42 with H-6 and H-10, respectively, showed C-6 and C-10 to be hydroxylated. The NOE relationships between H-3 (δ_{H} 2.21, m) and H-6 (δ_{H} 4.63, br d, J = 9.6 Hz) and from H-1b (δ_{H} 2.10, d, J = 15.4 Hz) to H₃-15 and H-10 (δ_{H} 3.98, d, J = 5.5 Hz) were evident from the spatial proximity of H-1b to H-10 and H₃-15, indicating the same orientation of H-10, OH-6, and H₃-15. On the basis of Mosher's method,²⁰ the absolute configuration of C-6 was assigned as S (Figure 3). Accordingly, the configurations of the remaining asymmetric centers were assigned as 3R and 10S.

The structure of capillosanane C (**3**) was determined as the C-10 epimer of **2** based on the close similarity of NMR spectroscopic data with the exception of an upfield shifted H-10 (δ_{H} 3.60) and a downfield shifted C-10 (δ_{C} 87.3) of **3** in comparison with those of **2**. The NOE correlations of H-1b (δ_{H} 2.31) to H₃-15 (δ_{H} 0.96) and OH-10 (δ_{H} 4.50) supported the structural assignment.

A comparison of the NMR data (Tables 1 and 3) in association with analyses of 2D NMR spectra established capillosanane D (**4**) to be a 10-ketone analogue of **2**. The similar NOE interactions and coupling constants of the two compounds indicated **4** had the same relative configuration as that of **2**. Adding Rh₂(OCOFCF₃)₄ to a solution of **4** in CHCl₃ formed a metal complex as an auxiliary chromophore. The sign of the E band reflected the absolute configuration of the secondary alcohol by applying the bulkiness rule.^{21,22} The positive E band sign at 350 nm observed in the CD spectrum (Figure 4) was in agreement with a 6S configuration. Transformation of **4** to **2** and **3** with NaBH₄ further supported the configurational assignment.

Capillosanane E (**5**) had a molecular formula of C₁₅H₂₂O₃, as deduced from the HRESIMS data, requiring five degrees of unsaturation. The ¹H and ¹³C NMR resonances of **5** were closely similar to those of **4**, except for the presence of two oxygenated carbons at δ_{C} 76.7 (C-2) and 67.1 (C-9) that replaced the olefinic carbons at C-2 and C-9 of **4**. These findings in addition to **5** possessing an extra oxygen atom compared to **4** assigned an epoxy group across C-2 and C-9. The olefin geometry was identified as Z on the basis of the NOE correlation between H-8 (δ_{H} 5.59, s) and H₃-14 (δ_{H} 1.69, s). The absolute configuration of C-6 in **5** was assigned as 6S by the modified Mosher's method (Figure 3). Thus, the NOE correlations from H-3 (δ_{H} 2.09, m) to H-6 (δ_{H} 4.34, dt, J = 12.0, 4.2, 3.6 Hz) assigned a 3R configuration. There are two possibilities for the epoxy orientation. The observation of NOE




Figure 1. Key NOE correlations of **1**, **2**, **5**, **9**, **11**, **12**, and **17**.




Figure 2. Chemical conversion of **1** to **1a** and the formation of MPA esters of **1a**.




Figure 3. $\Delta\delta^{RS}$ ($\Delta(\delta_R - \delta_S)$) data for the MPA esters.

interactions between H-3 and H-1 α (δ_H 1.90, d, J = 14.4 Hz), and between H₃-15 (δ_H 1.08, d, J = 6.0 Hz) and H-1 β (δ_H 2.22, d, J = 14.4 Hz) (Figure 1), as well as the absence of NOE interaction between H₂-1 and H₂-4, indicated H-3 was spatially proximal to H-1 α . These findings indicated the epoxy group to be β -oriented. Accordingly, the remaining stereogenic centers were determined as 2R and 9S.

Capillosanane F (**6**) was determined to be the 6-ketone analogue of **4** on the basis of 1D and 2D NMR data. The configuration of the C-3 stereogenic center was assumed to be the same as that of **4** based on biogenetic considerations.

The molecular formula of capillosanane G (**7**) was determined as C₁₅H₂₄O₃ on the basis of its HRESIMS and NMR data. Analyses of 1D and 2D NMR data led to the establishment of an asteriscane-type sesquiterpene, structurally related to **2**. The major distinction was attributed to the presence of two conjugated double bonds at $\Delta^{7,8}$ and $\Delta^{9,10}$ instead of $\Delta^{7,8}$ and $\Delta^{2,9}$, which were evident from the HMBC correlations from H₃-12/H₃-13 (δ_H 1.05, s) to C-10 (δ_C 140.8) and from H₃-14 (δ_H 1.60, s) to C-7 (δ_C 140.7) and C-8 (δ_C 120.8). Because an OH at C-6 was recognized by the COSY relationship between H-6 (δ_H 4.42, dd, J = 5.2, 11.5 Hz) and OH (δ_H 4.67) and the HMBC interaction between H₃-14 and C-6 (δ_C 68.8), the remaining O₂H unit was assigned as a

Table 1. ¹H NMR Data for Capillosananes A–H (1–8)^a

| no. | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|-------|----------------------|------------------|------------------|---------------------------|---------------------------|----------------------------|----------------------|----------------------|
| 1 | 1.61, dd (12.8, 8.5) | 2.10, d (15.4) | 2.31, d (15.9) | 2.54, d (18.2) | 2.22, d (14.4) | 2.71, d (19.6) | 2.06, d (14.1) | 1.65, br d (5.5) |
| | 1.29, dd (12.8, 7.5) | 1.89, d (15.4) | 1.79, d (15.9) | 2.35, d (18.2) | 1.90, d (14.4) | 2.57, d (19.6) | 1.76, d (14.1) | |
| 2 | 2.43, dd (8.5, 7.5) | | | | | | | |
| 3 | 1.58, m | 2.21, m | 2.19, m | 2.59, m | 2.09, m | 2.75, m | 1.65, m | 1.96, m |
| 4 | 1.89, m | 1.70, m | 1.74, m | 1.82, m | 1.20, m | 2.21, m | 1.91, m | 1.57, m |
| | | 1.34, m | 1.33, m | 1.51, m | 1.03, m | 1.89, m | 0.93, m | 1.29, m |
| 5 | 3.14, m | 1.66, m | 1.65, m | 1.73, m | 1.59, tt (12.4, 5.2) | 2.65, m | 1.57, m | 2.31, m |
| | 1.96, dd (11.6, 5.6) | 1.45, m | 1.44, m | 1.48, m | 1.28, m | 2.11, ddd (11.3, 4.3, 2.3) | 1.33, br t (12.6) | 1.46, m |
| 6 | | 4.63, br d (9.6) | 4.56, br d (9.6) | 4.37, dt (10.0, 4.2, 3.6) | 4.34, dt (12.0, 4.2, 3.6) | | 4.42, dd (11.5, 5.2) | 3.92, dd (10.5, 6.1) |
| 8 | 5.42, s | 5.63, s | 5.80, s | 5.80, s | 5.59, s | 6.68, s | 5.80, s | 5.81, s |
| 10 | 2.12, d (13.3) | 3.98, d (5.5) | 3.60, d (6.7) | | | | 5.23, d (1.3) | 5.36, d (1.7) |
| | 1.75, d (13.3) | | | | | | | |
| 12 | 1.03, s | 1.00, s | 0.94, s | 1.00, s | 1.04, s | 1.08, s | 1.05, s | 1.03, s |
| 13 | 0.73, s | 0.84, s | 0.91, s | 0.98, s | 0.98, s | 1.09, s | 1.05, s | 1.11, s |
| 14 | 1.20, s | 1.67, s | 1.65, s | 1.72, s | 1.69, s | 1.90, s | 1.60, s | 1.77, s |
| 15 | 0.85, d (6.9) | 0.91, d (6.9) | 0.96, d (6.8) | 1.12, d (6.9) | 1.08, d (6.9) | 1.11, d (6.9) | 0.95, d (6.9) | 0.77, s |
| 2-OOH | | | | | | | 10.71, s | 10.58, s |
| 6-OH | | 4.63, br s | 4.63, d (4.2) | 4.80, d (4.3) | 4.91, d (3.6) | 6.27, s | 4.67, s | 4.61, d (6.1) |
| 7-OH | 5.57, s | | | | | | | |
| 10-OH | | 4.52, d (5.8) | 4.50, d (6.7) | | | | | |

^aRecorded in DMSO-*d*₆ at 500 MHz.

Table 2. ^1H NMR Data for Capillosananes I–N (9–14)^a

| no. | 9 | 10 | 11 | 12 | 13 | 14 |
|-------|-----------------------------|-----------------------------|-----------------------------|----------------|----------------------|----------------------|
| 1 | 1.72, d (13.6) | 1.76, d (13.6) | 1.70, d (14.0) | 1.71, d (13.3) | 1.83, d (14.1) | 1.99, d (14.0) |
| | 1.38, d (13.6) | 1.35, d (13.6) | 1.50, d (14.0) | 1.24, d (13.3) | 1.73, d (14.1) | 1.22, d (14.0) |
| 3 | 1.50, m | 1.78, m | 1.93, m | 1.48, m | 1.61, m | |
| 4 | 1.70, m | 1.79, m | 1.97, m | 1.61, m | 1.58, br d (13.6) | 1.68, m |
| | 1.31, br d (11.2) | 1.28, dd (12.3, 4.0) | 1.16, dt (13.4, 4.0, 3.5) | 1.21, m | 1.32, qd (13.6, 3.6) | 1.39, m |
| 5 | 1.66, ddd (14.0, 12.8, 4.5) | 1.65, ddd (14.0, 13.5, 4.5) | 1.50, ddd (14.0, 12.5, 3.5) | 1.82, m | 1.58, dd (12.1, 3.6) | 1.71, m |
| | 1.49, m | 1.50, dt (12.9, 4.1, 2.5) | 1.31, ddd (13.3, 4.2, 2.9) | 1.47, m | 1.46, dt (3.6, 12.9) | 1.69, m |
| 6 | | | | 4.45, d (8.8) | | |
| 8 | 5.46, d (1.5) | 5.65, s | 5.54, s | 5.38, s | 6.14, br s | 5.51, d (1.7) |
| 10 | 2.13, dd (14.2, 1.7) | 3.61, d (2.0) | 1.95, d (13.3) | 2.13, d (13.2) | 5.25, s | 2.39, dd (12.5, 1.6) |
| | 1.88, d (14.2) | | 1.68, d (13.3) | 1.39, d (13.2) | | 1.72, d (12.5) |
| 12 | 1.05, s | 0.88, s | 1.04, s | 1.12, s | 1.10, s | 1.05, s |
| 13 | 0.99, s | 0.96, s | 1.06, s | 1.08, s | 1.08, s | 0.95, s |
| 14 | 1.05, s | 1.11, s | 1.52, d (1.0) | 1.72, s | 1.63, s | 1.10, s |
| 15 | 1.03, d (6.9) | 0.98, d (6.9) | 0.93, d (6.9) | 0.83, d (6.9) | 0.80, d (6.5) | 1.19, s |
| 2-OH | | | | 3.92, s | | |
| 3-OH | | | | | | 4.44, s |
| 6-OH | 5.07, s | 5.10, s | 5.96, s | | 6.27, s | 5.08, s |
| 7-OH | 4.88, s | 4.88, s | | | | 4.85, s |
| 9-OH | | | 4.54, s | | | |
| 10-OH | | 4.84, d (2.5) | | | | |

^aRecorded in DMSO-*d*₆, at 500 MHz.

hydroperoxy group, which was characterized by the presence of a hydroperoxy proton (OOH) at δ_{H} 10.71 (s).²³ This group was located at C-2 (δ_{C} 100.7) on the basis of the HMBC correlations from the OOH proton and H-10 (δ_{H} 5.23, d, J = 1.3 Hz) to C-2. The NOE correlation between H-3 and H-6 indicated both protons are oriented on the same face, while the NOE relationship between H-3 (δ_{H} 1.65, m) and H-1a (δ_{H} 1.76, d, J = 14.1 Hz) suggested that the hydroperoxy group is oriented opposite H-3.

The gross structure of capillosanane H (8) was determined to be the same as that of 7 on the basis of HRESIMS data and 1D and 2D NMR data analyses. The NOE correlation between H-3 (δ_{H} 1.96, m) and H-6 (δ_{H} 3.92, dd, J = 11.5, 6.1 Hz) in the NOESY spectrum indicated their spatial proximity. However, the presence of an NOE interaction from H₃-15 to H₂-1 and the absence of an interaction between H-3 and the protons at C-1 were indicative of a C-2 epimer. The absolute configuration at C-2 of 8 was established by Cotton effects in the CD spectrum. On the basis of the Beecham rule for allylic oxygen-containing dienes,^{24,25} the positive sign at 220 nm in 8 follows from the right-handed helix with the allylic oxygen and the first double bond of diene, indicating a 2S configuration. Accordingly, the negative sign at 220 nm in 7 reflected a 2R configuration. These assignments were confirmed by the calculated ECD data of 8 at the B3LYP-SCRF/6-311++G(d,2p) level (Figure 5).

Capillosanane I (9) had a molecular formula of C₁₅H₂₄O₃, as determined from the HRESIMS and NMR data, requiring four degrees of unsaturation. The ^1H and ^{13}C NMR data were closely related to those of 1, indicating an asteriscane-type analogue. Differences included an oxygenated quaternary carbon at δ_{C} 84.8 (C-2) of 9 in place of a methine in 1, whereas C-6 of 9 presented as a hemiacetal carbon (δ_{C} 98.2) based on a hydroxy proton at δ_{H} 5.07 (s) correlated to C-7 (δ_{C} 68.2), C-6, and C-5 (δ_{C} 26.6) in the HMBC spectrum. On the basis of the molecular formula, the remaining site of unsaturation was attributed to an ether bridge across C-2 and




C-6. The NOE interaction between H-3 (δ_{H} 1.50, m) and H-10a (δ_{H} 2.13, dd, J = 1.7, 14.2 Hz) (Figure 1) indicated that the oxygen bridge was oriented on the same face as H₃-15. Additional NOE interactions between H₃-14 (δ_{H} 1.05, s)/H-5a (δ_{H} 1.50) and H₃-15 (δ_{H} 1.03, d, J = 6.9 Hz)/H-5b (δ_{H} 1.66) clarified the orientation of H₃-14 as opposite of H₃-15. Upon addition of dimolybdenum tetraacetate [Mo₂(OAc)₄] to a solution of 9 in DMSO, a metal complex was generated. The sign of the induced CD bands at 310, 350, and 400 nm reflected the O–C–C–O torsion angle on the basis of the in situ dimolybdenum CD method.^{26,27} The negative CD effects observed at 300 and 380 nm (Figure 6) permitted assignment of the 6S and 7R configurations. Accordingly, the configurations of remaining asymmetric centers of 9 were assigned as 2R, 3R, 6S, and 7R.

Interpretation of 1D and 2D NMR data disclosed the structure of capillosanane J (10) as a 10-hydroxylated analogue of 9. The closely similar NOE interactions confirmed that 10 shares the same relative configuration as that of 9, while the NOE correlation between H-3 (δ_{H} 1.78, m) and 10-OH (δ_{H} 4.84, d, J = 2.5 Hz) revealed H-10 to be oriented on the same face as H₃-15. The absolute configuration of 10 was determined by the modified Mosher's method.²⁰ Analysis of the $\Delta\delta^{\text{RS}}$ ($\delta_{\text{R}} - \delta_{\text{S}}$) values (Figure 3) resulted in a 10S assignment. Thus, the other stereogenic centers in 10 were assigned as 2R, 3R, 6S, and 7R.

Capillosanane K (11) and 9 possessed the same molecular formula, while the ^1H NMR and APT data of 11 were mostly similar to those of 9. An obvious distinction was recognized for the double bond in 11 residing at C-7/C-8 instead of C-8/C-9, whereas C-9 (δ_{C} 79.6) of 11 was hydroxylated. These assignments were confirmed through HMBC correlations of the olefinic methyl protons H₃-14 (δ_{H} 1.52, d, J = 1.0 Hz) to C-6 (δ_{C} 97.0), C-7 (δ_{C} 132.1), and C-8 (δ_{C} 132.0), in addition to the correlations of OH-9 (δ_{OH} 4.54, s) to C-2 (δ_{C} 91.3), C-8, C-9, and C-10 (δ_{C} 57.9). The NOE correlations indicated the relative configurations of the stereogenic centers at C-2, C-3,

Table 3. ^{13}C NMR Data for Capillosananes A–N (1–14)^a

| no. | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
|-----|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 1 | 47.1, CH ₂ | 49.2, CH ₂ | 49.7, CH ₂ | 46.4, CH ₂ | 42.6, CH ₂ | 43.3, CH ₂ | 48.5, CH ₂ | 43.0, CH ₂ | 49.4, CH ₂ | 48.6, CH ₂ | 52.7, CH ₂ | 45.9, CH ₂ | 52.4, CH ₂ | 46.8, CH ₂ |
| 2 | 40.1, CH | 141.5, C | 144.7, C | 175.0, C | 76.7, C | 180.5, C | 100.7, C | 99.7, C | 84.8, C | 84.7, C | 91.3, C | 89.8, C | 91.2, C | 87.0, C |
| 3 | 36.9, CH | 37.3, CH | 37.3, CH | 38.6, CH | 35.8, CH | 35.0, CH | 43.9, CH | 32.3, CH | 31.0, CH | 33.3, CH | 28.4, CH | 39.2, CH | 38.8, CH | 70.6, C |
| 4 | 33.8, CH ₂ | 31.3, CH ₂ | 31.0, CH ₂ | 30.8, CH ₂ | 26.0, CH ₂ | 39.9, CH ₂ | 27.2, CH ₂ | 31.3, CH ₂ | 25.9, CH ₂ | 25.6, CH ₂ | 26.7, CH ₂ | 29.9, CH ₂ | 29.8, CH ₂ | 34.0, CH ₂ |
| 5 | 35.3, CH ₂ | 33.9, CH ₂ | 33.9, CH ₂ | 32.9, CH ₂ | 34.8, CH ₂ | 38.7, CH ₂ | 36.0, CH ₂ | 32.7, CH ₂ | 26.6, CH ₂ | 26.8, CH ₂ | 28.3, CH ₂ | 30.8, CH ₂ | 33.6, CH ₂ | 31.3, CH ₂ |
| 6 | 216.6, C | 69.6, CH | 69.4, CH | 69.5, CH | 68.6, CH | 205.6, C | 68.8, CH | 72.8, CH | 68.2, C | 98.2, C | 97.0, C | 82.7, CH | 98.8, C | 98.0, C |
| 7 | 76.5, C | 140.0, C | 139.3, C | 143.4, C | 146.8, C | 138.8, C | 140.7, C | 141.2, C | 68.2, C | 68.4, C | 132.1, C | 143.4, C | 138.8, C | 68.5, C |
| 8 | 128.8, CH | 120.9, CH | 120.8, CH | 115.7, CH | 112.5, CH | 129.0, CH | 120.8, CH | 120.2, CH | 125.1, CH | 128.6, CH | 132.0, CH | 124.6, CH | 121.3, CH | 126.8, CH |
| 9 | 141.5, C | 133.8, C | 133.1, C | 131.3, C | 67.1, C | 135.0, C | 140.3, C | 140.3, C | 145.6, C | 147.1, C | 79.6, C | 102.5, C | 136.4, C | 145.7, C |
| 10 | 53.1, CH ₂ | 85.4, CH | 87.3, CH | 212.7, C | 214.9, C | 210.7, C | 140.8, CH | 141.7, CH | 44.6, CH ₂ | 81.4, CH | 57.9, CH ₂ | 49.9, CH ₂ | 134.5, CH | 48.0, CH ₂ |
| 11 | 36.9, C | 41.2, C | 39.8, C | 42.5, C | 42.0, C | 42.7, C | 41.5, C | 42.4, C | 35.4, C | 40.8, C | 35.9, C | 35.5, C | 43.6, C | 36.4, C |
| 12 | 28.8, CH ₃ | 27.5, CH ₃ | 28.6, CH ₃ | 25.7, CH ₃ | 27.8, CH ₃ | 25.9, CH ₃ | 30.9, CH ₃ | 29.7, CH ₃ | 31.3, CH ₃ | 29.3, CH ₃ | 33.3, CH ₃ | 33.3, CH ₃ | 28.1, CH ₃ | 30.1, CH ₃ |
| 13 | 27.3, CH ₃ | 23.2, CH ₃ | 23.0, CH ₃ | 25.1, CH ₃ | 26.5, CH ₃ | 24.6, CH ₃ | 29.1, CH ₃ | 28.8, CH ₃ | 31.3, CH ₃ | 24.3, CH ₃ | 33.4, CH ₃ | 33.1, CH ₃ | 33.4, CH ₃ | 29.8, CH ₃ |
| 14 | 26.2, CH ₃ | 18.2, CH ₃ | 18.5, CH ₃ | 18.5, CH ₃ | 17.4, CH ₃ | 20.3, CH ₃ | 16.7, CH ₃ | 23.4, CH ₃ | 23.8, CH ₃ | 23.7, CH ₃ | 17.5, CH ₃ | 12.6, CH ₃ | 18.5, CH ₃ | 24.0, CH ₃ |
| 15 | 14.8, CH ₃ | 23.2, CH ₃ | 23.0, CH ₃ | 22.7, CH ₃ | 20.1, CH ₃ | 17.3, CH ₃ | 19.9, CH ₃ | 17.1, CH ₃ | 12.7, CH ₃ | 12.7, CH ₃ | 16.0, CH ₃ | 18.8, CH ₃ | 18.0, CH ₃ | 22.0, CH ₃ |

^aRecorded in DMSO-*d*₆ at 125 MHz.Figure 4. CD spectra for Rh₂(OCOCH₃)₄ complexes of 4 and 16.Figure 5. CD curves of 7 and 8 and the calculated CD spectrum of the 2*S*, 3*R*, 6*S* isomer of 8.Figure 6. ICD curves of 9 and 14 induced by Mo₂(OAc)₄ in DMSO-*d*₆.

and C-6 to be the same as those of 9 (Figure 1). Additional NOE correlations from 9-OH to H-3 (δ_{H} 1.93, m) and H-10 α (δ_{H} 1.68, d, $J = 13.3$ Hz) assigned 9-OH to the same face as H-3.

Capillosanane L (**12**) had a molecular formula of $C_{15}H_{24}O_2$, as determined through HRESIMS and NMR data. Analysis of the 1H NMR and ^{13}C NMR data disclosed **12** to be an asteriscane-type sesquiterpenoid, structurally related to **11**. The HMBC correlations from H_3 -14 (δ_H 1.72, s) to C-6 (δ_C 82.7), C-7 (δ_C 143.4), and C-8 (δ_C 124.6) ascertained a double bond at $\Delta^{7,8}$, whereas C-6 was oxygenated. In addition the quaternary carbons C-2 (δ_C 89.8) and C-9 (δ_C 102.5) were distinguished through HMBC correlations from H_3 -15 (δ_H 0.83, d, J = 6.9 Hz) to C-2. An additional HMBC correlation from H-6 (δ_H 4.45, d, J = 8.8 Hz) to C-9 confirmed the formation of an ether bridge across C-6 and C-9. Thus, a D_2O exchangeable proton at δ_{OH} 3.92 (s) was derived from a hydroxy group at C-2. The NOE correlations between H-8 (δ_H 5.38, s)/H-10a (δ_H 2.14), H-10b/H-1b (δ_H 1.24, d, J = 13.3 Hz), H-10b/OH-2, and H-1a (δ_H 1.71, d, J = 13.3 Hz)/H-3 (δ_H 1.48, m) located the bridging oxygen on the same face as that of OH-2.

The molecular formula of capillosanane M (**13**) was determined as $C_{15}H_{22}O_2$ according to the HRESIMS and NMR data. Interpretation of the 1D and 2D NMR data revealed the gross structure of **13** to be a $\Delta^{7,8}$, $\Delta^{9,10}$ analogue of **9**. The NOE observed from H_3 -15 to H_3 -12 (δ_H 1.10, s) and H-1b (δ_H 1.83, d, J = 14.1 Hz) and between H-1a (δ_H 1.74, d, J = 14.1 Hz) and H-3 (δ_H 1.61, m) established that the ether oxygen was oriented on the same face as H-3. In order to determine the absolute configuration of **13**, compound **9** was converted to **9a** and **9b** in an acidic solution (Figure 7). The




Figure 7. Acidic transformation of **9** and **11** to **9a** and **9b**.

absolute configurations of the stereogenic centers in **9b** were the same as those of **9**. The opposite CD effects of **13** and **9b** (Figure 8) led to the assignment of the configuration of **13** as 2*S*, 3*R*, and 6*R*.

Analyses of 1D and 2D NMR data and comparison of the NMR data revealed capillosanane N (**14**) to be a 3-hydroxylated analogue of **9**, which was supported by the methyl singlet H_3 -15 (δ_H 1.19, s) and an OH proton (δ_{OH} 4.44, s) correlated to C-3 (δ_C 70.6) and to C-2 (δ_C 87.0) and C-4 (δ_C 34.0). The ROESY spectrum showed NOE correlations from OH-3 to H-10 β (δ_H 2.39, dd, J = 12.5, 1.6 Hz) and H-1 β (δ_H 1.99, d, J = 14.0 Hz), enabling assignment of the same orientation of the ether oxygen as for that of **13**, whereas OH-3 was β -oriented. Based on the in situ dimolybdenum CD method, the positive Cotton effects around 310 and 400 nm (Figure 6) were in agreement with 6*R* and 7*R* configurations. Accordingly, the remaining asymmetric centers were assigned the 2*R* and 3*R* configurations.

Capillosanane O (**15**) was determined as a $\Delta^{9,10}$ analogue of (–)-sinularone A on the basis of the similar NMR data, except for the presence of two additional olefinic carbons at δ_C 168.3 (C-10, CH) and 140.2 (C-9, C) as well as the HMBC




Figure 8. CD curves of **9b** and **13**.

correlations of H-10 (δ_H 7.10, s) to C-1, C-2, C-11, C-12, and C-13.

Comparison of the NMR data between capillosanane P (**16**) and **15** and analyses of the 2D NMR data indicated the structure of **16** to be a 3-hydroxylated analogue of **15**, as evident by the presence of a hydroxymethine (δ_H 3.56, m; δ_C 65.8) replacing the ketone carbon at C-3 of **15**, in addition to the COSY relationship between H-3 and H_3 -15 (δ_H 1.03, d, J = 6.0 Hz). On the basis of the bulkiness rule, the $Rh_2(CF_3COO)_4$ complex of **16** presenting a positive CD effect at 350 nm was in agreement with a 3*S* configuration (Figure 4).

Capillosanane Q (**17**) had a molecular formula of $C_{15}H_{22}O_3$, as determined by HRESIMS data, requiring five degrees of unsaturation. The APT spectrum displayed four methyl, four methylene, two methine, and five quaternary carbons, including the resonances for two olefinic carbons, a ketone, and a lactone carbonyl. Thus, the structure of **17** should be a bicyclic nucleus. A dimethylcyclopentane ring was established on the basis of the HMBC interactions from the methyl singlets at δ_H 1.09 (s, H_3 -12) and 1.01 (s, H_3 -13) to C-10 (δ_C 51.3), C-1 (δ_C 47.8), and C-11 (δ_C 36.7) in addition to the methylene protons H_2 -1 (δ_H 1.97, 1.59) and H_2 -10 (δ_H 2.22, 2.18) to C-11, C-2 (δ_C 92.7) and C-9 (δ_C 159.5). An olefinic proton at δ_H 6.41 (1H, s, H-8) correlating to C-2 and C-10, as well as a ketone (δ_C 198.4, C-7) and a methyl carbon C-14, and correlations from H_3 -14 (δ_H 2.17, s) to C-7 and C-8 (δ_C 125.8) in the HMBC spectrum supported an ylideneprop-2-one unit fused to C-9. Additional HMBC and COSY correlations established a spirocyclic 3-methyl- δ -lactone linked to C-2, which was evidenced by the HMBC interactions from H_2 -1 to C-3 and from H-3 (δ_H 2.84, m) to C-2 and C-1 in association with the molecular unsaturation. The NOE interaction between H_2 -10 and H-8 was indicative of an 8*Z* geometry, while the NOE correlations from H_3 -15 (δ_H 0.85, d, J = 6.9 Hz) to H-1a (δ_H 1.97) and H-10a (δ_H 2.18) clarified C-1 to be oriented on the same face as H_3 -15. The absolute configuration of C-2 was determined as *R* according to the chemical conversion of **9** to **17** by $NaIO_4$ oxidation (Figure 9). This assignment was also supported by the computed ECD data of the 2*R* cognate at the B3LYP/aug-cc-pVDZ+6-31+G(d) level using a TD-DFT/B3LYP calculation, which showed a similar CD curve as compared to the measured CD spectrum (Figure 10).

Capillosanane R (**18**) had the same molecular formula as that of **17** based on HRESIMS data. The 1H and ^{13}C NMR features




Figure 9. Transformation from 9 to 17.




Figure 10. CD and computed ECD spectra of 17.

were very similar to those of 17, implying 18 to be structurally related to 17. However, 2D NMR data distinguished different structural patterns. The HMBC correlations established a dimethyl cyclopentene. The olefinic doublet H-1 (δ_H 5.56, d, J = 1.5 Hz) was attributed to a long-range coupling with H-9 (δ_H 3.27, m), indicating C-2 to be substituted. COSY correlations gave rise to a segment from H₂-8 to H₂-10, while H-9, H₂-8, and H₃-14 (δ_H 2.10, s) correlated to a ketone carbon at δ_C 208.0 (C-7) in the HMBC spectrum, revealing a propanone unit linked to C-9. A γ -methyl- γ -lactone ring was assembled based on the COSY coupling between H₂-4 and H₂-5 and the HMBC correlations from a carbonyl carbon at δ_C 176.6 and an oxygenated sp^3 quaternary carbon at δ_C 85.7 (C-3) to H₂-4 and H₂-5 in addition to the correlation between H₃-15 (δ_H 1.43, s) and C-3. The linkage of the γ -lactone to the cyclopentene ring through a C-2/C-3 bond was deduced by the HMBC interaction from H₃-15 to C-2 (δ_C 144.5). The presence of an NOE interaction between H-9 and H₃-15 and the absence of an interaction between H₃-15 and H₂-8 supported the same orientation for H-9 and H₃-15. The measured CD curve of 18 was in accordance with that calculated for the 3S,9S-cognate at the TD-DFT/B3LYP-6-31+G(d) level using the B3LYP/aug-cc-pVDZ method and opposite of that calculated for the 3R,9R-cognate (Figure 11).

In addition, (–)-sinularone A and sinularone A²⁷ were separated on a chiral-phase column by semipreparative HPLC. The structure of (–)-sinularone A was identified on the basis of the identical NMR data compared to those of sinularone A, except for the specific rotation $[\alpha]_D^{15} -23.4$ (c 0.2, MeOH) and CD effects (see SI) opposite those of sinularone A $[\alpha]_D^{15} +7.3$ (c 0.27, MeOH).²⁷ Because C-5 of sinularone A has the S configuration, the same stereogenic center in (–)-sinularone A has the SR configuration.

During the separation and storage of the new compounds, degradation and/or rearrangement of some of the compounds occurred. Capillosanane I (9) was partially converted to 9a and




Figure 11. CD spectrum of 18 and calculated CD spectra of 3R,9R and 3S,9S isomers.

9b during storage in CDCl₃ solution for two weeks (Figure 7). HRESIMS and 2D NMR data established the structures of 9a and 9b, which are the dehydrated analogues of 9. Adding a trace of TFA to a solution of 9 rapidly induced the conversion. The absolute configuration of C-6 in 9b was determined via the CD data of the in situ formed [Rh₂(OCOCF₃)₄] complex, where the negative E band (363 nm) sign in the CD spectrum determined a 6S configuration (see SI). The remaining asymmetric centers were determined as 2R and 3R, accordingly. Treatment of 11 with a trace of TFA also produced 9a and 9b (Figure 7). Thus the absolute configuration of 11 was determined to be 2R, 3R, 6S, and 9R. Capillosanane G (7) was partially converted to derivative 7a during HPLC separation using MeCN/H₂O (Figure 12). Compound 7a




Figure 12. Autotransformation of 7 to 7a and 4.

had the same molecular formula as that of 7 as determined by HRESIMS data, while the ¹H NMR resonances in 7a were mostly similar to those of 7 with the exception of H-10 in 7a being shifted upfield to δ_H 3.96 (s). This finding suggested 7a to be a 10-hydroperoxy- $\Delta^{2(9)}$ -analogue of 7. The pathway is depicted as an allylic rearrangement to give the isomeric hydroperoxide.^{28,29} Further conversion of 7a to 4 during ¹³C NMR measurement in CDCl₃ confirmed the structural assignment (Figure 12).

All compounds were inactive against human tumor cell lines HCT-8, HePG2, BGC-823, A549, and SKOV3 with IC₅₀ values of >10 μ M/mL. Compounds 2, 9, and (–)-sinularone A, representing different structural patterns, were selected for a bioassay to inhibit inflammation-related TNF- α . They showed weak activity with inhibitory rates of 16%, 21%, and 23% at 10 μ M, respectively. Nevertheless, compound 1 showed potent activity for antifouling against *Balanus amphitrite*, with an IC₅₀ value of 9.7 μ M, whereas the IC₅₀ values of 9 and (–)-sinularone A were 54 and 74 μ M. In addition, sinularone A was previously reported to potently inhibit *B. amphitrite* with the EC₅₀ of 13.9 μ g/mL,²⁷ indicating the configuration at C-9

plays a key role in the inhibitory activity. These results suggest that asteriscane-based sesquiterpenoids in *S. capillosa* play a chemoeological role such as antifouling without toxicity to higher organisms.

The soft coral *S. capillosa* has been reported to contain various terpenoids with unique skeletons. Furanos sesquiterpenes are a group of typical sesquiterpenes isolated from this specimen collected from an Australian coral reef.³⁰ The unprecedented sesquiterpenoid capillosanol,¹⁶ the furanos sesquiterpenoid capillofuranocarboxylate, the furanobenzos sesquiterpenoids capillobenzopyranol and capillobenzofuranol, and other sesquiterpenoids were isolated from the same species inhabiting Dongsha Atoll.¹⁸ However, the same species from Sanya Bay of Hainan Island produced cembranoids.¹⁹ The present work dramatically enriches the numbers of the asteriscane-type family, while it represents the first isolation of asteriscane-based sesquiterpenoids from soft corals. These findings support the depiction that an asteriscane-type sesquiterpenoid from the aeolid nudibranch *Phyllodesmium magnum* originates in the *Sinularia* soft coral, which is the prey for this organism. The *seco*-derivatives (**15**–**18**) are derived from asteriscane-type sesquiterpenoids through oxidation, while compounds **17** and **18** present an unusual carbon skeleton.

EXPERIMENTAL SECTION

General Experimental Procedures. Optical rotations were measured using an Autopol III automatic polarimeter (Rudolph Research Co.). UV spectra were detected by a UVIS-205 detector (Alltech). CD spectra were measured on a JASCO J-810/J-815 spectropolarimeter. IR spectra were recorded on a Thermo Nicolet Nexus 470 FT-IR spectrometer. NMR spectra were measured on a Bruker Avance-500 FT NMR spectrometer (500 MHz) using TMS as the internal standard. HRESIMS spectra were obtained on a PEQ-STAR ESI-TOF-MS/MS spectrometer. Column chromatography was performed on silica gel (200–300 mesh). The HF₂₅₄ silica gel for TLC was provided by Sigma Co. Ltd. Sephadex LH-20 (18–110 μ m) was obtained from Pharmacia Co., and ODS (50 μ m) was provided by YMC Co. HPLC was performed on an Alltech 426 apparatus with a UV detector. A YMC-Pack C8 column (ODS, 10 mm \times 250 mm, for reversed phase) and a chiral-phase column (OD-RH, 4.6 mm \times 150 mm) were used in the HPLC separation.

Animal Material. The fresh soft coral *Sinularia capillosa* was collected from the inner coral reef at a depth of around 10 m in Sanya Bay, Hainan Province, People's Republic of China, in May 2004, and the sample was immediately frozen after collection. The specimen was identified by one of the authors (L.v.O.). A voucher specimen (HSE-14) was deposited at the State Key Laboratory of Natural and Biomimetic Drugs, Peking University.

Extraction and Isolation. The frozen soft coral was homogenized and extracted with EtOH. The concentrated extract was desalted by dissolving in MeOH to obtain a residue that was further partitioned between H₂O and EtOAc. The EtOAc fraction (20.0 g) was subjected to VLC (vacuum liquid chromatography) using 200–300 mesh Si gel and eluted with a gradient of petroleum ether/acetone (50:1, 20:1, 10:1, 5:1, 3:1, 1:1) to get six fractions (F1–F6). Each fraction was examined by ¹H NMR spectroscopy, and sesquiterpenoids were recognized to exist in fractions F3–F5. F3 was further fractionated on an ODS column with a gradient of MeOH/H₂O (from 53% to 93%) followed by preparative RP-C8 HPLC to give **3** (0.7 mg), **4** (0.8 mg), **6** (0.9 mg), **7** (5.2 mg), and **8** (1.1 mg). F4 was subjected to flash chromatography on an ODS column with a gradient of MeOH/H₂O (from 60% to 80%) followed by semipreparative RP-C8 HPLC to give **1** (9.0 mg), **2** (4.6 mg), **5** (1.7 mg), **9** (10.6 mg), **13** (4.8 mg), **11** (6.4 mg), **12** (1.6 mg), **15** (0.7 mg), **16** (0.8 mg), **17** (4.5 mg), **18** (2.2 mg), and racemic sinularone A (9.6 mg). The enantiomers were separated on an OD-RH chiral-phase column using a mobile phase of H₂O/MeOH (2:8) to yield (–)-sinularone A (4.6 mg) and sinularone

A (5.0 mg). F5 was fractionated by repeated ODS column chromatography and preparative HPLC to give **10** (1.4 mg) and **14** (2.6 mg).

Capillosanane A (1): colorless oil; $[\alpha]_D^{26}$ –39 (c 0.5, MeOH); UV (MeOH) λ_{\max} (log ϵ) 204 (3.81), 289 (2.64) nm; IR (KBr) ν_{\max} 3488, 1700, 1470, 1432, 1361 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 259.1666 [M + Na]⁺ (calcd for C₁₅H₂₄O₂Na, 259.1674).

Capillosanane B (2): colorless oil; $[\alpha]_D^{24}$ –160 (c 0.8, MeOH); UV (MeOH) λ_{\max} (log ϵ) 214 (3.17), 236 (3.94) nm; IR (KBr) ν_{\max} 3330, 1652, 1458, 1370 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 259.1666 [M + Na]⁺ (calcd for C₁₅H₂₄O₂Na, 259.1674).

Capillosanane C (3): colorless oil; $[\alpha]_D^{27}$ –120 (c 0.1, MeOH); UV (MeOH) λ_{\max} (log ϵ) 217 (3.28), 234 (3.75) nm; IR (KBr) ν_{\max} 3427, 1635, 1457, 1380 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 259.1669 [M + Na]⁺ (calcd for C₁₅H₂₄O₂Na, 259.1674).

Capillosanane D (4): colorless oil; $[\alpha]_D^{27}$ –44 (c 0.2, MeOH); UV (MeOH) λ_{\max} (log ϵ) 213 (3.91), 265 (2.94) nm; IR (KBr) ν_{\max} 3437, 1694, 1655, 1460, 1370 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 257.1512 [M + Na]⁺ (calcd for C₁₅H₂₂O₂Na, 257.1518).

Capillosanane E (5): colorless oil; $[\alpha]_D^{26}$ –31 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 206 (4.21) nm, 303 (2.78); IR (KBr) ν_{\max} 3262, 1736, 1593, 1457, 1371 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 273.1460 [M + Na]⁺ (calcd for C₁₅H₂₂O₃Na, 273.1467).

Capillosanane F (6): colorless oil; $[\alpha]_D^{27}$ –10 (c 0.02, MeOH); UV (MeOH) λ_{\max} (log ϵ) 217 (3.89), 243 (3.34), 287 (3.16), 322 (2.42) nm; IR (KBr) ν_{\max} 3449, 1717, 1691, 1453, 1377, 1282 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 233.1535 [M + H]⁺ (calcd for C₁₅H₂₁O₂, 233.1541).

Capillosanane G (7): colorless oil; $[\alpha]_D^{18}$ –76 (c 0.5, MeOH); UV (MeOH) λ_{\max} (log ϵ) 219 (3.61) nm; IR (KBr) ν_{\max} 3450, 1642, 1457, 1378, 1043 cm^{–1}; CD (MeOH) λ (mdeg) 220 (–37) nm; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 275.1618 [M + Na]⁺ (calcd for C₁₅H₂₄O₃Na, 275.1623).

Capillosanane H (8): colorless oil; $[\alpha]_D^{27}$ –9 (c 0.2, MeOH); UV (MeOH) λ_{\max} (log ϵ) 217 (3.86) nm; IR (KBr) ν_{\max} 3446, 1653, 1459, 1369, 1163 cm^{–1}; CD (MeOH) λ (mdeg) 220(+30), 250(–10) nm; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 1 and 3; HRESIMS m/z 275.1618 [M + Na]⁺ (calcd for C₁₅H₂₄O₃Na, 275.1623).

Capillosanane I (9): colorless oil; $[\alpha]_D^{26}$ –55 (c 0.6, MeOH); UV (MeOH) λ_{\max} (log ϵ) 206 (4.21) nm; IR (KBr) ν_{\max} 3431, 1635, 1385, 1164 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 2 and 3; HRESIMS m/z 275.1612 [M + Na]⁺ (calcd for C₁₅H₂₄O₃Na, 275.1623).

Capillosanane J (10): colorless oil; $[\alpha]_D^{24}$ –51 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 206 (3.81) nm; IR (KBr) ν_{\max} 3440, 1653, 1464, 1360, 1064 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 2 and 3; HRESIMS m/z 291.1564 [M + Na]⁺ (calcd for C₁₅H₂₄O₄Na, 291.1573).

Capillosanane K (11): colorless oil; $[\alpha]_D^{22}$ +20 (c 1.0, MeOH); UV (MeOH) λ_{\max} (log ϵ) 205 (3.94) nm; IR (KBr) ν_{\max} 3421, 1647, 1454, 1376 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 2 and 3; HRESIMS m/z 275.1615 [M + Na]⁺ (calcd for C₁₅H₂₄O₃Na, 275.1623).

Capillosanane L (12): colorless oil; $[\alpha]_D^{21}$ –2 (c 0.16, MeOH); UV (MeOH) λ_{\max} (log ϵ) 206 (4.24) nm; IR (KBr) ν_{\max} 3418, 1639, 1475, 1387 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see Tables 2 and 3; HRESIMS m/z 259.1668 [M + Na]⁺ (calcd for C₁₅H₂₄O₂Na, 259.1674).

Capillosanane M (13): colorless oil; $[\alpha]_D^{23}$ –49 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 212 (2.96), 239 (3.85) nm; CD (c 0.3, MeOH) λ ($\Delta\epsilon$) 242(–2.0) nm; IR (KBr) ν_{\max} 3432, 1637, 1454, 1382, 1108 cm^{–1}; ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) data, see

Table 4. ^1H and ^{13}C NMR Data for Capillosananes O–R (15–18)^a

| no. | 15 | | 16 | | 17 | | 18 | |
|------|-----------------------|----------------------------------|-----------------------|-------------------------------|-----------------------|--|-----------------------|--|
| | δ_{C} | δ_{H} (J in Hz) | δ_{C} | δ_{H} (J in Hz) | δ_{C} | δ_{H} (J in Hz) | δ_{C} | δ_{H} (J in Hz) |
| 1 | 50.3, CH ₂ | 2.23, s | 50.3, CH ₂ | 2.23, s | 47.8, CH ₂ | 1.97, d (14.4) 1.59, dd (14.4, 2.0) | 138.3, CH | 5.56, d (1.50) |
| 2 | 208.6, C | | 208.5, C | | 92.7, C | | 144.5, C | |
| 3 | 65.8, CH | 3.56, m | 208.5, C | | 34.6, CH | 2.84, m | 85.7, C | |
| 4 | 39.7, CH ₂ | 1.33, m | 43.2, CH ₂ | 2.46, t (7.3) | 24.9, CH ₂ | 1.65, m 1.50, m | 32.9, CH ₂ | 2.37, m 2.06, m |
| 5 | 24.5, CH ₂ | 2.00, m | 22.4, CH ₂ | 2.17, q (7.3) | 30.0, CH ₂ | 2.51, m | 29.0, CH ₂ | 2.63, ddd (17.0, 9.0, 6.7) 2.46, ddd (16.9, 9.5, 7.0) |
| 6 | 127.2, CH | 5.23, t (7.1) | 125.9, CH | 5.17, t (7.2) | 170.3, C | | 176.6, C | |
| 7 | 131.9, C | | 132.7, C | | 198.4, C | | 208.0, C | |
| 8 | 26.8, CH ₂ | 2.76, d (15.5) 2.72, d (15.5) | 26.8, CH ₂ | 2.76, s | 125.8, CH | 6.41, s | 49.4, CH ₂ | 2.89, dd (17.0, 4.3) 2.36, dd (17.0, 9.8) |
| 9 | 140.4, C | | 140.2, C | | 159.5, C | | 39.7, CH | 3.27, m |
| 10 | 168.1, CH | 7.10, s | 168.3, CH | 7.13, s | 51.3, CH ₂ | 2.22, br d (15.4) 2.18, br d (15.4) | 46.8, CH ₂ | 1.97, dd (13.0, 8.4) 1.26, dd (13.0, 6.7) |
| 11 | 38.8, C | | 38.9, C | | 36.7, C | | 43.2, C | |
| 12 | 28.4, CH ₃ | 1.14, s | 28.4, CH ₃ | 1.15, s | 28.8, CH ₃ | 1.09, s | 29.8, CH ₃ | 1.06, s |
| 13 | 28.4, CH ₃ | 1.14, s | 28.4, CH ₃ | 1.15, s | 27.8, CH ₃ | 1.01, s | 28.7, CH ₃ | 0.98, s |
| 14 | 23.8, CH ₃ | 1.60, s | 23.7, CH ₃ | 1.58, s | 32.1, CH ₃ | 2.17, s | 30.6, CH ₃ | 2.10, s |
| 15 | 24.1, CH ₃ | 1.03, d (6.0) | 30.2, CH ₃ | 2.08, s | 16.5, CH ₃ | 0.85, d (6.9) | 26.8, CH ₃ | 1.43, s |
| 3-OH | | 4.37, d (4.7) | | | | | | |

^aRecorded in DMSO-*d*₆, at 500 MHz.

Tables 2 and 3; HRESIMS m/z 257.1509 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{22}\text{O}_2\text{Na}$, 257.1518).

Capillosanane N (14): colorless oil; $[\alpha]_{\text{D}}^{24} -5$ (c 0.5, MeOH); UV (MeOH) λ_{max} (log ϵ) 202 (3.78) nm; IR (KBr) ν_{max} 3435, 1634, 1461, 1384, 1077 cm^{-1} ; ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) data, see Tables 2 and 3; HRESIMS m/z 291.1563 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{24}\text{O}_4\text{Na}$, 291.1573).

Capillosanane O (15): colorless oil; UV (MeOH) λ_{max} (log ϵ) 221 (4.12) nm; IR (KBr) ν_{max} 1709, 1678, 1646, 1474, 1376 cm^{-1} ; ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) data, see Table 4; HRESIMS m/z 257.1511 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{22}\text{O}_2\text{Na}$, 257.1518).

Capillosanane P (16): colorless oil; $[\alpha]_{\text{D}}^{21} -1.8$ (c 0.14, MeOH); UV (MeOH) λ_{max} (log ϵ) 222 (4.32) nm; $\text{Rh}_2(\text{CF}_3\text{COO})_4$ -induced CD (c 0.4, CHCl_3) λ ($\Delta\epsilon$) 353 (+0.7) nm; IR (KBr) ν_{max} 3441, 1704, 1623, 1454, 1383, 1123 cm^{-1} ; ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) data, see Table 4; HRESIMS m/z 259.1669 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{24}\text{O}_2\text{Na}$, 259.1674).

Capillosanane Q (17): colorless oil; $[\alpha]_{\text{D}}^{27} -6.2$ (c 0.4, MeOH); UV (MeOH) λ_{max} (log ϵ) 211 (3.12), 237 (3.92) nm; IR (KBr) ν_{max} 1737, 1702, 1633, 1461, 1380, 1052 cm^{-1} ; CD (c 0.3, MeOH) λ ($\Delta\epsilon$) 220 (−27) nm; ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) data, see Table 4; HRESIMS m/z 273.1462 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{22}\text{O}_3\text{Na}$, 273.1467).

Capillosanane R (18): colorless oil; $[\alpha]_{\text{D}}^{27} -8.6$ (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 208 (3.83) nm; IR (KBr) ν_{max} 1723, 1634, 1440, 1382, 1113 cm^{-1} ; CD (c 0.5, MeOH) λ ($\Delta\epsilon$) 223 (−5) nm; ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) data, see Table 4; HRESIMS m/z 273.1461 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{22}\text{O}_3\text{Na}$, 273.1467).

(−)-Sinularone A: colorless oil; $[\alpha]_{\text{D}}^{15} -23.4$ (c 0.2, MeOH); CD (c 0.3, MeOH) λ ($\Delta\epsilon$) 300 (−0.56) nm. $[\alpha]_{\text{D}}^{15} +25.5$ (c 0.2, MeOH) is for the freshly separated sinularone A. The reported $[\alpha]_{\text{D}}^{23} +7.26$ (c 0.27, MeOH)²⁷ is attributed to a mixture of enantiomers.

Compound 7a: colorless oil; $[\alpha]_{\text{D}}^{27} -70$ (c 0.2, MeOH); IR (KBr) ν_{max} 3437, 1642, 1459, 1368, 1160 cm^{-1} ; ^1H NMR (500 MHz, DMSO-*d*₆) δ_{H} 11.31 (1H, s, 10-OH), 5.80 (1H, s, H-8), 4.70 (1H, d, J = 4.3 Hz, 6-OH), 4.55 (1H, m, H-6), 3.96 (1H, s, H-10), 2.36 (1H, d, J = 16.3 Hz, H-1a), 2.22 (1H, m, H-3), 1.80 (1H, d, J = 16.3 Hz, H-1b), 1.66 (3H, s, H₃-14), 1.09 (3H, s, H₃-12), 0.99 (3H, s, H₃-13), 0.95

(3H, d, J = 6.9 Hz, H₃-15); HRESIMS m/z 275.1616 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{24}\text{O}_3\text{Na}$, 275.1623).

Compound 9a: colorless oil; $[\alpha]_{\text{D}}^{23} +8.1$ (c 0.1, MeOH); IR (KBr) ν_{max} 3436, 1657, 1464, 1365 cm^{-1} ; ^1H NMR (500 MHz, DMSO-*d*₆) δ_{H} 6.19 (1H, s, 6-OH), 6.10 (1H, s, H-8), 5.00 (1H, s, H-14a), 4.81 (1H, s, H-14b), 2.23 (1H, d, J = 14.4 Hz, H-10a), 1.97 (1H, d, J = 14.4 Hz, H-10b), 1.78 (1H, m, H-4a), 1.77 (1H, m, H-5a), 1.75 (1H, d, J = 14.3 Hz, H-1a), 1.45 (1H, m, H-3), 1.41 (1H, d, J = 14.3 Hz, H-1b), 1.35 (1H, m, H-4b), 1.34 (1H, m, H-5b), 1.07 (3H, s, H₃-12), 1.03 (3H, d, J = 6.9 Hz, H₃-15), 0.94 (3H, s, H₃-13); ^{13}C NMR (125 MHz, DMSO-*d*₆) δ_{C} 149.0 (C, C-9), 146.6 (C, C-7), 122.1 (CH, C-8), 109.4 (CH₂, C-14), 97.6 (C, C-6), 85.2 (C, C-2), 50.2 (CH, C-1), 45.5 (CH₂, C-10), 36.4 (C, C-11), 33.4 (CH₂, C-5), 30.9 (CH, C-3), 30.6 (CH₃, C-13), 30.2 (CH₃, C-12), 26.1 (CH₂, C-4), 13.3 (CH₃, C-15); HRESIMS m/z 257.1510 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{22}\text{O}_2\text{Na}$, 257.1518).

Compound 9b: colorless oil; $[\alpha]_{\text{D}}^{23} +20$ (c 0.1, MeOH); IR (KBr) ν_{max} 3435, 1634, 1436, 1377, 1061 cm^{-1} ; ^1H NMR (500 MHz, DMSO-*d*₆) δ_{H} 6.25 (1H, s, 6-OH), 6.14 (1H, s, H-8), 5.26 (1H, s, H-10), 2.04 (1H, d, J = 13.1 Hz, H-1a), 1.47 (1H, d, J = 13.1 Hz, H-1b), 1.66 (3H, s, H₃-14), 1.62 (1H, m, H-3), 1.07 (3H, s, H₃-12), 1.08 (3H, d, J = 6.9 Hz, H₃-15), 1.05 (3H, s, H₃-13); ^{13}C NMR (125 MHz, DMSO-*d*₆) δ_{C} 141.1 (C, C-9), 139.0 (C, C-7), 132.4 (CH, C-10), 121.1 (CH, C-8), 98.8 (C, C-6), 88.3 (C, C-2), 50.3 (CH, C-1), 42.4 (C, C-11), 34.8 (CH, C-3), 31.5 (CH₃, C-13), 28.3 (CH₃, C-12), 27.6 (CH₂, C-4), 27.1 (CH₂, C-5), 18.6 (CH₃, C-14), 16.0 (CH₃, C-15); HRESIMS m/z 257.1510 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{22}\text{O}_2\text{Na}$, 257.1518).

Conversion of 9 to 18. To a MeOH/H₂O (3:2, 3 mL) solution of 9 (5.0 mg) was added NaIO₄ (10 mg), and the reaction mixture was stirred for 7 h at room temperature (rt). The solution was then diluted with H₂O (5 mL) and extracted with CHCl₃ (10 mL) to yield 18 (1.0 mg) after Si gel column chromatography. ^1H NMR (400 MHz, CDCl₃) δ 6.29 (1H, s, H-8), 3.09 (1H, m, H-3), 2.75 (1H, m), 2.59 (1H, dd, J = 17.3, 6.2 Hz) 2.24 (3H, s, H₃-14), 1.13 (6H, s, H₃-12, 13), 0.92 (3H, d, J = 6.9 Hz, H₃-15).

Reduction of Capillosanane A (1) to 1a. Compound 1 (5.1 mg) was treated with NaBH₄ (10 mg) in MeOH (2 mL) by stirring at rt for 60 min. The reaction was quenched by the addition of H₂O (2 mL), and the mixture was evaporated in vacuo. The residue was purified by

reversed-phase semipreparative HPLC using MeCN/H₂O (65:35) as the mobile phase to obtain **1a** (3.1 mg).

Compound 1a: colorless oil; $[\alpha]_D^{25} +40$ (c 0.1, MeOH); ¹H NMR (CD₃OD, 500 MHz) δ_H 5.55 (1H, s, H-8), 3.49 (1H, dd, *J* = 10.5, 2.5 Hz, H-6), 3.09 (1H, dd, *J* = 13.5, 6.4 Hz, H-2), 2.20 (1H, d, *J* = 13.4 Hz, H-10a), 1.96 (1H, m, H-4a), 1.89 (1H, t, *J* = 13.4 Hz, H-5a), 1.78 (1H, d, *J* = 13.4 Hz, H-10b), 1.66 (1H, m, H-3), 1.65 (1H, m, H-5b), 1.64 (1H, m, H-1a), 1.61 (1H, m, H-1b), 1.36 (3H, s, Me-14), 1.20 (1H, m, H-4b), 1.11 (3H, s, Me-12), 0.92 (3H, s, Me-13), 0.83 (3H, d, *J* = 6.8 Hz, Me-15); ¹H NMR (CDCl₃, 400 MHz) δ_H 5.58 (1H, s, H-8), 3.51 (1H, dd, *J* = 10.5, 2.5 Hz, H-6), 2.96 (1H, m, H-2), 2.17 (1H, d, *J* = 13.4 Hz, H-10a), 1.79 (1H, d, *J* = 13.4 Hz, H-10b), 1.41 (3H, s, Me-14), 1.13 (1H, m, H-4b), 1.11 (3H, s, Me-12), 0.91 (3H, s, Me-13), 0.81 (3H, d, *J* = 6.8 Hz, Me-15); ¹³C NMR (CD₃OD, 125 MHz) δ_C 143.2 (C, C-9), 130.2 (CH, C-8), 78.7 (CH, C-6), 75.9 (C, C-7), 53.0 (CH₂, C-10), 43.8 (CH₂, C-1), 39.8 (CH, C-2), 36.3 (C, C-11), 35.3 (CH, C-3), 31.9 (CH₂, C-4), 28.2 (CH₂, C-5), 28.0 (CH₃, C-13), 27.9 (CH₃, C-12), 20.4 (CH₃, C-14), 16.6 (CH₃, C-15).

Preparation of (R)- and (S)-MPA Esters of 1a. To a CDCl₃ solution (0.5 mL) containing **1a** (1.4 mg, 0.0059 mmol) were added (R)-MPA (2.0 mg, 0.012 mmol), DMAP (0.8 mg, 0.006 mmol), and *N,N'*-dicyclohexylcarbodiimide (DCC, 2.5 mg, 0.012 mmol). After reacting at rt for 24 h, the crude products were separated by silica gel column chromatography eluting with petroleum ether/acetone (7:1) to afford the (R)-MPA ester (1.5 mg). By the same protocol, the (S)-MPA ester (1.3 mg) was prepared from (S)-MPA.

(R)-MPA ester of 1a: ¹H NMR (CDCl₃, 400 MHz) δ_H 5.382 (1H, s, H-8), 4.784 (1H, dd, *J* = 9.2, 2.2 Hz, H-6), 2.935 (1H, m, H-2), 2.154 (1H, d, *J* = 13.6 Hz, H-10a), 1.942 (1H, m, H-4a), 1.770 (1H, d, *J* = 13.7 Hz, H-10b), 1.277 (1H, m, H-4b), 1.124 (3H, s, Me-14), 1.086 (3H, s, Me-12), 0.871 (3H, s, Me-13), 0.800 (3H, d, *J* = 6.8 Hz, Me-15); ESIMS *m/z* 387.25 [M + H]⁺.

(S)-MPA ester of 1a: ¹H NMR (CDCl₃, 400 MHz) δ_H 5.443 (1H, s, H-8), 4.874 (1H, dd, *J* = 9.2, 2.2 Hz, H-6), 2.769 (1H, m, H-2), 2.154 (1H, d, *J* = 13.6 Hz, H-10a), 1.828 (1H, m, H-4a), 1.799 (1H, d, *J* = 13.2 Hz, H-10b), 1.180 (1H, m, H-4b), 1.338 (3H, s, Me-14), 1.077 (3H, s, Me-12), 0.888 (3H, s, Me-13), 0.741 (3H, d, *J* = 6.8 Hz, Me-15); ESIMS *m/z* 387.25 [M + H]⁺.

Preparation of (R)- and (S)-MPA Esters of 2. Following the same protocol as that for **1a**, the (R)-MPA ester (1.3 mg) and the (S)-MPA ester (1.2 mg) were obtained from **2** (each 0.8 mg, 0.0032 mmol).

Preparation of (R)- and (S)-MPA Esters of 5. Following the same protocol as that for **1a**, the (R)-MPA ester (0.9 mg) and the (S)-MPA ester (0.8 mg) were obtained from **5** (each 0.8 mg, 0.0032 mmol).

(R)-MPA ester of 2: ¹H NMR (CDCl₃, 400 MHz) δ_H 5.865 (1H, m, H-6), 5.701 (1H, s, H-8), 5.336 (1H, s, H-10), 2.258 (1H, m, H-3), 2.158 (1H, d, *J* = 16.7 Hz, H-1a), 2.097 (1H, d, *J* = 16.7 Hz, H-1b), 1.574 (3H, s, Me-14), 0.983 (3H, d, *J* = 6.8 Hz, Me-15), 0.931 (3H, s, Me-12), 0.591 (3H, s, Me-13); ESIMS *m/z* 385.23 [M + H]⁺.

(S)-MPA ester of 2: ¹H NMR (CDCl₃, 400 MHz) δ_H 5.862 (1H, m, H-6), 5.346 (1H, s, H-10), 5.116 (1H, s, H-8), 2.318 (1H, m, H-3), 2.261 (1H, d, *J* = 15.8 Hz, H-1a), 2.151 (1H, d, *J* = 15.8 Hz, H-1b), 1.129 (3H, s, Me-12), 1.108 (3H, s, Me-14), 1.021 (3H, s, Me-13), 0.976 (3H, d, *J* = 6.8 Hz, Me-15); ESIMS *m/z* 385.23 [M + H]⁺.

(R)-MPA ester of 5: ¹H NMR (CDCl₃, 400 MHz) δ_H 5.849 (1H, s, H-8), 5.572 (1H, dd, *J* = 12.1, 4.2 Hz, H-6), 2.292 (1H, d, *J* = 14.3 Hz, H-1a), 2.069 (1H, m, H-3), 1.835 (1H, d, *J* = 14.3 Hz, H-1b), 1.707 (3H, s, Me-14), 1.161 (3H, d, *J* = 6.8 Hz, Me-15), 1.152 (3H, s, Me-12), 1.104 (3H, s, Me-13); ESIMS *m/z* 399.23 [M + H]⁺.

(S)-MPA ester of 5: ¹H NMR (CDCl₃, 400 MHz) δ_H 5.775 (1H, s, H-8), 5.603 (1H, dd, *J* = 12.3, 4.7 Hz, H-6), 2.292 (1H, d, *J* = 14.3 Hz, H-1a), 2.086 (1H, m, H-3), 1.835 (1H, d, *J* = 14.3 Hz, H-1b), 1.381 (3H, s, Me-14), 1.183 (3H, d, *J* = 6.8 Hz, Me-15), 1.144 (3H, s, Me-12), 1.108 (3H, s, Me-13); ESIMS *m/z* 399.23 [M + H]⁺.

Reduction of Capillosanane D (4) to 2 and 3. Compound **4** (0.6 mg) was stirred with NaBH₄ (1.0 mg) in MeOH (2 mL) at rt for 45 min. The reaction was quenched by addition of H₂O (2 mL), and the mixture was evaporated in vacuo. The residue was separated by

reversed-phase HPLC using MeCN/H₂O (61:39) as eluent to yield **2** (0.3 mg) and **3** (0.25 mg).

Preparation of (R)- and (S)-MPA Esters of 10. Following the same protocol as that for **1a**, the (R)-MPA ester (2.3 mg) and the (S)-MPA ester (2.0 mg) were obtained from **10** (each 1.8 mg).

(R)-MPA ester of 10: ¹H NMR (CDCl₃, 400 MHz) δ_H 6.072 (1H, s, H-8), 4.944 (1H, s, H-10), 1.254 (3H, s, Me-14), 0.978 (3H, s, Me-12), 0.741 (3H, s, Me-13); ESIMS *m/z* 417.22 [M + H]⁺.

(S)-MPA ester of 10: ¹H NMR (CDCl₃, 400 MHz) δ_H 6.047 (1H, s, H-8), 4.914 (1H, s, H-10), 1.039 (3H, s, Me-12), 1.009 (3H, s, Me-14), 0.873 (3H, s, Me-13); ESIMS *m/z* 417.22 [M + H]⁺.

Bioassay against Larval Settlement (refs 31, 32). Adults of the barnacle *Balanus amphitrite* Darwin were exposed to air for more than 6 h and then were placed in a container filled with fresh 0.22 μ m filtered seawater to release nauplii. The collected nauplii were reared to the cyprid stage according to the method described by Thiyagarajan et al.²⁶ When kept at 26–28 °C and fed with *Chaetoceros gracilis*, larvae developed to cyprids within four days. Fresh cyprids were used in the tests.

Assay for the Production of Pro-inflammatory Cytokines (TNF- α) (ref 33). Compounds were dissolved in DMSO, and NK-007 was used as a positive control.²⁸ Suspensions of BALB/c mouse splenocytes were cultured in complete RPMI 1640 medium (Hyclone Laboratories) containing 10% FBS (Gibco). Raw 264.7 cells were grown in DMEM (Hyclone Laboratories) supplemented with 10% FBS. Cultured cells were treated in the presence or absence of NK-007 (100 nM) and tested compounds (10 μ M) for 24 h followed by LPS stimulation (1 μ g/mL). Supernatants were collected for analysis of cytokine levels. For cytokine determinations in culture supernatants or serum, cytometric bead array kits (BD Biosciences) were used.

Cytotoxic Assay. Cell viability in the presence or absence of tested compounds was determined using the standard MTT assay as described.³⁴

■ ASSOCIATED CONTENT

⑤ Supporting Information

NMR spectroscopic data for the new compounds (**1–18**) including ¹H, ¹³C, and 2D NMR spectra, IR, and ESIMS/MS data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Tel: ++86-10-82806188. Fax: ++86-10-82806188. E-mail: whlin@bjmu.edu.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by grants from NSFC (No. 30672607), the National Hi-Tech 863-Projects (2010DFA31610, 2011AA090701, 2013AA092902), and Sino-German Project GZ816.

■ REFERENCES

- (1) San Feliciano, A.; Barrero, A. F.; Medarde, M.; de Corral, J. M.; Aramburu, A.; Perales, A.; Fayos, J. *Tetrahedron Lett.* **1985**, 26, 2369–2372.
- (2) El-Dahmy, S.; Jakupovic, J.; Bohlmann, F.; Sarg, T. M. *Tetrahedron* **1985**, 41, 309–316.
- (3) Catalan, C. A. N.; de Lampasona, M. E. P.; Cerda-Garcia-Rojas, C. M.; Joseph-Nathan, P. J. *Nat. Prod.* **1995**, 58, 1713–1717.
- (4) Fricke, C.; Hardt, I. H.; Koenig, W. A.; Joulain, D.; Zygadlo, J. A.; Guzman, C. A. *J. Nat. Prod.* **1999**, 62, 694–696.
- (5) Mao, S.-C.; Gavagnin, M.; Mollo, E.; Guo, Y.-W. *Biochem. Syst. Ecol.* **2011**, 39, 408–411.

- (6) Su, J.-H.; Huang, C.-Y.; Li, P.-J.; Lu, Y.; Wen, Z.-H.; Kao, Y.-H.; Sheu, J.-H. *Arch. Pharm. Res.* **2012**, *35*, 779–784.
- (7) Huang, C.-Y.; Su, J.-H.; Liu, Y.-C.; Wen, Z.-H.; Hsu, C.-H.; Chiang, M. Y.; Sheu, J.-H. *Bull. Chem. Soc.* **2010**, *83*, 678–682.
- (8) Su, J.-H.; Chiang, M. Y.; Wen, Z.-H.; Dai, C.-F.; Hsu, C.-H.; Sheu, J.-H. *Chem. Pharm. Bull.* **2010**, *58*, 250–253.
- (9) Jin, P.; Deng, Z.-W.; Pei, Y.-H.; Lin, W.-H. *Zhongguo Haiyang Yaowu* **2011**, *30*, 11–14.
- (10) Ramesh, P.; Ravikanth, V.; Venkateswarlu, Y. *Indian J. Chem.* **2001**, *40B*, 867–868.
- (11) Zhang, C.-X.; Zhu, C.-C.; Yan, S.-J.; Li, J.; Su, J.-Y.; Liang, Y.-J.; Yang, X.-P.; Zheng, K.-C.; Zeng, L.-M. *J. Asian Nat. Prod. Res.* **2008**, *10*, 277–280.
- (12) Braekman, J. C.; Daloze, D.; Tursch, B.; Hull, S. E.; Declercq, J. P.; Germain, G.; Van Meerssche, M. *Experientia* **1980**, *36*, 893–893.
- (13) Tseng, Y.-J.; Shen, K.-P.; Lin, H.-L.; Huang, C.-Y.; Dai, C.-F.; Sheu, J.-H. *Mar. Drugs* **2012**, *10*, 1572–1581.
- (14) Su, J.-H.; Hsieh, C.-H.; Lo, C.-L.; Huang, C.-Y.; Dai, C.-F.; Kuo, Y.-H.; Sheu, J.-H. *J. Chin. Chem. Soc.* **2008**, *55*, 1286–1289.
- (15) Chao, C.-H.; Hsieh, C.-H.; Chen, S.-P.; Lu, C.-K.; Dai, C.-F.; Sheu, J.-H. *Tetrahedron Lett.* **2006**, *47*, 5889–5891.
- (16) Cheng, S.-Y.; Huang, K.-J.; Wang, S.-K.; Wen, Z.-H.; Hsu, C.-H.; Dai, C.-F.; Duh, C.-Y. *Org. Lett.* **2009**, *11*, 4830–4833.
- (17) Cheng, S.-Y.; Huang, K.-J.; Wang, S.-K.; Duh, C.-Y. *Mar. Drugs* **2011**, *9*, 1469–1476.
- (18) Cheng, S.-Y.; Huang, K.-J.; Wang, S.-K.; Wen, Z.-H.; Chen, P.-W.; Duh, C.-Y. *J. Nat. Prod.* **2010**, *73*, 771–775.
- (19) Su, J.; Yang, R.; Kuang, Y.; Zeng, L. *J. Nat. Prod.* **2000**, *63*, 1543–1545.
- (20) Freire, F.; Calderon, F.; Seco, J. M.; Fernandez-Mayoralas, A.; Quinoa, E.; Riguera, R. *J. Org. Chem.* **2007**, *72*, 2297–2301.
- (21) Gerards, M.; Snatzke, G. *Tetrahedron: Asymmetry* **1990**, *1*, 221–236.
- (22) Frelek, J.; Szczepek, W. *J. Tetrahedron: Asymmetry* **1999**, *10*, 1507–1520.
- (23) Ono, M.; Tsuru, T.; Abe, H.; Eto, M.; Okawa, M.; Abe, F.; Kinjo, J.; Ikeda, T.; Nohara, T. *J. Nat. Prod.* **2006**, *69*, 1417–1420.
- (24) Beecham, A. F.; Mathieson, A. McL.; Johns, S. R.; Lamberton, J. A.; Sioumis, A. A.; Batterham, T. J.; Young, I. G. *Tetrahedron* **1971**, *27*, 3725–3738.
- (25) Bari, L. D.; Pescitelli, G.; Pratelli, C.; Pini, D.; Salvadori, P. *J. Org. Chem.* **2001**, *66*, 4819–4825.
- (26) Chen, D.; Yu, S.; van Ofwegen, L.; Proksch, P.; Lin, W. *J. Agric. Food Chem.* **2012**, *60*, 112–123.
- (27) Shi, H.; Yu, S.; Liu, D.; van Ofwegen, L.; Proksch, P.; Lin, W. *Mar. Drugs* **2012**, *10*, 1331–1344.
- (28) Porter, N. A.; Wujek, J. S. *J. Org. Chem.* **1987**, *52*, 5085–5089.
- (29) Brill, W. F. *J. Chem. Soc., Perkin Trans.* **1984**, *11*, 621–627.
- (30) Bowden, B. D.; Coil, J. C.; Silva, E. D.; Costa, M. S. L.; Djura, P. J.; Mahendran, M.; Tapiolas, D. M. *Aust. J. Chem.* **1983**, *36*, 371–376.
- (31) Thiyagarajan, V.; Harder, T.; Qian, P. *J. Exp. Mar. Biol. Ecol.* **2003**, *287*, 223–236.
- (32) Dobretsov, S.; Xiong, H.; Xu, Y.; Levin, L. A.; Qian, P. *Mar. Biotechnol.* **2007**, *9*, 388–397.
- (33) Ti, W.; Li, Y.; Wu, M.; Sun, X.; Bao, X.; Lin, Y.; Hao, J.; Han, L.; Cao, G.; Wang, Z.; Liu, Y.; Wu, Z.; Hong, Z.; Wang, P.; Zhao, L.; Li, Z.; Wang, Q.; Yin, Z. *Arthritis Rheum.* **2012**, *64*, 2896–2906.
- (34) Li, F.; Awale, S.; Tezuka, Y.; Kadota, S. *Bioorg. Med. Chem.* **2008**, *16*, 5434–5440.