

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/233740831>

Alkaloids of Anatolian *Thalictrum* sultanabadense

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · JULY 1985

Impact Factor: 3.8 · DOI: 10.1021/np50040a036

CITATIONS

2

READS

10

3 AUTHORS, INCLUDING:



K. Husnu Can Baser

Near East University

1,074 PUBLICATIONS 10,098 CITATIONS

SEE PROFILE

ALKALOIDS OF ANATOLIAN *THALICTRUM SULTANABADENSE*

KEMAL HÜSNÜ CAN BAŞER,* MUZAFFER ÖĞÜTVEREN

Anadolu University, Faculty of Pharmacy, Department of Pharmacognosy, Eskişehir, Turkey

and NORMAN G. BISSET

Chelsea College, Department of Pharmacy, Pharmacognosy Research Laboratories, University of London, Manresa Rd., London SW3 6LX, U.K.

Thalictrum sultanabadense Stapf (Ranunculaceae) is a small, perennial plant found growing near the town of Kemaliye (Erzincan) in Eastern Turkey. The occurrence of thalbadensine, hernandezine, hernandezine *N*-oxide, and thalidezine in *T. sultanabadense* growing in the USSR has been reported (1, 2). Here, we report on the isolation of thalbadensine, hernandezine, thalictine, thalifoline, and two more alkaloids—for which uv and tlc comparison with authentic samples strongly suggest their identification as berberine and magnoflorine—from the roots and above ground parts of *T. sultanabadense* of Anatolian origin (Table 1). The last four alkaloids are reported to be isolated for the first time from this species.

TABLE 1. Alkaloids Isolated from Anatolian *Thalictrum sultanabadense*

Alkaloid	Above ground parts	Roots	Identified by
Thalbadensine . .	+ ^a	+	uv, ¹ H nmr, ms
Thalictine	+	+	uv, ¹ H nmr, ms
Thalifoline	+	+	uv, ¹ H nmr, ms, tlc comparison
Hernandezine . . .	+	—	uv, ¹ H nmr, ms
Berberine	—	+	uv, tlc comparison
Magnoflorine . . .	+	+	uv, tlc comparison

^a+ = detected, — = not detected.

EXPERIMENTAL

PLANT MATERIAL.—Whole plants were collected from Sandık Village near Kemaliye (Erzincan) in Eastern Turkey in June 1983. A voucher specimen is kept at the Herbarium of the Faculty of Pharmacy, Anadolu University (Turkey).

EXTRACTION AND ISOLATION.—Dried roots (170 g) and above ground parts (245 g) were separately extracted and worked up by a previously reported procedure (3). The alkaloids were isolated and purified by column and thin layer chromatographic techniques. Details of the isolation and identification of the alkaloids are available on request to the first author.

ACKNOWLEDGMENTS

This work was supported by NATO research grant No. 029.81. We thank Dr. R. W. Doskotch for a sample of thalifoline.

LITERATURE CITED

1. S. Abdizhabbarova, S. Kh. Maekh, S. Yu. Yunusov, *Khim. Prir. Soedin*, 139 (1978).
2. S. Abdizhabbarova, S. Kh. Maekh, S. Yu. Yunusov, *Khim. Prir. Soedin*, 250 (1981).
3. K.H.C. Başer, *Doğa, seri A*, 5, 163 (1981).

Received 4 February 1985