

# THE SURFACE TENSIONS OF TERNARY SOLUTIONS. PART I. THE SURFACE TENSIONS OF AQUEOUS SOLUTIONS OF (a) SODIUM AND POTASSIUM CHLORIDES, (b) SODIUM CHLORIDE AND HYDROCHLORIC ACID.

BY J. W. BELTON.

*Received 7th June, 1935.*

The addition of an electrolyte to water usually produces an increase in surface tension which is given by the relation

$$\gamma - \gamma_0 = km, \quad (1)$$

where  $\gamma$  and  $\gamma_0$  are the surface tensions of the solution and of water respectively,  $k$  is a constant, and  $m$  is the number of moles of solute present per 1000 gms. of water. When two electrolytes are present the change in surface tension is usually taken as equal to the sum of the changes each would produce separately, and is given by

$$\gamma_{12} - \gamma_0 = k_1 m_1 + k_2 m_2, \quad (2)$$

where  $\gamma_{12}$  is the surface tension of the solution and the subscripts 1 and 2 refer to the two solutes. This relation may be conveniently tested by keeping the concentration of one solute constant and varying that of the other. A plot of  $\gamma_{12}$  against  $m$  should give

$$\left( \frac{\partial \gamma_{12}}{\partial m_1} \right)_{m_2} = k_1. \quad (3)$$

$$\left( \frac{\partial \gamma_{12}}{\partial m_2} \right)_{m_1} = k_2. \quad (4)$$

These considerations are applied below to solutions of sodium and potassium chlorides and of hydrochloric acid and sodium chloride.

## Experimental.

The surface tensions of the solutions were obtained by the maximum bubble pressure method, the apparatus used being similar to that of Brown<sup>1</sup> to which reference should be made for complete details. Dry air was stored under pressure in a reservoir over mercury and was allowed to pass slowly over phosphorus pentoxide, through a glass wool trap, through a finely adjustable valve and into the experimental cell (Fig. 1) at A, and thence via B to a large reservoir kept at constant temperature. C was connected with a reservoir of butyl phthalate, the level of which could be easily changed, thus giving a fine adjustment of the level of the liquid in the cell. The cell at A and B was also in communication with the limbs of a manometer containing butyl phthalate, by means of which the pressure required to force a single bubble through the surface of the liquid could be measured with a cathetometer placed at a suitable distance. Both cell and manometer were kept in a thermostat at 25° (± 0.02). At

<sup>1</sup> *Phil. Mag.*, 1932, 13, 578.

## 1414 THE SURFACE TENSIONS OF TERNARY SOLUTIONS

each measurement the jet was arranged to be just coincident with the surface of the liquid, and when the bubble had passed through the surface the contact was broken. Some difficulty was experienced in the cleaning of the cell and jet. Before each fresh solution was examined, it was washed out with hot chromic acid and then with a large quantity of water; steam was then blown into it for about five minutes and finally it was dried by evacuation at 100°. The jet was made by drawing down a piece of quill tubing, breaking off at the constricted part and examining under a microscope until one satisfactory was obtained.

The surface tension was calculated from

$$\gamma = \frac{rg}{2} \left( h\rho - \frac{2}{3}r\rho_1 \right) \quad (5)$$

where  $r$  is the radius of the jet = 0.0102 cm.




FIG. 1.

$g = 981$  cm./sec.<sup>2</sup>

$h$  = maximum bubble pressure in cm. butyl phthalate.

$\rho$  = density of butyl phthalate at 25° = 1.0434 gms./c.c.

$\rho_1$  = density of liquid under observation, found by a Westphal balance.

The surface tension in dyne/cm. of pure water found in this way was 72.01 at 25°, and that of pure benzene at the same temperature 28.20. These values are in good agreement with those generally accepted.

### Binary Solutions.

The results obtained for binary solutions containing sodium chloride, potassium chloride and hydrochloric acid are given in Table I. The final column gives the value of the constant,  $k$ , calculated from equation (1). The surface tensions of hydrochloric acid solutions decrease with increasing concentration, while those of sodium and potassium chlorides increase.

TABLE I.—SURFACE TENSIONS OF BINARY SOLUTIONS.

	$m$ .	$h$ .	$\gamma$ .	$\Delta\gamma$ .	$k$ .
Sodium Chloride Solutions	1.021	14.15	73.84	1.75	1.73
	2.080	14.48	75.59	3.58	1.72
	3.197	14.84	77.44	5.43	1.70
	4.360	15.22	79.43	7.42	1.70
	5.608	15.62	81.54	9.53	1.70
Potassium Chloride Solutions	1.000	14.12	73.68	1.67	1.67
	2.010	14.43	75.34	3.33	1.65
	3.051	14.70	76.73	4.72	1.55
	4.015	14.96	78.14	6.13	1.53
Hydrochloric Acid Solutions.	1.00	13.77	71.87	—0.14	—0.14
	3.00	13.73	71.64	—0.37	—0.12
	5.00	13.68	71.43	—0.58	—0.11

### Ternary Solutions.

(a) **Aqueous Solutions of Sodium Chloride and Potassium Chloride.**—The mixtures were made up by weight from B.D.H. AR. chemicals and pure water of surface tension 72.01 dynes/cm. The surface tensions measured are given in Table II. together with the molal concentrations of the salts. The fifth and sixth columns give ( $\gamma_{12} - \gamma_0$ ) observed, and calculated from (2). The agreement is good, but there is a discrepancy which is greater than the experimental error.

TABLE II.—SOLUTIONS OF SODIUM AND POTASSIUM CHLORIDES.

$m_1$ .	$m_2$ .	$h$ .	$\gamma_{12}$ .	$\Delta\gamma_{12}$ obs.	$\Delta\gamma_{12}$ calc.
1.01	1.05	14.435	75.35	3.34	3.55
1.01	2.07	14.735	76.91	4.90	5.24
1.01	3.09	15.015	78.37	6.36	6.51
1.01	4.12	15.24	79.54	7.53	8.09
2.02	1.00	14.725	76.86	4.85	5.15
2.02	2.01	14.98	78.46	6.45	6.81
2.02	3.00	15.26	79.97	7.96	8.03
3.01	1.01	15.05	78.55	6.54	6.80
3.01	2.01	15.39	80.15	8.34	8.44
3.00	3.00	15.62	81.54	9.53	9.67
4.02	1.00	15.38	80.30	8.29	8.50
4.02	2.01	15.72	82.08	10.07	10.17

Plots of  $\gamma_{12}$  against  $m$  for constant sodium chloride concentration and increasing potassium chloride, and for constant potassium chloride and increasing sodium chloride gave a series of nearly parallel straight lines. According to equations (3) and (4) a series of parallel straight lines should be obtained on both graphs, of slope  $k_2$  and  $k_1$ . The slopes ( $k'_1$  and  $k'_2$ ) for the various mixtures are given in Table III.; these values should be compared with those given in Table I. for sodium and potassium chloride alone. The curves for the mixtures with one component constant appear

TABLE III.—SOLUTIONS OF SODIUM AND POTASSIUM CHLORIDES.

$m_2 = 1.$			$m_2 = 2.$			$m_2 = 3.$		
$m_1$ .	$\Delta\gamma$ .	$k'_1$ .	$m_1$ .	$\Delta\gamma$ .	$k'_1$ .	$m_1$ .	$\Delta\gamma$ .	$k'_1$ .
1	1.67	1.67	1	1.81	1.81	1	1.64	1.64
2	3.18	1.59	2	3.36	1.68	2	3.24	1.62
3	4.87	1.62	3	5.05	1.68	3	4.81	1.61
4	6.62	1.65	4	6.98	1.75		Mean	1.62
	Mean	1.63		Mean	1.73			


  

$m_1 = 1.$			$m_1 = 2.$			$m_1 = 3.$		
$m_2$ .	$\Delta\gamma$ .	$k'_2$ .	$m_2$ .	$\Delta\gamma$ .	$k'_2$ .	$m_2$ .	$\Delta\gamma$ .	$k'_2$ .
1	1.61	1.61	1	1.41	1.41	1	1.44	1.44
2	3.17	1.59	2	3.01	1.50	2	3.04	1.52
3	4.63	1.54	3	4.52	1.51	3	4.43	1.48
	Mean	1.57		Mean	1.50		Mean	1.48

## 1416 THE SURFACE TENSIONS OF TERNARY SOLUTIONS

TABLE IV.—SOLUTIONS OF SODIUM CHLORIDE AND HYDROCHLORIC ACID.

$m_1$ .	$m_2$ .	$h$ .	$\gamma_{12}$ .	$\Delta\gamma_{12}$ .	$\Delta\gamma_{12}$ calc. from (2).	$\Delta\gamma_{12}$ calc. from (6).
0.99	1.0	14.01	73.12	1.11	1.63	1.13
1.98	1.0	14.22	74.23	2.22	3.27	2.28
2.95	1.0	14.45	75.42	3.41	4.87	3.40
3.98	1.0	14.69	76.68	4.67	6.63	4.64
4.96	1.0	14.86	77.57	5.56	8.29	5.81
1.06	2.0	13.96	72.86	0.85	1.64	0.89
2.00	2.0	14.14	73.78	1.77	3.16	1.75
3.01	2.0	14.31	74.68	2.67	4.84	2.72
4.00	2.0	14.47	75.54	3.53	6.55	3.83
1.00	3.0	13.91	72.59	0.58	1.44	0.57
2.01	3.0	14.07	73.43	1.42	3.10	1.36
3.02	3.0	14.23	74.28	2.27	4.76	2.12
0.99	4.0	13.88	72.43	0.42	1.29	0.30
1.99	4.0	14.02	73.15	1.14	2.95	0.96
1.00	5.0	13.84	72.26	0.25	1.24	0.12

FIG. 2.—Plot of Molality HCl —  $\gamma_{12}$  for NaCl — HCl mixtures.

to be slightly less steep than those with one component alone, although in one case they are equally steep (NaCl in 2MKCl). The contribution of each solute to the surface tension is thus a little less than it would have been in a separate solution.

(b) **Aqueous Solutions of Sodium Chloride and Hydrochloric Acid.**—The solutions were made up by weight from pure chemicals and the same water as before. The strength of the hydrochloric acid was verified by standardisation. Table IV. gives the molal concentrations of sodium chloride ( $m_1$ ) and hydrochloric acid ( $m_2$ ) and the corresponding surface tension. The fifth and sixth columns give the surface tension increments observed and calculated from (2). Comparison of these shows that the simple additive relation is only very approximate and that the discrepancies are much greater than those obtained for solutions of sodium and potassium chlorides. The plot of  $\gamma_{12}$  against  $m_1$  ( $m_2$  constant) gives a series of straight lines, but they are not parallel as would be expected from (3); the plot of  $\gamma_{12}$  against  $m_2$  ( $m_1$  constant) gives a series of curves again showing no agreement with (4) (Fig. 2). Examination of the data showed

that  $\gamma_{12}$  could be represented by the relation

$$\gamma_{12} - \gamma_0 = k_1 m_1 + k_2 m_2 + k_3 m_1 m_2^{\frac{1}{2}} \quad (6)$$

in which  $k_3$  is an empirical constant equal to  $-0.5$ . The values of  $\Delta\gamma_{12}$  so calculated are given in the final column of Table IV. The agreement with the observed values is very good—a difference of one in the first place represents an error in the surface tension of one in seven hundred. The slopes of the curves for  $\gamma_{12}$  against  $m_1$  ( $m_2$  constant) and  $\gamma_{12}$  against  $m_2$  ( $m_1$  constant) are thus given by

$$\left(\frac{\partial \gamma_{12}}{\partial m_1}\right)_{m_2} = k_1 + k_3 m_2^{\frac{1}{2}} \quad \left(\frac{\partial \gamma_{12}}{\partial m_2}\right)_{m_1} = k_2 + \frac{1}{2} k_3 m_1 m_2^{-\frac{1}{2}} \quad (7)$$

### Application of the Gibbs Equation.

The values of  $(\partial \gamma_{12}/\partial m_1)_{m_2}$  and  $(\partial \gamma_{12}/\partial m_2)_{m_1}$ , which are calculable from (2) and from (6), may also be derived from the Gibbs equation, according to which

$$d\gamma = -\Gamma_1 d\mu_1 - \Gamma_2 d\mu_2 - \Gamma_3 d\mu_3, \quad (8)$$

where  $\Gamma_1$ ,  $\Gamma_2$ ,  $\Gamma_3$  are the surface concentrations of the components and  $\mu_1$ ,  $\mu_2$ ,  $\mu_3$  their chemical potentials. If the surface is chosen so that  $\Gamma_3 = 0$ , then

$$d\gamma = -\Gamma_1 d\mu_1 - \Gamma_2 d\mu_2 = -\Gamma_1 2RT d \log f_1 m_1 - \Gamma_2 2RT d \log f_2 m_2, \quad (9)$$

where  $f_1$  and  $f_2$  are the appropriate activity coefficients of the solutes. If the concentration of one is kept constant and the other varied, then this may be written

$$-\frac{1}{2RT} \left(\frac{\partial \gamma_{12}}{\partial m_1}\right)_{m_2} = \Gamma_1 \left(\frac{1}{m_1} + \left(\frac{\partial \log f_1}{\partial m_1}\right)_{m_2}\right) + \Gamma_2 \left(\frac{\partial \log f_2}{\partial m_1}\right)_{m_2} \quad (10)$$

$$-\frac{1}{2RT} \left(\frac{\partial \gamma_{12}}{\partial m_2}\right)_{m_1} = \Gamma_2 \left(\frac{1}{m_2} + \left(\frac{\partial \log f_2}{\partial m_2}\right)_{m_1}\right) + \Gamma_1 \left(\frac{\partial \log f_1}{\partial m_2}\right)_{m_1} \quad (11)$$

For only one solute (subscript 0)

$$-\frac{1}{2RT} \frac{d\gamma}{dm} = -\frac{k}{2RT} = {}_0\Gamma \left(\frac{1}{m} + \frac{d \log f}{dm}\right). \quad (12)$$

Equations (3) and (4) and consequently (2) are thus special cases of a more general equation. A complete solution of equations (10) and (11) requires a knowledge of the variations of the activity coefficient of each component of the system, data not usually available. In certain cases, as below, approximations may be made and the adsorptions calculated.

#### (a) Aqueous Solutions of Sodium and Potassium Chlorides.

It may be assumed for this system that  $(\partial \log f_1/\partial m_1)_{m_2} = d \log f_1/dm_1$ . Combining (10) and (12), and (11) and (12) and putting  $k_1$  for the slope in the mixed solution, we have

$$-\frac{k'_1}{2RT} = -\frac{\Gamma_1}{{}_0\Gamma_1} \cdot \frac{k_1}{2RT} + \Gamma_2 \left(\frac{\partial \log f_2}{\partial m_1}\right)_{m_2} \quad (13)$$

$$-\frac{k'_2}{2RT} = -\frac{\Gamma_2}{{}_0\Gamma_2} \cdot \frac{k_2}{2RT} + \Gamma_1 \left(\frac{\partial \log f_1}{\partial m_2}\right)_{m_1} \quad (14)$$

## 1418 THE SURFACE TENSIONS OF TERNARY SOLUTIONS

For the lower concentrations the  $\partial \log f / \partial m$  terms are small and may be neglected. Hence

$$\Gamma_1 = \frac{k'_1}{k_1} {}_0\Gamma_1 \quad (15)$$

Even for 5*M* solutions the error introduced is only a few per cent. Similarly.

$$\Gamma_2 = \frac{k'_2}{k_2} {}_0\Gamma_2 \quad (16)$$

The approximate values of  $\Gamma_1$  and  $\Gamma_2$  calculated in this way are given in Table V. The sum  $\Gamma_1 + \Gamma_2$  gives the total negative adsorption at the surface. The values of  ${}_0\Gamma_1 + {}_0\Gamma_2$  for separate solutions, given for comparison, have been calculated from (12), the values of  $d \log f / dm$  being obtained from the empirical equations of Harned.<sup>2</sup>

It appears that for this system each solute is hardly affected by the presence of the other, and that each is adsorbed to the same extent in the mixed solution as it would be in a separate solution. If the layer of water at the surface is assumed to be unimolecular, its thickness,  $\delta$ , is given by  $1000 (\Gamma_1 + \Gamma_2) / (m_1 + m_2)$ . These values are given in the final column of Table V. and the result, as for separate solutions of sodium and potassium chlorides, is about the diameter of a water molecule, with a tendency to become less at higher concentrations, probably due to a change in orientation.

TABLE V.—SOLUTIONS OF SODIUM AND POTASSIUM CHLORIDES.

$m_1$	$m_2$	$\Gamma_1$	$\Gamma_2$	$\Gamma_1 + \Gamma_2$	${}_0\Gamma_1 + {}_0\Gamma_2$	$\delta(\text{\AA}^\circ)$
1	1	0.39	0.35	0.74	0.79	3.7
1	2	0.41	0.65	1.06	1.10	3.5
1	3	0.38	0.89	1.27	1.29	3.2
1	4	0.38	1.07	1.45	1.46	2.9
2	1	0.60	0.34	0.94	1.01	3.1
2	2	0.63	0.63	1.26	1.32	3.1
2	3	0.59	0.85	1.44	1.51	2.9
3	1	0.78	0.34	1.12	1.19	2.8
3	2	0.82	0.62	1.44	1.50	2.9
3	3	0.77	0.83	1.60	1.69	2.7
4	1	0.92	0.34	1.26	1.34	2.5
4	2	0.97	0.62	1.59	1.64	2.6

$\Gamma$  in moles/sq. cm.  $\times 10^{10}$ .

**(b) Aqueous Solutions of Sodium Chloride and Hydrochloric Acid.**

For this system  $\partial \log f_2 / \partial m_1$  may be computed, but not  $\partial \log f_1 / \partial m_2$ . In order, then, to calculate either  $\Gamma_1$  or  $\Gamma_2$  the other must be known. Now it appears that the relative values of the terms in (10), calculated below, are such that changes in  $\Gamma_2$  may be neglected. It may thus be assumed that  $\Gamma_2 = {}_0\Gamma_2$  and  $\Gamma_1$  calculated. A very large change in  $\Gamma_2$  would be required to produce only a small error in  $\Gamma_1$  calculated in this way. Further it must be assumed that  $d \log f / dm$  is the same in both mixed and separate solutions; this will introduce a small error.

<sup>2</sup> *J. Amer. Chem. Soc.*, 1920, **42**, 1808; 1922, **44**, 252.

Equation (10) may now be written (using (7) and (12))

$$-\frac{1}{2RT}(k_1 + k_3 m_2^{\frac{1}{2}}) = -\frac{k_1}{2RT} \cdot \frac{\Gamma_1}{\Gamma_1} + \Gamma_2 \left( \frac{\partial \log f_2}{\partial m_1} \right)_{m_2},$$

or

$$1 + \frac{k_3 m_2^{\frac{1}{2}}}{k_1} + \Gamma_2 \frac{2RT}{k_1} \cdot \left( \frac{\partial \log f_2}{\partial m_1} \right)_{m_2} = \frac{\Gamma_1}{\Gamma_1}. \quad (18)$$

The values of  $k_1$  (taken as 1.7) and  $k_3$  are known,  $\Gamma_2$  is put equal to  $\Gamma_2$  calculated from the data in Table I. The values of  $(\partial \log f_2 / \partial m_1)_{m_2}$  were calculated from the data of Hawkins<sup>3</sup> who found for mixtures of sodium chloride and hydrochloric acid up to total concentrations of 6M

$$\log f = -\frac{0.354\sqrt{2C_T}}{1 + A\sqrt{2C_T}} + B(2C_1) + B'(2C_T - 2C_1) - \log(1 - 0.002wm_T),$$

where  $C_T$  is the total molar concentration,  $C_1$  the molar acid concentration, and  $A$ ,  $B$  and  $B'$  are determined constants. The values of  $\Gamma_1/\Gamma_1$  calculated in this way are given in Table VI. The term involving  $\Gamma_2$  is quite small throughout.

TABLE VI.—SOLUTIONS OF SODIUM CHLORIDE AND HYDROCHLORIC ACID.

$m_1$	$m_2$	$\Gamma_2$	$\left( \frac{\partial \log f_2}{\partial m_1} \right)_{m_2}$	$\frac{\Gamma_1}{\Gamma_1}$	$\delta$	$\delta_{12}$
1	1	$2.44 \times 10^{-12}$	0.0646	0.72	4.1	3.0
2	1	$2.44 \times 10^{-12}$	0.0697	0.72	3.2	2.3
3	1	$2.44 \times 10^{-12}$	0.0689	0.72	2.7	1.9
4	1	$2.44 \times 10^{-12}$	0.0662	0.72	2.4	1.7
5	1	$2.44 \times 10^{-12}$	0.0627	0.72	2.1	1.5
1	2	$3.48 \times 10^{-12}$	0.0755	0.60	4.1	2.4
2	2	$3.48 \times 10^{-12}$	0.0745	0.60	3.2	1.9
3	2	$3.48 \times 10^{-12}$	0.0714	0.60	2.7	1.6
4	2	$3.48 \times 10^{-12}$	0.0679	0.60	2.4	1.4
1	3	$3.86 \times 10^{-12}$	0.0799	0.51	4.1	2.1
2	3	$3.86 \times 10^{-12}$	0.0769	0.51	3.2	1.6
3	3	$3.86 \times 10^{-12}$	0.0728	0.51	2.7	1.4
1	4	$4.25 \times 10^{-12}$	0.0826	0.44	4.1	1.8
2	4	$4.25 \times 10^{-12}$	0.0783	0.44	3.2	1.4
1	5	$4.14 \times 10^{-12}$	0.0838	0.37	4.1	1.5

The thickness of the adsorbed water layer, assuming it to be uni-molecular, is given under  $\delta$  for the corresponding pure sodium chloride solution, and in the last column under  $\delta_{12}$  for the mixtures ( $\delta = 1000\Gamma/m$ ). The effect of the addition of hydrochloric acid is to reduce the thickness of this layer. The increased adsorption of HCl molecules seems to displace water molecules from the surface. An increase in the concentration of sodium chloride reduces the thickness of the water layer, but HCl is more effective than NaCl.

### Summary.

The surface tensions of aqueous solutions of sodium and potassium chlorides, and of sodium chloride and hydrochloric acid, have been measured. The variation of the surface tension with the concentration of each component is discussed, and the surface adsorption of water calculated by means of the Gibbs equation.

Physical Chemistry Department,  
The University, Leeds.

<sup>3</sup> J. Amer. Chem. Soc., 1932, 54, 4480.