

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7892538>

Sesquiterpenoids from *Pulicaria canariensis* and Their Cytotoxic Activities

ARTICLE in JOURNAL OF NATURAL PRODUCTS · MAY 2005

Impact Factor: 3.8 · DOI: 10.1021/np0496183 · Source: PubMed

CITATIONS

16

READS

40

8 AUTHORS, INCLUDING:



Francisco J. Pérez

University of Chile

50 PUBLICATIONS 1,046 CITATIONS

SEE PROFILE



Javier Gonzalez-Platas

Universidad de La Laguna

129 PUBLICATIONS 911 CITATIONS

SEE PROFILE



José Quintana

Universidad de Las Palmas de Gran Canaria

53 PUBLICATIONS 814 CITATIONS

SEE PROFILE



Francisco León

University of Mississippi

106 PUBLICATIONS 790 CITATIONS

SEE PROFILE

Sesquiterpenoids from *Pulicaria canariensis* and Their Cytotoxic Activities[#]

Jorge Triana,^{†,‡} Mariana López,^{†,‡} Francisco J. Pérez,^{†,‡} Javier González-Platas,[§] José Quintana,^{‡,⊥} Francisco Estévez,^{‡,⊥} Francisco León,^{‡,||} and Jaime Bermejo^{*,||}

Departamento de Química, Universidad de Las Palmas de Gran Canaria, Campus de Tafira, 35017 Las Palmas de Gran Canaria, Spain, Instituto Canario de Investigación del Cáncer, Avenida Astrofísico F. Sánchez 2, 38206 La Laguna, Tenerife, Spain, Instituto Universitario de Bio-Organica "Antonio González", Servicio de Difracción de Rayos X (SIDIX), Departamento Física Fundamental II, Universidad de La Laguna, Avenida Astrofísico F. Sánchez, E-38204 La Laguna, Tenerife, Spain, Departamento de Bioquímica y Biología Molecular, Centro de Ciencias de la Salud, Universidad de Las Palmas de Gran Canaria, Avenida S. Cristóbal, 35016 Las Palmas de Gran Canaria, Spain, and Instituto de Productos Naturales y Agrobiología, C.S.I.C., Instituto Universitario de Bio-Organica "Antonio González", Avenida Astrofísico F. Sánchez 3, 38206 La Laguna, Tenerife, Spain

Received November 23, 2004

Thirteen new sesquiterpenes, pulicanadiene A (**1**), B (**2**), and C (**3**), pulicanone (**4**), pulicanol (**5**), pulicanarals A (**6**), B (**7**), and C (**8**), pulicanadienals A (**9**) and B (**10**), pulicanadienol (**11**), and pulioplopanones A (**12**) and B (**13**), and seven known compounds, stigmaterol, ergosterol peroxide, calenduladiol, 7,4'-di-*O*-methylhydrokaempferol, 5,7-dihydroxy-3,3',4'-trimethoxyflavone, dihydroquercetin 7,3'-dimethyl ether, and 6,15 α -epoxy-1 β ,4 β -dihydroxyeudesmane, were isolated from *Pulicaria canariensis*. Compound **4a** showed cytotoxicity on the human myeloid leukemia cell line HL-60. The cytotoxicity was caused by induction of apoptosis as determined by microscopy of nuclear changes, activation of caspases, and the cleavage of poly(ADP-ribose) polymerase-1.

The genus *Pulicaria* (Compositae) is represented in the Canary Islands by three species distributed in the more easterly islands, namely, *P. crispa* Forssk., *P. burchardii* Hutch., and *P. canariensis* Bolle. The last two are endemic, with *P. canariensis* having two subspecies, *canariensis* and *lanata*.¹ Several species of this genus have been used as insect repellents² and in the treatment of dysentery.^{3,4} This genus has been the subject of several chemical investigations, giving rise to the isolation of flavonoids,⁵ sesquiterpenes,^{6,7} and triterpenes.⁸ As a part of our continuing search for novel, plant-derived cancer chemotherapeutic agents,^{9,10} we have investigated an ethanolic extract of the aerial part of *P. canariensis* Bolle subsp. *canariensis*, leading to the isolation of 13 new sesquiterpenes, pulicanadienes A (**1**), B (**2**), and C (**3**), pulicanone (**4**), pulicanol (**5**), pulicanarals A (**6**), B (**7**), and C (**8**), pulicanadienals A (**9**) and B (**10**), pulicanadienol (**11**), and pulioplopanones A (**12**) and B (**13**), and seven known compounds, stigmaterol,¹¹ ergosterol peroxide,¹² calenduladiol,¹³ 7,4'-di-*O*-methylhydrokaempferol,¹⁴ 5,7-dihydroxy-3,3',4'-trimethoxyflavone,¹⁵ dihydroquercetin 7,3'-dimethyl ether,¹⁶ and 6,15 α -epoxy-1 β ,4 β -dihydroxyeudesmane.¹⁷ The structures of the known compounds were confirmed by comparison of their spectroscopic properties with published data. The structure elucidation of the new compounds was carried out by extensive spectral data interpretation as well as by chemical transformation. To assess whether these compounds display cytotoxic properties, we also studied their effects on the viability of the human promyelocytic cell line HL-60, using the colorimetric MTT procedure.

Results and Discussion

Pulicanadiene A (**1**) was isolated as an amorphous powder, $[\alpha]_D^{25}$ 21.4° (c 0.014, CHCl₃), and its molecular formula was determined as C₁₉H₂₈O₆ by HREIMS (*m/z* 352.1877 [M]⁺). The ¹H and ¹³C NMR spectra of compound **1** (Table 1) carried out at 25 °C presented poor resolution owing to the existence of a conformational equilibrium,¹⁸ but heating to 50 °C sharpened most lines and allowed assignment of the signals. Thus, characteristic signals for three methyl groups were observed at δ_H 0.97 (3H, d, *J* = 4.5 Hz, CH₃-13), 1.06 (3H, d, *J* = 5.6 Hz, CH₃-12), and 1.84 (3H, s, CH₃-15), two acetoxy groups at δ_H 2.00 (3H, s, OAc) and 2.05 (3H, s, OAc), two olefinic protons at δ_H 5.18 (1H, d, *J* = 8.9 Hz, H-5) and 6.78 (1H, t, *J* = 8.0 Hz, H-1), and two signals at δ_H 5.77 (1H, d, *J* = 9.1 Hz, H-6) and 5.63 (1H, br s, H-8) corresponding to the geminal protons to the acetoxy groups. The ¹³C NMR and DEPT spectra of **1** disclosed 19 carbons, which were indicative of an α,β -unsaturated carboxylic acid (δ_C 173.0, s, C-14; 137.4, s, C-10; 141.6, d, C-1), two oxygenated methines (δ_C 70.0, d, C-6; 70.1, d, C-8), an isopropyl group (δ_C 26.7, d, C-11; 22.4, q, C-12; 21.8, q, C-13), in addition to two acetoxy groups, three methylenes, two other methines, and one other quaternary carbon. These NMR data were closely comparable to those of the known germacrane diacetate tovarol (**14**)¹⁹ and suggested that compound **1** is also a germacrane.¹⁵ The IR absorption at 3200–3067 and 1693 cm⁻¹ confirmed the presence of an α,β -unsaturated carboxylic acid. In fact, methylation with diazomethane yielded **1a**, in the ¹H NMR spectrum (Table 1) of which appears a new singlet at δ_H 3.73 (3H, s, COOMe). Comparison of **1a** and **14** led us to conclude that acetoxy groups could be placed at C-6 and C-8 in **1a**. The ¹H and ¹³C NMR spectral features found in both compounds suggest that they are similar. This was deduced from the following 2D NMR data of **1a**, in which the α,β -unsaturated acid group was supported by the HMBC correlations H-1 with C-10 and C-14; H₂-9 with C-1, C-8, C-10, and C-14; CH₃-15 with C-3, C-4, and C-5; CH₃-12 with C-7; and CH₃-13 with C-7.

[#] Dedicated to the memory of Professor Antonio G. González.

^{*} To whom correspondence should be addressed. Tel: (34)-922-318583. Fax: (34)-922-318571. E-mail: jbermejo@ull.es.

[†] Departamento de Química, Universidad de Las Palmas de Gran Canaria.


[‡] Instituto Canario de Investigación del Cáncer.

[§] Servicio de Difracción de Rayos X (SIDIX).

[⊥] Departamento de Bioquímica, Universidad de Las Palmas de Gran Canaria.

^{||} Instituto de Productos Naturales y Agrobiología.

Chart 1


**Table 1.** ¹H and ¹³C NMR Data for Compounds 1, 1a, 2, and 3 (benzene-*d*₆ at 50 °C, *J* values (Hz) in parentheses)

position	1		1a		2		3	
	δ _C	δ _H	δ _C	δ _H	δ _C	δ _H	δ _C	δ _H
1	141.6	6.78 t (8.0)	138.6	6.60 t (8.0)	150.4	5.86 t (8.4)	126.7	5.34 t (8.5)
2	26.7	2.10 m 2.35 m	29.6	2.12 m 2.32 m	27.1	1.70 m 1.80 m	26.0	1.70 m 2.01 m
3	36.5	1.80 m 2.10 m	37.0	1.80 m 2.12 m	36.8	1.70 m	31.0	1.40 m
4	132.2		132.8		136.6		136.1	
5	123.7	5.18 d (8.9)	123.3	5.17 d (9.1)	124.3	5.11 d (8.6)	127.0	5.11 m
6	70.0	5.77 d (9.1)	69.9	5.77 d (9.2)	70.0	6.02 dd (1.8, 8.8)	71.0	6.13 d br (5.9)
7	52.2	1.80 m	53.0	1.80 m	52.6	1.80 m	50.0	1.92 dd br (3.3, 13.4)
8	70.1	5.63 s br	70.6	5.60 s br	70.1	6.09 s br	71.1	5.76 t (9.2)
9	31.6	2.35 m 2.66 d (11.2)	32.0	2.32 m 2.66 dd (4.0, 13.6)	30.2	2.34 s br 2.61 dd (5.3, 14.0)	30.0	2.50 m 2.68 m
10	137.3		137.4		143.6		135.0	
11	26.7	2.10m	26.5	2.12 m	26.7	1.80 m	27.0	2.12 t (6.3)
12	22.4	1.06 d (5.6)	22.2	1.02 d (5.3)	22.6	1.28 d (6.3)	19.9	1.13 d (6.8)
13	21.8	0.97 d (4.5)	21.7	0.97 d (5.3)	21.6	1.09 d (6.3)	21.4	0.97 d (6.9)
14	173.0		168.7		193.6	9.26 s	69.7	4.12 d br (17.8)
15	20.6	1.84 s	20.5	1.83 s	20.6	1.98 s	20.0	1.52 s
OMe			50.8	3.73 s				
OAc	18.0	2.00 s	20.4	1.99 s	17.9	1.77 s	21.5	1.78
	20.6	2.05 s	20.5	2.05 s	20.5	1.72 s	21.5	1.85
	169.1		168.7		169.0		170.2	
	169.1		168.7		169.0		170.2	

NOESY correlations between H-6 (δ_H 5.77) and CH₃-15, H-7 clearly showed that these protons were on the same face. Correlations were also observed between H-8 (δ_H 5.63) and CH₃-12, CH₃-13, and H-5. Thus, the C-6 acetoxy and C-7 isopropyl groups were deduced to be α correlated. This was also confirmed when similar correlations were observed in a NOESY experiment at −50 °C. From the above-mentioned spectra it could be deduced that the preferred conformers in solution maintain a similar orientation in space, and therefore compound **1** was assigned the structure (1*E*,4*E*)-6α,8β-diacetoxy-7β*H*-germacra-1(10),4-dien-14-oic acid, which was unequivocally confirmed by an X-ray crystallographic analysis of **1a** (Figure 1).

Pulicanadiene B (**2**) was obtained as a colorless oil, with [α]_D²⁵ 35.7° (c 0.042, CHCl₃). The molecular formula of **2** was determined as C₁₉H₂₈O₅, one oxygen atom less than obtained for **1**, on the basis of its HRFABMS data (*m/z* 359.1874 [M + Na]⁺). Both the ¹H and ¹³C NMR (Table 1) spectral data of compound **2** were close to those of **1**, but differences were evident between these compounds since the carboxyl group at C-14 of **1** is replaced with an aldehyde group in **2**, δ_H 9.26 (1H, s, H-14). Therefore, the structure of **2** was elucidated as (1*E*,4*E*)-6α,8β-diacetoxy-7β*H*-germacra-1(10),4-dien-14-al.


Pulicanadiene C (**3**) was also isolated as a colorless oil with [α]_D²⁵ −75.0° (c 0.008, CHCl₃). The molecular formula

**Figure 1.** ORTEP drawing of compound **1a**.**Table 2.** ^1H and ^{13}C NMR Data for Compound **4a** (CDCl_3 , J values (Hz) in parentheses)

position	δ_{C}	δ_{H}
1	131.9	5.59 dd (5.7, 11.4)
2	23.3	2.00 m
3	32.6	2.07 m
		2.94 d br, (12.7)
4	145.4	
5	198.7	
6	72.9	5.95 s
7	45.5	2.54 dd (3.4, 10.5)
8	69.6	5.41 dt (3.5, 10.6)
9	30.0	2.32 dd (3.5, 8.9)
		2.44 td (4.3, 12.6)
10	132.6	
11	27.4	2.27 m
12	21.6	1.12 d (6.9)
13	17.6	0.75 d (6.9)
14	69.3	4.51 s
15	124.1	5.66 s
OAc	20.8	2.03 s
	20.9	2.04 s
	21.1	2.19 s
	169.7	
	169.9	
	170.6	

$\text{C}_{19}\text{H}_{30}\text{O}_5$ was determined from EIMS and HREIMS analysis. The IR spectrum of **3** showed absorption bands at 3414 and 1728 cm^{-1} due to hydroxyl and acetoxy functions. The ^1H NMR (Table 1) spectrum of **3** showed signals analogous to those of **2** with the exception that the C-14 is replaced with a hydroxymethyl group, δ_{H} 4.12 (2H, d, J = 11 Hz, H_2 -14), in **3**. Thus, **2** was treated with sodium borohydride (NaBH_4) in methanol to give **3** as the major product. The IR and ^1H NMR spectra of both compounds were superimposable. On the basis of the above spectral and chemical evidence, **3** was assigned as (1*E*,4*E*)-6 α ,8 β -diacetoxy-7 β *H*-germacra-1(10),4-dien-14-ol.

Pulicanone (**4**) was purified as its triacetate (**4a**) by treatment with acetic anhydride (Ac_2O) in pyridine: **4a** [colorless oil, $[\alpha]_{\text{D}}^{25}$ 4.9° (c 0.081, CHCl_3)]. The IR spectrum showed the presence of acetoxy functions (1741 cm^{-1}) and a conjugated carbonyl group (1694 cm^{-1}). In the UV spectrum of **4a**, an absorption maximum was observed at 225 nm, suggestive of an α,β -unsaturated carbonyl group. The ^1H and ^{13}C NMR (Table 2) spectra of **4a** showed signals assignable to the methyl groups of an isopropyl group at δ_{H} 0.75 (3H, d, J = 6.9 Hz, H_3 -13) and 1.12 (3H, d, J = 6.9 Hz, H_3 -12) and three olefinic protons at δ_{H} 5.59 (1H, dd, J = 5.7, 11.4 Hz, H-1) and 4.51 (2H, s br, H_2 -14). In the ^1H - ^1H COSY spectrum of **4a**, H-6 (δ_{H} 5.95) correlated to H-7 (δ_{H} 2.54) and the latter to H-8 (δ_{H} 5.41), suggesting that the acetoxy groups are located at C-6 and C-8. To determine the position of the carbonyl group, the HMBC spectrum of

**Figure 2.** Selected ROESY correlations for compound **4a**.**Table 3.** ^1H and ^{13}C NMR Data for Compound **5a** (CDCl_3 , J values (Hz) in parentheses)

position	δ_{C}	δ_{H}
1	135.0	5.64 t (7.5)
2	24.4	2.36 m
3	38.4	2.19 m
		1.25 m
4	58.8	
5	66.5	2.87 d (6.6)
6	73.5	4.89 d (6.8)
7	47.5	1.63 d (9.2)
8	72.1	5.46 dd (4.7, 12.1)
9	29.6	2.63 m
		1.27 m
10	129.3	
11	26.2	1.86 m
12	22.9	1.12 d (6.5)
13	21.2	0.92 d (6.6)
14	63.5	4.62 d (12.26)
		4.57 d (12.12)
15	16.3	1.17 s
OAc	20.8	2.00 s
	21.0	2.07 s
	21.1	2.09 s
	169.8	
	169.8	
	170.8	

4a was useful. Thus, connectivities observed in the signals at δ_{H} 4.51 (H_2 -14) and 5.95 (H-6) correlated with the carbonyl and helped situate that carbonyl group at C-5. The relative stereochemistry of **4a** was determined by a ROESY experiment (Figure 2) in which correlations were observed between H-7 (δ 2.54), H-6 (δ 5.95), and H_2 -14 (δ 4.51). Thus, the C-6 acetoxy and C-7 isopropyl groups, as in **1**, possess an α -orientation. Therefore, the structure of **4** was determined to be (1*E*)-6 α ,8 β ,14-trihydroxy-7 β *H*-germacra-1(10),4(15)-dien-5-one.

Pulicanol (**5**) was isolated as a colorless oil. The EIMS of **5** showed a molecular ion at m/z 312 $[\text{M}]^+$. The molecular formula $\text{C}_{17}\text{H}_{28}\text{O}_5$ of **5** was determined from molecular ion measurement. The IR spectrum of **5** showed absorption bands at 3404, 1736, 1647, and 897 cm^{-1} assigned to hydroxyl, acetoxy, and olefinic functions, respectively. The ^1H NMR spectrum of **5** showed the presence of three methyls at δ_{H} 0.95 (3H, d, J = 6.4 Hz, CH_3 -13), 1.12 (3H, d, J = 6.4 Hz, CH_3 -12), and 1.16 (3H, s, CH_3 -15), one acetoxy group at δ_{H} 2.15 (3H, s, OAc), an olefinic proton at δ_{H} 5.45 (1H, t, J = 6.8 Hz, H-1), one hydroxy methylene group at δ_{H} 4.29 (1H, d, J = 11.8 Hz, H_a -14) and 4.12 (1H, d, J = 11.8 Hz, H_b -14), and three oxygenated methine protons at δ_{H} 2.82 (1H, d, J = 7.7 Hz, H-5), 3.51 (1H, m, H-6), and 5.18 (1H, dd, J = 5.2, 12.0 Hz, H-8). To obtain a better resolution in the ^1H and ^{13}C NMR spectra, acetylation (Ac_2O -pyridine) of **5** yielded an acetoxy derivative (**5a**). The ^{13}C NMR spectrum (Table 3) of **5a** disclosed 21 carbons, which were indicative of an oxygenated methylene (δ_{C} 63.5, t, C-14), an epoxide functional group (δ_{C} 66.5, d, C-5; 58.8, s, C-4), two oxygenated methines (δ_{C} 73.5, d, C-6; 72.1, d, C-8), and an isopropyl group (δ_{C} 26.2, d, C-11; 22.9, q, C-12; 21.2, q, C-13). An analysis of 2D NMR data revealed that **5a** possesses the same skeleton as **3**, with a




Figure 3. Selected HMBC (H to C) correlations for compound **5a**.

methyl-bearing trisubstituted epoxide occurring in **5a** instead of the methyl-bearing trisubstituted olefin in **3**. HMBC correlations (Figure 3) of CH₃-15 with C-4 and C-5, H-5 with C-3 and C-4, and H-7 with C-5, C-6, C-8, C-12, and C-13 helped position the methyl-bearing trisubstituted epoxide. The relative stereochemistry of **5a** was deduced from a 2D ROESY experiment, which indicated correlations of CH₃-15 (δ_{H} 1.17) with H-7 and H-6, clearly showing that these protons were on the same face, while H-5 (δ_{H} 2.87) correlated with H-8 on the opposite side of the molecule, indicating that the 4,5-epoxide ring was in *trans* form. Due to its large size, the conformational movement of this 10-membered ring is probably restricted.²⁰ The above confirmed the structure of **5** as (1*E*)-4 α ,5 β -epoxy-8 β -acetoxy-7 β *H*-germacra-1(10)-ene-6 α ,14-diol.

Pulicanaral A (**6**) was isolated as colorless needles, mp 183–184 °C and $[\alpha]_{\text{D}}^{25}$ 11.1° (*c* 0.108, CHCl₃), and its molecular formula was determined as C₁₉H₂₈O₆ by HREIMS (m/z 352.1905 [M]⁺). The ¹³C NMR spectrum (Table 4) gave a total of 19 separate carbon resonances (five methyls, three methylenes, seven methines, and four quaternary carbons), in agreement with the molecular formula. The ¹H NMR spectrum (Table 4) of compound **6** displayed signals for three methyl groups at δ_{H} 0.97 (3H, d, *J* = 6.6 Hz, CH₃-13), 1.32 (3H, d, *J* = 6.6 Hz, CH₃-12), and 1.45 (3H, s, CH₃-15), two acetoxy groups at δ_{H} 2.00 (3H, s, OAc) and 2.08 (3H, s, OAc), one olefinic proton at δ_{H} 5.43 (1H, d, *J* = 7.2 Hz, H-5), and one aldehyde proton at δ_{H} 9.28 (1H, s, H-14). The ¹H NMR spectra of **6** and **2** were similar with regard to their functional group analysis. However, compound **6** has only one olefinic proton. This led us to conclude that one of the two double bonds is epoxidized. In fact, its ¹³C NMR spectrum showed signals (δ_{C} 63.9, d, C-1; 61.1, s, C-10) corresponding to an epoxide

ring. In the HMBC spectrum of **6**, the signals at δ_{H} 9.28 (H-14) correlated with the oxygenated quaternary carbon (δ_{C} 61.1, C-10), which indicated that an epoxide ring was attached at C-1 (10). To establish the relative stereochemistry of **6**, a ROESY experiment was carried out. A correlation was observed between H-1 and H-8, while correlations observed between H-6, H-7 and CH₃-15 suggested that they have an opposite orientation. From these data together with an X-ray diffraction analysis (Supporting Information) the stereochemistry of the substituents was observed as well as the *trans* relationship between the methyl group at C-4 and the H at C-5 and of the aldehyde group at C-10 and the H at C-1. From these data the new compound **6** was deduced to be (4*E*)-1 β ,10 α -epoxy-6 α ,8 β -diacetoxy-7 β *H*-germacra-4-en-14-al.

Compound **7**, pulicanaral B, was purified as colorless needles by recrystallization from chloroform and methanol and has the molecular formula C₁₉H₂₈O₇ as determined by HRFABMS (m/z 391.1738 [M + Na]⁺) and NMR spectroscopy. The ¹H NMR data of **7** were very similar to those of **6**, suggesting that the two compounds are closely related in structure. The major difference is that **7** has an additional epoxide functional group. A detailed comparison of the ¹H and ¹³C NMR signals of **6** and **7** (Table 4) revealed that the signals due to H-5 (δ_{H} 5.43, d, *J* = 7.2 Hz), C-4 (δ_{C} 133.6), and C-5 (δ_{C} 129.7) of **6** have been replaced by signals due to an epoxide group [H-5 (δ_{H} 3.04, d, *J* = 6.7 Hz), C-4 (δ_{C} 58.0), C-5 (δ_{C} 66.2)]. The structure was shown to be 1 β ,10 α ;4 α ,5 β -diepoxy-6 α ,8 β -diacetoxy-7 β *H*-germacra-14-al, and again, only the relative configuration could be determined by an X-ray diffraction analysis (Supporting Information).

Compound **8**, pulicanaral C, was obtained as a colorless oil. The ¹H and ¹³C NMR spectra (Table 4) were similar to those of **7**, with the exception of the absence of an acetoxy signal since the oxygenated methine proton of **7** at δ_{H} 4.87 (1H, dd, *J* = 1.0, 6.7 Hz, H-6) was displaced to δ_{H} 3.51 (1H, dd, *J* = 1.2, 6.7 Hz, H-6) in **8** and one of two acetyl methyl signals in **7** disappeared in **8**. Acetylation (Ac₂O–pyridine) of **8** gave a diacetoxy derivative whose physical and spectroscopical constants were identical with those of **7**.

Table 4. ¹H and ¹³C NMR Data for Compounds **6**–**8** (CDCl₃, *J* values (Hz) in parentheses)



position	6		7		8	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}
1	63.9	3.34 d (11.6)	62.3	3.50 d (11.2)	62.5	3.45 d (11.5)
2	23.7	1.75 m	23.1	1.70 m	23.0	1.75 m
		2.38 m		2.38 m		2.40 m
3	35.9	2.33 m	35.7	1.29 m	35.8	1.30 m
		2.44 m		2.22 ddd (3.5, 7.1, 13.8)		2.20 ddd (3.5, 7.0, 13.5)
4	133.6		58.0		58.0	
5	129.7	5.43 d (7.2)	66.2	3.04 d (6.7)	67.7	3.03 d (6.7)
6	71.2	5.54 d (6.1)	72.5	4.87 dd (1.0, 6.7)	70.2	3.51 dd (1.2, 6.7)
7	51.4	1.58 d br (9.3)	47.9	1.52 m	49.7	1.54 m
8	68.8	5.40 dd (4.2, 12.6)	69.2	5.44 dd (4.6, 12.1)	70.0	5.20 dd (4.6, 12.6)
9	34.6	2.79 t (12.8)	33.3	2.84 t (12.6)	34.5	3.01 t (12.6)
		1.43 d (4.20)		1.55 m		1.54 m
10	61.1		61.8		61.5	
11	26.3	1.96 m	26.3	1.95 m	26.3	1.90 m
12	22.0	1.32 d (6.6)	21.9	1.29 d (6.7)	22.1	1.31 d (6.6)
13	21.1	0.97 d (6.6)	21.0	0.96 d (6.7)	21.1	0.99 d (6.6)
14	199.7	9.28 s	199.8	9.38 s	199.3	9.40 s
15	16.4	1.45 s	16.7	1.10 s	16.1	1.13 s
OAc	20.9	2.00 s	20.8	2.06 s	21.0	2.12 s
	21.1	2.08 s	20.9	2.06 s	173.0	
	169.8		169.7			
	170.4		169.8			

Table 5. ^1H and ^{13}C NMR Data for Compounds **9**–**11** (CDCl_3 , J values (Hz) in parentheses)

position.	9		10		11	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}
1	163.7	6.58 dd (4.4, 11.9)	161.6	6.41 dd (5.8, 10.5)	134.0	5.36 dd (2.5, 9.0)
2	25.0	2.11 m	25.2	2.04 m	23.1	1.80 m
		2.64 q (12.0)		2.53 m		2.26 d (10.2)
3	41.0	1.67 dd (11.0, 14.0)	39.9	1.74 m	41.2	1.50 dd (10.8, 12.8)
		2.01 m		1.98 m		1.78 m
4	73.6		73.6		74.1	
5	139.7	5.00 d (15.3)	138.6	5.20 d (15.3)	140.7	5.17 d (15.3)
6	122.2	5.22 dd (10.7, 15.3)	121.4	5.25 dd (10.0, 15.4)	120.0	5.22 dd (9.4, 15.3)
7	55.7	1.56 m	52.8	1.60 m	55.7	1.98 m
8	66.7	3.76 m	70.3	5.00 m	68.6	3.85 dd (3.3, 6.2)
9	32.1	2.83 dd (4.8, 15.3)	31.9	2.79 dd (3.4, 15.0)	35.6	2.10 m
		2.40 d (15.3)		2.51 dd (3.4, 15.0)		2.88 d br (14.9)
10	139.7		157.6		150.0	
11	26.6	2.17 ddd (2.7, 6.8, 6.8)	27.5	2.16 m	27.3	2.01 m
12	21.7	0.81 d (6.9)	21.4	0.80 d (6.8)	21.8	0.87 d (6.4)
13	16.4	0.79 d (6.9)	16.8	0.79 d (6.8)	16.7	0.85 d (6.4)
14	199.5	9.39 s	194.8	9.42 s	69.8	4.03 d (12.0)
						4.10 d (12.0)
15	29.8	1.32 s	29.6	1.34 s	30.0	1.29 s
OAc			171.2			
			21.2	2.12 s		

On the basis of the above evidence, we assigned **8** as 1β -, 10α -, 4α -, 5β -diepoxy- 8β -acetoxy- 6α -hydroxy- 7β H-germacra-14-al.

Pulicanadienal A (**9**), obtained as a colorless oil, has a molecular formula of $\text{C}_{15}\text{H}_{24}\text{O}_3$ as determined by HR-FABMS (m/z 275.1645 $[\text{M} + \text{Na}]^+$). The IR absorption of **9** (3410 and 1660 cm^{-1}) indicated the presence of a hydroxyl and an α,β -unsaturated carbonyl functional group. The ^1H NMR spectrum (Table 5) of **9** showed signals for an oxygenated methine proton at δ_{H} 3.76 (1H, m, H-8), two methyl groups on a methine carbon at δ_{H} 0.81 (3H, d, $J = 6.9$ Hz, CH_3 -12) and 0.79 (3H, d, $J = 6.9$ Hz, CH_3 -13), an angular methyl group at δ_{H} 1.32 (3H, s, CH_3 -15), three olefinic protons at δ_{H} 6.58 (1H, dd, $J = 4.4, 11.9$ Hz, H-1), 5.22 (1H, dd, $J = 10.7, 15.3$ Hz, H-6), and 5.00 (1H, d, $J = 15.3$ Hz, H-6), and one aldehyde proton at δ_{H} 9.39 (1H, s, H-14). The ^{13}C NMR spectrum and the corresponding DEPT spectrum exhibited 15 carbon signals, consisting of three methyl, three methylene, seven methine, and two quaternary carbons atoms, suggesting the presence of a germacrane-type sesquiterpene skeleton. The ^{13}C NMR spectrum also established that two olefinic bonds (one disubstituted and one trisubstituted) were present and a hydroxyl group was located on the carbon totally substituted (a singlet carbon at δ_{C} 73.6). The COSY experiment disclosed two partial structures, CHCH_2CH_2 and CHCH-CHCHCH_2 , corresponding to the C-1, C-2, C-3 and C-5, C-6, C-7, C-8, C-9 fragments. HMBC correlations of H-1 with C-10, C-14, and C-3; H-5 with C-4 and C-6; H-6 with C-5, C-4, and C-8; H-7 with C-8, C-12, and C-13; H-14 with C-1, C-9, and C-10; and CH_3 -15 with C-4 and C-5 were observed. The stereochemistry of the C-5, C-6 double bond was determined to be *E* by considering the coupling constant ($J_{5,6} = 15.3$ Hz). The relative stereochemistry of **9** was determined from the NOESY spectrum (Figure 4). Important correlations were observed between H-8, H-6, CH_3 -12, and CH_3 -13 as well as between H-14, H-1, and CH_3 -15, leading to the conclusion that the stereochemistry of the C-1, C-10 double bond is *E*. Correlations were also observed between CH_3 -15, H-5, and H-7. The absolute configuration of C-8 (Figure 5) was determined using the modified Mosher ester procedure.²¹ Compound **9** was treated with (S)-(+)- and (R)-(-)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride in anhydrous pyridine at room

**Figure 4.** Selected NOESY correlations for compound **9**.**Figure 5.** $\Delta\delta$ values [$\Delta\delta$ (in ppm) = $\delta_S - \delta_R$] obtained for (S)- and (R)-MTPA esters (**9a** and **9b**, respectively) of compound **9**.

temperature for 24 h, to afford the (R)- and (S)-MTPA ester derivatives **9a** and **9b**, respectively. Negative values ($\Delta\delta_{S-R}$) were obtained for H-7, H-11, H-6, and H-5, indicating that the absolute configuration at C-8 is *R*. Thus the structure of **9** may be formulated as (1*E*,5*E*,8*R*)- 4α , 8β -dihydroxy- 7β H-germacra-1(10),5-dien-14-al.

Pulicanadienal B (**10**) was isolated as an amorphous solid, $[\alpha]_{\text{D}}^{25} 49.1^\circ$ (c 0.059, CHCl_3), and its molecular formula was determined as $\text{C}_{17}\text{H}_{26}\text{O}_4$ by HRFABMS data (m/z 317.1718 $[\text{M} + \text{Na}]^+$). The ^1H and ^{13}C NMR (Table 5) spectral data of **10** were similar to those of **9** with the exception of signals for an acetoxy group. Thus, treatment of **9** with Ac_2O –pyridine gave a monoacetyl derivative whose physical and spectroscopic constants were identical with those of **10** and was determined as (1*E*,5*E*)- 8β -acetoxy- 4α -hydroxy- 7β H-germacra-1(10),5-dien-14-al.

Pulicanadienol (**11**) was obtained as an amorphous solid, $[\alpha]_{\text{D}}^{25} 10.2^\circ$ (c 0.049, CHCl_3). The IR spectrum showed the presence of hydroxyl groups (3335 cm^{-1}). The ^1H NMR spectrum (Table 5) of **11** showed signals analogous to those of **9** with the difference that C-14 is replaced by a hydroxymethyl group, δ_{H} 4.03 (1H, d, $J = 12.0$ Hz, H-14a), 4.10 (1H, d, $J = 12.0$ Hz, H-14b). The relative stereochemistry of **11** was assigned as in **9** based upon the very close

Table 6. ^1H and ^{13}C NMR Data for Compounds **12** and **13** (CDCl_3 , J values (Hz) in parentheses)

position	12		13	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}
1	51.7	α , 1.95 m	57.0	α , 1.50 m
2	27.3	α , 1.73 m	25.4	α , 1.50 m
		β , 1.83 m		β , 1.86 m
3	29.4	α , 1.95 m	29.5	α , 1.96 m
		β , 1.60 m		β , 1.60 m
4	212.4		212.4	
5	50.8	α , 2.66 ddd (4.0, 5.8, 9.7)	50.8	α , 2.59 ddd (5.0, 6.3, 10.6)
6	52.0	β , 1.86 m	47.0	β , 1.92 m
7	49.0	β , 1.25 m	49.2	β , 1.10 m
8	26.3	α , 1.70 m	22.8	1.60 m
		β , 1.10 dq (4.2, 12.9)		1.10 m
9	35.2	α , 2.01 (m)	42.0	1.80 br d (15.6)
		β , 2.37 dt (4.0, 6.0)		1.41 m
10	150.4		73.0	
11	30.2	1.33 dt (2.8, 6.9)	30.0	1.29 m
12	21.9	0.89 d (6.8)	21.9	0.89 d (6.7)
13	15.4	0.61 d (6.8)	15.3	0.66 d (6.7)
14	103.9	4.68 d (1.0)	20.4	1.21 s
		4.57 d (1.0)		
15	67.7	4.31 s	68.0	4.30 t (4.3)

^{13}C NMR (Table 5) correlations between these compounds. This confirmed the structure of **11** as (1*E*,5*E*)-7 β H-germacra-1(10),5-dien-4 α ,8 β ,14-triol.

Puliplopanone A (**12**) exhibited the molecular formula $\text{C}_{15}\text{H}_{24}\text{O}_2\text{Na}$ by HRFABMS (m/z 259.1702 [$\text{M} + \text{Na}$] $^+$). It showed IR bands at 3489, 1710, and 1660 cm^{-1} due to hydroxyl and carbonyl groups and a double bond. Its ^1H NMR spectrum (Table 6) showed signals that corresponded to an exocyclic double bond at δ_{H} 4.57 (1H, d, $J = 1.1$ Hz, H-14a) and 4.68 (1H, d, $J = 1.1$ Hz, H-14b), a hydroxymethyl group at δ_{H} 4.31 (2H, s, H₂-15), and an isopropyl moiety at δ_{H} 0.61 (3H, d, $J = 6.8$ Hz, H₃-12), 0.89 (3H, d, $J = 6.8$ Hz, H₃-12), and 1.33 (1H, m, H-11), suggesting an oplopene skeleton.²² From these data, the new compound **12** was deduced to be 15-hydroxy-10(14)-oplopen-4-one, and the structural model could be determined by a single-crystal X-ray diffraction analysis (Supporting Information).

Puliplopanone B (**13**) exhibited a molecular formula of $\text{C}_{15}\text{H}_{26}\text{O}_3$, established from HREIMS (m/z 254.1901 [M] $^+$). Its ^1H and ^{13}C NMR spectra (Table 6) were similar to those of **12** except for the presence of a hydroxyl group attached to C-10. NOE effects between H-5 and H₂-15, H-7, and H-1 and between H-6 and H₃-14, observed in a NOESY experiment, suggested that OH-10 in **13** has an α -orientation and pointed to the same relative stereochemistry for **12** and **13**, leading to the conclusion that **13** is 10 α ,15-dihydroxyoplopen-4-one. A cinnamoyl derivative has been reported in the literature.²³

Next, we investigated whether these sesquiterpenes possess cytotoxic activities against the HL-60 human tumor cell line. As determined by the MTT assay, all compounds tested were weakly cytotoxic to this cell line with 50% inhibitory concentration (IC_{50}) values (in μM) of 20 ± 7 , 298 ± 22 , 115 ± 18 , 264 ± 6 , and 242 ± 80 , for **4a**, **5a**, **6**, **7**, and **12**, respectively. The triacetate of pulicanone (**4a**) was the most potent compound, with an IC_{50} value of 6- and 15-fold lower than those of **6** and **5a**, respectively. To




Figure 6. Induction of apoptosis in HL-60 cells in response to triacetoxypulicanone **4a**. (A) Cells were cultured for 12 h in the absence (control) or presence of the indicated doses of compound **4a** and then stained with Hoeschst 33258 to evaluate nuclear chromatin condensation, i.e., apoptosis. (B) Quantitative analysis of apoptotic cells by fluorescence microscopy; etoposide (Eto, 10 μM) was used as a positive control. The results of a representative experiment are shown, and each point represents the average (\pm SE) of triplicate determinations.




Figure 7. Triacetoxypulicanone **4a** induces caspase-3/8 activation. HL-60 cells were cultured in the presence of increasing concentrations of compound **4a** and harvested at 12 h. Total cell lysates were obtained and used to determine caspase-3 (left panel) and caspase-8 (right panel) activities. The results of a representative experiment performed in duplicate are shown and expressed as x -fold increase of caspase activity compared with control.

know if compound **4a** decreases cell viability through apoptosis activation, morphological changes characteristic of apoptotic cells (i.e., condensed and fragmented chromatin) were analyzed and quantified by fluorescent microscopy. The percentage of apoptotic cells increased in a dose-response manner in response to **4a**, as demonstrated in Figure 6. Furthermore, flow cytometric analysis of HL-60 cells incubated with **4a** (30 μM , 12 h) indicates that transition of cells from the diploid to hypodiploid (i.e., apoptotic cells) region occurs with the accumulation of cells in the G1 phase of the cell cycle (data not shown). In concert with induction of apoptosis, HL-60 cells also responded to **4a** with activation of caspase-3 and caspase-8 (Figure 7) and cleavage of PARP-1 (Figure 8).

Experimental Section

General Experimental Procedures. Melting points were determined on a Büchi B-540 apparatus and are uncorrected. Optical rotations were recorded in a Perkin-Elmer model 343 polarimeter, and UV spectra were recorded using a JASCO model V-560 spectrophotometer. ^1H NMR and ^{13}C NMR spectra were obtained on Bruker model AMX-400 and AMX-500 spectrometers with standard pulse sequences operating at 400 and 500 MHz in ^1H NMR and 100 and 125 MHz in ^{13}C NMR, respectively. CDCl_3 , $\text{C}_5\text{D}_5\text{N}$, and $\text{DMSO}-d_6$ were used as solvents. EIMS, HREIMS, and HRFABMS were taken on a Micromass model Autospec (70 eV) spectrometer. Column chromatography was carried out on silica gel 60 (Merck 230–




Figure 8. Triacetoxypulicanone **4a** induces poly(ADP-ribose)polymerase-1 (PARP-1) cleavage in HL-60 cells. The indicated concentrations of compound **4a** were added to the cells for 12 h, and total cell lysates were then analyzed by immunoblotting with an anti-PARP-1 antibody. Etoposide (Eto, 10 μ M) was included as a positive control.

400 mesh), and preparative TLC on silica gel 60 PF₂₅₄₊₃₆₆ plates (20 \times 20 cm, 1 mm thickness).

Plant Material. *Pulicaria canariensis* was collected in Playa Quemada (Lanzarote, Canary Islands) in August 1998. The plant material was identified by Professor Rosa Febles, and a voucher specimen has been deposited at the Herbarium of the Viera y Clavijo Botanical Garden in Gran Canaria (No. 20146).

Extraction and Isolation. The aerial parts of the plant (1.5 kg) were extracted with EtOH (3 L, four times). The solvent was concentrated under reduced pressure. The residue (90 g) obtained after removing EtOH was fractionated by silica gel flash column chromatography using hexane and EtOAc mixtures of increasing polarity to yield four fractions. Fraction 1 (500 mg) eluted with hexane–EtOAc (4:1) was concentrated under reduced pressure and recrystallized from hexane–EtOAc to obtain the known compounds stigmasterol (180 mg), 7,4'-di-*O*-methylhydrokaempferol (26 mg), and calenduladiol (41 mg), respectively; the liquid supernatant was chromatographed by medium-pressure liquid chromatography (MPLC) with the mobile phase hexane–EtOAc (4:1), furnishing after purification by preparative TLC with hexane–EtOAc (9:1) the new sesquiterpenes **6** (45 mg), **2** (48 mg), **7** (35 mg), and **1** (24 mg). Fraction 2 (425 mg) eluted with hexane–EtOAc (7:3) and treated in the same way as fraction 1 was then recrystallized from hexane–EtOAc to yield 5,7-dihydroxy-3,3',4'-trimethoxyflavone (150 mg). The liquid supernatant was chromatographed on MPLC using hexane–EtOAc (7:3) as eluant, and subsequent purification by preparative TLC using benzene–EtOAc (4:1) gave **3** (9 mg), ergosterol peroxide (178 mg), **10** (12 mg), and a mixture which, after acetylation with Ac₂O–pyridine and MPLC in hexane–EtOAc (9:1), yielded **4a** (30 mg). Fraction 3 (200 mg), eluted with hexane–EtOAc (3:2), was subjected to column chromatography using hexane–EtOAc (4:1) to give dihydroquercetin 7,3'-dimethyl ether (18 mg) and the semipure compound **9**. This was finally purified by preparative TLC developed with benzene–acetonitrile (4:1), to afford pure compound **9** (42 mg). Fraction 4 (300 mg) eluted with hexane–EtOAc (3:1) and finally purified by preparative TLC developed with hexane–EtOAc (1:1) to afford compounds 6,15 α -epoxy-1 β ,4 β -dihydroxyeudesmane (45 mg), **11** (9 mg), **13** (8 mg), **8** (25 mg), **5** (40 mg), and **12** (68 mg).

Pulicanadiene A (1): amorphous solid; $[\alpha]_D^{25}$ 21.4° (c 0.014, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 206 (3.2), 250 (2.3), 273 (1.8) nm; IR (KBr) ν_{max} 3067, 2937, 1731, 1693, 1554, 1430, 1372, 1245, 1023, 960 cm⁻¹; ¹H and ¹³C NMR, see Table 1; EIMS m/z 352 (M⁺, 1), 316 (1), 292 (1), 281 (0.5), 274 (1), 250 (12), 233 (15), 232 (66), 217 (24), 203 (12), 189 (100), 187 (41), 176 (16), 171 (17), 161 (19), 145 (89), 143 (44), 131 (40), 123 (56), 105 (34), 91 (49), 81 (42), 69 (30), 55 (30); HREIMS m/z 352.1877 [M]⁺ (calcd for C₁₉H₂₈O₆ 352.1886).

Methylation of Compound 1. To compound **1** (4 mg) in diethyl ether (5 mL) was added an excess of freshly prepared diazomethane in diethyl ether, and the mixture was kept at room temperature for 12 h. Concentration of the reaction mixture under vacuum furnished **1a** (4 mg), as colorless needles: mp 162–163 °C; $[\alpha]_D^{25}$ 3.2° (c 0.062, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 212 (2.0), 251 (2.1) nm; IR (KBr) ν_{max} 2930, 2360, 2341, 1732, 1435, 1371, 1246, 1215, 1021, 963 cm⁻¹; ¹H and ¹³C NMR, see Table 1, ¹H NMR (CDCl₃, at –50 °C, 500

MHz) δ 6.64 (1H, t, J = 7.9 Hz, H-1), 5.69 (1H, d br, J = 11.0 Hz, H-6), 5.65 (1H, m, H-8), 5.14 (1H, d, J = 10.9 Hz, H-5), 3.74 (3H, s, OMe), 2.49 (1H, dd, J = 5.2, 13.9 Hz, H-9a), 2.32 (2H, m, overlapped signals, H-2a and H-3a), 2.11 (3H, s, OAc), 2.08 (1H, dd, J = 13.9, 10.8 Hz, H-9b), 2.00 (3H, s, OAc), 1.98 (2H, m, overlapped signals, H-2b and H-3b), 1.93 (3H, s, CH₃-15), 1.72 (1H, d, J = 11.4 Hz, H-7), 1.57 (1H, m, H-11), 0.97 (3H, d, J = 6.2 Hz, CH₃-12), 0.95 (3H, d, J = 6.2 Hz, CH₃-13); EIMS m/z 366 (M⁺, 2), 307 (2), 306 (2), 274 (4), 264 (12), 247 (13), 246 (58), 231 (23), 214 (16), 203 (100), 187 (78), 171 (30), 145 (37), 143 (59), 131 (37), 123 (59), 105 (28), 91 (34), 81 (38), 79 (25), 69 (20), 55 (22); HREIMS m/z 366.2019 [M]⁺ (calcd for C₂₀H₃₀O₆ 366.1733).

Pulicanadiene B (2): colorless oil; $[\alpha]_D^{25}$ 35.7° (c 0.042, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 209 (3.2), 222 (2.7), 251 (2.1) nm; IR (KBr) ν_{max} 2932, 1732, 1687, 1433, 1372, 1243, 1141, 1023, 960 cm⁻¹; ¹H and ¹³C NMR, see Table 1; EIMS m/z 277 (35), 235 (10), 217 (74), 216 (72), 199 (79), 187 (82), 173 (93), 150 (62), 145 (94), 131 (41), 123 (83), 109 (38), 107 (100), 95 (39), 93 (33), 91 (33), 84 (45), 55 (49); HRFABMS m/z 359.1874 [M + Na]⁺ (calcd for C₁₉H₂₈O₅Na 359.1834).

Pulicanadiene C (3): colorless oil; $[\alpha]_D^{25}$ –75.0° (c 0.008, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 205 (3.2), 271 (1.9) nm; IR (KBr) ν_{max} 3414, 3020, 2928, 2854, 1728, 1502, 1411, 1381, 1332, 1219, 1159, 1107, 1028, 766 cm⁻¹; ¹H and ¹³C NMR, see Table 1; EIMS m/z 338 (M⁺, 1), 290 (10), 279 (10), 241 (13), 218 (48), 203 (25), 200 (78), 193 (11), 187 (100), 178 (59), 175 (99), 161 (44), 157 (71), 149 (45), 143 (46), 135 (38), 131 (35), 122 (28), 109 (23), 105 (30), 97 (33), 95 (26), 91 (25), 83 (36), 81 (30), 69 (33), 57 (45), 55 (95); HREIMS m/z 338.2019 [M]⁺ (calcd for C₁₉H₃₀O₅ 338.2093).

Reduction of 2. Compound **2** (5.0 mg) in methanol (5 mL) was treated with excess NaBH₄ at room temperature for 30 min. The reaction was worked up by addition of H₂O and was extracted with CH₃Cl. The concentrated dried CH₃Cl extract was separated by preparative TLC (benzene–AcOEt, 80:20) and furnished **3** (1.2 mg).

Triacetoxypulicanone (4a): colorless oil; $[\alpha]_D^{25}$ 4.9° (c 0.081, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 205 (3.2), 223 (2.7), 271 (3.1) nm; IR (KBr) ν_{max} 2960, 2926, 2851, 1741, 1694, 1463, 1438, 1373, 1233, 1123, 1053, 1025, 935, 852, 666 cm⁻¹; ¹H and ¹³C NMR, see Table 2; EIMS m/z 394 (M⁺, 0.5), 334 (12), 292 (17), 274 (78), 249 (24), 232 (62), 214 (85), 205 (94), 189 (95), 171 (42), 161 (47), 149 (68), 137 (41), 136 (44), 107 (74), 96 (65), 69 (61), 57 (100); HREIMS m/z 394.1982 [M]⁺ (calcd for C₂₁H₃₀O₇ 394.1991).

Pulicanol (5): colorless oil; IR (KBr) ν_{max} 3404, 2929, 1736, 1647, 1470, 1375, 1243, 1088, 1023, 963, 897 cm⁻¹; EIMS m/z 312 (M⁺, 2), 279 (11), 252 (4), 234 (5), 216 (4), 173 (14), 171 (16), 167 (37), 149 (87), 133 (32), 129 (33), 123 (46), 121 (50), 111 (56), 105 (48), 97 (87), 95 (91), 91 (57), 85 (70), 83 (100), 57 (89), 55 (96); ¹H NMR (CDCl₃, 400 MHz) δ 5.45 (1H, t, J = 6.8 Hz, H-1), 5.18 (1H, dd, J = 5.2, 12.0 Hz, H-8), 4.29 (1H, d, J = 11.8 Hz, H-14a), 4.12 (1H, d, J = 11.6 Hz, H-14b) 3.51 (1H, m, H-6), 2.82 (1H, d, J = 7.7 Hz, H-5), 2.53 (2H, d, J = 12.4 Hz, H-2), 2.34 (2H, m, H-2-3), 2.15 (3H, s, OAc), 2.10 (1H, m, H-2-3), 1.80 (1H, m, H-11), 1.59 (1H, m, H-7), 1.16 (3H, s, CH₃-15), 1.12 (3H, d, J = 6.4 Hz, CH₃-12), 0.95 (3H, d, J = 6.4 Hz, CH₃-13).

Acetylation of 5. Compound **5** (20 mg) was dissolved in pyridine (3 mL) and acetic anhydride (3 mL), and the solution was stirred for 16 h at room temperature. The product was dried under vacuum to furnish **5a** (15 mg) as a colorless oil; $[\alpha]_D^{25}$ –6.9° (c 0.188, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 214 (2.4) nm; IR (KBr) ν_{max} 2930, 2874, 1739, 1653, 1597, 1458, 1371, 1238, 1130, 1078, 1026, 962, 817 cm⁻¹; ¹H and ¹³C NMR, see Table 3; EIMS m/z 336 (6), 294 (8), 276 (5), 234 (21), 216 (32), 194 (26), 191 (28), 175 (45), 173 (37), 163 (25), 150 (29), 145 (33), 139 (46), 131 (50), 119 (33), 107 (49), 91 (47), 84 (100), 81 (40), 69 (28), 67 (28), 57 (22), 55 (35); HRFABMS m/z 419.2009 [M + Na]⁺ (calcd for C₂₁H₃₂O₇Na 419.2046).

Pulicanaral A (6): colorless needles; mp 183–184 °C; $[\alpha]_D^{25}$ 11.1° (c 0.108, CHCl₃); UV (EtOH) λ_{max} (log ϵ) 209 (3.1), 272 (1.9) nm; IR (KBr) ν_{max} 2928, 1732, 1716, 1375, 1243, 1025, 935 cm⁻¹; ¹H and ¹³C NMR, see Table 4; EIMS m/z 352 (M⁺,

0.5), 337 (0.6), 292 (6), 249 (10), 232 (68), 203 (21), 189 (66), 175 (31), 171 (20), 166 (28), 161 (45), 153 (47), 145 (33), 143 (100), 135 (22), 129 (58), 109 (46), 107 (28), 105 (28), 95 (42), 84 (56), 81 (61), 69 (36), 55 (50); HREIMS m/z 352.1905 $[M]^+$ (calcd for $C_{19}H_{28}O_6$, 352.1886).

Pulicanaral B (7): colorless needles; mp 205–210 °C; $[\alpha]_D^{25}$ –60.0° (c 0.005, $CHCl_3$); UV (EtOH) λ_{max} (log ϵ) 208 (2.3) nm; IR (KBr) ν_{max} 2932, 1737, 1715, 1432, 1373, 1244, 1130, 1090, 1033, 894, 767 cm^{-1} ; 1H and ^{13}C NMR, see Table 4; EIMS m/z 309 (3), 308 (1), 297 (4), 279 (3), 266 (6), 255 (27), 248 (4), 237 (12), 223 (15), 220 (18), 219 (28), 209 (10), 207 (13), 205 (27), 195 (34), 191 (14), 187 (11), 177 (44), 166 (27), 163 (28), 159 (14), 151 (24), 149 (47), 141 (30), 139 (66), 127 (50), 123 (44), 109 (59), 99 (76), 97 (100), 95 (74), 85 (58), 83 (100), 81 (83), 71 (53), 69 (60), 55 (90); HRFABMS m/z 391.1738 $[M + Na]^+$ (calcd for $C_{19}H_{28}O_7Na$ 391.1733).

Pulicanaral B (8): colorless oil; IR (KBr) ν_{max} 3462, 2964, 2925, 2856, 1719, 1597, 1455, 1382, 1255, 1074 cm^{-1} ; 1H and ^{13}C NMR, see Table 4; EIMS m/z 326 (M^+ , 0.2), 279 (15), 251 (6), 237 (6), 223 (8), 219 (10), 205 (7), 195 (5), 191 (13), 177 (13), 167 (27), 163 (19), 149 (90), 139 (28), 137 (31), 123 (42), 111 (41), 109 (68), 97 (100), 95 (79), 83 (72), 81 (66), 69 (61), 55 (77).

Acetylation of 8. Compound **8** (4.0 mg) was dissolved in pyridine (1 mL) and acetic anhydride (1 mL), and the solution was stirred for 18 h at room temperature. The product was dried under vacuum to furnish **7** (3.5 mg).

Pulicanadienal A (9): colorless oil; $[\alpha]_D^{25}$ 210° (c 0.020, $CHCl_3$); UV (EtOH) λ_{max} (log ϵ) 212 (2.9), 253 (2.2) nm; IR (KBr) ν_{max} 3406, 2957, 2926, 2870, 1728, 1660, 1623, 1440, 1370, 1272, 1142, 1060, 987, 669 cm^{-1} ; 1H and ^{13}C NMR, see Table 5; EIMS m/z 234 (28), 219 (9), 216 (15), 205 (10), 201 (8), 194 (7), 191 (15), 187 (9), 176 (26), 173 (14), 169 (11), 161 (11), 151 (76), 145 (27), 133 (22), 131 (22), 123 (100), 122 (62), 119 (18), 117 (15), 107 (53), 105 (39), 95 (43), 93 (62), 91 (43), 81 (53), 69 (54), 67 (46), 55 (58); HRFABMS m/z 275.1645 $[M + Na]^+$ (calcd for $C_{15}H_{24}O_3Na$ 275.1623).

Preparation of the (R)- and (S)-MTPA Ester Derivatives of 9. Two portions (1.5 mg) of compound **9** were treated with (R)-(+)- and (S)-(–)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (8 μ L) in anhydrous pyridine (0.5 mL) at room temperature overnight. When the reaction was completed, the residue was concentrated and purified on silica gel by preparative TLC eluting with $CHCl_3$ – Et_2O – $MeOH$ (14:6:1), affording the (S)- and (R)-MTPA ester derivatives (**9a** and **9b**) of **9**, respectively. 1H NMR data of **9a** (500 MHz, $CDCl_3$): δ_H 9.420 (1H, s, H-14), 6.484 (1H, t, J = 7.8 Hz, H-1), 5.360 (1H, m, H-8), 5.322 (1H, d, J = 15.4 Hz, H-5), 5.120 (1H, dd, J = 11.1, 15.4 Hz, H-6), 2.684 (1H, dd, J = 6.7, 14.3 Hz, H-9a), 2.610 (1H, dd, J = 4.0, 14.3 Hz, H-9b), 2.170 (1H, dd, J = 3.0, 8.9 Hz, H-7), 1.471 (1H, td, J = 2.8, 6.9 Hz, H-11), 1.310 (3H, s, CH_3 -15), 0.730 (3H, d, J = 6.9 Hz, CH_3 -13), 0.632 (3H, d, J = 6.8 Hz, CH_3 -12); 1H NMR data of **9b** (500 MHz, $CDCl_3$): δ_H 9.347 (1H, s, H-14), 6.466 (1H, t, J = 8.3 Hz, H-1), 5.352 (1H, d, J = 15.1 Hz, H-5), 5.320 (1H, m, H-8), 5.151 (1H, dd, J = 10.7, 15.1 Hz, H-6), 2.680 (1H, dd, J = 7.7, 14.7 Hz, H-9a), 2.518 (1H, d br, J = 14.1 Hz, H-9b), 2.251 (1H, m, H-7), 1.765 (1H, m, H-11), 1.321 (3H, s, CH_3 -15), 0.815 (3H, d, J = 7.0 Hz, CH_3 -13), 0.784 (3H, d, J = 6.7 Hz, CH_3 -12).

Pulicanadienol B (10): amorphous solid; $[\alpha]_D^{25}$ 49.1° (c 0.059, $CHCl_3$); UV (EtOH) λ_{max} (log ϵ) 214 (3.0), 251 (2.1) nm; IR (KBr) ν_{max} 3459, 2922, 1732, 1684, 1630, 1375, 1256, 1026, 880, 803 cm^{-1} ; 1H and ^{13}C NMR, see Table 5; EIMS m/z 317 (M^+ , 2), 165 (1), 154 (8), 136 (13), 109 (12), 97 (17), 95 (31), 81 (35), 71 (31), 69 (66), 57 (81), 55 (100); HRFABMS m/z 317.1718 $[M + Na]^+$ (calcd for $C_{17}H_{26}O_4Na$ 317.1729).

Acetylation of 9. Compound **9** (5.0 mg) was dissolved in pyridine (1 mL) and acetic anhydride (1 mL), and the solution was stirred for 18 h at room temperature. The product was dried under vacuum to furnish **10** (4.0 mg).

Pulicanadienol (11): amorphous solid; $[\alpha]_D^{25}$ 10.2° (c 0.049, $CHCl_3$); UV (EtOH) λ_{max} (log ϵ) 212 (3.0) nm; IR (KBr) ν_{max} 3335, 2957, 2926, 2871, 1661, 1625, 1449, 1384, 1368, 1244, 1174, 1037, 984, 891 cm^{-1} ; 1H and ^{13}C NMR, see Table 5; EIMS m/z 236 (5), 218 (26), 203 (11), 202 (5), 201 (10), 189 (12), 187

(12), 185 (9), 175 (56), 160 (17), 159 (25), 157 (30), 145 (44), 133 (42), 131 (41), 123 (39), 107 (58), 105 (69), 95 (55), 93 (64), 91 (72), 81 (79), 69 (79), 57 (84), 55 (100); HREIMS $[M - H_2O]^+$ m/z 236.2155 (calcd for $C_{15}H_{24}O_2$ 236.1776).

Puliooplopanone A (12): colorless needles; mp 141–142 °C; $[\alpha]_D^{25}$ –40.0° (c 0.005, $CHCl_3$); UV (EtOH) λ_{max} (log ϵ) 209 (2.7), 274 (1.2) nm; IR (KBr) ν_{max} 3489, 2960, 2939, 2876, 1710, 1660, 1602, 1405, 1280, 1230, 1055, 893, 748 cm^{-1} ; 1H and ^{13}C NMR, see Table 6; EIMS m/z 236 (M^+ , 12), 205 (100), 187 (74), 177 (35), 135 (17), 121 (55), 107 (18), 95 (39), 93 (27), 91 (18), 79 (24), 69 (17), 67 (15), 55 (13); HRFABMS m/z 259.1702 $[M + Na]^+$ (calcd for $C_{15}H_{24}O_2Na$ 259.1703).

Puliooplopanone B (13): colorless oil; $[\alpha]_D^{25}$ 33.3° (c 0.012, $CHCl_3$); IR (KBr) ν_{max} 3401, 2930, 2871, 2360, 2342, 1718, 1459, 1369, 1263, 1126, 1053 cm^{-1} ; 1H and ^{13}C NMR, see Table 6; EIMS m/z 254 (M^+ , 2), 236 (5), 224 (15), 223 (96), 205 (35), 187 (43), 177 (100), 169 (20), 147 (9), 135 (13), 121 (33), 107 (19), 95 (34), 81 (29), 69 (25), 55 (15); HRFABMS $[M]^+$ m/z 254.1901 (calcd for $C_{15}H_{26}O_3$ 254.1882).

Cell Culture and Cytotoxicity Assays. HL-60 cells were cultured in suspension in RPMI-1640 medium supplemented with 10% (v/v) heat-inactivated fetal bovine serum, penicillin (100 units/mL), and streptomycin (100 μ g/mL) in a humidified atmosphere of 95% air and 5% CO_2 at 37 °C. Cells were maintained at a density of $<1 \times 10^6$ cells/mL. Cytotoxicity studies were performed using the MTT assay as described.⁹

Apoptosis Quantitation by Fluorescence Microscopy. Quantitation of apoptotic cells was performed as described previously.¹⁰

Assay of Caspase Activity. Activity of caspase was determined as described previously²⁴ from cytosolic lysates, using the specific colorimetric substrate DEVD-pNA (caspase-3) or IETD-pNA (caspase-8).

Analysis of PARP-1 Hydrolysis by Immunoblotting. PARP-1 cleavage was analyzed by Western blot assay as described¹⁰ using a PARP-1 monoclonal antibody which recognizes the native form (116 kDa) and the hydrolyzed form (85 kDa).

Acknowledgment. This work was supported in part by a grant from the Instituto Canario de Investigación del Cáncer (G-04-09 to J.B. and 08/2004 to J.Q.) and Spanish MEC funds (SAF 2004-07928 to F.E.). We thank J. Estévez (Hospital Universitario Insular de Gran Canaria) for his collaboration in the Western blot assays and S. Rubio for excellent technical assistance.

Supporting Information Available: 1H NMR spectra of compound **1**, **1a** (at 50 °C, 25 °C, –50 °C), **5a**, **6**, **7**, **9**, and **12**. Enlargement of the NOESY experiment performed at 50 and –50 °C, and X-ray diffraction data for compounds **1a**, **6**, **7**, and **12**. These materials are available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Bramwell, D.; Bramwell, Z. I. *Flores Silvestres de las Islas Canarias*; Rueda S. L., Ed.; Madrid, 2001.
- Grieve, M. A. *Modern Herbal*; Penguin Books: Harmondsworth, UK, 1980; p 32.
- Awadh, A. N. A.; Julich, W. D.; Kusnick, C.; Lindequist, U. J. *Ethnopharmacol.* **2001**, *74*, 173–179.
- Mahfouz, M.; Ghazal, A.; El-Dakhakhny, M.; Ghoneim, M. T. *J. Drug Res.* **1973**, *5*, 151–172.
- Williams, C. A.; Harborne, J. B.; Greenham, J. *Biochem. Syst. Ecol.* **2000**, *28*, 679–687.
- Marco, J. A.; Sanz, J. F.; Albiach, R. *Phytochemistry* **1992**, *31*, 2409–2413.
- Dendougui, H.; Benayache, S.; Benayache, F.; Connolly, J. D. *Fito-terapia* **2000**, *71*, 373–378.
- Eshbakova, K. A.; Saidkhodzhaev, A. I. *Chem. Nat. Prod.* **2001**, *37*, 196–197.
- Triana, J.; Lopez, M.; Rico, M.; González-Platas, J.; Quintana, J.; Estevez, F.; León, F.; Bermejo, J. *J. Nat. Prod.* **2003**, *66*, 943–948.
- Rivero, A.; Quintana, J.; Eiroa, J. L.; López, M.; Triana, J.; Bermejo, J.; Estévez, F. *Eur. J. Pharmacol.* **2003**, *482*, 77–84.
- Shopee, C. W. *Chemistry of the Sterols*; Butterworth: London, 1964; pp 62–63.
- González, A. G.; Bermejo, J. B.; Rodríguez, P. E. M.; Hernández, H. C. *Planta Med.* **1992**, *58*, 214–218.
- Datanarayana, A. P.; Kumar, N. S.; Sultanbawa, M. U.; Balasubramaniam, S. *J. Chem. Soc., Perkin Trans. 1* **1981**, 2717–2723.
- Islam, M. T.; Tahara, S. *Phytochemistry* **2000**, *54*, 901–907.

- (15) González, A. G.; Bermejo, J.; Triana, J.; Eiroa, J. L.; Lopez, M. *J. Nat. Prod.* **1995**, *58*, 432–437.
- (16) Balza, F.; Towers, G. H. N. *Phytochemistry* **1984**, *23*, 2333–2337.
- (17) Fang, N.; Yu, S.; Mabry, T. J.; Abboud, K. A.; Simonsen, S. H. *Phytochemistry* **1988**, *27*, 3187–3196.
- (18) Barrero, A. F.; Herrador, M. M.; Quilez, J. F.; Alvarez-Manzaneda, R.; Portal, D.; Gavin, J. A.; Gravalos, D. G.; Simmonds, M. S. J.; Blaney, W. M. *Phytochemistry* **1999**, 529–541.
- (19) Pascual, J. T.; Moran, J. R.; Hernández, J. M.; Grande, M. *Phytochemistry* **1985**, *24*, 1779–1783.
- (20) Wong, H. F.; Brown, G. D. *Phytochemistry* **2002**, *59*, 529–536.
- (21) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4096.
- (22) Joseph-Nathan, P.; Villagomez, J. R.; Roman, L. U.; Hernandez, J. D. *Phytochemistry* **1990**, *29*, 977–979.
- (23) Tamayo-Castillo, G.; Jakupovic, J.; Bohlmann, F.; Rojas, A.; Castro, V.; King, R. M. *Phytochemistry* **1988**, *27*, 2893–2897.
- (24) Benyahia, S.; Benayache, S.; Benayache, F.; Quintana, J.; López, M.; León, F.; Hernández, J. C.; Estévez, F.; Bermejo, J. *J. Nat. Prod.* **2004**, *67*, 527–531.

NP0496183