

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/41011503>

Mass Spectrometric Evidence for the Existence of Distinct Modifications of Different Proteins by 2(E),4(E)-Decadienal

ARTICLE *in* CHEMICAL RESEARCH IN TOXICOLOGY · MARCH 2010

Impact Factor: 3.53 · DOI: 10.1021/tx900379a · Source: PubMed

CITATIONS

4

READS

34

6 AUTHORS, INCLUDING:



Xiaochun Zhu

Amgen

21 PUBLICATIONS 468 CITATIONS

SEE PROFILE



Jianye Zhang

Case Western Reserve University

23 PUBLICATIONS 330 CITATIONS

SEE PROFILE



Vernon Emmett Anderson

National Institutes of Health

159 PUBLICATIONS 4,169 CITATIONS

SEE PROFILE

Mass Spectrometric Evidence for the Existence of Distinct Modifications of Different Proteins by 2(*E*),4(*E*)-Decadienal

Xiaochun Zhu,^{*,†} Xiaoxia Tang,[†] Jianye Zhang,[†] Gregory P. Tochtrop,[†]
Vernon E. Anderson,^{§,‡} and Lawrence M. Sayre^{†,||}

Departments of Chemistry and Biochemistry, Case Western Reserve University, Cleveland, Ohio 44106

Received October 16, 2009

2(*E*),4(*E*)-Decadienal (DDE), a lipid peroxidation product, was found to covalently modify Lys residues of different proteins by different reactions using mass spectrometry (MALDI-TOF-MS and LC-ESI-MS). DDE mainly formed Lys Schiff base adducts with cytochrome *c* and ribonuclease A at 10 min, but these reversibly formed adducts almost disappeared after 24 h. In contrast, β -lactoglobulin (β -LG) was highly modified by DDE after 24 h. In addition to the Lys Schiff base adducts, DDE formed novel Lys pyridinium adducts as well as Cys Michael adducts with β -LG.

Introduction

Lipid peroxidation generates various aldehydes, including reactive α,β -unsaturated aldehydes. Characterization of protein modification by these aldehydes has been widely investigated using mass spectrometry in the last two decades. Among them, 4-hydroxy-2(*E*)-nonenal (HNE), acrolein, and malondialdehyde (MDA) are the most studied. HNE has been reported to form Cys/His/Lys Michael adducts, Lys Schiff base adducts, Lys pyrrole, Cys/His-Lys reversible cross-link adducts, and Lys-Lys fluorescent cross-link adduct (1–8). Acrolein has been reported to form the Cys/His Michael adduct, Lys Schiff base adduct, Lys 3-formyldihydropyridine adduct, Lys 3-methylpyridinium adduct, and His-Lys cross-link adduct (9–11). MDA has been reported to form the Lys-Lys bis-Schiff base cross-link adduct, Lys fluorescent adduct, and Arg-Lys cross-link adduct (12–16). More recently, 4-oxo-2(*E*)-nonenal (ONE) was discovered as a lipid peroxidation product (17, 18), which has been found to be a more reactive electrophile with proteins and to exhibit greater neurotoxicity than HNE (3, 19). ONE was reported to form the Cys/His/Lys Michael adduct, Lys Schiff base adduct, Lys pyrrolinone adduct, Lys 4-ketoamide adduct, and His/Cys-Lys pyrrole cross-link adduct with proteins (3, 4, 7, 19–21).

2(*E*),4(*E*)-Decadienal (DDE) is a widespread dialdehyde. It occurs in various foods and has been detected in fish (22, 23), chicken (24), meat (25), bread (26, 27), and heated oil (28). DDE has been shown to inhibit cell proliferation and induce the apoptosis of cells (29). It has also been reported to be carcinogenic and associated with lung adenocarcinoma in women (28). Additionally, DDE has been shown to covalently modify DNA (30, 31). However, protein modification by DDE has been rarely studied, and the nature of chemistry between

DDE and protein nucleophiles is not completely clear. The Medeiros group has shown that the Schiff base is the almost exclusive product found in DDE treated cytochrome *c* by mass spectrometry (32).

By incubating DDE with β -lactoglobulin (β -LG), cytochrome *c*, and ribonuclease A (RNase A), this study shows that DDE has a unique modification on the Lys residue of β -LG, although the other two are also Lys-rich proteins. This finding of differential selective modification of specific proteins by DDE strongly suggests that lipid peroxidation products can and will differentially modify specific proteins. To determine why some proteins are especially sensitive to specific modifications can help identify and/or predict which proteins will be sensitive toward each type of modification *in vivo*. To our knowledge, this is the first report on the unusual phenomenon that distinct modifications of the same type of amino acid were observed in different proteins exposed to a lipid peroxidation-derived aldehyde.

Materials and Methods

General. Chymotrypsin, cytochrome *c*, β -LG, and RNase A were purchased from Sigma. α -Cyano-4-hydroxycinnamic acid, DDE, iodoacetamide (IAM), and sinapinic acid were purchased from Acros Organics. Sequencing grade modified trypsin was received as a kit with buffer from Promega. PD-10 columns (Sephadex G-25) were obtained from GE Healthcare. ZipTips were purchased from Millipore. Dialysis tubing (Nominal MW 6000–8000) was purchased from Fisher Scientific. All other reagents were of the highest grade commercially available. The protein solution was concentrated through centrifugal evaporation under reduced pressure with a Savant Speed Vac SC110 system (Farma Scientific, Inc.). MALDI samples were spotted on the stainless steel target and analyzed with MALDI-TOF-MS as previously reported (33).

Proteolytic Digestion. Modified protein (1 mg) was dissolved in 100 μ L of pH 8.0 buffer containing 6 M guanidine hydrochloride and 50 mM Tris. After the addition of dithiothreitol (200 mM, 2.5 μ L), the sample was vortexed and incubated at 37 °C for 1 h. Then IAM (400 mM, 5.0 μ L) was added, vortexed, and allowed to stand at 25 °C in a dark place for 1 h. Dithiothreitol (200 mM, 7.5 μ L) was added again, vortexed, and allowed to consume the unreacted IAM at 25 °C for another 1 h. Twenty-five microliters of this denatured sample was added to 50 μ L of trypsin (0.1 μ g/ μ L) suspension buffer or 10 μ L of α -chymotrypsin (2.0 μ g/ μ L) solution. Then the digestion mixture was diluted to 200 μ L with pH 7.8

* To whom correspondence should be addressed. Xiaochun Zhu, Ph.D., Pharmacokinetics and Drug Metabolism, Amgen Inc., One Amgen Center Drive, 30E-2-C, Thousand Oaks, CA 91320, Tel: (805) 447-8385, E-mail: xiaochun@amgen.com.

[†] Department of Chemistry.

[§] Department of Biochemistry.

^{||} Current address: National Institute of General Medical Sciences, 45 Center Drive, Bethesda, MD 20892.

^{||} This work was performed in Dr. Sayre's laboratory. Dr. Sayre was a respectable mentor and good friend. He passed away on May 8, 2009. This paper is dedicated to him.




Figure 1. MALDI-TOF mass spectra of 0.25 mM cytochrome *c* (A) or RNase A (B) modified by 20 equivalents of DDE (5 mM) in pH 7.4 phosphate buffer at 37 °C for various times. The mass increment of 134 Da is due to the DDE-Lys Schiff base formation.

NH₄HCO₃ solution (50 mM), and the mixture was incubated at 37 °C for 24 h. The solution was stored at −20 °C for LC-MS or MALDI-TOF-MS analysis.

HPLC-ESI-MS/MS. ESI-MS was performed with a Thermo LCQ DecaXP MAX or Advantage instrument in the positive mode using nitrogen as the sheath and auxiliary gas. The capillary temperature was 300 °C, the capillary voltage was 35.00 V, and the source voltage was 4.50 kV. Typically, two scan events were used: (1) *m/z* 300–2000 full scan MS; (2) data dependent scan MS/MS on the most intense ion from event 1 or from the predefined parent mass list. The spectra were recorded using the dynamic exclusion of previously analyzed ions for 0.5 min with three repeats and a repeat duration of 0.5 min. The MS/MS normalized collision energy was set to 35%. Reversed-phase HPLC was performed with a Surveyor LC system equipped with a 5 μ m 2.1 \times 250 mm Grace Vydac C18 column with a gradient elution program at a flow rate of 200 μ L/min. Eluent A was a mixture of 95% H₂O, 5% MeOH, and 0.1% formic acid. Eluent B was a mixture of 95% MeOH, 5% H₂O, and 0.1% formic acid. The gradient program was 0–70 min, 80% A to 20% A; 70–75 min, 20% A to 80% A; 75–80 min, 80% A.

Preparation of Cys121-Carbamidomethylated β -LG. GT buffer was prepared as a pH 8.0 mixture of 6.0 M guanidine hydrochloride and 50 mM Tris. A solution of β -LG (2.5 mM, 200 μ L) added to GT buffer (1000 μ L) and water (1284.4 μ L) was incubated with IAM (400 mM, 15.6 μ L) at 25 °C in the dark for 20 h. A 2 μ L aliquot was subjected to MALDI-TOF-MS. The remainder was dialyzed against pH 7.4 sodium phosphate buffer (50 mM, 500 mL) to remove excess IAM three times with 6 h/time.

Incubation of Proteins with DDE for MALDI-TOF-MS. A solution of β -LG, Cys121-carbamidomethylated β -LG, cytochrome *c*, or RNase A (2.5 mM, 20 μ L) added to pH 7.4 phosphate buffer (50 mM, 160 μ L) was incubated with a solution of DDE in EtOH (50 mM, 20 μ L). The molar ratio of DDE to the protein was 20/1. After 10, 30 min, 1, 2, 6, and 24 h, 10 μ L of each sample was mixed with matrix and deposited onto the MALDI stainless steel target for analysis.

Preparation of DDE-Modified Proteins with or without NaBH₄ Quenching. A solution of β -LG or cytochrome *c* (2.5 mM, 100 μ L) added to pH 7.4 phosphate buffer (50 mM, 800 μ L) was incubated with a solution of DDE in EtOH (50 mM, 100 μ L) at 37 °C. The molar ratio of DDE to the protein was 20/1. After 24 h, a 500 μ L aliquot was diluted to 2.5 mL and subjected to a PD-10 column eluting with 3.5 mL of water to remove the unbound DDE and the buffer salts. The eluted solution was concentrated by centrifugal evaporation prior to proteolytic digestion. Another 500 μ L aliquot was treated with NaBH₄ (3.8 mg) at 25 °C overnight prior to PD-10 column elution.

Results and Discussion

Characterization of Intact DDE-Modified Protein by MALDI-TOF-MS. The Medeiros group reported that DDE




Figure 2. MALDI-TOF mass spectra of 0.25 mM β -LG modified by 20 equivalents of DDE (5 mM) in pH 7.4 phosphate buffer at 37 °C for various times. The mass increments of 134, 152, and 268 Da are due to the DDE-Lys Schiff base, DDE-Cys Michael adduct, and the DDE-Lys pyridinium adduct, respectively. Arrows a, b, and c indicate that the pyridinium adduct was gradually formed after 2 to 24 h on the β -LG A chain.

modified cytochrome *c* mainly through Lys Schiff base formation with a mass increment of 134 Da (32), which was also observed in this study (Figure 1A). In agreement with this result, when RNase A was incubated with DDE, the major product was also the Lys Schiff base adduct (Figure 1B). Moreover, the spectra indicate that DDE-Lys Schiff base formation is not stable during incubation under our conditions as the prominent +134 Da adduct gradually disappeared at longer incubation times, while apparently unmodified protein remained after 24 h.

However, when β -LG was incubated with DDE under the same conditions, the spectra observed revealed a totally different pattern of modification (Figure 2). The protein, composed of an A chain and B chain (33), was modified extensively by DDE over a period of 24 h. Figure 2 shows the formation of the DDE-Lys Schiff base (M+134) and DDE Michael adduct (M + 152) after 10 min. Under longer incubation times, the native β -LG peak decreases concomitant with an increase in the modified β -LG peaks. After 24 h, β -LG was heavily modified by DDE. In contrast, the spectra of DDE-modified cytochrome *c* and RNase A after 24 h are almost identical to those of their controls (Figure 1). Therefore, there are obvious distinctions in modification by DDE between β -LG and cytochrome *c* or RNase A.

β -LG contains a free Cys residue, i.e., Cys121, while neither cytochrome *c* nor RNase A has any free Cys. It is possible that




Figure 3. MALDI-TOF mass spectra of 0.25 mM Cys12-carbamidomethylated β -LG modified by 20 equivalents of DDE (5 mM) in pH 7.4 phosphate buffer at 37 °C at various times. The mass increments of 134 and 269 Da are attributed to the DDE-Lys Schiff base and DDE-Lys pyridinium adduct, respectively.

the above difference observed is attributable to the reaction between DDE and Cys121 in β -LG. To investigate this possibility, Cys121 of β -LG is protected by carbamidomethylation after the reaction with IAM. MALDI mass spectra show that β -LG has a mass increment of 57 Da after the treatment of IAM in the guanidine hydrochloride and Tris buffer after 20 h (Figure S1, Supporting Information), suggesting that Cys121 is effectively carbamidomethylated (Scheme S1, Supporting Information). Figure 3 displays MALDI-TOF mass spectra of DDE-modified carbamidomethylated β -LG at various times. It indicates that after 10 min some Schiff base adducts were formed on the protein ($M + n \times 134$), but Michael adducts were not observed ($M + 152$), suggesting that Michael adducts formed on DDE-modified β -LG (Figure 2) were ascribed to the Cys121 reaction. Longer incubation results in a more abundant native Cys121-carbamidomethylated β -LG (M) and less abundant Schiff base adducted peak ($M + 134$), presumably due to the reversibility of Schiff base formation. However, other than this, carbamidomethylated β -LG is still intensely modified by DDE after 24 h. The most abundant modified protein has a mass shift of around 269 Da, which is apparently equal to the addition of two Schiff base adducts (2×134). If this peak does arise from the independent formation of two Schiff base adducts, the mono-Schiff base bound protein ($M + 134$) should also be observed since the native protein (M) is as abundant as the bis-Schiff base bound protein ($M + 269$). However, the peak $M + 134$ is not evident after 24 h. Therefore, the mass shift of 269 Da must be attributed to an alternative modification(s). As Cys121 was protected, this special modification was obviously not ascribed to the reaction of Cys121.

Characterization of DDE-Modified β -LG by Mass Spectrometry after Proteolytic Digestion. To characterize the unique adduct of β -LG, proteolytic digests of DDE-modified β -LG were analyzed by MALDI-TOF-MS and LC-ESI-MS/MS. MALDI-TOF mass spectra of chymotryptic digests of the control and DDE-modified β -LG are shown in Figure 4. After 10 min, the mass spectra of the modified and control digests showed minimal differences. However, after 24 h, the mass spectrum profile changes substantially in comparison to those of 10 min as well as the control, revealing that a large number of novel modified peptides were generated after the long-time incubation. This dramatic change in the peptide fingerprint was




Figure 4. MALDI-TOF mass spectra of chymotryptic digests of 0.25 mM β -LG control (A) and β -LG modified by 20 equivalents of DDE (5 mM) after 10 min (B) or 24 h (C). The peaks marked with an asterisk are peptides modified by DDE through Lys pyridinium formation.

Table 1. Modified Peptides Detected by LC-ESI-MS/MS from Chymotryptic Digest of DDE Treated β -LG at 10 min or 24 h


| peptides ^a | position | <i>m/z</i> | modified <i>m/z</i> | assignment | incubation time |
|-----------------------|----------|------------|------------------------|------------|--------------------|
| LIVTQ | 1–5 | 573.4 | 841.4 | pyridinium | 24 h |
| KGL | 8–10 | 317.2 | 585.2 | pyridinium | 24 h |
| KGLDIQ | 8–13 | 673.3 | 941.4 | pyridinium | 24 h |
| DIQKVAGTW | 11–19 | 1017.5 | 1285.5 | pyridinium | 24 h |
| VEELKPTPEGDLEIL | 43–57 | 1681.9 | 975.5 ^b | pyridinium | 24 h |
| LQKW | 58–61 | 574.3 | 842.3 | pyridinium | 24 h |
| KIPAVF | 77–82 | 674.4 | 942.4 | pyridinium | 24 h |
| KIDAL | 83–87 | 559.3 | 827.3 | pyridinium | 24 h |
| KIDALNENKVL | 83–93 | 1256.7 | 763.1 ^b | pyridinium | 24 h |
| NENKVL | 88–93 | 716.3 | 984.3 | pyridinium | 24 h |
| KVL | 91–93 | 359.3 | 627.3 | pyridinium | 24 h |
| KKY | 100–102 | 438.3 | 706.3 | pyridinium | 24 h |
| VRTPEVDDEALEKF | 123–136 | 1647.8 | 958.7 ^b | pyridinium | 24 h |
| DKAL | 137–141 | 446.3 | 714.3 | pyridinium | 24 h |
| KALPM | 141–145 | 559.3 | 827.3 | pyridinium | 24 h |

^a Bold and underlined residue was modified. ^b Doubly charged peak.

observed in the tryptic digest as well (Figure S2, Supporting Information).

In order to locate the modified residues as well as their mass shifts, all of these unique peaks were separated and subjected

Scheme 1. Proposed Reaction Mechanism for the Formation of DDE-Lys Pyridinium Adducts 1a and 1b






Figure 5. Characteristic fragment ions Py^+ and M-285 of DDE-Lys pyridinium adducts.

to collision-induced dissociation (CID) by LC-ESI-MS/MS. The resulting tandem mass spectra indicate that all of the peptides marked with an asterisk in Figure 4 contain a modified Lys residue with a mass increment of 268 Da (Table 1). The addition of 268 Da suggests that the modification incorporates portions of at least two DDE molecules, whose molecular weight is 152. Arrows a, b, and c in Figure 2 demonstrate that this adduct was gradually produced after 2 to 24 h on the β -LG A chain. In accordance with this observation, this adduct was not detected after 10 min of incubation as shown in Figure 4B.

All of the modified peptides ($M + 268$) have unusually high intensities in MALDI-TOF mass spectra (Figures 4 and S2 (Supporting Information)), which are comparable to or even higher than that of the most abundant native peptide. In other words, they are at least either more abundant in the solution or much more sensitive to MALDI-TOF-MS than native peptides. However, even if all of the protein modifications occur through this reaction with a mass shift of 268 Da, Figure 2 shows that at most 5 Lys residues are modified in a β -LG molecule after 24 h. Since a β -LG chain contains 15 Lys residues and most of the protein molecules have less than 5 modified Lys residues, native Lys containing peptides should be more abundant than the modified ones in the sample solution. Therefore, these Lys modified peptides must be more sensitive to MALDI-TOF-MS than normal peptides. It is well known that the peptides containing a fixed positive charge are extremely sensitive to MALDI-TOF-MS (34–36). Various Lys pyridinium adducts involving two or more alkenal molecules have been detected

and characterized by MS and/or NMR upon incubation of alkenals with a model peptide or a protein (37–41). DDE belongs to the alkenal class, but with one more conjugated double bonds. Hence it may also form a pyridinium adduct with Lys residues. The extraordinary sensitivity to MALDI-TOF-MS of these modified peptides implicates the formation of the pyridinium derivative, which has a fixed positive charge. On the basis of the pyridinium product formed between alkenals and Lys (37, 38, 40), the proposed structures **1a** and **1b** of these DDE adducts as well as their formation mechanism are shown in Scheme 1. These two isomeric adducts were both detected in the LC-MS. Examples of extracted ion chromatograms of modified peptides KGL, KVL, DKAL, and DIQKVAGTW are shown in Figure S3 (Supporting Information). Each of them has two peaks with different intensity and retention time. However, their tandem mass spectra are identical.

Additionally, two prominent characteristic fragment ions (Figure 5) of these modified peptides were always observed, which corroborated the presence of the pyridinium moiety. One is m/z $M - 285$ resulting from a neutral loss of the pyridine moiety from the precursor ion. The other product ion is the protonated pyridine moiety m/z 286 except for peptides such as DIQKVAGTW whose tandem mass spectrum does not cover this ion because of the low mass cutoff rule for the ion trap. These two kinds of ions are known to be featured product ions of pyridinium cations from CID (39, 42, 43). Figure 6 shows four examples of tandem mass spectra of DDE-modified peptides, i.e., KVL, KIDAL, KGLDIQ, and DIQKVAGTW. All of them have these characteristic product ions in the spectra.

Additional chemical evidence that suggests the presence of the pyridinium moiety came from the reduction by sodium borohydride (40), which results in an additional mass increment of 3 Da (Table 2) and a dramatically reduced intensity in the MALDI-TOF mass spectra. The reduction of pyridinium **1a** ($M+268$) by NaBH_4 leads to the intermediate dihydropyridinium





Figure 6. Tandem mass spectra of modified peptides KVL (m/z 627.32), KIDAL (m/z 827.34), KGLDIQ (m/z 941.41), and DIQKVAGTW (m/z 1285.49) through DDE-Lys pyridinium formation. Fragment ions Py^+ and M-285 are shown in Figure 5. Asterisks denote the modified residue.

Table 2. Modified Peptides Detected by LC-ESI-MS/MS from a Chymotryptic Digest of DDE Treated β -LG at 10 min or 24 h after Reduction

| peptides ^a | position | <i>m/z</i> | modified <i>m/z</i> | assignment | incubation time |
|-----------------------|----------|------------|---------------------|----------------------------|-----------------|
| LIVTQ | 1–5 | 573.4 | 709.4 | reduced Lys Schiff base | 10 min, 24 h |
| KGLDIQ | 8–13 | 673.3 | 809.3 | reduced Lys Schiff base | 10 min, 24 h |
| VEELKPTPEGDLEIL | 43–57 | 1681.9 | 909.8 ^b | reduced Lys Schiff base | 10 min, 24 h |
| LQKW | 58–61 | 574.3 | 710.3 | reduced Lys Schiff base | 10 min, 24 h |
| KIPAVF | 77–82 | 674.4 | 810.4 | reduced Lys Schiff base | 10 min, 24 h |
| KIDAL | 83–87 | 559.3 | 695.3 | reduced Lys Schiff base | 10 min, 24 h |
| KIDALNENKVL | 83–93 | 1256.7 | 1392.7 | reduced Lys Schiff base | 10 min, 24 h |
| NENKVL | 88–93 | 716.3 | 852.3 | reduced Lys Schiff base | 10 min, 24 h |
| VRTPEVDDEALEKF | 123–136 | 1647.8 | 893.0 ^b | reduced Lys Schiff base | 10 min, 24 h |
| DKAL | 137–140 | 446.3 | 582.3 | reduced Lys Schiff base | 10 min, 24 h |
| KALPM | 141–145 | 559.3 | 695.3 | reduced Lys Schiff base | 10 min, 24 h |
| AC ^c QCL | 118–122 | 594.3 | 748.3 | reduced Cys Michael adduct | 10 min |
| VC ^c QCL | 118–122 | 622.3 | 776.3 | reduced Cys Michael adduct | 10 min |
| CL | 121–122 | 235.1 | 389.1 | reduced Cys Michael adduct | 10 min |
| KGL | 8–10 | 317.2 | 589.2 | reduced pyridinium | 24 h |
| VEELKPTPEGDLEIL | 43–57 | 1681.9 | 978.0 ^b | reduced pyridinium | 24 h |
| LQKW | 58–61 | 574.3 | 846.3 | reduced pyridinium | 24 h |
| KIDAL | 83–87 | 559.3 | 831.3 | reduced pyridinium | 24 h |
| KIDALNENKVL | 83–93 | 1256.7 | 1528.7 | reduced pyridinium | 24 h |
| KVL | 91–93 | 359.27 | 631.3 | reduced pyridinium | 24 h |
| VRTPEVDDEALEKF | 123–136 | 1647.8 | 961.0 ^b | reduced pyridinium | 24 h |
| DKAL | 137–140 | 446.3 | 718.3 | reduced pyridinium | 24 h |
| KALPM | 141–145 | 559.3 | 831.3 | reduced pyridinium | 24 h |

^a Bold and underlined residues were modified. ^b Doubly charged peak. ^c Cys was *S*-carbamidomethylated with IAM.

Scheme 2. Reduction of DDE-Lys Pyridinium Adduct 1a by NaBH₄

2 (M+269) (Scheme 2). After protonation and rearrangement to iminium, dihydropyridinium is further reduced to produce tetrahydropyridinium 3 (M + 271) (40). Compound 3 is a neutral amine, which requires protonation to be detected by mass spectrometry resulting in a *m/z* of M + 272. Therefore, all of the reduced pyridinium adducts listed in Table 2 have an apparent mass shift of 272 Da instead of 271 Da. Since sodium borohydride was in large excess in our experiment, every pyridinium adduct should be converted to tetrahydropyridinium. As tetrahydropyridinium has lost the fixed positive charge, it is not highly sensitive to MALDI-TOF-MS anymore.

Other than pyridinium adducts, DDE-Lys Schiff base adducts were detected after borohydride treatment (Table 2). They are not observed without reduction because they are reversibly formed and hydrolyze after standard denaturation and 24 h digestion treatment. In addition, DDE-Cys Michael adducts were observed in the reduced sample after 10 min of incubation. However, they were not observed after 24 h, suggesting that they were either reversible adducts or that they evolved into more advanced products. Considering that intensities of the native Cys containing peptides in LC-MS at 24 h decreased dramatically compared to that at 10 min (data not shown), it was possible that these Michael adducts were finally converted into other advanced products. Although both the reduced DDE-Lys Schiff base adduct and the DDE-Cys Michael adduct were

observed, a reduced Lys-DDE-Cys Schiff base–Michael adduct cross-link was not detected with our conditions and analysis.

Modification of cytochrome *c* by DDE is significantly different from that of β -LG. As a control to β -LG, cytochrome *c* was incubated with DDE for 10 min and 24 h and subjected to proteolytic digestion and mass spectrometry. DDE-Lys Schiff base was observed at 10 min as expected (32). However, no pyridinium adduct was detected after 10 min or 24 h of incubation by either MALDI-TOF-MS or LC-ESI-MS.

It is not clear what makes this difference in modifications between β -LG and cytochrome *c* by DDE. In fact, every Lys residue of β -LG was found to form the pyridinium adduct from combined results of both chymotryptic and tryptic digests. In contrast, none of the cytochrome *c* Lys residues were observed to form this adduct. The environment of Lys residues in X-ray structures of both proteins was investigated, but no obvious difference was observed. Further study is needed to answer this question.

Conclusions

DDE has been reported to exclusively form the Lys Schiff base adducts with cytochrome *c* (32), which to our knowledge is the first and only literature investigating the protein modification by DDE. In our study, we found that DDE not only forms

the Lys Schiff base adduct with β -LG but also forms other Lys products, pyridinium adducts, as well as DDE-Cys Michael adducts. After 24 h, β -LG was substantially modified by DDE, while cytochrome *c* and RNase A were minimally modified. The pyridinium adduct was observed when the DDE concentration was as low as 500 μ M with a DDE/protein molar ratio of 2/1 (data not shown). At physiologic conditions, where the DDE concentration may be much lower, the formation of the pyridinium adduct may be minimal. Further work is required to identify the conditions required to generate this pyridinium modification, as they may be physiologically relevant. Furthermore, these results indicate that DDE displays different behaviors with different proteins, and suggest that different proteins may have different modifications by the same reactive aldehyde. MDA, acrolein, HNE, and ONE etc. have been reported to modify proteins through various reactions. Although incubation of these aldehydes with different proteins has been investigated previously (44), none of them has been reported to modify different proteins in distinct reactions. Our results with DDE suggest that the surface nature of individual proteins can affect the observed products of modification by bifunctional lipid peroxidation products.

Acknowledgment. This work was supported by NIH Grants HL 53315 and AG 15885.

Supporting Information Available: MALDI-TOF mass spectra of β -LG control (A), β -LG incubated in guanidine hydrochloride and tris buffer for 20 h (B), and β -LG incubated with IAM in guanidine hydrochloride and tris buffer for 20 h (C); MALDI-TOF mass spectra of tryptic digests of 0.25 mM of β -LG control (A) and β -LG modified by 20 equivalents of DDE (5 mM) after 10 min (B) or 24 h (C); extracted ion chromatograms of modified peptides KGL, KVL, DKAL, and DIQKVAGTW through DDE-Lys pyridinium formation; carbamidomethylation of Cys residue by IAM. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Szveda, L. I., Uchida, K., Tsai, L., and Stadtman, E. R. (1993) Inactivation of glucose-6-phosphate dehydrogenase by 4-hydroxy-2-nonenal. Selective modification of an active-site lysine. *J. Biol. Chem.* 268, 3342–3347.
- (2) Nadkarni, D. V., and Sayre, L. M. (1995) Structural definition of early lysine and histidine adduction chemistry of 4-hydroxynonenal. *Chem. Res. Toxicol.* 8, 284–291.
- (3) Doorn, J. A., and Petersen, D. R. (2002) Covalent modification of amino acid nucleophiles by the lipid peroxidation products 4-hydroxy-2-nonenal and 4-oxo-2-nonenal. *Chem. Res. Toxicol.* 15, 1445–1450.
- (4) Liu, Z., Minkler, P. E., and Sayre, L. M. (2003) Mass spectroscopic characterization of protein modification by 4-hydroxy-2-(E)-nonenal and 4-oxo-2-(E)-nonenal. *Chem. Res. Toxicol.* 16, 901–911.
- (5) Cohn, J. A., Tsai, L., Friguet, B., and Szveda, L. I. (1996) Chemical characterization of a protein-4-hydroxy-2-nonenal cross-link: immunochemical detection in mitochondria exposed to oxidative stress. *Arch. Biochem. Biophys.* 328, 158–164.
- (6) Xu, G., and Sayre, L. M. (1998) Structural characterization of a 4-hydroxy-2-alkenal-derived fluorophore that contributes to lipoperoxidation-dependent protein crosslinking in aging and degenerative disease. *Chem. Res. Toxicol.* 11, 247–251.
- (7) Sayre, L. M., Lin, D., Yuan, Q., Zhu, X., and Tang, X. (2006) Protein adducts generated from products of lipid oxidation: focus on HNE and ONE. *Drug Metab. Rev.* 38, 651–675.
- (8) Szpac, M. E., Kim, H.-Y. H., Porter, N. A., and Liebler, D. C. (2008) Identification of proteins adducted by lipid peroxidation products in plasma and modifications of apolipoprotein A1 with a novel biotinylated phospholipid probe. *J. Proteome Res.* 7, 4237–4246.
- (9) Uchida, K., Kanematsu, M., Morimitsu, Y., Osawa, T., Noguchi, N., and Niki, E. (1998) Acrolein is a product of lipid peroxidation reaction. Formation of free acrolein and its conjugate with lysine residues in oxidized low density lipoproteins. *J. Biol. Chem.* 273, 16058–16066.
- (10) Furuhashi, A., Ishii, T., Kumazawa, S., Yamada, T., Nakayama, T., and Uchida, K. (2003) N^ε-(3-methylpyridinium)lysine, a major antigenic adduct generated in acrolein-modified protein. *J. Biol. Chem.* 278, 48658–48665.
- (11) Ishii, T., Yamada, T., Mori, T., Kumazawa, S., Uchida, K., and Nakayama, T. (2007) Characterization of acrolein-induced protein cross-links. *Free Radical Res.* 41, 1253–1260.
- (12) Kikugawa, K., and Beppu, M. (1987) Involvement of lipid oxidation products in the formation of fluorescent and cross-linked proteins. *Chem. Phys. Lipids* 44, 277–296.
- (13) Nair, V., Offerman, R. J., Turner, G. A., Pryor, A. N., and Baenziger, N. C. (1988) Fluorescent 1,4-dihydropyridines. The malondialdehyde connection. *Tetrahedron* 44, 2793–2803.
- (14) Slatter, D. A., Murray, M., and Bailey, A. J. (1998) Formation of a dihydropyridine derivative as a potential cross-link derived from malondialdehyde in physiological systems. *FEBS Lett.* 421, 180–184.
- (15) Itakura, K., Uchida, K., and Osawa, T. (1996) A novel fluorescent malondialdehyde-lysine adduct. *Chem. Phys. Lipids* 84, 75–79.
- (16) Slatter, D. A., Avery, N. C., and Bailey, A. J. (2004) Identification of a new cross-link and unique histidine adduct from bovine serum albumin incubated with malondialdehyde. *J. Biol. Chem.* 279, 61–69.
- (17) Lee, S. H., and Blair, I. A. (2000) Characterization of 4-oxo-2-nonenal as a novel product of lipid peroxidation. *Chem. Res. Toxicol.* 13, 698–702.
- (18) Spiteller, P., Kern, W., Reiner, J., and Spiteller, G. (2001) Aldehydic lipid peroxidation products derived from linoleic acid. *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids* 1531, 188–208.
- (19) Lin, D., Lee, H.-g., Liu, Q., Perry, G., Smith, M. A., and Sayre, L. M. (2005) 4-Oxo-2-nonenal is both more neurotoxic and more protein reactive than 4-hydroxy-2-nonenal. *Chem. Res. Toxicol.* 18, 1219–1231.
- (20) Oe, T., Arora, J. S., Lee, S. H., and Blair, I. A. (2003) A novel lipid hydroperoxide-derived cyclic covalent modification to Histone H4. *J. Biol. Chem.* 278, 42098–42105.
- (21) Zhang, W.-H., Liu, J., Xu, G., Yuan, Q., and Sayre, L. M. (2003) Model studies on protein side chain modification by 4-oxo-2-nonenal. *Chem. Res. Toxicol.* 16, 512–523.
- (22) Milo, C., and Grosch, W. (1996) Changes in the odorants of boiled salmon and cod as affected by the storage of the raw material. *J. Agric. Food Chem.* 44, 2366–2371.
- (23) Cha, Y. J., and Cadwallader, K. R. (1998) Aroma-active compounds in skipjack tuna sauce. *J. Agric. Food Chem.* 46, 1123–1128.
- (24) Farkas, P., Sadecka, J., Kovac, M., Siegmund, B., Leitner, E., and Pfannhauser, W. (1997) Key odorants of pressure-cooked hen meat. *Food Chem.* 60, 617–621.
- (25) Brewer, M. S., and Vega, J. D. (1995) Detectable odor thresholds of selected lipid oxidation compounds in a meat model system. *J. Food Sci.* 60, 592–595.
- (26) Schieberle, P., and Grosch, W. (1991) Potent odorants of the wheat bread crumb. Differences from the crust and effect of a longer dough fermentation. *Z. Lebensm.-Unters. Forsch.* 192, 130–135.
- (27) Vermeulen, N., Czerny, M., Gaenzle, M. G., Schieberle, P., and Vogel, R. F. (2007) Reduction of (E)-2-nonenal and (E, E)-2,4-decadienal during sourdough fermentation. *J. Cereal Sci.* 45, 78–87.
- (28) Chang, L. W., Lo, W.-S., and Lin, P. (2005) Trans,trans-2,4-decadienal, a product found in cooking oil fumes, induces cell proliferation and cytokine production due to reactive oxygen species in human bronchial epithelial cells. *Toxicol. Sci.* 87, 337–343.
- (29) Hansen, E., Even, Y., and Genevieve, A.-M. (2004) The alpha, beta, gamma, delta-unsaturated aldehyde 2-trans-4-trans-decadienal disturbs DNA replication and mitotic events in early sea urchin embryos. *Toxicol. Sci.* 81, 190–197.
- (30) Loureiro, A. P. M., Di Mascio, P., Gomes, O. F., and Medeiros, M. H. G. (2000) trans,trans-2,4-Decadienal-induced 1, N2-etheno-2'-deoxyguanosine adduct formation. *Chem. Res. Toxicol.* 13, 601–609.
- (31) Carvalho, V. M., Asahara, F., Di ascio, P., de Campos, I. P., Cadet, J., and Medeiros, M. H. G. (2000) Novel 1, N6-etheno-2'-deoxyadenosine adducts from lipid peroxidation products. *Chem. Res. Toxicol.* 13, 397–405.
- (32) Sigolo, C. A. O., Di Mascio, P., and Medeiros, M. H. G. (2007) Covalent modification of cytochrome c exposed to trans,trans-2,4-decadienal. *Chem. Res. Toxicol.* 20, 1099–1110.
- (33) Zhu, X., Gallogly, M. M., Mielay, J. J., Anderson, V. E., and Sayre, L. M. (2009) Covalent cross-linking of glutathione and carnosine to proteins by 4-oxo-2-nonenal. *Chem. Res. Toxicol.* 22, 1050–1059.
- (34) Liao, P.-C., and Allison, J. (1995) Enhanced detection of peptides in matrix-assisted laser desorption/ionization mass spectrometry through the use of charge-localized derivatives. *J. Mass Spectrom.* 30, 511–512.
- (35) Naven, T. J. P., and Harvey, D. J. (1996) Cationic derivatization of oligosaccharides with Girard's T reagent for improved performance in matrix-assisted laser desorption/ionization and electrospray mass spectrometry. *Rapid Commun. Mass Spectrom.* 10, 829–834.

- (36) Zhu, X., Anderson, V. E., and Sayre, L. M. (2009) Charge-derivatized amino acids facilitate model studies on protein side-chain modifications by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* 23, 2113–2124.
- (37) Zidek, L., Dolezel, P., Chmelik, J., Baker, A. G., and Novotny, M. (1997) Modification of horse heart cytochrome c with trans-2-hexenal. *Chem. Res. Toxicol.* 10, 702–710.
- (38) Baker, A., Zidek, L., Wiesler, D., Chmelik, J., Pagel, M., and Novotny, M. V. (1998) Reaction of N-acetylglycyllysine methyl ester with 2-alkenals: an alternative model for covalent modification of proteins. *Chem. Res. Toxicol.* 11, 730–740.
- (39) Baker, A. G., Wiesler, D., and Novotny, M. V. (1999) Tandem mass spectrometry of model peptides modified with Trans-2-hexenal, a product of lipid peroxidation. *J. Am. Soc. Mass Spectrom.* 10, 613–624.
- (40) Pournamdari, M., Saadi, A., Ellis, E., Andrew, R., Walker, B., and Watson, D. G. (2009) Development of a derivatisation method for the analysis of aldehyde modified amino acid residues in proteins by Fourier transform mass spectrometry. *Anal. Chim. Acta* 633, 216–222.
- (41) Ichihashi, K., Osawa, T., Toyokuni, S., and Uchida, K. (2001) Endogenous formation of protein adducts with carcinogenic aldehydes. Implications for oxidative stress. *J. Biol. Chem.* 276, 23903–23913.
- (42) Katritzky, A. R., Watson, C. H., Dega-Szafran, Z., and Eyler, J. R. (1990) Kinetics and mechanisms of nucleophilic displacements with heterocycles as leaving groups. Part 27. Collisionally activated dissociation of N-alkylpyridinium cations to pyridinium cation and olefins in the gas phase. *J. Am. Chem. Soc.* 112, 2479–2484.
- (43) Katritzky, A. R., Watson, C. H., Dega-Szafran, Z., and Eyler, J. R. (1990) Kinetics and mechanism of nucleophilic displacements with heterocycles as leaving groups. 26. Collisionally activated dissociation of N-alkylpyridinium cations to pyridine and alkyl cations in the gas phase. *J. Am. Chem. Soc.* 112, 2471–2478.
- (44) Codreanu, S. G., Zhang, B., Sobecki, S. M., Billheimer, D. D., and Liebler, D. C. (2009) Global analysis of protein damage by the lipid electrophile 4-hydroxy-2-nonenal. *Mol. Cell. Proteomics* 8, 670–680.

TX900379A