

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/23221006>

Combination of Perfluoroalkyl and Triazole Moieties: A New Recovery Strategy for TEMPO

ARTICLE in ORGANIC LETTERS · SEPTEMBER 2008

Impact Factor: 6.36 · DOI: 10.1021/ol801555f · Source: PubMed

CITATIONS

44

READS

46

5 AUTHORS, INCLUDING:



Tamilselvi Chinnusamy

Universität Regensburg

7 PUBLICATIONS 84 CITATIONS

SEE PROFILE



Petra Hilgers

Universität Regensburg

20 PUBLICATIONS 142 CITATIONS

SEE PROFILE



Oliver Reiser

Universität Regensburg

371 PUBLICATIONS 5,495 CITATIONS

SEE PROFILE

Combination of Perfluoroalkyl and Triazole Moieties: A New Recovery Strategy for TEMPO


Alexandru Gheorghe, Tamilselvi Chinnusamy, Erick Cuevas-Yañez, Petra Hilgers, and Oliver Reiser*

Institut für Organische Chemie, Universität Regensburg,
Universitätsstr. 31, 93053 Regensburg, Germany

oliver.reiser@chemie.uni-regensburg.de

Received July 9, 2008

ABSTRACT



The attachment of multiple triazole moieties and perfluoroalkyl chains to TEMPO promotes emulsion formation in dichloromethane/water, giving a highly active and easily recoverable catalyst for the oxidation of alcohols.

The use of perfluorinated alkyl chains as tags has been broadly applied to impose fluorous properties onto a given molecule. Arguably the most important application of this strategy is to separate a perfluorinated compound from a complex reaction mixture by fluorous biphasic separations, for example, by extraction or retention of the tagged molecule with a perfluorinated solvent, fluorous silica, or perfluorinated polymers.¹ As this property provides an attractive method for catalyst recovery and recycling, perfluorinated tags have been attached to several metal-based catalysts² and organo-catalysts.³ Thus, homogeneous catalysis is achieved in an

organic solvent due to the good solubility perfluorinated tagged compounds still have, but easy and selective recovery of the catalyst by a phase switch to a perfluorinated medium is possible.

A well-known alternative approach for the development of recyclable catalysts is their attachment to a polymeric support that is insoluble in the reaction medium, allowing its recovery by simple filtration. An inherent disadvantage of this strategy is the generally reduced reactivity of such modified catalysts due to the heterogeneous nature of the reaction.⁴

We report here the combination of multiple perfluoroalkyl chains and triazole moieties, leading to a new kind of building block that can be used for the tagging of TEMPO. The resulting reagent promotes the oxidation of alcohols to aldehydes in organic solvent/water mixtures with reaction


(1) Leading reviews: (a) Gladysz, J. A.; Curran, D. P. *Tetrahedron* **2002**, 58, 3823. (b) Tzschucke, C. C.; Markert, C.; Bannwarth, W.; Roller, S.; Hebel, A.; Haag, R. *Angew. Chem., Int. Ed.* **2002**, 41, 3964. (c) Curran, D. P. In *Handbook of Fluorous Chemistry*; Gladysz, J. A., Curran, D. P., Horváth, I. T., Eds.; Wiley-VCH, Weinheim, 2004; p 101.

(2) Leading examples: (a) Dinh, L. V.; Gladysz, J. A. *Angew. Chem., Int. Ed.* **2005**, 44, 4095. (b) Matsugi, M.; Curran, D. P. *J. Org. Chem.* **2005**, 70, 1636. (c) Contel, M.; Izuel, C.; Laguna, M.; Villuendas, P. R.; Alonso, P. J.; Fish, R. H. *Chem. Eur. J.* **2003**, 9, 4168. (d) Tzschucke, C. C.; Markert, C.; Glatz, H.; Bannwarth, W. *Angew. Chem., Int. Ed.* **2002**, 41, 4500.

(3) Leading examples: (a) Zu, L.; Wang, J.; Li, H.; Wang, W. *Org. Lett.* **2006**, 8, 3077. (b) Dalicsek, Z.; Pollreis, F.; Gömöry, A.; Soós, T. *Org. Lett.* **2005**, 7, 3243.

(4) Leading reviews: (a) De Vos, D. E.; Sels, B. F.; Jacobs, P. A. *Adv. Catal.* **2001**, 46, 1. (b) Gladysz, J. A. *Pure Appl. Chem.* **2001**, 73, 1319. (c) Oehme, G. In *Comprehensive Asymmetric Catalysis I-III*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: Berlin, Germany, 1999; Vol 3, p 1377.

rates comparable to homogeneous TEMPO reagents, but can be easily recovered by liquid/emulsion filtration.




Based on our interest in developing immobilized reagents on organic and inorganic supports,⁵ we recently utilized the copper-catalyzed⁶ alkyne azide cycloaddition⁷ (CuAAC) reaction as a versatile method for the ligation of TEMPO to polymers⁸ such as **1** or **2**.¹⁰ In this context, we discovered that **1** showed low solubility in perfluorinated, but good solubility in organic solvents, and more strikingly, showed little affinity to fluorosilica but high affinity to conventional silica.⁹

To further investigate the combination of perfluorinated alkyl chains and triazole moieties, we attempted to synthesize azides **8** and **9-F**, which should be suitable building blocks for the attachment of catalysts and reagents modified by an alkynyl group.

The benzyl chlorides **5**¹¹ and **6** could be readily synthesized in a three-step procedure starting from commercially available benzoic acid esters **3** and **4**, respectively (Scheme


Scheme 1. Synthesis of Benzyl Azides with Multiple Triazole and Perfluoroalkyl Moieties



1). Ligation with azide **7-R** using catalytic amounts of CuI¹² followed by chloride azide exchange resulted in the target compounds **8** and **9** in high yields.

The fluorosilica tagged azides **8** or **9-F** were only slightly soluble in organic solvents, nevertheless, they smoothly underwent the CuAAC with **10** in high yields, again using catalytic amounts of CuI in THF (Scheme 2). **11** and **12** precipitated from the reaction mixture and could be conve-

Scheme 2. Synthesis of Novel TEMPO Reagents with Multiple Triazole and Perfluoroalkyl Moieties



(5) (a) Glos, M.; Reiser, O. *Org. Lett.* **2000**, *2*, 2045. (b) Fraile, J. M.; García, J. I.; Herrerías, C. I.; Mayoral, J. A.; Reiser, O.; Socuélamos, A.; Werner, H. *Chem. Eur. J.* **2004**, *10*, 2997. (c) Gissibl, A.; Finn, M. G.; Reiser, O. *Org. Lett.* **2005**, *7*, 2325. (d) Werner, H.; Herrerías, C. I.; Glos, M.; Gissibl, A.; Fraile, J. M.; Pérez, I.; Mayoral, J. A.; Reiser, O. *Adv. Synth. Catal.* **2006**, *348*, 125. (e) Gissibl, A.; Padié, C.; Hager, M.; Jaroschik, F.; Rasappan, R.; Cuevas-Yañez, E.; Turrin, C.-O.; Caminade, A.-M.; Majoral, J.-P.; Reiser, O. *Org. Lett.* **2007**, *9*, 2895.

(6) (a) Tornøe, C. W.; Meldal, M. In *17th American Peptide Symposium*; Lebl, M.; Houghten, R. A., Eds.; American Peptide Society and Kluwer Academic Publishers: San Diego, CA, 2001; p 263. (b) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596. (c) Tornøe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057.

(7) Huisgen, R. *Pure Appl. Chem.* **1989**, *61*, 613.

(8) Immobilized TEMPO reagents: (a) Bolm, C.; Fey, T. *Chem. Commun.* **1999**, 1795. (b) Dijkman, A.; Arends, I. W. C. E.; Sheldon, R. A. *Chem. Commun.* **2000**, 271. (c) Weik, S.; Nicholson, G.; Jung, G.; Rademann, J. *Angew. Chem., Int. Ed.* **2001**, *40*, 1436. (d) Fey, T.; Fischer, H.; Bachmann, S.; Albert, K.; Bolm, C. *J. Org. Chem.* **2001**, *66*, 8154. (e) Brunel, D.; Fajula, F.; Nagy, J. B.; Deroide, B.; Verhoef, M. J.; Veum, L.; Peters, J. A.; van Bekkum, H. *Appl. Catal. A* **2001**, *213*, 73. (f) Ciriminna, R.; Pagliaro, M.; Blum, J.; Avnir, D. *Chem. Commun.* **2000**, 1441. (g) Ciriminna, R.; Bolm, C.; Fey, T.; Pagliaro, M. *Adv. Synth. Catal.* **2002**, *344*, 159. (h) Pozzi, G.; Cavazzini, M.; Quici, S.; Benaglia, M.; Dell'Anna, G. *Org. Lett.* **2004**, *6*, 441. (i) Ferreira, P.; Hayes, W.; Phillips, E.; Rippon, D.; Tsang, S. C. *Green Chem.* **2004**, *6*, 310. (j) Benaglia, M.; Puglisi, A.; Holczknecht, O.; Quici, S.; Pozzi, G. *Tetrahedron* **2005**, *61*, 12058. (k) Tanyeli, C.; Gümmüş, A. *Tetrahedron Lett.* **2003**, *44*, 1639. (l) Holczknecht, O.; Cavazzini, M.; Quici, S.; Shepperson, I.; Pozzi, G. *Adv. Synth. Catal.* **2005**, *347*, 677. (m) Pozzi, G.; Cavazzini, M.; Holczknecht, O.; Quici, S.; Shepperson, I. *Tetrahedron Lett.* **2004**, *45*, 4249. (n) Luo, J.; Pardin, C.; Lubell, W. D.; Zhu, X. X. *Chem. Commun.* **2007**, 2136. (o) Gilhespy, M.; Lok, M.; Baucherel, X. *Chem. Commun.* **2005**, 1085. (p) Gilhespy, M.; Lok, M.; Baucherel, X. *Catal. Today* **2006**, *117*, 114. (q) Karimi, B.; Biglari, A.; Clark, J. H.; Budarin, V. *Angew. Chem., Int. Ed.* **2007**, *46*, 7210. (r) Mason, B. P.; Bogdan, A. R.; Goswami, A.; McQuade, D. T. *Org. Lett.* **2007**, *9*, 3449.

(9) Gheorghe, A.; Cuevas-Yañez, E.; Horn, J.; Bannwarth, W.; Narsaiah, B.; Reiser, O. *Synlett* **2006**, 2767.

(10) Gheorghe, A.; Matsuno, A.; Reiser, O. *Adv. Synth. Catal.* **2006**, *348*, 1016.

Table 1. Oxidation of Alcohols with **11** and **12**^a

| entry | alcohol | catalyst | conversion ^b (%) | yield (%) | purity ^b (%) |
|-------|---------------------------|-----------|--------------------------------|-----------|----------------------------|
| 1 | 4-bromobenzyl alcohol | 11 | >98 | 95 | >98 |
| 2 | | 12 | >98 | 95 | >98 |
| 3 | 4-methylbenzyl alcohol | 11 | >98 | 98 | >98 |
| 4 | | 12 | >98 | 98 | >98 |
| 5 | 1-octanol | 11 | >98 | 95 | >97 |
| 6 | | 12 | >98 | 96 | >96 |
| 7 | 1-decanol | 11 | >98 | 98 | >96 |
| 8 | | 12 | >98 | 92 | >95 |
| 9 | cyclohexanol ^c | 11 | >98 | 92 | >98 |
| 10 | | 12 | >98 | 91 | >98 |

^a Alcohol (1.0 mmol) in CH₂Cl₂ (2 mL), KBr (0.2 mmol), catalyst (1.0 mol %), NaOCl (0.8 mL, 1.1 mmol), NaHCO₃ (0.5 mmol), 0 °C. Reaction time 15 min. ^b Determined by ¹H NMR. ^c Reaction time 1 h.

niently isolated by simple filtration and purified from residual copper by washing with aqueous EDTA (0.1 M) solution. The absence of the typical azide band around 2100 cm⁻¹ in the IR-spectrum further indicated that the products were obtained free from the starting materials **8** or **9-F**, respectively.

The new TEMPO derivatives **11** and **12** showed remarkable properties: They are largely insoluble in organic or fluorinated solvents (≤ 1 mg/10 mL CH₂Cl₂ or perfluoro-(methylcyclohexane) at 25 °C) as well as in water; nevertheless, oxidation of alcohols using Anelli's conditions,¹³ calling for KBr as the cocatalyst and NaOCl as the terminal oxidant, or using Minisci's conditions,¹⁴ calling for Mn(II) and Co(II) nitrate as cocatalysts in acetic acid and oxygen as the terminal oxidant, proceeded with high yields and selectivity. Under both protocols, **12** could be recovered with high efficiency and subsequently reused as such without loss of activity.

Oxidations of alcohols were carried out at 0 °C in the presence of 1 mol % of catalyst and 1.1 equiv of NaOCl (Table 1-3). Under these conditions, most alcohols were quantitatively oxidized within 15 min, with the exception of the more challenging cyclohexanol, which was smoothly oxidized within 1 h. In all cases, the purity of the isolated carbonyl derivatives was >95%.

Table 2. Activity Comparison of Heterogeneous (**2**, **12**) and Homogeneous (TEMPO, **1**, **13**) TEMPO Catalysts in the Anelli Oxidation of 4-Methylbenzyl Alcohol^a

| entry | catalyst (mol %) | time (min) | yield ^b (%) | TON | TOF (h ⁻¹) |
|-------|------------------|------------|------------------------|-----|------------------------|
| 1 | TEMPO (0.2) | 40 | ≥ 98 | 500 | ≥ 750 |
| 2 | TEMPO (0.2) | 15 | 60 | 300 | 1200 |
| 3 | 1 (0.2) | 40 | ≥ 98 | 500 | ≥ 750 |
| 4 | 2 (2.5) | 15 | 68 | 29 | 116 |
| 5 | 2 (0.2) | 40 | 58 | 300 | 450 |
| 6 | 12 (1.0) | 15 | ≥ 98 | 100 | ≥ 400 |
| 7 | 12 (0.2) | 40 | 97 | 500 | ≥ 750 |
| 8 | 12 (0.2) | 15 | 74 | 370 | 1480 |
| 9 | 13 (0.2) | 40 | 96 | 500 | ≥ 750 |

^a Alcohol (25 mmol, 1.0 equiv) in CH₂Cl₂ (50 mL), KBr (0.2 equiv), TEMPO catalyst, NaOCl (1.1 equiv, 20 mL), NaHCO₃ (0.5 equiv), 0 °C. ^b Purity >98% in all cases.

Table 3. Oxidation of 4-Methylbenzyl Alcohol with TEMPO **12**. Recovery and Recycling Experiment^a

| run | conversion ^b (%) | yield (%) | purity ^b (%) |
|-----|-----------------------------|-----------|-------------------------|
| 1 | >98 | 98 | >98 |
| 2 | >98 | 99 | >98 |
| 3 | >98 | 97 | >98 |
| 4 | >98 | 92 | >98 |

^a Alcohol (2.5 mmol), CH₂Cl₂ (5 mL), KBr (0.5 mmol), **12** (1.0 mol %), NaOCl (2.0 mL, 2.8 mmol), NaHCO₃ (1.3 mmol), 0 °C. Reaction time 15 min. **12** was recovered by filtration and reused after washing once with water and twice with CH₂Cl₂. ^b Determined by ¹H NMR.

Notably, the activity of **12** (Table 2, entries 6–8) in the oxidation of 4-methylbenzylalcohol on a 25 mmol scale in CH₂Cl₂/water is about four times higher than polystyrene supported TEMPO **2** (entries 4–5) and at least comparable to that of TEMPO or fluorine tagged TEMPO **1** (entries 1–3), both being soluble in the solvent system employed for the reaction.

However, **12** can be recovered in a highly efficient manner by simple filtration of the reaction mixture through a standard sintered glass funnel (16–40 μ m pore size) and reused without further purification. For example, the oxidation of 4-methylbenzyl alcohol to the corresponding aldehyde was carried out in four runs in 92–98% yield with no observable loss of activity using an initial amount of 1 mol % of catalyst **12** (Table 3). Analysis of the filtrates by NMR gave no indication that **12** or fragments thereof due to degradation were present. Moreover, the catalytic activity of the filtrates for alcohol oxidations under the reaction conditions was in the limits of the known^{13,15} background reaction in the absence of any catalyst (see Supporting Information). In contrast, **11**, although also insoluble but apparently more highly dispersed in CH₂Cl₂/water, was not held back by a sintered glass funnel as described above for **12**. Therefore, its recovery by filtration was not possible, but could be readily achieved by chromatography due to its high affinity to silica in a similar way as already noticed for **1**.⁹ For practical purposes, **12** therefore seems to be the catalyst of choice.

The high activity of **12** can be most likely attributed to its behavior to act like a surfactant by forming an emulsion between the organic and aqueous phase, suggesting a detergent effect that enhances catalysis (Figure 1).¹⁶ The combination of multiple polar triazole and perfluoroalkyl

(11) Wu, P.; Feldman, A. K.; Nugent, A. K.; Hawker, C. J.; Scheel, A.; Voit, B.; Pyun, J.; Frechet, J. M. J.; Sharpless, K. B.; Fokin, V. V. *Angew. Chem., Int. Ed.* **2004**, *43*, 3928.

(12) Löber, S.; Rodriguez-Loaiza, P.; Gmeiner, P. *Org. Lett.* **2003**, *5*, 1753.

(13) Anelli, P. L.; Biffi, C.; Montanari, F.; Quici, S. *J. Org. Chem.* **1987**, *52*, 2559.

(14) (a) Cecchetto, A.; Fontana, F.; Minisci, F.; Recupero, F. *Tetrahedron Lett.* **2001**, *42*, 6651. (b) Minisci, F.; Recupero, F.; Rodino, M.; Sala, M.; Schneider, A. *Org. Proc. Res. Dev.* **2003**, *7*, 794.

(15) Huang, J.-Y.; Li, S.-J.; Wang, Y.-G. *Tetrahedron Lett.* **2006**, *47*, 5637.

(16) Dwars, T.; Paetzold, E.; Oehme, G. *Angew. Chem., Int. Ed.* **2005**, *44*, 7174.




Figure 1. Reaction mixture containing **12** (left; Table 2, entry 7) or **13** (right; Table 2, entry 9).

chains seems to be decisive for this property since both, **1** or **13**, the latter being analogous to **12** but having the perfluoroalkyl chains replaced by benzyl groups, are readily soluble in dichloromethane. Those display similar activities as **12** (Table 2, entries 3 and 9) but cannot be recovered by simple filtration from the reaction mixture.

Alternatively, the TEMPO derivative **12** was investigated under Minisci's conditions, using Mn(II)- and Co(II)-nitrate as cocatalysts in acetic acid oxygen as the terminal oxidant.¹⁴ Under these conditions benzylic alcohols were oxidized within 2–4 h, while 1-octanol, cyclohexyl alcohol and cinnamyl alcohol required 7–20 h (Table 4) to reach conversions $\geq 90\%$.

Table 4. Aerobic Oxidation of Alcohols with TEMPO **12**^a

| entry | alcohol | time (h) | conversion ^b (%) | yield (%) |
|----------------|-------------------------|----------|-----------------------------|-----------|
| 1 | benzyl alcohol | 2 | 90 | 89 |
| 2 | 4-methylbenzyl alcohol | 2 | >98 | 98 |
| 3 | 4-methoxybenzyl alcohol | 3 | >98 | 96 |
| 4 | 4-bromobenzyl alcohol | 3 | 95 | 92 |
| 5 | cinnamyl alcohol | 20 | >98 | 90 |
| 6 | cyclohexyl alcohol | 12 | >98 | 93 |
| 7 ^c | 1-octanol | 7 | 90 | 82 |
| 8 | thiophen-2-yl-methanol | 4 | >98 | 91 |

^a Alcohol (1.0 mmol) in AcOH (1 mL), Mn(NO₃)₂·4H₂O (2.0 mol %), Co(NO₃)₂·6 H₂O (2.0 mol %), **12** (1.0 mol %), oxygen atmosphere, 40 °C. ^b Determined by ¹H NMR. ^c 5.0 mol % of catalyst used.

In contrast to unmodified TEMPO, **12** can be also recovered by simple filtration of the reaction mixture through a standard sintered glass funnel (see above) and reused without further purification. For example, the oxidation of 4-methylbenzyl alcohol to the corresponding aldehyde was carried out in three runs in 96–98% yield with consistent catalytic activity (Table 5).

Again, the activity of **12** (Table 6) in the oxidation of 4-methyl benzyl alcohol is about three to four times higher

Table 5. Oxidation of 4-Methylbenzyl Alcohol with TEMPO **12**. Recovery and Recycling Experiment^a

| run | conversion ^b (%) | yield (%) | purity ^b (%) |
|-----|-----------------------------|-----------|-------------------------|
| 1 | >98 | 98 | >98 |
| 2 | >98 | 99 | >98 |
| 3 | 97 | 96 | >98 |

^a Alcohol (3.0 mmol) in AcOH (3 mL), for conditions see Table 4, reaction time = 2 h. **12** was recovered by filtration and reused after washing once with water and twice with CH₂Cl₂. ^b Determined by ¹H NMR.

than polystyrene supported TEMPO **2** and only slightly reduced to that of unmodified TEMPO.

Table 6. Activity Comparison of Heterogeneous (**2**, **12**) and Homogeneous TEMPO Catalysts in the Aerobic Oxidation of 4-Methylbenzyl Alcohol^a

| catalyst | conversion ^b (%) | yield (%) |
|-----------|-----------------------------|-----------|
| TEMPO | 97 | 92 |
| 2 | 26 | 25 |
| 12 | 84 | 82 |

^a For conditions, see Table 4, reaction time = 1 h. ^b Determined by ¹H NMR.

In conclusion, we have developed a new TEMPO derivative that retains the activity of homogeneous TEMPO catalysts for the oxidation of alcohols, but nevertheless allows recovery by simple filtration. The fluororous/triazole building blocks developed here should also be suitable for other catalysts and reagents that are carried out in aqueous/organic two-phase media, which is under investigation in our laboratories.

Acknowledgment. This work was supported by the Degussa foundation (fellowship for C.T.), the DAAD (fellowship for E. C.-Y.), the DFG (HI-1329/1-1 and SPP 1179 Organokatalyse), and the Fonds der Chemischen Industrie. We thank Dr. Marianne Engeser, University of Bonn, for carrying out high-resolution mass spectra analyses on compounds **11** and **12**. Helpful comments from Prof. John Gladysz, Texas A&M University, and the reviewers during the preparation of this manuscript are gratefully acknowledged.

Supporting Information Available: Experimental procedures and characterization data for compounds **8**, **9**, **11**, **12**, and **13**; details on catalyzes performed and catalyst recovery. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL801555F