

Erratum: Force Constants and Normal Vibrations of the Propargyl Halides

G. Zerbi and M. Gussoni

Citation: The Journal of Chemical Physics 46, 2450 (1967); doi: 10.1063/1.1841065

View online: http://dx.doi.org/10.1063/1.1841065

View Table of Contents: http://scitation.aip.org/content/aip/journal/jcp/46/6?ver=pdfcov

Published by the AIP Publishing

Articles you may be interested in

Normal coordinates calculation and valence force field for propyne, propargyl halides, butyne-2, halobutyne-2, and 1,4-dihalobutyne-2

J. Chem. Phys. 70, 4786 (1979); 10.1063/1.437246

Erratum: Force constants of the metaborate ion in alkali halides

J. Chem. Phys. 58, 1782 (1973); 10.1063/1.1679431

Fundamental Vibrations and Force Constants in the Partially Deuterated Methyl Halides

J. Chem. Phys. 44, 745 (1966); 10.1063/1.1726756

Force Constants and Normal Vibrations of the Propargyl Halides

J. Chem. Phys. 41, 456 (1964); 10.1063/1.1725889

Normal Vibrations and Urey-Bradley Force Constants of Methyl Amine

J. Chem. Phys. 35, 2060 (1961); 10.1063/1.1732209



Errata

presented in my paper will serve as a useful supplement to the paper of Dobryakov.

¹ R. H. Silsbee, J. Chem. Phys. **45**, 1710 (1966). ² S. N. Dobryakov, Zh. Strukt. Khim. **6**, 39 (1965) [English transl.: J. Struct. Chem. (USSR) **6**, 30 (1965)].

Erratum: Force Constants and Normal Vibrations of the Propargyl Halides

[J. Chem. Phys. 41, 456 (1964)]

G. ZERBI AND M. GUSSONI

Istituto di Chimica Industriale del Politecnico—Piazza L. da Vinci, 32 Milano, Italy

THE value of the force constant $F_{\rm HX}$ for propargyl fluoride in Table III, p. 459, should read 1.052 mdyn/Å instead of 1.52 as erroneously reported. This is only a typographical error and does not affect the calculations and the content of the paper.

Erratum: Flash Photolysis of H₂O₂ Vapor in the Presence of D₂, Ar, and H₂¹⁸O

[J. Chem. Phys. 45, 99 (1966)]

N. R. GREINER

University of California and Los Alamos Scientific Laboratory
Los Alamos, New Mexico

CONTINUED work on the H_2O_2 -Ar system at room temperature has shown that $k_e = k_{10} = 5.2 \times 10^{11}$ cc mole⁻¹·sec⁻¹ = 8.5×10^{-13} cc molecule⁻¹·sec⁻¹, rather than 5.8×10^{-14} cc molecule⁻¹·sec⁻¹ as reported in this

paper. It is believed that something introduced into the cell during repairs caused the rapid dark decomposition of H_2O_2 resulting in a low observed OH disappearance rate. All the other experiments reported were done before the repairs when the dark decomposition was known to be insignificant.

Erratum: Nuclear Quadrupole Resonance of Antimony Tribromide and Its Molecular Complexes

[J. Chem. Phys. 45, 1076 (1966)]

HISAO NEGITA, TSUTOMU OKUDA, AND MIKITO KASHIMA
Department of Chemistry, Hiroshima University, Hiroshima, Japan

THE chemical formulas of the complexes and the lowest resonance frequency of antimony tribromide (2) in Table I should be corrected as follows:

2SbBr₃·C₆H₆, 2SbBr₃·C₁₄H₁₀, and 135. 112 Mc/sec

instead of

SbBr₃·C₆H₆, SbBr₃·C₁₄H₁₀, and 133. 112 Mc/sec.

Three resonance frequencies of antimony tribromide (2) are not completely consistent with those of the form I by Ogawa, indicating that the former crystal structure is slightly different from the latter. Therefore, there should be at least four forms in the crystal of antimony tribromide at a given temperature between room and liquid-nitrogen temperatures.

¹ S. Ogawa, J. Phys. Soc. Japan 13, 618 (1958).