

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5900216>

Mercurial-Induced Hydrogen Peroxide Generation in Mouse Brain Mitochondria: Protective Effects of Quercetin

ARTICLE *in* CHEMICAL RESEARCH IN TOXICOLOGY · JANUARY 2008

Impact Factor: 3.53 · DOI: 10.1021/tx7002323 · Source: PubMed

CITATIONS

70

READS

45

11 AUTHORS, INCLUDING:



Jeferson Franco

Universidade Federal do Pampa (Unipampa)

65 PUBLICATIONS 1,305 CITATIONS

[SEE PROFILE](#)



Thaís Posser

Universidade Federal do Pampa (Unipampa)

54 PUBLICATIONS 797 CITATIONS

[SEE PROFILE](#)



Adair Roberto Soares Santos

Federal University of Santa Catarina

303 PUBLICATIONS 7,810 CITATIONS

[SEE PROFILE](#)



Marcelo Farina

Federal University of Santa Catarina

161 PUBLICATIONS 3,395 CITATIONS

[SEE PROFILE](#)

Mercurial-Induced Hydrogen Peroxide Generation in Mouse Brain Mitochondria: Protective Effects of Quercetin

Jeferson L. Franco,^{†,‡} Hugo C. Braga,[†] James Stringari,[†] Fabiana C. Missau,[§] Thais Posser,[†] Beatriz G. Mendes,[§] Rodrigo B. Leal,[†] Adair R. S. Santos,[‡] Alcir L. Dafre,[‡] Moacir G. Pizzolatti,[§] and Marcelo Farina^{*,†}

Departamento de Bioquímica, Centro de Ciências Biológicas, Departamento de Ciências Fisiológicas, Centro de Ciências Biológicas, and Departamento de Química, Centro de Ciências Físicas e Matemáticas, Universidade Federal de Santa Catarina, 88040-900, Florianópolis, SC, Brazil

Received June 28, 2007

Plants of the genus *Polygala* have been shown to possess protective effects against neuronal death and cognitive impairments in neurodegenerative disorders related to excitotoxicity. Moreover, previous reports from our group have shown the neuroprotective effects of the plant *Polygala paniculata* against methylmercury (MeHg)-induced neurotoxicity. In this work, we have examined the potential protective effects of three compounds (7-prenyloxy-6-methoxycoumarin, quercetin, and 1,5-dihidroxi-2,3-dimethoxy xanthone) from *Polygala* species against MeHg- and mercuric chloride (HgCl₂)-induced disruption of mitochondrial function under in vitro conditions using mitochondrial-enriched fractions from mouse brain. MeHg and HgCl₂ (10–100 μ M) significantly decreased mitochondrial viability; this phenomenon was positively correlated to mercurial-induced glutathione oxidation. Among the isolated compounds, only quercetin (100–300 μ M) prevented mercurial-induced disruption of mitochondrial viability. Moreover, quercetin, which did not display any chelating effect on MeHg or HgCl₂, prevented mercurial-induced glutathione oxidation. The present results suggest that the protective effects of quercetin against mercurial-induced mitochondrial dysfunction is related to the removal of oxidant species generated in the presence of either MeHg or HgCl₂. Reinforcing this hypothesis, MeHg and HgCl₂ increased the production of hydrogen peroxide in the brain mitochondria, as well as the levels of malondialdehyde. These oxidative phenomena were prevented by co-incubation with quercetin or catalase. These results are the first to show the involvement of hydrogen peroxide as a crucial molecule related to the toxic effects of both organic and inorganic mercurials in brain mitochondria. In addition, the study is the first to show the protective effect of quercetin against mercurial-induced toxicity, pointing to its capability to counteract mercurial-dependent hydrogen peroxide generation as a potential molecular mechanism of protection. Taken together, these data render quercetin a promising molecule for pharmacological studies with respects to mercurials' poisoning.

Introduction

Methylmercury (MeHg) is a highly neurotoxic compound leading to neurological and developmental deficits in animals and humans (1). Although MeHg-induced neurotoxicity is an extensively reported phenomenon, the molecular mechanisms underlying its toxicity are still under debate. The major mechanisms involved in MeHg neurotoxicity currently being explored are the impairment of intracellular calcium homeostasis (2), the alteration of glutamate homeostasis (3–6), and oxidative stress (7). The last seems to be related to the direct oxidative properties of MeHg on endogenous thiols (8) and to its effects toward antioxidant enzymes (9–11). In addition, it is noteworthy that MeHg-induced oxidative stress and MeHg-induced glutamate dyshomeostasis appear to be connected phenomena affecting each other; for review, see Aschner et al (12).

Notwithstanding the massive efforts in the search for new drugs that counteract mercurial toxicity, there are no effective treatments available that completely abolish its toxic effects. In MeHg poisoning, supportive care is given when necessary to maintain vital functions. Moreover, the use of chelating agents assists the body's ability to eliminate mercury from the tissues. However, these drugs are of limited use because of their adverse side effects (13).

Several studies have focused their efforts on the protective effects of plants and their isolated compounds on diverse neuropathological conditions (14, 15). In this regard, plants of the genus *Polygala* have been shown to possess protective effects against neuronal death and cognitive impairments in neurodegenerative disorders related to excitotoxicity (16, 17).

A previous study from our group (10) demonstrated that the hydroalcoholic extract of *Polygala paniculata* protected against MeHg-induced neurotoxicity in mice. Although data examined in that study have showed that both behavioral and neurochemical changes induced by MeHg were prevented by *Polygala paniculata* extract co-administration, data about the potential compound(s) responsible for the observed neuroprotective effects as well as its/their potential molecular mechanisms were lacking.

* Corresponding author. Telephone: 55 48 3721 9589. Fax: 55 48 3721 9672. E-mail: farina@ccb.ufsc.br.

[†] Departamento de Bioquímica, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina.

[‡] Departamento de Ciências Fisiológicas, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina.

[§] Departamento de Química, Centro de Ciências Físicas e Matemáticas, Universidade Federal de Santa Catarina.




Figure 1. Chemical structures of *Polygala*-derived compounds.

As already mentioned, oxidative stress is a central phenomenon related to MeHg-induced neurotoxicity. In this regard, mitochondria appear to be important cellular organelles targeted by the pro-oxidative effects of MeHg (18). Apoptosis under mitochondrial control has been shown to have an important role in the neuronal death process (19, 20). It usually involves the reduction of mitochondrial transmembrane potential and the release of cytochrome C into the cytoplasm, followed by caspase activation (19, 21). Therefore, disturbances of mitochondrial integrity may initiate cellular death processes. Mercury is known to accumulate inside mitochondria, and it can thus change mitochondrial membrane permeability and cause reduction of mitochondrial transmembrane potential (22–24). Consequently, several studies have pointed to apoptosis as a critical phenomenon involved with MeHg-induced neurotoxicity (25–27).

Taking into account the protective effects of *Polygala paniculata* against MeHg-induced neurotoxicity in mice (10) and the involvement of mitochondrial dysfunction during MeHg-induced neurotoxicity, the aim of this study was to investigate the potential protective effects of three compounds (7-prenyloxy-6-methoxycoumarin, quercetin, and 1,5-dihydroxy-2,3-dimethoxy xanthone) isolated from two plants of genus *Polygala* (*Polygala paniculata* and *Polygala sabulosa*) against mercurial-induced mitochondrial dysfunction under in vitro conditions, using mitochondrial-enriched fractions from mouse brain. Mercuric chloride was also investigated as mitochondrial challenge because of the degradation of MeHg into inorganic mercury in mammalian tissues (28).

Materials and Methods

Materials. Adult Swiss Albino male mice (2 months old) were bred in the animal facilities of the Universidade Federal de Santa Catarina. The mice were maintained according to the Animal Care Guidelines from the National Institutes of Health of the United States of America, and all experiments were approved by our ethic committee for animal use (313/CEUA; 23080.026023/2004-39/UFSC). The animals were maintained at 23°C on a 12 h light/dark cycle with free access to water and food (Nuvital, PR, Brazil). Methylmercury (II) chloride, mercuric chloride, 5,5'-dithiobis-(2-nitrobenzoic acid), and 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) were purchased from Sigma-Aldrich (St Louis, MO). All other chemicals were analytical grade.

Plant Material. *Polygala paniculata* L. (Polygalaceae) was collected in Florianópolis (SC, Brazil) and identified by comparison with the voucher UPCNB 26027 by Prof. Dr. Olavo de Araújo Guimarães of the Botany Department of the Universidade Federal do Paraná, Brazil. *Polygala sabulosa* A.W. Bennett was collected in Rancho Queimado (SC, Brazil) and was also identified by Dr. Guimarães by comparison with the voucher UPCNB 19640.

Extraction and Isolation. 1. *Polygala paniculata*. Air-dried whole plant (3.500 g) was degreased with hexane and extracted with ethanol:water (4:1, v/v). The hydroalcoholic extract was subjected to chromatography fractionation on silica gel column using mixtures of hexane:ethyl acetate:methanol of increasing polarity to give 75 fractions. Fractions 29–35 were combined on

the basis TLC analysis in hexane:ethyl acetate (3:4, v/v) and purified by flash chromatography to give the 1,5-dihydroxy-2,3-dimethoxy xanthone.

2. *Polygala sabulosa*. The dried and powdered whole plant (500 g) was extracted exhaustively with ethanol (96%) at room temperature. The crude extract (135 g) obtained was partitioned into hexane, CH₂Cl₂, ethyl acetate (EtOAc), and water to give four distinct hexane-soluble (16.1 g), CH₂Cl₂-soluble (28 g), AcOEt-soluble (16.8 g), and aqueous (64.6 g) fractions. The CH₂Cl₂-soluble fraction (16 g) was chromatographed on a silica gel column eluted with increasing amounts of ethyl acetate in *n*-hexane to give 36 fractions. The combined fractions 21–26 were dissolved in EtOAc (50 mL) and treated under heating and shaking with activated charcoal (1 g). After filtration, drying (anhydrous NaSO₄), and removal of the solvent, the white residue was purified by recrystallization from hexane:EtOAc (3:1) to obtain 530 mg of the 7-prenyloxy-6-methoxycoumarin (29). The EtOAc-soluble fraction (10 g) was further subjected to column chromatography over silica gel eluted with a hexane:EtOAc gradient system. The polar fractions were further subjected to Sephadex LH-20 eluted with MeOH to yield quercetin (87 mg). The compounds isolated were identified by comparison of their physical and spectral (IR, ¹H, and ¹³C NMR) data with authentic samples, and their chemical structures are depicted in Figure 1.

Preparation of Mitochondrial Enriched Fractions. Mitochondrial-enriched fractions were prepared essentially as described previously (30). Briefly, adult (2 months old) male Swiss mice were sacrificed by decapitation. The whole brains (less cerebellum) were removed and homogenized on ice in 10 volumes of isolation medium (10 mM HEPES buffer pH 7.0 containing 220 mM mannitol, 68 mM sucrose, 10 mM KCl, and 0.1% serum albumin), and the homogenate was centrifuged at 4°C for 10 min at 1000 g. The supernatant was then centrifuged at 17 500 g for 10 min at 4°C, resulting in a myelin-rich supernatant and a pellet (P2) consisting of synaptosomes and free mitochondria. The supernatant was discarded, and the pellet was resuspended in the isolation medium but without albumin. The mitochondrial-enriched fractions were kept on ice for 10–15 min until the experiments were performed.

Incubations. P2 (approximately 2 mg of protein) was incubated with different concentrations of mercurials (0, 10, 30, and 100 μM) and/or the *Polygala*-derived compounds (7-prenyloxy-6-methoxycoumarin, quercetin, and 1,5-dihydroxy-2,3-dimethoxy xanthone at 0, 100, and 300 μM) in a medium containing 10 mM HEPES buffer (pH 7.0), 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). Incubations were carried out for 30 min at 25 °C. *Polygala*-derived compounds were dissolved in dimethyl sulfoxide, of which the final concentration did not exceed 0.5% and did not display any effect itself (data not shown). After incubations, mitochondrial function or glutathione content were determined. Parallel experiments were also carried out in the presence of catalase (200 units) or hydrogen peroxide (100 μM) in order to elucidate molecular mechanisms of toxicity and protection.

Assessment of Mitochondrial Function. Mitochondrial function was assessed by the conversion of the dye methylthiazolyl-diphenyl-tetrazolium bromide (MTT) to formazan (18). This assay is based on the ability of the mitochondrial enzyme succinate dehydrogenase to metabolize MTT into formazan, a reaction that takes place only in functionally intact mitochondria. Briefly, after the preincubation of P2 with mercurials and/or *Polygala*-derived compounds (30 min

at 25 °C), the reaction medium (300 μ L) was incubated with 300 μ L of 1.2 mM MTT for 30 min at 25 °C. The purple formazan crystals were pelleted by centrifugation, and the supernatant was discarded. The pellets were dissolved in DMSO and the formazan was quantified spectrophotometrically at 550 nm. Data were expressed as percentage of control.

Assessment of Glutathione Content. Glutathione content was measured as nonprotein thiols according to a method previously described (31), with minor modifications. Briefly, after the preincubation of P2 with mercurials and/or *Polygala*-derived compounds (30 min at 25 °C), 300 μ L of trichloroacetic acid 10% was added to the reaction medium (300 μ L). After centrifugation (4000 g at 4 °C for 10 min), the protein pellet was discarded and free thiol groups were determined in the clear supernatant (which was neutralized with 0.1 M NaOH) by the method of Ellman (31).

Assessment of Hydrogen Peroxide Content and Lipid Peroxidation. The levels of hydrogen peroxide produced by mitochondria were evaluated through the xylenol orange assay (32), with minor modifications. In short, after the preincubation of P2 with mercurials and/or quercetin/catalase (30 min at 25 °C), the reaction medium was centrifuged at 17 500 g for 10 min at 4 °C and the supernatant was incubated for 30 min in a reaction medium containing 250 mM perchloric acid, 2.5 mM ammonium iron (II) sulfate hexahydrate, and 1 mM xylenol orange. Hydroperoxide levels were determined at 560 nm using a hydrogen peroxide curve as standard. Lipid peroxidation levels were measured as thiobarbituric acid reactive substances (TBARS) according to the method described by Ohkawa and collaborators (33). Briefly, samples were incubated in a reaction media containing 0.28 % 2-thiobarbituric acid, 1.2 % SDS, and 0.45 M/0.12 M acetic acid/HCl buffer (pH 3.4). After incubation at 95 °C for 60 min, TBARS were measured at 532 nm and compared to a standard curve of malondialdehyde.

In vitro Experiments To Detect Possible Chelating Effects of Quercetin. Taking into account the protective role of quercetin on mercurial-induced mitochondrial dysfunction and glutathione oxidation, the potential chelating effects of quercetin towards mercurials were analyzed. Such methodological approach was based on the indirect determination of free mercurial using reduced glutathione (GSH), where free mercurial is available to oxidize GSH. Briefly, different concentrations of MeHg or $HgCl_2$ (0, 10, 25, 50, and 100 μ M) were incubated with GSH (100 μ M) in the presence or in the absence of quercetin (100 μ M) at 25 °C (volume total of reaction = 1 mL). After incubation for 30 min, the amount of GSH was determined by reaction with 55'-dithiobis-(2-nitrobenzoic acid) (31).

Assessment of Protein Content. Protein concentration was determined according to Bradford (34), using a bovine serum albumin as standard.

Statistical Analysis. Statistical differences among groups were analyzed by one-way or two-way analysis of variance, followed by Duncan's multiple range test when appropriate. Pearson analysis was used to correlate variables. Differences were considered statistically significant when $P < 0.05$.

Results

Figure 2 shows that both MeHg (A) and $HgCl_2$ (B) caused a significant and dose-dependent decrease in mitochondrial viability in mitochondrial-enriched fractions of mouse brain. In addition, both mercurials induced a significant oxidation of endogenous glutathione in a dose-dependent manner (Figure 3). Correlation analyses showed significant positive correlations between mitochondrial viability and glutathione content for MeHg (Pearson coefficient = 0.933; $P < 0.01$) and or $HgCl_2$ (Pearson coefficient = 0.854; $P < 0.01$). It is important to state that even the lowest mercurial concentration (10 μ M) was able to significantly decrease mitochondrial viability and glutathione levels when either organic or inorganic mercury was used. This concentration is closely related to that (around 2 μ g/g) found




Figure 2. Effects of MeHg or $HgCl_2$ on mitochondrial viability. Mitochondrial-enriched fractions from mouse brain were incubated with different concentrations (0, 10, 30, and 100 μ M) of (A) MeHg or (B) $HgCl_2$ in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μ L). After preincubation (30 min at 25 °C), mitochondrial viability was assessed by the MTT reduction assay (see Materials and Methods). Data are expressed as mean \pm standard deviation and represented as percent control for four independent assays. *Statistically different from control ($P < 0.05$).

in the cortex of humans exposed to mercury because of the consumption of MeHg-contaminated food (35).

The potential protective effects of *Polygala*-derived compounds against mercurial-induced mitochondrial dysfunction are depicted in Figures 4 and 5. Figures 4A and 5A show that quercetin, at 100 and 300 μ M, completely prevented MeHg- and $HgCl_2$ -induced mitochondrial dysfunction, respectively. However, 7-prenyloxy-6-methoxycoumarin and 1,5-dihydroxy-2,3-dimethoxy xanthone did not display protective effects (Figures 4B, 4C, 5B, and 5C). Indeed, 7-prenyloxy-6-methoxycoumarin and 1,5-dihydroxy-2,3-dimethoxy xanthone alone decreased mitochondrial viability.

Taking into account the positive correlation between mitochondrial viability and glutathione content after mercurial exposure and the protective effects of quercetin against mercurial-induced mitochondrial dysfunction, we evaluated glutathione levels after simultaneous exposure to both MeHg/ $HgCl_2$ and quercetin. The flavonoid completely prevented MeHg- and $HgCl_2$ -induced decrease in reduced glutathione levels (Figure 6). In an attempt to discard the possible direct chemical interaction between quercetin and mercurials, we performed an in vitro assay in order to elucidate if this flavonoid is able to decrease the amount of "free" mercurial in the reaction medium. As expected, Table 1 showed that quercetin was unable to bind MeHg or $HgCl_2$. In fact, the presence of quercetin in the reaction medium did not change the oxidative capability of both MeHg and $HgCl_2$ to reduced glutathione.

Taking the aforementioned results into consideration, we hypothesized that quercetin could be capable of detoxifying reactive/oxidant molecules generated in the presence of mer-




Figure 3. Effects of the MeHg or HgCl_2 on glutathione content. Mitochondrial-enriched fractions from mouse brain were preincubated with different concentrations of (A) MeHg or (B) HgCl_2 (0, 10, 30, and 100 μM) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). After preincubation (30 min at 25 $^\circ\text{C}$), glutathione content was quantified using 5,5'-dithiobis-(2-nitrobenzoic acid) (see Materials and Methods). Data are expressed as mean \pm standard deviation and represented as percent control for four independent assays. *Statistically different from control ($P < 0.05$). Glutathione content in basal condition was 1.0 ± 0.04 nmol/mg of protein.

curials in the brain synaptosomal medium. This idea was based on the study of Lund and collaborators (36), which points to increased hydrogen peroxide production in rat kidney mitochondria during mercuric chloride exposure. Figure 7 shows that catalase, an enzyme involved in the detoxification of hydrogen peroxide, completely prevented MeHg-induced mitochondrial dysfunction. In addition, catalase partially prevented HgCl_2 -induced mitochondrial dysfunction.

Because the previous data (Figure 7) have pointed to hydrogen peroxide as an important molecule involved with mercurial-induced mitochondrial dysfunction, the levels of hydrogen peroxide generated during MeHg or HgCl_2 exposure were evaluated. Figure 8 shows that both MeHg (Figure 8A) and HgCl_2 (Figure 8B) enhanced the levels of hydrogen peroxide. Interestingly, such phenomena were abolished by the addition of either catalase (200 units) or quercetin (100 μM) in the reaction medium. Moreover, both MeHg (Figure 9A) and HgCl_2 (Figure 9B) enhanced the levels of thiobarbituric acid reactive substances (a marker for lipid peroxidation), and these phenomena were also abolished by the addition of catalase (200 units) or quercetin (100 μM) in the reaction medium.

Discussion

A previous study from our group (10) reported significant beneficial effects of the hydroalcoholic extract of *Polygala paniculata* against MeHg-induced neurotoxicity in mice. In that study, the hydroalcoholic extract of *Polygala paniculata* was able to prevent behavioral and neurochemical changes induced by the oral exposure to MeHg, pointing to such a plant as a




Figure 4. Effects of the *Polygala*-derived compounds on MeHg-induced mitochondrial dysfunction. Mitochondrial-enriched fractions from mouse brain were preincubated with MeHg (30 μM) and/or (A) quercetin, (B) 7-prenyloxy-6-methoxycoumarin, or (C) 1,5-dihydroxy-2,3-dimethoxy xanthone (0, 100, and 300 μM) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). After pre- (30 min at 25 $^\circ\text{C}$), mitochondrial viability was assessed by the MTT reduction assay (see Materials and Methods). Data are expressed as mean \pm standard deviation and represented as percent control for four independent assays. *Statistically different from control ($P < 0.05$).

potential therapeutic agent for the treatment of pathological conditions related to excitotoxicity and oxidative stress, including MeHg poisoning. In agreement, plants of the genus *Polygala* have been reported to display neuroprotective effects in several neuropathological conditions related to excitotoxicity and oxidative stress (16, 17).

In the present study, we sought potential compounds derived from *Polygala* that might be responsible for protection against mercurial-induced neurotoxicity, using mitochondrial-enriched fractions from mouse brain. Three molecules, from different chemical groups, a flavonoid (quercetin), a coumarin (7-prenyloxy-6-methoxycoumarin), and a xanthone (1,5-dihydroxy-2,3-dimethoxy xanthone), were selected for this proposal. Taking into account the possible degradation of MeHg into inorganic mercury in mammalian tissues (28), we investigated both




Figure 5. Effects of the *Polygala*-derived compounds on $HgCl_2$ -induced mitochondrial dysfunction. Mitochondrial-enriched fractions from mouse brain were preincubated with $HgCl_2$ (0 or 30 μM) and/or (A) quercetin, (B) 7-prenyloxy-6-methoxycoumarin, or (C) 1,5-dihydroxy-2,3-dimethoxy xanthone (0, 100, and 300 μM) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). After preincubation (30 min at 25 $^{\circ}C$), mitochondrial viability was assessed by the MTT reduction assay (see Materials and Methods). Data are expressed as mean \pm standard deviation and represented as percent control for four independent assays. *Statistically different from control ($P < 0.05$).

organic (MeHg) and inorganic ($HgCl_2$) mercurials as potential mitochondrial challenges.

It was noteworthy that both mercurials (MeHg and $HgCl_2$) caused a similar and dose-dependent decrease in mitochondrial viability, which was positively correlated to glutathione oxidation. From a molecular point of view, these results represent important findings. In fact, considering the differential number of potential electrophilic regions of mercury atom in the organic-methylated form (one positive charge) and in the mercuric mercury form (two positive charges), when dissociated in aqueous medium, the similar profiles of mitochondrial glutathione oxidation and dysfunction observed after MeHg and $HgCl_2$ exposure were unexpected. Such results indicate that MeHg- and $HgCl_2$ -induced mitochondrial dysfunction depend on phenomena other than the simple direct interaction between




Figure 6. Effects of the quercetin on MeHg- or $HgCl_2$ -induced glutathione oxidation. Mitochondrial-enriched fractions from mouse brain were preincubated with (A) MeHg or (B) $HgCl_2$ (30 μM) and/or quercetin (0, 100 or 300 μM) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). After preincubation (30 min at 25 $^{\circ}C$), glutathione content was quantified using 5,5'-dithiobis-(2-nitrobenzoic acid) (see Materials and Methods). Data are expressed as mean \pm standard deviation and represented as percent control for four independent assays. *Statistically different from control ($P < 0.05$). Glutathione content in basal condition was 1.0 ± 0.028 nmol/mg protein.

Table 1. Effects of Quercetin on the Mercurial-Induced Glutathione (GSH) Oxidation^a

	remaining reduced GSH (nmol)	
	without quercetin	with quercetin
MeHg (nmol/mL)		
0	100 \pm 3.9	96.8 \pm 5.2
10	92.0 \pm 2.5	88.7 \pm 3.4
25	81.2 \pm 1.7	81.7 \pm 2.3
50	65.0 \pm 8.4	60.7 \pm 8.8
100	20.3 \pm 0.6	18.2 \pm 2.7
$HgCl_2$ (nmol/mL)		
0	98.9 \pm 4.0	97.1 \pm 2.2
10	87.7 \pm 4.4	87.3 \pm 6.5
25	62.1 \pm 2.4	57.2 \pm 5.2
50	18.5 \pm 2.0	16.9 \pm 2.1
100	1.3 \pm 0.1	1.5 \pm 0.3

^a MeHg or $HgCl_2$ (0, 10, 25, 50, and 100 μM) were incubated with GSH (100 μM) in the presence or in the absence of quercetin (100 μM) at 25 $^{\circ}C$ (volume total of reaction = 1 mL). After incubation for 30 min, the amount of remaining GSH was determined using 5,5'-dithiobis-(2-nitrobenzoic acid). Data are expressed as mean \pm standard deviation and represented as nmol of reduced glutathione for three independent assays. Two-way analysis of variance (5 mercurial doses vs quercetin) showed no significant interactions of quercetin with MeHg [$F(4,29) = 0.624$, $p = 0.651$] or with $HgCl_2$ [$F(4,29) = 0.695$, $p = 0.604$] toward glutathione oxidation.

mercury and sulfhydryl groups from biomolecules. Taking into account the degradation of MeHg into inorganic mercury in mammalian tissues, one could suppose that the similar potency of MeHg and $HgCl_2$ toward mitochondrial viability and glutathione oxidation could be related, at least in part, to the




Figure 7. Effects of catalase on MeHg- or HgCl₂-induced mitochondrial dysfunction. Mitochondrial-enriched fractions from mouse brain were preincubated with MeHg or HgCl₂ (30 μM) and/or catalase (200 units) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). After preincubation (30 min at 25 °C), mitochondrial viability was assessed by the MTT reduction assay (see Materials and Methods). Data are expressed as mean ± standard deviation and represented as percent control (dotted line) for three independent assays. *Statistically different from the corresponding condition with catalase ($P < 0.05$). [#]Statistically different from control ($P < 0.05$).

instability of MeHg and its conversion to inorganic mercury. Although this hypothesis could not be ruled out, it is important to state that the degradation of MeHg into inorganic mercury depends on the activity of NADPH-cytochrome P-450 reductase, an enzyme present only in microsomal preparations (28).

Herein, the incubation of mitochondrial enriched fractions with MeHg or HgCl₂ enhanced by around 80% the hydrogen peroxide (H₂O₂) production and such effects were reduced by the addition of catalase, strongly pointing to H₂O₂ as a crucial molecule involved in mercurials-induced mitochondrial dysfunction. This result is confirmed by the fact that both mercurials increased the levels of thiobarbituric acid reactive substances (marker of lipid peroxidation) and such phenomena were also prevented by the addition of catalase, an enzyme that specifically detoxifies H₂O₂.

Therefore, the increase in H₂O₂ formation by mouse brain mitochondria exposed to mercurials may represent a significant finding of our study, considering the oxidative role of H₂O₂ toward sulphydryl groups. This observation is in agreement with Lund et al. (36), who effectively demonstrated that HgCl₂ is able to induce renal mitochondrial dysfunction and pointed to H₂O₂ as a key molecule in mercury toxicity to renal tissue. Furthermore, an increase in reactive oxygen species generation in brain preparations after mercurials (organic or inorganic mercury) exposure has been also demonstrated (8, 18). However, those studies were based on the 2',7'-dichlorofluorescein methodology, which can not discriminate between hydroxyl radicals, peroxynitrite, or H₂O₂ (37).

Among the three Polygala-derived compounds, only quercetin displayed protective effects against mercurial-induced mitochondrial dysfunction. In this regard, several studies have shown the protective effects of quercetin in different models of chemical injury under in vitro and in vivo conditions (38–40). Particularly important: quercetin has been reported to display beneficial effects against metal-induced toxicity (41, 42). Moreover, two recent and interesting studies have reported the anti-peroxidative action of quercetin under in vitro conditions (43, 44). However, to the best of our knowledge, there are no data in the literature showing protective effects of quercetin against mercurial-induced neurotoxicity.




Figure 8. Effects of mercurials on hydrogen peroxide production. Mitochondrial-enriched fractions from mouse brain were preincubated with (A) 100 μM MeHg or (B) HgCl₂ and/or catalase (200 units) or quercetin (100 μM) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μL). After preincubation (60 min at 25 °C), hydrogen peroxide formation was assessed by the xylenol orange assay (see Materials and Methods). Data are expressed as mean ± standard deviation and represented as percent of control (dotted line) for four independent assays. Different letters indicate significant difference ($P < 0.05$) by one-way analysis of variance. Hydrogen peroxide level in basal condition was 0.14 ± 0.021 nmol/mg of protein.

Even though quercetin prevented mercurial-induced mitochondrial dysfunction and glutathione oxidation, our results demonstrate that such protective effect is not directly related to chelating effects. Indeed, quercetin did not change the oxidative capability of both MeHg and HgCl₂ regarding reduced glutathione, indicating that the “free” amounts of both mercurials are not changed by the presence of quercetin. Because the mitochondrial dysfunction observed after mercurial exposure was related to H₂O₂ production, we hypothesized that quercetin could be capable of detoxifying the H₂O₂ generated in the presence of mercurials in the brain mitochondria. Our results indeed showed that quercetin decreased mercurial-induced H₂O₂ production and such data are in agreement with the fact that quercetin also abolished mercurial-induced lipid peroxidation in the brain mitochondrial-enriched fraction. In conformity, the protective effect of quercetin against H₂O₂-induced toxicity has been shown under in vitro conditions (45).

Even though quercetin prevented mercurial-induced mitochondrial toxicity under in vitro conditions, it is difficult to affirm that quercetin is directly and exclusively involved with the beneficial effects (10, 16, 17) of plants of the genus *Polygala* observed under in vivo conditions. Several compounds—aurapten, phebalosin, murrangatin, and 7-methoxy-8-(1,4-dihydroxy-3-methyl-2-butenyl)coumarin (46); 1-hydroxy-2,3,5-trimethoxyxanthone, 1-hydroxy-5-methoxy-2,3-methylenedioxyxanthone, 1,5-dihydroxy-2,3-dimethoxyxanthone, coumarin murragatin, rutin, spinasterol, and delta-25-spinasterol (29)—were isolated from *Polygala paniculata*, and the knowledge about their biological effects are scarce. So, it is possible that the beneficial effects displayed by *Polygala paniculata* under in vivo condi-




Figure 9. Effects of mercurials on lipid peroxidation. Mitochondrial-enriched fractions from mouse brain were pre-incubated with 100 μ M (A) MeHg or HgCl₂ (B) and/or catalase (200 units) or quercetin (100 μ M) in 10 mM HEPES buffer (pH 7.0) containing 220 mM mannitol, 68 mM sucrose, and 10 mM KCl (total incubation volume = 300 μ L). After preincubation (60 min at 25 °C), thiobarbituric acid reactive substances (TBARS) levels were assessed (see Materials and Methods). Data are expressed as mean \pm standard deviation and represented as percent control (dotted line) for four independent assays. *Statistically different from control ($P < 0.05$) by one-way analysis of variance. TBARS levels in basal condition were 7.8 ± 0.14 nmol MDA/mg of protein.

tions (10) are not necessarily related to a single compound. This hypothesis is reinforced by the fact that 100 and 300 μ M are high quercetin concentrations that practically could not be reached under in vivo conditions.

In summary, this study shows that thiol oxidation is a crucial phenomenon involved with MeHg- and HgCl₂-induced mitochondrial dysfunction and that the protective effect of quercetin on such process is not directly related to chelating effects. Moreover, our results are the first to show that H_2O_2 is involved with the toxic effects of both organic and inorganic mercurials in brain mitochondria. The study also shows the protective effect of quercetin against mercurial-induced toxicity, pointing to its capability to counteract mercurial-dependent H_2O_2 generation as a potential molecular mechanism of protection. Taken together, these data add new information about molecular mechanisms associated to mercurial toxicity and render quercetin a promising molecule for pharmacological studies with respect to mercurials' poisoning.

Acknowledgment. This study was supported by grants from CNPq to M. Farina (475329/2004-0 and 474796/2006-0) and by the FINEP research grant "Rede Instituto Brasileiro de Neurociência (IBN-Net)" 01.06.0842-00.

References

- (1) Clarkson, T. W., Magos, L., and Myers, G. J. (2003) The toxicology of mercury—current exposures and clinical manifestations. *New Engl. J. Med.* 349, 1731–1737.
- (2) Sirois, J. E., and Atchison, W. D. (2000) Methylmercury affects multiple subtypes of calcium channels in rat cerebellar granule cells. *Toxicol. Appl. Pharmacol.* 167, 1–11.
- (3) Aschner, M., Yao, C. P., Allen, J. W., and Tan, K. H. (2000) Methylmercury alters glutamate transport in astrocytes. *Neurochem. Int.* 37, 199–206.
- (4) Farina, M., Dahm, K. C., Schwalm, F. D., Brusque, A. M., Frizzo, M. E., Zeni, G., Souza, D. O., and Rocha, J. B. 2003a Methylmercury increases glutamate release from brain synaptosomes and glutamate uptake by cortical slices from suckling rat pups: modulatory effect of ebselen. *Toxicol. Sci.* 73, 135–140.
- (5) Manfro, C. B., Schwalm, F. D., Cereser, V., Abreu, F., Oliveira, A., Bizarro, L., Rocha, J. B., Frizzo, M. E., Souza, D. O., and Farina, M. (2004) Maternal milk as methylmercury source for suckling mice: neurotoxic effects involved with the cerebellar glutamatergic system. *Toxicol. Sci.* 81, 172–178.
- (6) Fonfria, E., Vilaro, M. T., Babot, Z., Rodriguez-Farre, E., and Sunol, C. (2005) Mercury compounds disrupt neuronal glutamate transport in cultured mouse cerebellar granule cells. *J. Neurosci. Res.* 79, 545–553.
- (7) Ou, Y. C., White, C. C., Krejsa, C. M., Ponce, R. A., Kavanagh, T. J., and Faustman, E. M. (1999) The role of intracellular glutathione in methylmercury-induced toxicity in embryonic neuronal cells. *Neurotoxicology* 20, 793–804.
- (8) Shanker, G., Syversen, T., Aschner, J. L., and Aschner, M. (2005) Modulatory effect of glutathione status and antioxidants on methylmercury-induced free radical formation in primary cultures of cerebral astrocytes. *Brain Res. Mol. Brain Res.* 137, 11–22.
- (9) Farina, M., Brandão, R., Lara, F. S., Soares, F. A., Souza, D. O., and Rocha, J. B. 2003b Mechanisms of the inhibitory effects of selenium and mercury on the activity of delta-aminolevulinic dehydratase from mouse liver, kidney and brain. *Toxicol. Lett.* 139, 55–66.
- (10) Farina, M., Franco, J. L., Ribas, C. M., Meotti, F. C., Missau, F. C., Pizzolatti, M. G., Dafre, A. L., and Santos, A. R. (2005) Protective effects of Polygala paniculata extract against methylmercury-induced neurotoxicity in mice. *J. Pharm. Pharmacol.* 57, 1503–1508.
- (11) Franco, J. L., Teixeira, A., Meotti, F. C., Ribas, C. M., Stringari, J., Garcia Pomblum, S. C., Moro, A. M., Bohrer, D., Bairos, A. V., Dafre, A. L., Santos, A. R., and Farina, M. (2006) Cerebellar thiol status and motor deficit after lactational exposure to methylmercury. *Environ. Res.* 102, 22–28.
- (12) Aschner, M., Syversen, T., Souza, D. O., Rocha, J. B., and Farina, M. (2007) Involvement of glutamate and reactive oxygen species in methylmercury neurotoxicity. *Braz. J. Med. Biol. Res.* 40, 285–291.
- (13) Tchounwou, P. B., Ayensu, W. K., Ninashvili, N., and Sutton, D. (2003) Environmental exposure to mercury and its toxicopathologic implications for public health. *Environ. Toxicol.* 18, 149–175.
- (14) Scalbert, A., and Williamson, G. (2000) Dietary intake and bioavailability of polyphenols. *J. Nutr.* 130, 2073S–2085S.
- (15) Walle, T. (2004) Absorption and metabolism of flavonoids. *Free Radical Biol. Med.* 36, 829–837.
- (16) Lee, H. J., Ban, J. Y., Koh, S. B., Seong, N. S., Song, K. S., Bae, K. W., and Seong, Y. H. (2004) Polygalae radix extract protects cultured rat granule cells against damage induced by NMDA. *Am. J. Chin. Med.* 32, 599–610.
- (17) Park, J. H., Kim, J. S., Jang, D. S., and Lee, S. M. (2006) Effect of Polygala tenuifolia root extract on cerebral ischemia and reperfusion. *Am. J. Chin. Med.* 34, 115–123.
- (18) Dreiem, A., Gertz, C. C., and Seegal, R. F. (2005) The effects of methylmercury on mitochondrial function and reactive oxygen species formation in rat striatal synaptosomes are age-dependent. *Toxicol. Sci.* 87, 156–162.
- (19) Ghribi, O., DeWitt, D. A., Forbes, M. S., Herman, M. M., and Savory, J. (2001) Co-involvement of mitochondria and endoplasmic reticulum in regulation of apoptosis: changes in cytochrome c, Bcl-2 and Bax in the hippocampus of aluminum-treated rabbits. *Brain Res.* 903, 66–73.
- (20) Fiskum, G., Starkov, A., Polster, B. M., and Chinopoulos, C. (2003) Mitochondrial mechanisms of neural cell death and neuroprotective interventions in Parkinson's disease. *Ann. N.Y. Acad. Sci.* 991, 111–119.
- (21) Kluck, R. M., Bossy-Wetzel, E., Green, D. R., and Newmeyer, D. D. (1997) The release of cytochrome c from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science* 275, 1132–1136.
- (22) InSug, O., Datar, S., Koch, C. J., Shapiro, I. M., and Shenker, B. J. (1997) Mercuric compounds inhibit human monocyte function by inducing apoptosis: evidence for formation of reactive oxygen species, development of mitochondrial membrane permeability transition and loss of reductive reserve. *Toxicology* 124, 211–224.
- (23) Shenker, B. J., Guo, T. L., O, I., and Shapiro, I. M. (1999) Induction of apoptosis in human T-cells by methyl mercury: temporal relationship between mitochondrial dysfunction and loss of reductive reserve. *Toxicol. Appl. Pharmacol.* 157, 23–35.

- (24) Araragi, S., Kondoh, M., Kawase, M., Saito, S., Higashimoto, M., and Sato, M. (2003) Mercuric chloride induces apoptosis via a mitochondrial-dependent pathway in human leukemia cells. *Toxicology* 184, 1–9.
- (25) Nishioku, T., Takai, N., Miyamoto, K., Murao, K., Hara, C., Yamamoto, K., and Nakanishi, H. (2000) Involvement of caspase 3-like protease in methylmercury-induced apoptosis of primary cultured rat cerebral microglia. *Brain Res.* 871, 160–164.
- (26) Dare, E., Gotz, M. E., Zhivotovsky, B., Manzo, L., and Ceccatelli, S. (2000) Antioxidants J811 and 17 β -estradiol protect cerebellar granule cells from methylmercury-induced apoptotic cell death. *J. Neurosci. Res.* 62, 557–565.
- (27) Dare, E., Li, W., Zhivotovsky, B., Yuan, X., and Ceccatelli, S. (2001) Methylmercury and H₂O₂ provoke lysosomal damage in human astrocytoma D384 cells followed by apoptosis. *Free Rad. Biol. Med.* 30, 1347–1356.
- (28) Suda, I., and Hirayama, K. (1992) Degradation of methyl and ethyl mercury into inorganic mercury by hydroxyl radical produced from rat liver microsomes. *Arch. Toxicol.* 66, 398–402.
- (29) Cristiano, R., Pizzolatti, M. G., Delle Monache, F., Rezende, C. M., and Branco, A. (2003) Two xanthenes from *Polygala paniculata* and confirmation of the 1-hydroxy-2,3,5-trimethoxy-xanthone at trace level by HRGC-MS. *Z. Naturforsch., C: J. Biosci.* 58, 490–494.
- (30) Andersen, J. M., Myhre, O., and Fonnum, F. (2003) Discussion of the role of the extracellular signal-regulated kinase-phospholipase A₂ pathway in production of reactive oxygen species in Alzheimer's disease. *Neurochem. Res.* 28, 319–326.
- (31) Ellman, G. L. (1959) Tissue sulfhydryl groups. *Arch. Biochem. Biophys.* 82, 70–77.
- (32) Nourooz-Zadeh, J., Tajaddini-Sarmadi, J., and Wolff, S. P. (1994) Measurement of plasma hydroperoxide concentrations by the ferrous oxidation-xylenol orange assay in conjunction with triphenylphosphine. *Anal. Biochem.* 220, 403–409.
- (33) Ohkawa, H., Ohishi, N., and Yagi, K. (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 95, 351–358.
- (34) Bradford, M. M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- (35) Davis, L. E., Kornfeld, M., Mooney, H. S., Fielder, K. J., Haaland, K. Y., Orrison, W. W., Cernichiari, E., and Clarkson, T. W. (1994) Methylmercury poisoning: long-term clinical, radiological, toxicological, and pathological studies of an affected family. *Ann. Neurol.* 35, 680–688.
- (36) Lund, B. O., Miller, D. M., and Woods, J. S. (1993) Studies on Hg(II)-induced H₂O₂ formation and oxidative stress in vivo and in vitro in rat kidney mitochondria. *Biochem. Pharmacol.* 45, 2017–2024.
- (37) Myhre, O., Andersen, J. M., Aarnes, H., and Fonnum, F. (2003) Evaluation of the probes 2',7'-dichlorofluorescein diacetate, luminol, and lucigenin as indicators of reactive species formation. *Biochem. Pharmacol.* 65, 1575–1582.
- (38) Naidu, P. S., Singh, A., and Kulkarni, S. K. (2003) Reversal of haloperidol-induced orofacial dyskinesia by quercetin, a bioflavonoid. *Psychopharmacology* 167, 418–423.
- (39) Heo, H. J., and Lee, C. Y. (2004) Protective effects of quercetin and vitamin C against oxidative stress-induced neurodegeneration. *J. Agric. Food Chem.* 52, 7514–7517.
- (40) Lu, J., Zheng, Y. L., Luo, L., Wu, D. M., Sun, D. X., and Feng, Y. J. (2006) Quercetin reverses D-galactose induced neurotoxicity in mouse brain. *Behav. Brain Res.* 171, 251–260.
- (41) Meng, F., Liu, R., Bai, H., Liu, B. W., and Liu, Y. (2004) Inhibitory effect of quercetin, rutin and puerarin on HDL oxidation induced by Cu²⁺. *Sichuan Daxue Xuebao Yi Xue Ban* 35, 836–838.
- (42) Morales, A. I., Vicente-Sanchez, C., Sandoval, J. M., Egido, J., Mayoral, P., Arevalo, M. A., Fernandez-Tagarro, M., Lopez-Novoa, J. M., and Perez-Barriocanal, F. (2006) Protective effect of quercetin on experimental chronic cadmium nephrotoxicity in rats is based on its antioxidant properties. *Food Chem. Toxicol.* 44, 2092–2100.
- (43) Dufour, C., and Loonis, M. (2007) Flavonoids and their oxidation products protect efficiently albumin-bound linoleic acid in a model of plasma oxidation. *Biochim. Biophys. Acta* 1770, 958–965.
- (44) Dufour, C., Loonis, M., and Dangles, O. (2007) Inhibition of the peroxidation of linoleic acid by the flavonoid quercetin within their complex with human serum albumin. *Free Radical Biol. Med.* 43, 241–252.
- (45) Chow, J. M., Shen, S. C., Huan, S. K., Lin, H. Y., and Chen, Y. C. (2005) Quercetin, but not rutin and quercitrin, prevention of H₂O₂-induced apoptosis via anti-oxidant activity and heme oxygenase 1 gene expression in macrophages. *Biochem. Pharmacol.* 69, 1839–1851.
- (46) Hamburger, M., Gupta, M., and Hostettmann, K. (1985) Coumarins from *Polygala paniculata*. *Planta Med.* 51, 215–217.

TX7002323