

A Cytotoxic Sesquiterpene Alkaloid from the South China Sea Gorgonian *Subergorgia suberosa*


Shu-Hua Qi,^{*,†} Si Zhang,^{†,‡} Xiang Li,[†] and Qing-Xin Li[†]

Guangdong Key Laboratory of Marine Materia Medica, South China Sea Institute of Oceanology, The Chinese Academy of Sciences, 164 West Xingang Road, Guangzhou 510301 Guangdong, People's Republic of China, and Hainan Key Lab of Tropical Marine Biology and Technology, South China Sea Institute of Oceanology, The Chinese Academy of Sciences, People's Republic of China

Received March 29, 2005

A new sesquiterpene alkaloid, 6-(9'-purine-6',8'-diolyl)-2 β -suberosanone (**1**), together with three known sesquiterpenes, suberosenol A (**2**), subergorgic acid (**3**), and subergorgiol (**4**), was isolated from the EtOH/CH₂Cl₂ extracts of the South China Sea gorgonian *Subergorgia suberosa*. The structure of **1** was determined through spectroscopic methods. Compound **1** showed moderate cytotoxicity against the human breast carcinoma MDA-MB-231 cell line with an IC₅₀ of 8.87 μ g/mL.

Previous studies on the chemical constituents of *Subergorgia suberosa* have led to the isolation of several sesquiterpenes^{1–5} and several 9,11-seco steroids.^{6–8} Some of these sesquiterpenes showed cytotoxicity toward several cancer cell lines.^{3–5} During the course of further searching for novel active compounds from gorgonians,^{9,10} we undertook the investigation of the South China Sea gorgonian *S. suberosa*. A new suberosane-type sesquiterpene alkaloid, 6-(9'-purine-6',8'-diolyl)-2 β -suberosanone (**1**), together with three known sesquiterpenes, suberosenol A (**2**),¹¹ subergorgic acid (**3**),¹ and subergorgiol (**4**),⁴ was isolated from the EtOH/CH₂Cl₂ extracts of *S. suberosa*. In the cytotoxicity assays, we observed that **1** showed moderate cytotoxicity against the human breast carcinoma MDA-MB-231 cell line with an IC₅₀ of 8.87 μ g/mL and potential cytotoxicity toward the MCF cell line at a concentration of 50 μ M. This paper deals with the isolation, structural elucidation, and cytotoxic activity of **1**.



Compound **1** had the molecular formula C₂₀H₂₆N₄O₃ as deduced from NMR spectra and HRESIMS. Thus, 10 degrees of unsaturation was determined for the molecule of **1**. Its UV spectrum exhibited maximum absorption at 212 and 264 nm (aromatic group), while the IR spectrum showed absorption bands for hydroxyls (3500, 3115 cm⁻¹), carbonyl groups (1740, 1710 cm⁻¹), and an aromatic ring




Figure 1. Key HMBC correlations of **1**.

(1670, 1600 cm⁻¹). The ¹³C NMR spectrum showed the presence of 15 basic skeleton carbons, including three methyls (δ_C 16.7, 27.0, 34.4), five methylenes (δ_C 40.8, 41.3, 27.0, 27.9, 48.5), four methines (δ_C 44.0, 52.4, 36.6, 49.7), two quaternary carbons (δ_C 56.7, 39.7), and a ketone carbon (δ_C 216.5, s), along with five low-field carbons [δ_C 108.1 (s), 140.6 (d), 150.2 (s), 151.9 (s), 155.8 (s)]. The ¹H NMR spectrum displayed three methyl groups at δ_H 0.80 (3H, d, J = 7.0 Hz), 1.14 (3H, s), 1.16 (3H, s) and an olefin proton at δ_H 7.91 (1H, s). These NMR spectral data showed similarity with those of suberosanone¹¹ and suberosenone⁵ with the exception of five additional low-field carbons. On the basis of the above data, **1** should be a suberosanone-type sesquiterpene linked with a five-carbon aromatic group.

The five low-field carbon signals [δ_C 108.1 (s), 140.6 (d), 150.2 (s), 151.9 (s), 155.8 (s)] with only one corresponding proton (δ_H 7.91, 1H, s) of **1** were similar to those of 3,7,9-tri-Me-6,8-purinediol that had been found in the South China Sea gorgonian *Echinogorgia pseudosapio*¹² and other analogues.¹³ When the measuring solvent was changed from CDCl₃ to pyridine-*d*₅, two additional signals [δ_H 12.9, 13.7 (each 1H, s, OH)] appeared in the ¹H NMR spectrum. According to the above NMR spectral data, the correlations of δ_H 7.91 (1H, s) with δ_C 150.2 (s), 108.1 (s), and 155.8 (s) in the HMBC spectrum (Figure 1) and the molecular formula of C₂₀H₂₆N₄O₃, the five-carbon aromatic group should be 6',8'-purinediol. In the HMBC spectrum, correlations of δ_H 4.46 (1H, dd, J = 4.6, 14.1 Hz, H-6a), 4.25 (1H, dd, J = 9.0, 14.1 Hz, H-6b) with δ_C 150.2 (s, C-4'), 151.9 (s, C-8') suggested the link of the 6',8'-purinediol moiety with the suberosanone moiety by a C(6)–N(9') bond. The relative stereochemistry of **1** was deduced from a 2D NOE experiment. In the NOESY spectrum of **1** (Figure 2), correlations of H-2 (δ_H 2.40, 1H, overlap) with Me-15 (δ_H

* Corresponding author. Tel: +86-20-89023105. Fax: +86-20-84458964. E-mail: shuhuqi2001@yahoo.com.

[†] Guangdong Key Laboratory of Marine Materia Medica.

[‡] Hainan Key Lab of Tropical Marine Biology and Technology.




Figure 2. Selective NOE correlations of **1**.

1.16, 3H, s) and H-12a (δ_{H} 1.65, 1H, d, $J = 14.7$ Hz) indicated that H-2 and Me-15 had a β -orientation, because the C-12 methylene was the β -substituent at C-1. NOE correlations of H-9a (δ_{H} 1.19, 1H, m) with Me-7 (δ_{H} 0.80, 3H, d, $J = 7.0$ Hz) and H-11 (δ_{H} 1.77, 1H, m), and H-5 with Me-7, and no NOE correlation of H-11 with H-12 [δ_{H} 1.69, 1.65 (each 1H, d, $J = 14.7$ Hz)], suggested the α -orientation of H-5, H-11, and Me-7. On the basis of the above data, the structure of **1** was elucidated as shown. The H-2 β and positive rotation $\{[\alpha]_{\text{D}}^{20} + 28^\circ$ (c 0.2, CHCl_3) of **1** were different from the H-2 α and negative rotation of suberosanone $\{[\alpha]_{\text{D}}^{25} - 60^\circ$ (c 0.1, CHCl_3) and other suberosanone-type sesquiterpenes.¹¹

The cytotoxicity of compound **1** toward the MDA-MB-231 and MCF cancer cell lines was evaluated quantitatively and qualitatively, respectively. It was found that compound **1** showed moderate cytotoxicity against the human breast carcinoma MDA-MB-231 cell line with an IC_{50} of $8.87 \mu\text{g/mL}$ and potential cytotoxicity toward the MCF cell line at a concentration of $50 \mu\text{M}$.

Experimental Section

General Experimental Procedures. Optical rotations were measured with a Horiba SEAP-300 spectropolarimeter. UV spectra were measured with a Shimadzu double-beam 210A spectrophotometer in MeOH solution. IR (KBr) spectra were obtained on a Bio-Rad FTS-135 infrared spectrophotometer. ^1H , ^{13}C NMR and 2D NMR spectra were recorded on a Bruker AV-500 MHz NMR spectrometer with TMS as internal standard. MS spectral data were obtained on an LCQ^{DECA} XP HPLC/MSⁿ spectrometer for ESIMS. Si gel (200–300 mesh) for column chromatography and GF₂₅₄ for TLC were obtained from the Qindao Marine Chemical Factory, Qindao, People's Republic of China.

Animal Material. The South China Sea gorgonian coral *S. suberosa* (3.5 kg, wet weight) was collected in Sanya, Hainan Province, China, in October 2003 and identified by Prof. R. L. Zou, the South China Sea Institute of Oceanology, Academia Sinica. A voucher specimen (No. 0312) was deposited in the South China Sea Institute of Oceanology, Academia Sinica, Guangzhou, China.

Extraction and Isolation. The frozen specimen was extracted with EtOH/ CH_2Cl_2 (2:1) three times at room temperature, and the solution was evaporated in vacuo. The residue was suspended in H_2O and extracted with CHCl_3 three

times. The CHCl_3 layer was concentrated in vacuo to afford 40 g of residue. The CHCl_3 extract was subjected to column chromatography (CC) on silica, using $\text{CHCl}_3/\text{Me}_2\text{CO}$ (from 10:0 to 0:10) as eluent. By combining the fractions with TLC (GF₂₅₄) monitoring, eight fractions were obtained. Fraction 2 was subjected to CC on silica gel, eluted with petroleum ether/EtOAc (from 10:0 to 10:1), to afford **2** (6 mg). Fraction 4 was subjected to CC on silica gel, eluted with petroleum ether/EtOAc (from 10:1 to 8:2), to yield **3** (48 mg) and **4** (5 mg). Fraction 6 was chromatographed over Sephadex LH-20 eluting with $\text{CHCl}_3/\text{MeOH}$ (1:1), then subjected to CC on silica gel, eluted with $\text{CHCl}_3/\text{MeOH}$ (from 12:1 to 10:2), to yield **1** (8 mg).

6-(9'-Purine-6',8'-diolyl)-2 β -suberosanone (1**):** white powder; $[\alpha]_{\text{D}}^{20} + 28^\circ$ (c 0.2, CHCl_3); UV (MeOH) λ_{max} 212, 264 nm; IR (KBr) 3500, 3115, 1740, 1710, 1670, 1600 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 2.40 (1H, overlap, H-2 β), 2.47 (1H, m, H-3 β), 2.41 (1H, overlap, H-3 α), 3.32 (1H, dd, $J = 4.4$, 8.3 Hz, H-5 α), 4.46 (1H, dd, $J = 4.4$ Hz, 14.1, H-6a), 4.25 (1H, dd, $J = 8.3$, 14.1 Hz, H-6b), 0.80 (3H, d, $J = 7.0$ Hz, Me-7), 1.54 (1H, m, H-8 α), 1.87 (1H, m, H-9 β), 1.16–1.19 (1H, overlap, H-9 α), 1.61 (1H, m, H-10 β), 1.53 (1H, m, H-10 α), 1.77 (1H, m, H-11 α), 1.69 (1H, d, $J = 14.7$ Hz, H-12 β), 1.65 (1H, d, $J = 14.7$ Hz, H-12 α), 1.14 (3H, s, Me-14), 1.16 (3H, s, Me-15), 7.91 (1H, s, H-2'), ^{13}C NMR (125 MHz, CDCl_3) δ 56.7 (C, C-1), 44.0 (CH, C-2), 40.8 (CH₂, C-3), 216.5 (C, C-4), 52.4 (CH, C-5), 41.3 (CH₂, C-6), 16.7 (C, C-7), 36.6 (CH, C-8), 27.0 (CH₂, C-9), 27.9 (CH₂, C-10), 49.7 (CH, C-11), 48.5 (CH₂, C-12), 39.7 (C, C-13), 27.0 (CH₃, C-14), 34.4 (CH₃, C-15), 140.6 (CH, C-2'), 150.2 (C, C-4'), 108.1 (C, C-5'), 155.8 (C, C-6'), 151.9 (C, C-8'); HRESIMS m/z 369.1920 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{20}\text{H}_{25}\text{N}_4\text{O}_3$ 369.1926).

Biological Assays. Human breast carcinoma MDA-MB-231 and MCF cell lines were purchased from the American Type Culture Collection (ATCC, Rockville, MD). Cytotoxicity assays were measured by MTT methods as described previously.¹⁴

Acknowledgment. The authors are grateful to the Knowledge Innovation Program of Chinese Academy of Science (grant: KZCX3-SW-216) and Guangdong Natural Science Foundation (2003)11 for financial support.

References and Notes

- Groweiss, A.; Fenical, W.; He, C. H.; Clardy, J.; Wu, Z. D.; Yiao, Z. N.; Long, K. H. *Tetrahedron Lett.* **1985**, 26, 2379–2382.
- Parameswaran, P. S.; Naik, C. G.; Kamat, S. Y.; Puar, M. S.; Das, P.; Hegde, V. R. *J. Nat. Prod.* **1998**, 61, 832–834.
- Wang, G. H.; Ahmed, A. F.; Sheu, J. H.; Duh, C. Y.; Shen, Y. C.; Wang, L. T. *J. Nat. Prod.* **2002**, 65, 887–891.
- Wang, G. H.; Ahmed, A. F.; Kuo, Y. H.; Sheu, J. H. *J. Nat. Prod.* **2002**, 65, 1033–1036.
- Bokesch, H. R.; McKee, T. C.; Cardellina, J. H.; Boyd, M. R. *Tetrahedron Lett.* **1996**, 37, 3259–3262.
- Anjaneyulu, A. S. R.; Rao, K.; Venkateswara, G. *Indian J. Chem.* **1997**, 36B, 418–423.
- Aknin, M.; Costantino, V.; Mangoni, A. *Steroids* **1998**, 63, 575–578.
- Subrahmanyam, C.; Kumar, S. R.; Reddy, G. D. *Indian J. Chem.* **2003**, 42B, 219–220.
- Qi, S. H.; Zhang, S.; Huang, H.; Huang, J. S.; Xiao, Z. H. *J. Nat. Prod.* **2004**, 67, 1907–1910.
- Qi, S. H.; Zhang, S.; Xiao, Z. H.; Huang, J. S. *Magn. Reson. Chem.* **2005**, 43, 266–268.
- Sheu, J. H.; Hung, K. C.; Wang, G. H.; Duh, C. Y. *J. Nat. Prod.* **2000**, 63, 1603–1607.
- Su, J. Y.; Long, K. H.; Jian, Z. G. *Acta Sci. Nat. Univ. Sunyatseni* **1984**, 1, 97–101.
- Yu, D. Q.; Yang, J. D. *Analytical Chemistry Book*, Vol. VII; Chemical Industry Press: Beijing, 1999; p 743.
- Bednarski, P. J. *Biochem. Pharmacol.* **1992**, 43, 2609–2620.

NP050105L