

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/231715654>

Myricetin, Dihydromyricetin, and Quercetin Glycosides from *Catha edulis*

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · JUNE 2004

Impact Factor: 3.8 · DOI: 10.1021/np50043a032

CITATIONS

5

READS

44

3 AUTHORS, INCLUDING:



Mohamed S Hifnawy

Cairo University

48 PUBLICATIONS 380 CITATIONS

SEE PROFILE

MYRICETIN, DIHYDROMYRICETIN, AND QUERCETIN GLYCOSIDES
FROM *CATHA EDULIS*

IBRAHIM A. AL-MESHAL, MOHAMED S. HIFNAWY, and MOHAMMAD NASIR

Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia

As a continuation of our studies of the chemistry and biology of "khat," *Catha edulis* Forsk. (Celastraceae) (1-5), we report the isolation and characterization of four flavonoid glycosides, three of which were not previously reported in "khat."

EXPERIMENTAL

PLANT MATERIAL.—Plants of *C. edulis* were collected during April-May 1984, from the Fifa area in the southern region of Saudi Arabia. A voucher specimen representing the collection is deposited in the herbarium of the Research Centre, College of Pharmacy, King Saud University.

EXTRACTION AND ISOLATION OF FLAVONOIDS.—Air-dried leaves and young shoots of *C. edulis* (1.5 kg) were exhausted with C_6H_6 followed by $CHCl_3$ and then EtOAc. Evaporation of EtOAc in vacuo afforded 20 g of a dark-brown solid residue. Chromatographic separation of 6 g using Sephadex LH-20, polyamide and cellulose columns, and a Me_2CO/H_2O mixture for elution yielded myricetin-3-O- β -D-galactoside (80 mg), dihydromyricetin-3-O-rhamnoside (1.004 g), myricetin-3-O-rhamnoside (40 mg), and quercetin-3-O- β -D-galactoside (10 mg).

Identity of these glycosides was based on the standard spectral data of the original, hydrolytic, and methyl derivatives as well as by comparison with those reported in the literature (6-10). The structure of myricetin-3-O- β -D-galactoside was further confirmed by ^{13}C nmr. Details of the isolation and identification are available from the major author.

ACKNOWLEDGMENTS

This work was supported by the Saudi Arabia National Center for Science and Technology (SANCST). We are grateful to the Department of Chemistry, College of Science, King Saud University, for 1H -nmr and ^{13}C -nmr services.

LITERATURE CITED

1. I. A. Al-Meshal, M. S. Hifnawy, A. G. Mekki, M. Tariq, and F. J. Muhtadi, in: "Characterization and Evaluation of Saudi Arabian Khat," Proceedings of the International Conference on Khat, Antananarivo, Madagascar, January 17-21, 1983, pp. 110-134.
2. M. Tariq and I. A. Al-Meshal, in: "Toxic Studies on *Catha edulis*," Third International Congress on Toxicology, San Diego, California, 1983, p. 100.
3. I. A. Al-Meshal, A. M. Ageel, M. Tariq, and N. S. Parmar, *Res. Comm. Substance Abuse*, **4**, 143 (1983).
4. M. Tariq, N. S. Parmar, A. M. Ageel, I. A. Al-Meshal, and A. R. Abu-Jayyab, *Res. Comm. Substance Abuse*, **4**, 143 (1984).
5. I. A. Al-Meshal, M. Tariq, N. S. Parmar, A. M. Ageel, in: "The antiinflammatory activity of the flavonoid fraction of khat (*Catha edulis* Forsk.)," Seventh European Workshop on Inflammation, Capri (Italy), 1985, p. 52.
6. M. Gellert, K. Szendrei, and J. Reisel, *Phytochemistry*, **20**, 1759 (1981).
7. T. J. Mabry, K. R. Markham, and M. F. Thomas, "The Systematic Identification of Flavonoids," Springer-Verlag, Berlin, 1970, pp. 65, 67, 102.
8. T. K. Devon and A. T. Scott, "Handbook of Naturally Occurring Compounds," vol. 1, Academic Press, New York, 1975, p. 152.
9. E. Lamar and T. Bodalski, *Diss. Pharm. Pharmacol.*, **20**, 623 (1968).
10. H. Thiem, *Pharmazie*, **24**, 783 (1969).
11. K. R. Markham, B. Ternai, R. Stanelly, H. Geiger, and T. J. Mabry, *Tetrahedron*, **34**, 1389 (1978).

Received 18 June 1985