

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/248738551>

Sesquiterpene Lactones, Chromans, and Other Constituents of *Ophryosporus* *piquerioides*

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · AUGUST 1992

Impact Factor: 3.8 · DOI: 10.1021/np50086a022

CITATIONS

14

READS

7

4 AUTHORS, INCLUDING:



E. Elizabeth Sigstad

National Scientific and Tech...

22 PUBLICATIONS 199 CITATIONS

SEE PROFILE




Cesar Catalan

National University of Tucum...

215 PUBLICATIONS 2,475

CITATIONS

SEE PROFILE



ment (M45 pump, U6K injector with 2 ml loop and R-401 differential refractometer) and also a Konik KNC-500A liquid chromatograph (RI detector and Rheodyne injector with 3 ml loop) were used. Columns employed were a Phenomenex Maxsil 10C8 (10 × 500 mm) and a Phenomenex Ultramex 5C18 (10 × 250 mm). Rt's were measured from the solvent peak. Nmr spectra were obtained by means of Varian 300 and 500 MHz spectrometers and ms with a Finnigan 6C/MS system Model 4510 at 70 eV. Known compounds were identified by ms and ¹H-nmr spectra.

PLANT MATERIAL.—Aerial parts of *O. piquerioides* were collected at the flowering stage on May 18, 1989 at Sierras de Medina, Tucumán Province, Argentina. A voucher specimen (C. Catalán No. 101) is on deposit in the herbarium of the Instituto Miguel Lillo, Tucumán, Argentina.

EXTRACTION AND ISOLATION.—Air-dried flowers and leaves (619 g) were extracted with CHCl₃ (2 × 5 liters) at room temperature for 3 days to give after evaporation 49.6 g (8.2%) of crude extract which was suspended in EtOH (566 ml) at 55° and diluted with H₂O (426 ml). The mixture was extracted successively with *n*-hexane (3 × 300 ml) and CHCl₃ (3 × 900 ml). The CHCl₃ extract on evaporation at reduced pressure furnished a residue (7.4 g) which was chromatographed over Si gel (200 g) using CHCl₃ with increasing amounts of EtOAc (0–100%), 58 fractions being collected which were monitored by tlc.

Fractions 15–26 (480 mg) were combined and subjected to cc on Si gel [CHCl₃-EtOAc (19:1)], 20 fractions being collected. Fractions 6–20 of the rechromatogram (296 mg) were combined; a 100-mg portion was processed by hplc [MeOH-H₂O (2:1), 2 ml/min] to give 0.9 mg of **1** (Rt 13 min), mixtures or unidentified material (Rt 14, 20.3, 22.5 min), 0.6 mg of **2** (Rt 25.1 min), 2.7 mg of **5** (Rt 16.3 min), 0.5 mg of **4** (Rt 21.1 min), and 1.5 mg of **3** (Rt 23.1 min). Fractions 27–52 from the original column were combined (1.41 g) and subjected to flash chromatography [Si gel, CHCl₃-EtOAc (3:1), 22 fractions]. Fractions 4–9 (884 mg) were combined, and a portion (100 mg) was processed by hplc [MeOH-H₂O (3:2), 1.6 ml/min] to give 29.2 mg of a mixture which showed four spots on tlc. Final purification by tlc on Si gel 60 HF [CHCl₃-EtOAc-MeOH (170:30:3)] gave 6.3 mg of **6** [Rt 0.13 min, CHCl₃-EtOAc (3:1)], 8.1 mg of **7** [Rt 0.21 min, CHCl₃-EtOAc (3:1)], 0.8 mg of syringaresinol [Rt 0.26 min, CHCl₃-EtOAc (3:1)], and 0.6 mg of scopoletin [Rt 0.42 min, CHCl₃-EtOAc (3:1)].

trans-3,4-Dihydroxy-2,2-dimethyl-6,7-dimethoxychroman [**6**].—Obtained only in the form of a gum: ir (film) 3401, 3009, 2966, 1612, 1503, 1448, 1409, 1379, 1364, 1327, 1266, 1220, 1188, 1170, 1144, 1122, 1005, 913, 852, 834, 793, 766, 665 cm⁻¹; ¹H-nmr (300 MHz, CDCl₃) δ 6.92 (s, H-8), 6.37 (s, H-5), 4.51 (brd, *J* = 8.5 Hz, H-4), 3.59 (d, *J* = 8.5 Hz, H-3), 3.85 and 3.82 (both s, 3H, -OMe's), 1.47 and 1.22 (s and brs, both 3H, Me's). The substance, kept in CHCl₃, had decomposed by the time an attempt at measuring the ms was made.

cis-3,4-Dihydroxy-2,2-dimethyl-6,7-dimethoxychroman [**7**].—Obtained only in the form of a gum: ir (film) 3403, 1619, 1510, 1451, 1413, 1384, 1369, 1335, 1265, 1244, 1201, 1148, 1128, 1106, 1012, 1040, 1012, 974, 940, 859, 833, 821, 790, 757, 665 cm⁻¹; ¹H-nmr (CDCl₃) δ 6.99 (s, H-8), 6.38 (s, H-5), 4.75 (br, H-4), 3.67 (br, H-3), 3.84 and 3.82 (both s, 3H, -OMe's), 1.46 and 1.27 (both s, 3H, Me's), 2.60 and 2.00 (both br, -OH's). On addition of D₂O the H-3 and H-4 signals sharpened to d's (*J* = 4) and the -OH signals disappeared. The substance, kept in CDCl₃, had decomposed by the time an attempt at measuring the ms was made.

ACKNOWLEDGMENTS

Work in Tucumán was supported by grants from CONICET (Argentina) and Fundación Antorchas. J. G. D. thanks the Dirección General de Investigación Científica y Técnica for a fellowship.

LITERATURE CITED

1. T. Bohlmann and C. Zdero, *Phytochemistry*, **18**, 145 (1979).
2. F. Bohlmann, M. Wallmeyer, R.M. King, and H. Robinson, *Phytochemistry*, **23**, 1513 (1984).
3. V.L. Ferracini, I. Roewer, F. Gao, and T.J. Mabry, *Phytochemistry*, **28**, 1463 (1989).
4. C. Zdero, F. Bohlmann, and H.M. Niemeyer, *Phytochemistry*, **29**, 3247 (1990).
5. A.B. Gutiérrez and W. Herz, *Phytochemistry*, **27**, 2225 (1988).
6. F. Bohlmann, R.K. Gupta, J. Jakupovic, R.M. King, and H. Robinson, *Phytochemistry*, **20**, 1635 (1981).
7. W. Herz and P. Kulanthaivel, *Phytochemistry*, **22**, 715 (1983).
8. W. Herz and P. Kulanthaivel, *Phytochemistry*, **23**, 435 (1984).

Received 27 January 1992