

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231110576>

Structure and Plant Growth Regulatory Activity of New Diterpenes from *Pterodon polygalaeflorus*

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · AUGUST 1996

Impact Factor: 3.8 · DOI: 10.1021/np960140f

CITATIONS

34

READS

25

4 AUTHORS, INCLUDING:



[Luiz Claudio De Almeida Barbosa](#)

Federal University of Minas Gerais

359 PUBLICATIONS 2,295 CITATIONS

SEE PROFILE

Structure and Plant Growth Regulatory Activity of New Diterpenes from *Pterodon polygalaeflorus*

Antonio Jacinto Demuner and Luiz Cláudio de Almeida Barbosa*

Departamento de Química, Universidade Federal de Viçosa, 36571-000 Viçosa-MG, Brazil

Dorila Piló Veloso and Dalton Luiz Ferreira Alves

Departamento de Química, Universidade Federal de Minas Gerais, 31270-901 Belo Horizonte-MG, Brazil

Oliver W. Howarth

Department of Chemistry, University of Warwick, Coventry CV4 7AL, U.K.

Received December 14, 1995*


Two new furanditerpenes, vouacapane-6 α ,7 β ,17 β -triol (**3**) and vouacapane-6 α ,7 β ,17 β ,19-tetraol (**4**), have been isolated from the seeds of *Pterodon polygalaeflorus* Benth. The selective effects of these two diterpenes and of 6 α ,7 β -dihydroxyvouacapane-17 β -oic acid (**1**) and 6 α -hydroxyvouacapane-7 β ,17 β -lactone (**2**) on the radicle growth of *Sorghum bicolor* and *Cucumis sativus* were evaluated.

The genus *Pterodon* (Leguminosae) is comprised of five native species in Brazil: *P. abruptus* Vog., *P. apparicioi* Pedersoli, *P. emarginatus* Benth, *P. polygalaeflorus* Benth, and *P. pubescens* Benth.^{1,2}

Oil from the fruits of *P. pubescens* inhibits skin penetration of *Schistosoma mansoni* cercaria, and this activity has been attributed to 14,15-dihydroxygeranylgeraniol.² An alcoholic infusion of the seeds of *P. polygalaeflorus* or *P. apparicioi* is traditionally used by the population of Goiás and Minas Gerais states in Brazil to treat throat infections. Further chemical investigation³ of the seed oil of *P. emarginatus* led to the isolation of 6 α ,7 β -dihydroxyvouacapane-17 β -oic acid (**1**) and 6 α -hydroxyvouacapane-7 β ,17 β -lactone (**2**). These two compounds were shown to have antiinflammatory and analgesic activity.⁴ Since this discovery we have been engaged in the preparation of several derivatives of the acid **1**,⁴⁻⁷ and polyols **3** and **4** were isolated during the routine isolation of quantities of **1** required for study of structure-activity relationships.

Compound **3** was isolated from *P. polygalaeflorus* Benth (Leguminosae) seeds as a white solid, mp 199–201 °C, and the molecular formula, C₂₀H₃₀O₄, was derived from HRMS and ¹³C-NMR spectra. Its IR spectrum showed hydroxyl absorptions bands at 3560 (sharp, free OH) and 3425 (broad, hydrogen bonded OH) cm⁻¹ and for a furan ring at 1678 and 1510 cm⁻¹ (C=C).

The ¹³C-NMR spectrum exhibited signals corresponding to four CH₃, four CH₂, seven CH, and five nonhydrogenated carbons. The presence of three hydroxyl groups was suggested by signals (CDCl₃) at δ 78.87 and 74.12 for CH and δ 72.46 for C and by the disappearance of three signals which were not observed in pyridine-d₅ solution ¹H-NMR at δ 4.02 (d, *J* = 1.3 Hz), 3.43 (s) and 2.18 (d, *J* = 4.6 Hz).



The ¹H-NMR spectrum confirmed the presence of four methyl groups by the signals at δ 1.69 (s, CH₃-17), 1.54 (s, CH₃-18), 1.29 (s, CH₃-19), and 1.03 (s, CH₃-20). Each assignment was confirmed by NOE difference experiments as shown in Figure 1. Stereochemistries at C-6 and C-7 were the same for all the furanditerpenes previously isolated from *Pterodon*, and this was clear from the large NOE observed at H-6 (3.4%) when CH₃-20 was irradiated and from the absence of any significant NOE at H-7 when H-8 was irradiated. The absence of NOE at the methyl attached to carbon-14 when H-8 was irradiated confirmed the proposed stereochemistry at C-14 as having the OH group in the β -position (Figure 1). Full ¹H- and ¹³C-NMR assignments were completed by analysis of HH-COSY, HC-HETCOR, DEPT, and NOE experiments.

Mahajan and Monteiro³ previously proposed structure **3** for a compound isolated from *P. emarginatus*. The structure was proposed on the basis of elemental

* To whom correspondence should be addressed. Fax: 31-899 2376. E-mail: LCAB@mail.UFV.BR.

* Abstract published in *Advance ACS Abstracts*, June 15, 1996.




Figure 1. Major NOE data for compounds 3 and 4.

analysis, IR, and degradation studies, but no NMR data were reported. The published melting point for compound 3 (218–222 °C) was higher than the value (199–201 °C) found by us. Also, the IR data reported³ showed only one stretching band for OH at 3413 cm^{-1} , while our data for compound 3 revealed two distinct OH stretching bands at 3560 and 3425 cm^{-1} . In view of these differences in spectroscopic and physical data we believe that the compound isolated by Mahajan and Monteiro³ could be an isomer of triol 3.

Furanditerpene (4)⁸ analyzed for $\text{C}_{20}\text{H}_{30}\text{O}_5$ by HRMS. Its IR spectrum showed a broad band at 3150–3550 cm^{-1} , with maxima at 3450, 3400, and 3280 cm^{-1} . The ^1H -NMR (CDCl_3) spectrum was very similar to that of compound 3, but only three methyl signals were present (δ 1.47, 1.25, and 1.08). However, four hydroxyl groups were observed in CDCl_3 at δ 4.15 (s), 3.90 (d, $J = 3.0$ Hz), 3.79 (s) and 2.24 (bt, $J_1 = 3.1$ Hz, $J_2 = 2.8$ Hz) that were absent in pyridine- d_5 . Also, hydrogens H-19 and H-19' appeared as a pair of double doublets at δ 4.27 ($J_1 = 10.3$ Hz, $J_2 = 3.1$ Hz) and δ 3.39 ($J_1 = 10.3$ Hz, $J_2 = 2.8$ Hz). In pyridine- d_5 , these signals appeared at δ 4.00 and 3.90 respectively.

The NOE observed at H-19 (2.0%) and H-19' (1.2%) when the methyl at C-10 was irradiated and at H-5 (2.2%) when CH_3 -18 was irradiated confirmed the β -(hydroxymethyl) stereochemistry at C-4 (Figure 1). ^1H -NMR and ^{13}C -NMR assignments presented were made on the basis of HH-COSY, HC-HETCOR, DEPT, and NOE experiments.

The antiinflammatory activity of compound 1 and some other furanditerpenes has been reported,⁴ but no report on the plant growth regulatory activity of this class of compound is available. Allelopathic activity of monoterpenes is widely reported,⁹ but little is known about the effect of diterpenes on plant growth.¹⁰ Compounds 1–4 were then submitted to a plant growth bioassay¹¹ to evaluate their effect on the radicle growth of *Sorghum bicolor* and *Cucumis sativus*. The experiments were carried out at two concentrations (100 and 1000 ppm) of each compound since it is known that some




Figure 2. Effect of compounds 1–4 on the radicular growth of *S. bicolor* at 100 and 1000 ppm after 3 days of incubation at 25 °C.




Figure 3. Effect of compounds 1–4 on the radicular growth of *C. sativus* at 100 and 1000 ppm after 3 days of incubation at 25 °C.

compounds exhibit both stimulatory and inhibitory effects on seedling growth depending on the concentration.¹⁰ Germination rate was not affected by any of these compounds. Diterpene 1 showed a 39% inhibitory effect on the radicular growth of *S. bicolor* at 100 ppm, and this effect was even higher (65%) at 1000 ppm. The other three compounds (2–4) showed a small inhibitory effect at 100 ppm, which increased to 13–40% at 1000 ppm (Figure 2). All compounds showed a stimulatory effect on radicular growth at 100 ppm for *C. sativus*. This effect decreased at 1000 ppm for compounds 2 and 3, while compounds 1 and 4 were significantly inhibitory (Figure 3).

Experimental Section

General Experimental Procedures. IR spectra were recorded with a Perkin-Elmer 881 double-beam grating spectrophotometer. NMR spectra were recorded with a Bruker WH 400 spectrometer (400 MHz), using TMS as an internal standard. Mass spectra were recorded on a V. G. Analytical ZAB-IF spectrometer. Flash chromatography was performed using Crosfield Sorbil C60 (40–60 μm), and the solvents used were purified according to Perrin and Armarego.¹² The melting points were corrected.

Plant Material. Seeds of *P. polygalaeiflorus* Benth were collected in Trés Marias-MG in September 1992. A voucher specimen (no. 10714) is deposited at the Herbarium of Department of Biology of Federal University of Viçosa.

Extraction and Isolation. Seeds were ground and extracted in a Soxhlet apparatus with hexane for 40 h. The hexane extract was submitted to column chromatography on silica gel,³ eluting with hexane:ether (1:10), to afford compounds 3 ($R_f = 0.40$) and 4 ($R_f = 0.22$).

Vouacapane-6 α ,7 β ,17 β -triol (3): white solid; mp 199–201 °C (lit.³ mp 218–220 °C); IR (KBr) ν_{\max} 3560, 3425, 3000, 2925, 1875, 1678, 1510, 1468, 1370, 1280, 1150, 1090, 910, and 725 cm⁻¹; ¹H-NMR (pyridine-*d*₅, 400 MHz) δ 1.03 (s, 20-CH₃), 1.03 (btd, $J_1 \approx J_2 = 12.8$ Hz, $J_3 = 3.5$ Hz, H-1 α), 1.04 (d, $J = 12.2$ Hz, H-5), 1.29 (s, 19-CH₃), 1.35 (btd, $J_1 \approx J_2 = 12.8$ Hz, $J_3 = 4.1$ Hz, H-1 β), 1.54 (s, 18-CH₃), 1.59–1.71 (m, H-3 α , H-3 β), 1.63–1.67 (m, H-9), 1.69 (s, 17-CH₃), 2.26 (dd, $J_1 = 12.5$ Hz, $J_2 = 10.1$ Hz, H-8), 2.47 (dd, $J_1 = 16.6$ Hz, $J_2 = 10.5$ Hz, H-11 β), 2.67 (dd, $J_1 = 16.6$ Hz, $J_2 = 6.5$ Hz, H-11 α), 4.02–4.09 (m, H-6 and H-7), 6.71 (d, 1.8 Hz, H-15), 7.47 (d, 1.8 Hz, H-16); ¹H-NMR (CDCl₃, 400 MHz) δ 0.95 (dt, $J_1 \approx J_2 = 12.7$ Hz, $J_3 = 4.0$ Hz, H-1 α), 0.99 (s, 20-CH₃), 1.03 (d, $J = 11.1$ Hz, H-5), 1.08 (s, 19-CH₃), 1.18 (s, 18-CH₃), 1.23 (ddt, $J_1 \approx J_2 = 12.7$ Hz, $J_3 = 4.1$ Hz, H-3 β), 1.37–1.43 (m, H-9), 1.40–1.68 (m, H-1 β , H-2 α , H-2 β , H-3 α), 1.45 (s, 17-CH₃), 1.96 (dd, $J_1 = 12.4$ Hz, $J_2 = 10.4$ Hz, H-8), 2.18 (d, $J = 4.6$ Hz, 6-OH), 2.41 (dd, $J_1 = 16.7$ Hz, $J_2 = 10.6$ Hz, H-11 β), 2.58 (dd, $J_1 = 16.7$ Hz, $J_2 = 6.6$ Hz, H-11 α), 3.43 (s, 14-OH), 3.67 (bt, $J_1 \approx J_2 = 9.1$ Hz, H-7), 3.82 (ddd, $J_1 = 11.1$ Hz, $J_2 = 9.1$ Hz, $J_3 = 4.6$ Hz, H-6), 4.02 (d, $J = 1.3$ Hz, 7-OH), 6.40 (d, 1.9 Hz, H-15), 7.23 (d, 1.9 Hz, H-16); ¹³C-NMR (pyridine-*d*₅, 100 MHz) δ 149.17 (C-12), 141.70 (C-16), 125.51 (C-13), 108.41 (C-15), 78.87 (C-7), 74.12 (C-6), 72.46 (C-14), 55.76 (C-5), 49.04 (C-8), 47.38 (C-9), 44.26 (C-3), 40.18 (C-1), 38.78 (C-10), 37.30 (C-18), 33.90 (C-4), 26.64 (C-17), 22.96 (C-19), 22.91 (C-11), 19.00 (C-2), 16.17 (C-20); HRMS (70 eV) m/z 334.2135 (M⁺, C₂₀H₃₀O₄, requires 334.2136, 2), 319 (79), 301 (100), 283 (31), 173 (9), 145 (22), 131 (7), 109 (26), 81 (16), 69 (20), 43 (25).

Vouacapane-6 α ,7 β ,17 β ,19-tetraol (4): white solid; mp 181–183 °C; IR (KBr) ν_{\max} 3500–3200, 3010, 2950, 2860, 1650, 1518, 1480, 1350, 1250, 1080, 1030, 1010, 900, and 690 cm⁻¹; ¹H-NMR (pyridine-*d*₅, 400 MHz) δ 0.98–1.02 (m, H-1 α), 0.99 (s, 20-CH₃), 1.17 (dt, $J_1 \approx J_2 = 12.8$ Hz, $J_3 = 4.1$ Hz, H-3 α), 1.35–1.40 (m, H-2 α), 1.46–1.48 (m, H-2 β), 1.43 (d, $J = 11.1$ Hz, H-5), 1.55 (bd, $J = 12.6$ Hz, H-1 β), 1.63 (ddd, $J_1 = 12.0$ Hz, $J_2 = 10.6$ Hz, $J_3 = 6.5$ Hz, H-9), 1.71 (s, 18-CH₃), 1.73 (s, 17-CH₃), 1.94 (bd, $J = 13.4$ Hz, H-3 β), 2.36 (dd, $J_1 = 12.0$ Hz, $J_2 = 10.5$ Hz, H-8), 2.39 (dd, $J_1 = 16.6$ Hz, $J_2 = 10.6$ Hz, H-11 β), 2.62 (dd, $J_1 = 16.6$ Hz, $J_2 = 6.5$ Hz, H-11 α), 3.90 (d, $J = 10.5$ Hz, H-19'), 4.00 (d, $J = 10.5$ Hz, H-19), 4.16 (dd, $J_1 = 10.5$ Hz, $J_2 = 8.7$ Hz, H-7), 4.37 (dd, $J_1 = 11.1$ Hz, $J_2 = 8.7$ Hz, H-6), 6.81 (d, 1.7 Hz, H-15), 7.54 (d, 1.7 Hz, H-16); ¹H-NMR (CDCl₃, 400 MHz) δ 1.00–1.08 (m, H-3 α), 1.08 (s, 20-CH₃), 1.13 (d, $J = 11.3$ Hz, H-5), 1.18–1.25 (m, H-2 α), 1.25 (s, 18-CH₃), 1.39–1.45 (m, H-1 α and H-9), 1.45–1.55 (m, H-1 β , H-2 β), 1.47 (s, 17-CH₃), 1.72 (dm, H-3 β), 2.02 (dd, $J_1 = 12.5$ Hz, $J_2 = 10.5$ Hz, H-8), 2.24 (bt, $J_1 = 3.1$ Hz, $J_2 = 2.8$ Hz, 19-OH), 2.42 (dd, $J_1 = 16.6$ Hz, $J_2 = 10.6$ Hz, H-11 β), 2.60 (dd, $J_1 = 16.6$ Hz, $J_2 = 6.5$ Hz, H-11 α), 3.39 (dd, $J_1 = 10.3$ Hz, $J_2 = 2.8$ Hz, H-19'), 3.69 (dd, $J_1 = 10.5$ Hz, $J_2 = 9.7$ Hz, H-7), 3.79 (s, 7-OH), 3.90 (d, $J = 3.0$ Hz, 6-OH), 4.04 (ddd, $J_1 = 11.3$ Hz, $J_2 = 9.7$ Hz,

$J_3 = 3.0$ Hz, H-6), 4.15 (s, 14-OH), 4.27 (dd, $J_1 = 10.3$ Hz, $J_2 = 3.1$ Hz, H-19), 6.44 (d, $J = 1.9$ Hz, H-15), 7.24 (d, $J = 1.9$ Hz, H-16); ¹³C-NMR (pyridine-*d*₅, 100 MHz) δ 148.94 (C-12), 141.72 (C-16), 125.71 (C-13), 108.73 (C-15), 78.36 (C-7), 74.21 (C-6), 72.39 (C-14), 66.49 (C-19), 56.08 (C-5), 49.20 (C-8), 47.59 (C-9), 40.26 (C-1), 39.29 (C-4), 39.25 (C-3), 38.16 (C-10), 32.15 (C-18), 26.94 (C-17), 23.03 (C-11), 18.82 (C-2), 16.98 (C-20); HRMS m/z 350.2084 (M⁺, C₂₀H₃₀O₅, requires 350.2085, 3), 335 (100), 317 (72), 299 (16), 269 (16), 131 (10), 109 (27), 81 (13), 43 (29).

Bioassays. Bioassays were carried out according to the method of Einhelling *et al.*¹⁰ with seeds of *S. bicolor* and *C. sativus*. CH₂Cl₂ solutions of compounds 1–4 were prepared at concentrations of 100 and 1000 ppm.

Assays were conducted in 100 × 15 mm glass Petri dishes lined with one sheet of Whatman no. 1 filter paper and sealed with Parafilm. To each dish was added 2 mL of each solution, and the solvent was evaporated before addition of 2 mL of water followed by 20 seeds of one of the two species. Assays were carried out at 25 °C under fluorescent light (8 × 40 W) in an incubator for 3 days. Radicle length was measured, and total germination was recorded. Seeds were considered to have germinated if a radicle protruded at least 1 mm. A control experiment was carried out under the same conditions described, using only water. Each bioassay was replicated four times in a complete randomized design.

Acknowledgment. We thank the Brazilian agencies CNPq, CAPES, and FAPEMIG for financial support.

References and Notes

- (1) Fascio, M.; Mors, W. B.; Gilbert, B.; Mahajan, J. R.; Monteiro, M. B.; Santos Filho, D.; Vichnewski, W. *Phytochemistry* **1976**, *15*, 201–203.
- (2) Mors, W. B.; Santos Filho, M. F.; Monteiro, H. J.; Gilbert, B. *Science* **1967**, *157*, 950–951.
- (3) Mahajan, J. R.; Monteiro, M. B. *J. Chem. Soc., Perkin Trans. 1* **1973**, 520–525.
- (4) Nunan, E. A. *Estudos da atividade antiinflamatória de furanoditerpenos isolados de Pterodon polygalaeiflorus Benth e de alguns de seus derivados*. MSc. Thesis, Departamento de Química (UFMG), Belo Horizonte, MG, 1985, 129 pp.
- (5) Moreira, J. M. *Síntese de derivados glicosilados do ácido 6 α ,7 β -dihidroxiouacapan-17 β -óico*. MSc Thesis, Departamento de Química (UFMG), Belo Horizonte, MG, 1990, 183 pp.
- (6) Rubinger, M. M. M.; Veloso, D. P. *J. Braz. Chem. Soc.* **1991**, *2*, 124–127.
- (7) Maltha, C. R. A.; Stefani, G. M.; Veloso, D. V. *J. Braz. Chem. Soc.* **1995**, *6*, 357–361.
- (8) After submitting this paper, we learned that compound 4 was presented at the following meeting: Arriaga, A. M. C.; Silveira, E. R.; Braz-Filho, R.; Castro, M. A. B.; Dourado, R. C. M.; Andrade, I. C. M. 18^a Reunião Anual da SBQ, Caxambu, 1995.
- (9) Vaughn, S. F.; Spencer, G. F. *Weed Sci.* **1993**, *41*, 114–119.
- (10) Einhelling, F. A.; Schan, M. K.; Rasmussen, J. A. *Plant Growth Regul.* **1983**, *1*, 251–258.
- (11) Elakovich, S. D. In *Terpenoids as Model for New Agrochemicals*; Cutler, H. G., Ed.; American Chemical Society: Washington, DC, 1988.
- (12) Perrin, D. D.; Armarego, W. L. F. *Purification of Laboratory Chemicals*, 3rd ed.; Pergamon Press: Oxford, 1988.

NP960140F