

# A general method for making bicyclic compounds with nitrogen at a bridgehead—application to the halichlorine spiro subunit†

Derrick L. J. Clive,\* Maolin Yu and Zhiyong Li


Received (in Corvallis, OR, USA) 26th August 2004, Accepted 7th October 2004

First published as an Advance Article on the web 23rd December 2004


DOI: 10.1039/b413481h

*N*-Protected  $\beta$ -amino aldehydes having the nitrogen in a ring are easily converted into Morita–Baylis–Hillman adducts; *O*-acetylation and *N*-deprotection result in spontaneous cyclization to bicyclic structures having nitrogen at a bridgehead.

A recent report from this laboratory<sup>1</sup> described the preparation of spiro amine **1**, which was made during studies on the total synthesis of the marine natural product halichlorine (**2**).<sup>2</sup> An obvious next step was to convert **1** into **3**, and this has now been achieved through the development of a new and general method for making compounds that contain a bicyclic subunit with nitrogen at a bridgehead. The approach (Scheme 1) is based on sequential formation of Morita–Baylis–Hillman (MBH) alcohols (**4**  $\rightarrow$  **5**) and intramolecular  $S_N2'$  displacement of the derived acetates (**6**  $\rightarrow$  **7**  $\rightarrow$  **8**). In the present work we have made alcohols of type **5** by MBH<sup>3</sup> condensation (and used only acrylates), but the same compounds should also be accessible by other<sup>3,4</sup> methods.



While *O*-acetates of MBH alcohols are known to undergo intermolecular  $S_N2'$  displacement,<sup>3,5</sup> the intramolecular 6-*endo* pathway<sup>6</sup> (Scheme 1, **7**  $\rightarrow$  **8**) requires that no competing




Scheme 1

† Electronic supplementary information (ESI) available: characterization data for compounds **3**, **21**, **26**, **32** and **38**. See <http://www.rsc.org/suppdata/cc/b413481h/>


\*derrick.clive@ualberta.ca

stereoelectronic or reactivity factors intervene to direct cyclization onto the  $CO_2R$  group<sup>7</sup> of **7** or to cause *O*  $\rightarrow$  *N* acetyl transfer; in the event, the desired ring closure (**7**  $\rightarrow$  **8**) occurs smoothly.

The lactam **9**,<sup>1</sup> an intermediate in the preparation of **1**, was deprotected ( $Me_3SiBr$ , 84%, **9**  $\rightarrow$  **10**) and oxidized (Swern, 85%) to aldehyde **11**. Wittig reaction with  $Ph_3P=CH(OMe)$  and hydrolysis (CSA, aqueous  $MeCN$ ) of the intermediate enol ethers then gave the expected aldehyde (**11**  $\rightarrow$  **12**  $\rightarrow$  **13**, 74% overall). When aldehyde **13**<sup>8</sup> was dissolved in methyl acrylate, condensation occurred on addition of DABCO and  $Sc(OTf)_3$ .<sup>9</sup> Although the resulting alcohols (**14a**, more polar) and **14b** (less polar) could be separated, it was more convenient to acetylate the mixture and separate the corresponding acetates **15a** (37% from **13**) and **15b** (34%). When each of the acetates **15a** and **15b** was treated with  $Me_3OBF_4$  and then with aqueous  $Na_2CO_3$ , the lactam ring was opened to amino esters **16a** and **16b**, respectively, and these cyclized *in situ* to afford the desired bis-ester **3** (77% for **15a** and 72% for **15b**).<sup>10</sup> Aldehyde **13** was recovered unchanged either after exposure to DABCO in  $CH_2Cl_2$  for 3 days, or when the MBH condensation was worked up before completion. These observations show that epimerization by retro-Michael elimination and re-addition does not occur.



**Scheme 2** Reagents and conditions: (i)  $Me_3SiBr$ ,  $CH_2Cl_2$ ,  $-10^\circ C$ , 2 h, 84%; (ii) Swern oxidation, 85%; (iii)  $MeOCH_2PPh_3Cl$ , *t*-BuOK, THF,  $0^\circ C$ , 2 h; (iv) camphorsulfonic acid,  $MeCN$ –water, 4 h, 74% over two steps; (v) methyl acrylate, DABCO,  $Sc(OTf)_3$ , 5 days; (vi)  $AcCl$ , pyridine,  $CH_2Cl_2$ ,  $0^\circ C$ , 1 h,  $25^\circ C$ , 1 h, over two steps 37% of **15a**, 34% of **15b**; (vii)  $Me_3OBF_4$ ,  $CH_2Cl_2$ , 1.5 h; (viii) 20% aqueous  $Na_2CO_3$ ,  $MeCN$ , 2 h, 77% from **15a**, 72% from **15b**.



**Scheme 3** Reagents and conditions: (i) References 10–12; (ii) methyl acrylate, DABCO, 3 days, 39% (**19a**), 34% (**19b**); (iii) AcCl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>, 1 h, 89% (**20a**), 85% (**20b**); (iv) CF<sub>3</sub>CO<sub>2</sub>H, CH<sub>2</sub>Cl<sub>2</sub>, 1 h; aq Na<sub>2</sub>CO<sub>3</sub>, MeCN, 1 h, 77%; (v) (a) (Boc)<sub>2</sub>O, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, 1 h, room temperature, 12 h, 87%; (b) Dess–Martin periodinane, CH<sub>2</sub>Cl<sub>2</sub>, 1.5 h, 88%; (vi) as in (ii), 46% (less polar), 38% (more polar); (vii) as in (iii), 94% for **24a**, 92% for **24b**; (viii) as in (iv), 77%; (ix) (a) NaN<sub>3</sub>, aq H<sub>2</sub>SO<sub>4</sub>, 0 °C, 16 h, 74%; (b) LiAlH<sub>4</sub>, dioxane, reflux, 24 h, 80%; (x) (a) (Boc)<sub>2</sub>O, EtOAc, 16 h, 98%; (b) Swern, 2 h, 87%; (xi) as in (ii), 5 days, 39% (less polar), 47% (more polar); (xii) as in (iii), 92% for **30a**, 94% for **30b**; (xiii) as in (iv) 90% (more polar), 87% (less polar); (xiv) (a) NaN<sub>3</sub>, aq H<sub>2</sub>SO<sub>4</sub>, 0 °C, 16 h, 80%; (b) LiAlH<sub>4</sub>, THF, reflux, 24 h, 71%; (xv) (a) as in (x), 83%; (b) as in (x), 80%; (xvi) as in (ix) 49% (more polar **36a**), 45% (less polar **36b**); (xvii) as in (iii), 78% for **37a**, 79% for **37b**; (xviii) as in (iv), 84% for **37a**, 93% for **37b**.

The approach of Scheme 2 appears to be general, and we have applied it to several other cases (Scheme 3).

L-Proline (**17**) was converted by literature methods<sup>11–13</sup> into aldehyde **18**, which underwent condensation with methyl acrylate, affording a separable mixture of **19a** (more polar, 39%) and **19b** (less polar, 34%). Acetylation produced the corresponding acetates **20a** (89%) and **20b** (85%). Finally, exposure of a mixture of both acetates to CF<sub>3</sub>CO<sub>2</sub>H resulted in *N*-deprotection, at which point, treatment with aqueous Na<sub>2</sub>CO<sub>3</sub> caused spontaneous cyclization to **21** (77% yield). HPLC analysis [Chiracel OD-H, 1% EtOH–hexane] showed the material to have an enantiomeric purity of 99.8%, indicating that little, if any, racemization occurs in the synthetic sequence.

In another series of experiments, commercial (2-hydroxyethyl)piperidine (**22**) was converted by *N*-protection (Boc<sub>2</sub>O, 87%) and Dess–Martin oxidation (88%) into aldehyde **23**, which underwent efficient MBH condensation [**23** → **24a** (more polar, 38%) and **24b** (less polar, 46%)]. Once again, acetylation (94% for **24a**, 92% for **24b**), *N*-deprotection (CF<sub>3</sub>CO<sub>2</sub>H), and treatment with aqueous Na<sub>2</sub>CO<sub>3</sub> resulted in ring closure, giving **26** (77% from a mixture of both acetates).

We also investigated two other ring sizes for the starting amine. Keto ester **27** was converted by Schmidt reaction and LiAlH<sub>4</sub> reduction into **28**,<sup>14</sup> which was protected on nitrogen [Boc<sub>2</sub>O, 98%], oxidized (Swern, 87%), and subjected to the MBH condensation [**29** → **30a** (more polar, 47%) and **30b** (less polar, 39%)]. Acetylation (92% for **31a**, 94% for **31b**) and *N*-deprotection (CF<sub>3</sub>CO<sub>2</sub>H) and basification gave **32** (90% from **31a**, 87% from **31b**). Similarly, keto ester **33** was converted into amino alcohol **34**

and then into aldehyde **35**. Our standard sequence (**35** → **36a,b**, → **37a,b**, → **38**) then proceeded in the expected way.

In summary, synthetic work related to halichlorine has led to the development of a method for generating bicyclic amines with nitrogen at a bridgehead. The process occurs with preservation of stereochemistry  $\alpha$  to the nitrogen.

All new compounds were fully characterized by spectroscopic methods, including high resolution mass spectrometry. We thank NSERC for financial support and C. Boucher [Boehringer Ingelheim (Canada)] for ee measurements. M.Y. holds a Province of Alberta Graduate Fellowship.

**Derrick L. J. Clive,\* Maolin Yu and Zhiyong Li**  
Chemistry Department, University of Alberta, Edmonton, Alberta,  
Canada T6G 2G2. E-mail: derrick.clive@ualberta.ca;  
Fax: 780-492-8231; Tel: 780-492-3251

## Notes and references

- M. Yu, D. L. J. Clive, V. S. C. Yeh, S. Kang and J. Wang, *Tetrahedron Lett.*, 2004, **45**, 2879–2881.
- D. Trauner, J. B. Schwarz and S. J. Danishefsky, *Angew. Chem., Int. Ed.*, 1999, **38**, 3542–3545.
- Review: D. Basavaiah, A. J. Rao and T. Satyanarayana, *Chem. Rev.*, 2003, **103**, 811–891 and reviews cited therein.
- e.g., (a) T. Tsuda, T. Yoshida and T. Saegusa, *J. Org. Chem.*, 1988, **53**, 1037–1040; (b) P. V. Ramachandran, M. V. R. Reddy, M. T. Rudd and J. R. de Alaniz, *Tetrahedron Lett.*, 1998, **39**, 8791–8794; (c) P. V. Ramachandran, M. V. R. Reddy and M. T. Rudd, *Chem. Commun.*, 1999, 1979–1980; (d) A. Bernardi, S. Cardani, L. Colombo, G. Poli, G. Schimperna and C. Scolastico, *J. Org. Chem.*, 1987, **52**, 888–891; (e) T. Kataoka, T. Iwama, H. Kinoshita, S. Tsujijima, Y. Tsurukami, T. Iwamura and S. Watanabe, *Synlett*, 1999, 197–198.

- 5 *e.g.*, (a) A. B. Charette, M. K. Janes and A. A. Boezio, *J. Org. Chem.*, 2001, **66**, 2178–2180; (b) C.-W. Cho, J.-R. Kong and M. J. Krische, *Org. Lett.*, 2004, **6**, 1337–1339; (c) D. Nilov, R. Räcker and O. Reiser, *Synthesis*, 2002, 2232–2242; (d) P. Bauchat, N. Le Bras, L. Rigal and A. Foucaud, *Tetrahedron*, 1994, **50**, 7815–7826.
- 6 *cf.* (a) D. K. O'Dell and K. M. Nicholas, *J. Org. Chem.*, 2003, **68**, 6427–6430; (b) M. L. Bode and P. T. Kaye, *J. Chem. Soc., Perkin Trans. I*, 1993, 1809–1813.
- 7 *cf.* (a) O. B. FAMILONI, P. T. Kaye and P. J. Klaas, *Chem. Commun.*, 1998, 2563–2564; (b) D. Basavaiah, R. M. Reddy, N. Kumaragurubaran and D. S. Sharada, *Tetrahedron*, 2002, **58**, 3693–3697.
- 8 Reduction of **11** gave back **10**, and ozonolysis of **14** gave **11**. These experiments establish that no epimerization occurs in the sequence leading to **12**.
- 9 *cf.* V. K. Aggarwal, A. Mereu, G. J. Tarver and R. McCague, *J. Org. Chem.*, 1998, **63**, 7183–7189.
- 10 *cf.* (a) I. Hayakawa, H. Arimoto and D. Uemura, *Chem. Commun.*, 2004, 1222–1223; (b) Y. Matsumura, S. Aoyagi and C. Kibayashi, *Org. Lett.*, 2004, **6**, 965–968.
- 11 G. R. Petti, S. B. Singh, D. L. Herald, P. Loyd-Williams, D. Kantoci, D. D. Burkett, J. Barkóczy, F. Hogan and T. R. Wardlaw, *J. Org. Chem.*, 1994, **59**, 6287–6295.
- 12 *cf.* M. F. Ansell, M. P. L. Caton and K. A. J. Stuttle, *J. Chem. Soc., Perkin Trans. I*, 1984, 1069–1077.
- 13 M. Ikeda, J. Shikaura, N. Maekawa, K. Daibuzono, H. Teranishi, Y. Teraoka, N. Oda and H. Ishibashi, *Heterocycles*, 1999, 31–34.
- 14 M. Psiorz, J. Heider, A. Bomhard, M. Reiffen, N. Huel, K. Noll, B. Narr, C. Lillie, W. Kobinger and J. Dammgen, *US Patent 5,175,157*, 1992.