

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/12159043>

Two New Stilbene Dimer Glucosides from Grape (*Vitis vinifera*) Cell Cultures

ARTICLE in JOURNAL OF NATURAL PRODUCTS · FEBRUARY 2001

Impact Factor: 3.8 · DOI: 10.1021/np000426r · Source: PubMed

CITATIONS

68

READS

29

6 AUTHORS, INCLUDING:



Pierre Waffo-Téguo

University of Bordeaux

75 PUBLICATIONS 1,515 CITATIONS

SEE PROFILE



Dongho Lee

Catholic University of Korea

191 PUBLICATIONS 2,235 CITATIONS

SEE PROFILE



Muriel Cuendet

University of Geneva

101 PUBLICATIONS 2,486 CITATIONS

SEE PROFILE



Jean-Michel Merillon

Université Victor Segalen Bordeaux 2

148 PUBLICATIONS 4,719 CITATIONS

SEE PROFILE

Two New Stilbene Dimer Glucosides from Grape (*Vitis vinifera*) Cell Cultures


Pierre Waffo-Teguo,[†] Dongho Lee,[†] Muriel Cuendet,[†] Jean-Michel Mérillon,[‡] John M. Pezzuto,[†] and A. Douglas Kinghorn^{*,†}

Program for Collaborative Research in the Pharmaceutical Sciences and Department of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, University of Illinois at Chicago, Chicago, Illinois 60612, and Groupe d'Etude des Substances Naturelles à Intérêt Thérapeutique, Université de Bordeaux 2, 3 Place de la Victoire, 33000 Bordeaux, France


Received August 31, 2000

Two new stilbene dimer glucosides, resveratrol (*E*)-dehydrodimer 11-*O*- β -D-glucopyranoside (**1**) and resveratrol (*E*)-dehydrodimer 11'-*O*- β -D-glucopyranoside (**2**), were isolated together with the known resveratrol (*E*)-dehydrodimer (**3**) and pallidol (**4**) from *Vitis vinifera* cell cultures. The structures and stereochemistry of the new compounds were determined on the basis of spectroscopic data analysis. Compounds **1** and **2** are dimers that belong to a new type of oligostilbene formed from a resveratrol unit and a resveratrol glucoside unit. Compounds **1** and **3** exhibited nonspecific inhibitory activity against cyclooxygenase-1 and -2, with IC₅₀ values in the range of 5 μ M, whereas compound **4** was approximately 10-fold less active.

Stilbenes occur naturally in several plant families, such as the Dipterocarpaceae, Vitaceae, Cyperaceae, and Gnetaceae,^{1,2} but grapes (*Vitis vinifera* L., Vitaceae) and products manufactured from grapes are considered the most important dietary sources of these substances.^{3,4} A previous study has shown that (*E*)-resveratrol (3,5,4'-trihydroxy-*E*-stilbene), a grapevine phytoalexin, inhibits cyclooxygenase-1 (COX-1).⁵ Earlier phytochemical studies on grape cell cultures have revealed that they biosynthesize resveratrol monomer derivatives.^{6–8} Resveratrol can be biotransformed by *Botrytis cinerea*, a fungal grapevine pathogen, into resveratrol (*E*)-dehydrodimer (**3**), pallidol (**4**), leachinol F, and restrytisols A–C.^{9,10}



- 1:** R₁ = β -glc, R₂ = H
2: R₁ = H, R₂ = β -glc
3: R₁ = H, R₂ = H



In our search for new cancer chemopreventive agents, a diverse group of natural products capable of mediating activities relevant to cancer chemoprevention has been isolated and characterized from plants.^{11,12} In further work

directed toward the search for novel natural product cancer chemopreventive agents, a fraction derived from an EtOAc-soluble extract of a grape cell culture was identified as a potent inhibitor of COX-1 activity (73% inhibition at 70 μ g/mL). Besides resveratrol (*E*)-dehydrodimer (**3**), which was synthesized more than 20 years ago,¹³ and pallidol (**4**),¹⁴ two new glucosides (**1** and **2**) of **3** were isolated by activity-guided fractionation using the COX-1 inhibitory assay, and their structures were determined on the basis of the spectroscopic data analysis.

Compound **1** was purified by semipreparative reversed-phase HPLC, and its molecular formula of C₃₄H₃₂O₁₁ was established by positive HRFABMS (m/z [M + Na]⁺, 639.1837). The ¹H and ¹³C NMR data (Table 1) of **1** indicated that it is a dimer consisting of a resveratrol unit and a resveratrol glucoside unit. Assignments of all ¹H and ¹³C NMR signals for **1** were made by the analysis of COSY, HMQC, and HMBC 2D NMR data. The ¹H NMR spectrum of **1** showed the presence of characteristic resonances in two distinct regions. The former, between δ_H 7.5 and δ_H 6.0, was constituted by a broad doublet at δ_H 7.43 (H-6'), a broad singlet at δ_H 7.26 (H-2'), and a doublet at δ_H 6.87 (H-5') of an ABX-spin system of ring C, two doublets at δ_H 7.25 (H-2 and H-6) and δ_H 6.86 (H-3 and H-5) of an AA'XX'-spin system of a 1,4-disubstituted aromatic ring A, a triplet at δ_H 6.25 (H-12') and a doublet at δ_H 6.53 (H-10' and H-14') of an AB₂-spin of ring D, three broad triplets at δ_H 6.52 (H-12), δ_H 6.44 (H-14), and δ_H 6.35 (H-10) of an ABC-spin system of ring B, and two coupled doublets at δ_H 7.06 (H-7') and δ_H 6.90 (H-8') with a large coupling constant (J = 16.3 Hz) for a trans olefinic proton system. The second region of the ¹H NMR spectrum of **1**, between δ_H 6.0 and δ_H 3.0, was characterized by two doublets at δ_H 5.52 (H-7) and δ_H 4.53 (H-8) for a dihydrobenzofuran moiety, one doublet at δ_H 4.94 for the anomeric proton of a glucose unit, and six protons of a glucosyl moiety between δ_H 4.0 and δ_H 3.0. Also, the ¹³C NMR spectrum showed six signals at δ_C 101.43, 77.87, 77.56, 74.58, 71.16, and 62.49, which are characteristic for a glucose unit.¹⁵ Moreover, enzymatic hydrolysis of **1** with β -glucosidase led to the generation of the aglycon, resveratrol (*E*)-dehydrodimer (**3**). The anomeric proton signal at δ_H 4.94 (1H, d, J = 7.6 Hz) is indicative of a β -configuration for the glucosyl bond.

* To whom correspondence should be addressed. Tel.: (312) 996-0914. Fax: (312) 996-7107. E-mail: Kinghorn@uic.edu.

[†] University of Illinois at Chicago.

[‡] Université de Bordeaux 2.

Table 1. ^1H and ^{13}C NMR Data of Compounds **1** and **2** in $\text{CD}_3\text{COCD}_3^a$

position	1		2	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1		132.51		132.19
2(6)	7.25 d (8.5)	128.55	7.25 d (8.6)	128.27
3(5)	6.86 d (8.5)	116.20	6.85 d (8.6)	115.81
4		158.44		158.99
7	5.52 d (7.8)	93.83	5.46 d (8)	93.75
8	4.53 d (7.8)	57.81	4.47 d (8)	57.49
9		145.27		144.90
10	6.35 t (2)	109.64	6.19 d (2.1)	107.06
11		160.15		159.88
12	6.52 t (2)	103.14	6.28 t (2.1)	102.00
13		159.60		159.88
14	6.44 t (2)	108.50	6.19 d (2.1)	107.06
1'		140.77		140.41
2'	7.26 brs	123.93	7.27 brs	123.60
3'		132.15		131.88
4'		160.57		160.61
5'	6.87 d (8.2)	110.21	6.88 d (8.2)	109.83
6'	7.43 brd (8.2)	128.65	7.44 brd (8.2)	128.76
7'	7.06 d (16.3)	129.10	7.14 d (16.3)	129.30
8'	6.90 d (16.3)	127.28	6.95 d (16.3)	128.20
9'		131.80		131.56
10'	6.53 d (2)	105.71	6.81 brs	107.83
11'		159.54		160.28
12'	6.25 t (2)	102.70	6.53 t (2)	103.50
13'		159.54		158.95
14'	6.53 d (2)	105.71	6.68 brs	106.03
Glc				
1	4.94 d (7.6)	101.43	4.94 d (7.4)	101.10
2		74.58		74.33
3		77.87		77.63
4		71.16		71.02
5		77.56		77.41
6		62.49		62.31

^a TMS was used as the internal standard; chemical shifts are shown in the δ scale with J values (Hz) in parentheses.

Therefore, compound **1** could be proposed as a glycoside of resveratrol (*E*)-dehydrodimer (**3**).

The position of the glucose unit in **1** was determined by the HMBC NMR experiment, which showed a long-range correlation between the anomeric proton signal at δ_{H} 4.94 (H-1 of Glc) and C-11 at δ_{C} 160.15 of ring B. Also, the presence of an ABC-spin system of ring B in the ^1H NMR spectrum of **1** instead of an AB₂-spin system confirmed the position of the glucose moiety at C-11. The relative configuration of the chiral centers of ring E of **1** was deduced from the coupling constant ($J = 7.8$ Hz) between the H-7 and H-8 benzofuran protons, which was identical to the coupling constant found for **3**.¹⁶ Therefore, the aromatic substituents of ring E were arranged in a trans configuration. Accordingly, the structure of **1** was assigned as resveratrol (*E*)-dehydrodimer 11-*O*- β -D-glucopyranoside.

Compound **2** was obtained as a minor constituent, and its molecular formula of $\text{C}_{34}\text{H}_{32}\text{O}_{11}$ was established by positive HRFABMS (m/z [$\text{M} + \text{Na}$]⁺, 639.1890), again corresponding to a dimer of a resveratrol unit and a resveratrol glucoside unit. The ^1H and ^{13}C NMR data (Table 1) of **2** were closely comparable to those of **1** except for the signals of rings B and D. This suggested that compound **2** is a regioisomer of **1** with the position of the glucose unit transferred from C-11 to C-11'. On enzymatic hydrolysis of **2** with β -glucosidase, compound **3** was generated. The position of the glucose unit was confirmed by the HMBC NMR technique, which showed a three-bond correlation between the anomeric proton signal at δ_{H} 4.94 (1H, d, $J = 7.4$ Hz, H-1 of Glc) and C-11' at δ_{C} 160.28 of ring D. Thus, the structure of **2** was concluded

to be resveratrol (*E*)-dehydrodimer 11'-*O*- β -D-glucopyranoside. The relative trans configuration of the chiral centers of ring E was deduced in the same manner as **1**.

Additionally, two known compounds, resveratrol (*E*)-dehydrodimer (**3**) and pallidol (**4**), were isolated and identified by comparison of the observed ^1H and ^{13}C NMR data with literature values.^{9,16,17} Using the ROESY NMR experiment and by J value comparison, the relative stereochemistry of resveratrol (*E*)-dehydrodimer (**3**) was confirmed. Thus, a ROE correlation between H-7 and H-10 (H-14) and the coupling constant ($J = 8$ Hz) between H-7 and H-8 clearly indicated a trans configuration of the chiral centers of ring E.¹⁷ Resveratrol (*E*)-dehydrodimer (**3**) and pallidol (**4**) have not been demonstrated previously as constituents of *V. vinifera* or its cell cultures. Resveratrol (*E*)-dehydrodimer (**3**) was reported recently as a natural product from the lianas of *Gnetum hainanense* C. Y. Cheng.¹⁶

Compounds **1**, **3**, and **4** were evaluated for their cyclooxygenase-1 and -2 (COX-1 and -2) inhibitory activity. Compound **4** was marginally active, with IC₅₀ values of 50 and 80 μM with COX-1 and -2, respectively. The isolated quantity of **2** was not sufficient to permit evaluation of biological activity. Compounds **1** and **3** demonstrated IC₅₀ values of 5.2 and 4.3 μM , respectively, when evaluated with COX-1, and 7.5 and 3.7 μM , respectively, when evaluated with COX-2. Thus, inhibitory potential is reasonably strong with these test agents, but specificity is lacking in both cases.

Experimental Section

General Experimental Procedures. Optical rotations were determined on a Perkin-Elmer 241 polarimeter. UV spectra were recorded on a Beckman DU-7 spectrometer. IR spectra were taken on a JASCO 410 FT-IR spectrometer. NMR spectra were measured on a Bruker DRX-500 MHz spectrometer using a 2.5 mm or a 5 mm sample tube. EIMS and FABMS were obtained using a Finnigan MAT-90 mass spectrometer, and HRFABMS were obtained on a VG 7070-HF instrument. HPLC was performed using a Waters 515 pump and a Waters 2487 UV detector.

Cell Culture Material. Cell cultures of *V. vinifera* (L.) cv. Gamay Freaux var. Tenturier were established in 1978 from pulp fragments of young fruits and provided by C. Ambid (ENSA, Toulouse, France). Suspension cultures of *V. vinifera* were maintained as described previously.¹⁸ Experiments were carried out by inoculating a 7-day-old cell suspension into an induction medium at a 1:8 (v/v) ratio, for each transfer.¹⁸

Extraction and Isolation. Frozen cells (2.5 kg, fresh weight) were extracted with acetone/water as reported previously.^{6,7} The aqueous mixture was partitioned with ethyl acetate. The ethyl acetate extract was chromatographed over a Dowex-50 \times 4-400 cation-exchange resin (Sigma, St. Louis, MO) and eluted by methanol/water. Crude polyphenols were eluted with 50% methanol. For further fractionation, the crude polyphenol-containing fraction was divided into three main subfractions by passage over Sephadex LH-20 and elution with MeOH/H₂O mixtures. Mixtures of (*Z*)-stilbenes and (*E*)-stilbenes were eluted by 20% MeOH and 30% MeOH, respectively, and were not investigated further. A mixture of stilbene dimers was eluted by 100% MeOH and further purified on Toyopearl HW-40S gel (Supelco, Bellefonte, PA), eluted with 100% MeOH, resulting in two main fractions. Final purification of the first fraction by HPLC resulted in the purification of compounds **1** (14 mg, 0.00056% w/w), **2** (0.6 mg, 0.00002% w/w), and **3** (13 mg, 0.00052% w/w) [column ODS-AQ Pack (YMC, Wilmington, NC), 20 \times 250 mm i.d., C₁₈, 5 μm , 120 Å; guard column ODS-AQ Guard Pack (YMC, Wilmington, NC), 20 \times 100 mm i.d.; linear gradient from 40% to 100% MeCN in H₂O (pH 2.4 with TFA), 30 min, 8 mL/min]. Pallidol (**4**,

12 mg, 0.00048% w/w) was purified from the second fraction using HPLC [50% MeCN in H₂O (pH 2.4 with TFA), 40 min, 8 mL/min].

Resveratrol (*E*)-dehydrodimer 11-*O*- β -D-glucopyranoside (1): powder; [α]_D²⁰ -18.9° (c 0.38, MeOH); UV (MeOH) λ_{\max} (log ϵ) 204 (4.73), 221 (4.58), 309 (4.43), 320 (4.39) nm; IR (neat) ν_{\max} 3352, 2924, 1698, 1597, 1515, 1487, 1451, 1356, 1236, 1202, 1152 cm⁻¹; ¹H and ¹³C NMR data of **1**, see Table 1; EIMS *m/z* 454 (15), 320 (40), 246 (28), 208 (45), 144 (100); FABMS *m/z* 639 [M + Na]⁺, 616 [M]⁺, 455 (15), 406 (20), 329 (27), 307 (77), 289 (56), 176 (100); HRFABMS *m/z* calcd for C₃₄H₃₂O₁₁Na 639.1833, found 639.1837.

Resveratrol (*E*)-dehydrodimer 11'-*O*- β -D-glucopyranoside (2): powder; [α]_D²⁰ -12.0° (c 0.05, MeOH); UV (MeOH) λ_{\max} (log ϵ) 206 (4.88), 309 (4.23), 322 (4.21) nm; IR (neat) ν_{\max} 3394, 2924, 1652, 1558, 1507, 1227, 1144, 1034 cm⁻¹; ¹H and ¹³C NMR data of **2**, see Table 1; FABMS *m/z* 639 [M + Na]⁺, 616 [M]⁺, 482 (72), 460 (58), 455 (28), 273 (95); HRFABMS *m/z* calcd for C₃₄H₃₂O₁₁Na 639.1833, found 639.1890.

Resveratrol *E*-dehydrodimer (3): oil; [α]_D²⁰ -1.7° (c 0.23, MeOH) [lit. [α]_D²⁵ -1.15° (c 7.3, acetone)];¹⁰ UV, IR, ¹H and ¹³C NMR, and EIMS data, consistent with literature values.¹⁰

Pallidol (4): [α]_D²⁰ 0° (c 0.45, MeOH) [lit. [α]_D²⁰ 0° (MeOH)];¹⁴ [α]_D²³ -36.3° (c 0.13, MeOH)¹⁶; UV, IR, ¹H and ¹³C NMR, and EIMS data, consistent with literature values.^{14,16}

Enzymatic Hydrolysis of 1 and 2. Separate solutions of **1** (1 mg) and **2** (0.3 mg) in acetate buffer (pH 6) were treated with β -glucosidase (1 mg/mL) for 24 h at room temperature. Each reaction solution was evaporated to dryness with the resultant residue analyzed by HPLC [column Nova-Pak (Waters, Milford, MA), 3.9 \times 300 mm i.d., C₁₈, 16 μ m, 60 Å; 30% MeCN in H₂O, 0.5 mL/min, *t*_R 28.0 min] and TLC (Si gel, CHCl₃/MeOH, 5:1, *R*_f 0.35) to afford **3**.

Cyclooxygenase-1 and -2 Inhibition Assay. The effect of test compounds on cyclooxygenase-1 and -2 (COX-1 and -2) was determined by measuring PGE₂ production. Reaction mixtures were prepared in 100 mM Tris-HCl buffer, pH 8.0, containing 1 μ M heme, 500 μ M phenol, 300 μ M epinephrine, sufficient amounts of COX-1 or COX-2 to generate 150 ng of PGE₂/mL, and various concentrations of test samples. The reaction was initiated by the addition of arachidonic acid (final concentration, 10 μ M) and incubated for 10 min at room temperature (final volume, 200 μ L). Then, the reaction was terminated by adding 20 μ L of the reaction mixture to 180 μ L of 27.8 μ M indomethacin, and PGE₂ was quantitated by an ELISA method. Samples were diluted to the desired concentration with 100 mM potassium phosphate buffer (pH 7.4) containing 2.34% NaCl, 0.1% bovine serum albumin, 0.01% sodium azide, and 0.9 mM Na₄EDTA. Following transfer to a 96-well plate (Nunc-Immuno Plate Maxisorp, Fisher) coated with a goat anti-mouse IgG (Jackson Immuno Research Laboratories), the tracer (PGE₂-acetylcholinesterase; Cayman Chemical, Ann Arbor, MI) and primary antibody (mouse anti PGE₂; Monsanto, St. Louis, MO) were added. Plates were then incubated at room temperature overnight, reaction mixtures

were removed, and wells were washed with a solution of 10 mM potassium phosphate buffer (pH 7.4) containing 0.01% sodium azide and 0.05% Tween 20. Ellman's reagent (200 μ L) was added to each well, and the plate was incubated at 37 °C for 3–5 h, until the control wells yielded OD = 0.5–1.0 at 412 nm. A standard curve with PGE₂ (Cayman Chemical, Ann Arbor, MI) was generated on the same plate, which was used to quantify the PGE₂ levels produced in the presence of test samples. Results were expressed as a percentage, relative to control (solvent-treated) samples, and dose–response curves were constructed for the determination of IC₅₀ values.

Acknowledgment. This work was supported by Program Project P01-CA-48112 funded by the National Cancer Institute, NIH, Bethesda, MD. P.W.-T. is grateful to Conseil Régional d'Aquitaine, France, for fellowship support. We thank Dr. K. Fagerquist, Mass Spectrometry Facility, Department of Chemistry, University of Minnesota, Minneapolis, MN, for the mass spectral data. We are grateful to the Research Resources Center, University of Illinois at Chicago, for the provision of certain spectroscopic equipment used in this investigation.

References and Notes

- (1) Hart, J. H. *Rev. Phytopathol.* **1981**, *19*, 437–458.
- (2) Sotheeswaran, S.; Pasupathy, V. *Phytochemistry* **1993**, *32*, 1083–1092.
- (3) Mattivi, F.; Reniero, F.; Korhammer, S. *J. Agric. Food Chem.* **1995**, *43*, 1820–1823.
- (4) Goldberg, D. M.; Ng, E.; Karumanchiri, A.; Diamandis, E. P.; Soleas, G. J. *Am. J. Enol. Vitic.* **1996**, *47*, 415–420.
- (5) Jang, M.; Cai, L.; Udeani, G. O.; Slowing, K. V.; Thomas, C. F.; Beecher, C. W. W.; Fong, H. H. S.; Farnsworth, N. R.; Kinghorn, A. D.; Mehta, R. G.; Moon, R. C.; Pezzuto, J. M. *Science* **1997**, *275*, 218–220.
- (6) Waffo-Teguo, P.; Decendit, A.; Vercauteren, J.; Deffieux, G.; Mérillon, J.-M. *Phytochemistry* **1996**, *42*, 1591–1593.
- (7) Waffo-Teguo, P.; Decendit, A.; Krisa, S.; Deffieux, G.; Vercauteren, J.; Mérillon, J.-M. *J. Nat. Prod.* **1996**, *59*, 1189–1191.
- (8) Waffo-Teguo, P.; Fauconneau, B.; Deffieux, G.; Huguet, F.; Vercauteren, J.; Mérillon, J.-M. *J. Nat. Prod.* **1998**, *61*, 655–657.
- (9) Breuil, A. C.; Adrian, M.; Pirio, N.; Meunier, P.; Bessis, R.; Jeandet, P. *Tetrahedron Lett.* **1998**, *39*, 537–540.
- (10) Cichewicz, R. H.; Kouzi, S. A.; Hamann, M. T. *J. Nat. Prod.* **2000**, *63*, 29–33.
- (11) Kinghorn, A. D.; Fong, H. H. S.; Farnsworth, N. R.; Mehta, R. G.; Moon, R. C.; Moriarty, R. M.; Pezzuto, J. M. *Curr. Org. Chem.* **1998**, *2*, 597–612.
- (12) Pezzuto, J. M. *Biochem. Pharmacol.* **1997**, *53*, 121–133.
- (13) Langcake, P.; Pryce, R. J. *Experientia* **1977**, *33*, 151–152.
- (14) Khan, M. A.; Nabi, S. G.; Prakash, S.; Zaman, A. *Phytochemistry* **1986**, *25*, 1945–1948.
- (15) Kim, N.-C.; Desjardins, A. E.; Wu, C. D.; Kinghorn, A. D. *J. Nat. Prod.* **1999**, *62*, 1379–1384.
- (16) Huang, K. S.; Wang, Y. H.; Li, R. L.; Lin, M. J. *J. Nat. Prod.* **2000**, *63*, 86–89.
- (17) Ohyama, M.; Tanaka, T.; Iinuma, M.; Goto, K. *Chem. Pharm. Bull.* **1994**, *42*, 2117–2120.
- (18) Decendit, A.; Mérillon, J.-M. *Plant Cell Rep.* **1996**, *15*, 762–765.

NP000426R