

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/24423594>

ChemInform Abstract: Daphenylline (I), a New Alkaloid with an Unusual Skeleton, from *Daphniphyllum longeracemosum*

ARTICLE *in* ORGANIC LETTERS · JUNE 2009

Impact Factor: 6.36 · DOI: 10.1021/ol9007958 · Source: PubMed

CITATIONS

26

READS

57

10 AUTHORS, INCLUDING:



Chun-Shun Li

Chinese Academy of Sciences

107 PUBLICATIONS 1,264 CITATIONS

SEE PROFILE



Xin Gui Fang

Peking University

84 PUBLICATIONS 1,040 CITATIONS

SEE PROFILE



Hehua Xu

Chinese Academy of Sciences

64 PUBLICATIONS 400 CITATIONS

SEE PROFILE



Hong-Ping He

Chinese Academy of Sciences

177 PUBLICATIONS 1,843 CITATIONS

SEE PROFILE

Daphenylline, a New Alkaloid with an Unusual Skeleton, from *Daphniphyllum longeracemosum*


Qiang Zhang,^{†,‡} Ying-Tong Di,[†] Chun-Shun Li,[†] Xin Fang,[†] Cheng-Jian Tan,[†]
Zhen Zhang,[†] Yu Zhang,[†] Hong-Ping He,[†] Shun-Lin Li,[†] and Xiao-Jiang Hao^{*,†}

State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming
Institute of Botany, Chinese Academy of Sciences, Kunming 650204, Yunnan, PRC,
and Graduate School of Chinese Academy of Sciences, Beijing 100039, PRC

haoxj@mail.kib.ac.cn

Received April 12, 2009

ABSTRACT



A new alkaloid, daphenylline (1), with an unprecedented rearranged 22-nor-calyciphylline skeleton, was isolated from the fruits of *Daphniphyllum longeracemosum*. Its structure and stereochemistry were elucidated on the basis of spectroscopic and computational approaches. A plausible biosynthetic pathway of 1 was also proposed.

Daphniphyllum alkaloids are a family of natural products with complex and diversified structures elaborated by trees of the genus *Daphniphyllum*.¹ In recent years, many new

Daphniphyllum alkaloids have been discovered from this genus, some of which possessed new carbon skeletons.² Those fused-heterocyclic systems have attracted great interest as challenging targets for total synthesis³ as well as biosynthesis.⁴ The previous work carried out by our group on the alkaloids of the genus *Daphniphyllum* led to a series of novel alkaloids with highly complex polycyclic systems.⁵ In a continuing search for structurally unique and biogenetically interesting alkaloids, the chemical constituents in *Daphniphyllum longeracemosum* Rosenth have been further in-

[†] Kunming Institute of Botany.

[‡] Graduate School of Chinese Academy of Sciences.


(1) For recent reviews of daphniphyllum alkaloids: (a) Morita, H.; Kobayashi, J. In *Modern Alkaloids Structure, Isolation, Synthesis, and Biology*; Fattorusso, E., Tagliatella-Scafati, O., Eds.; Wiley-VCH: Weinheim, 2008; pp 541–589. (b) Kobayashi, J.; Morita, H. In *The Alkaloids*; Cordell, G. A., Ed.; Academic Press: New York, 2003; Vol. 60, pp 165–205.

(2) (a) Saito, S.; Yahata, H.; Kubota, T.; Obara, Y.; Nakahata, N.; Kobayashi, J. *Tetrahedron* **2008**, *64*, 1901–1908. (b) Zhang, C.-R.; Yang, S.-P.; Yue, J.-M. *J. Nat. Prod.* **2008**, *71*, 1663–1668. (c) Saito, S.; Kubota, T.; Fukushima, E.; Kawabata, J.; Zhang, H.; Kobayashi, J. *Org. Lett.* **2007**, *9*, 1207–1209. (d) Li, Z.-Y.; Chen, P.; Xu, H.-G.; Yang, Y.-M.; Peng, S.-Y.; Zhao, Z.-Z.; Guo, Y.-W. *Org. Lett.* **2007**, *9*, 477–480. (e) Matsuno, Y.; Okamoto, M.; Hirasawa, Y.; Kawahara, N.; Goda, Y.; Shiro, M.; Morita, H. *J. Nat. Prod.* **2007**, *70*, 1516–1518. (f) Fan, C.-Q.; Yin, S.; Xue, J.-J.; Yue, J.-M. *Tetrahedron* **2007**, *63*, 115–119. (g) Saito, S.; Kubota, T.; Kobayashi, J. *Tetrahedron Lett.* **2007**, *48*, 3809–3812. (h) Saito, S.; Kubota, T.; Kobayashi, J. *Tetrahedron Lett.* **2007**, *48*, 5693–5695. (i) Saito, S.; Kubota, T.; Fukushima, E.; Kawabata, J.; Zhang, H.; Kobayashi, J. *Tetrahedron Lett.* **2007**, *48*, 1587–1589.

(3) (a) Ikeda, S.; Shibuya, M.; Kanoh, N.; Iwabuchi, Y. *Org. Lett.* **2009**, *11*, 1833–1836. (b) Cordero-Vargas, A.; Bradshaw, B.; Bonjoch, J. *Tetrahedron* **2008**, *64*, 8134–8140. (c) Denmark, S. E.; Baiazitov, R. Y. *J. Org. Chem.* **2006**, *71*, 593–605. (d) Wallace, G. A.; Heathcock, C. H. *J. Org. Chem.* **2001**, *66*, 450–454. (e) Heathcock, C. H. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 14323–14327. (f) Heathcock, C. H.; Joe, D. *J. Org. Chem.* **1995**, *60*, 1131–1142. (g) Heathcock, C. H.; Kath, J. C.; Ruggeri, R. B. *J. Org. Chem.* **1995**, *60*, 1120–1130. (h) Heathcock, C. H. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 665–681, and references therein.

(4) (a) Niwa, H.; Hirata, Y.; Suzuki, K. T.; Yamamura, S. *Tetrahedron Lett.* **1973**, *14*, 2129–2132. (b) Suzuki, K. T.; Okuda, S.; Niwa, H.; Toda, M.; Hirata, Y.; Yamamura, S. *Tetrahedron Lett.* **1973**, *14*, 799–802.

vestigated.⁶ A novel alkaloid, named daphenylline (**1**), was isolated from the fruits and possessed an expanded neo-hexamatomic ring with C-13 connected to C-1 instead of C-8 as usual. In this paper, the isolation and structure elucidation of **1** are described.



The fruits (60 kg) of *D. longracemosum* were extracted with 95% EtOH, and the crude extract was adjusted to pH 2 with 2% HCl. After extraction with petroleum ether and chloroform, the aqueous layer was then basified to pH 10 with 3% NaOH, followed by exhaustive extraction with chloroform and *n*-BuOH. The *n*-BuOH fraction was separated on silica gel eluted by acetone, methanol, and methanol/diethylamine (20:1) separately to afford three parts, the third of which was further chromatographed over a series of ion-exchange resin, C₁₈ silica gel, and silica gel to yield daphenylline (**1**, 44 mg, 0.00073%) and a known calyciphylline A-type alkaloid daphnilongeranin C (**2**, 1.5 g, 0.0025%).^{6c}

Daphenylline (**1**),⁷ a colorless oil, showed the pseudomolecular ion peak at m/z 294 $[M + H]^+$ in the FABMS and ESIMS, and the molecular formula of **1**, C₂₁H₂₇N, was established by HRESIMS $[m/z$ 294.2219, $(M + H)^+$, calcd. 294.2221] requiring nine degrees of unsaturation. ¹³C NMR and DEPT revealed 21 carbon signals due to four sp² quaternary carbons, two sp² methines, one sp³ quaternary carbon, five sp³

Table 1. ¹H, ¹³C, and DEPT NMR Data of Daphenylline (**1**) in CDCl₃

	δ_H (mult. Hz)	δ_C
1		143.8 (s)
2	2.75 (1H, m)	36.5 (d)
3 α	2.37 (1H, m)	17.8 (t)
3 β	1.95 (1H, m)	
4	3.73 (1H, m)	65.4 (d)
5		45.4 (s)
6	2.45 (1H, m)	47.2 (d)
7 α	3.74 (1H, m)	58.0 (t)
7 β	2.46 (1H, m)	
8		132.7 (s)
9		144.6 (s)
10	3.43 (1H, m)	42.9 (d)
11 α	2.04 (1H, m)	28.3 (t)
11 β	1.57 (1H, m)	
12 α	1.86 (1H, m)	27.7 (t)
12 β	1.30 (1H, m)	
13	6.87 (1H, d, 7.6 Hz)	127.0 (d)
14	7.05 (1H, d, 7.6 Hz)	124.1 (d)
15		137.4 (s)
16 α	2.76 (1H, m)	31.0 (t)
16 β	2.68 (1H, m)	
17 α	1.60 (1H, m)	36.0 (t)
17 β	2.33 (1H, m)	
18	1.87 (1H, m)	32.8 (d)
19 α	2.71 (1H, m)	50.0 (t)
19 β	2.97 (1H, dd, 13.2, 6.0)	
20	1.26 (1H, d, 7.2)	18.0 (q)
21	1.45 (1H, s)	26.0 (q)

methines, seven sp³ methylenes, and two methyl groups. Among them, two methylenes (δ_C 58.0 and δ_C 50.0) and one methine (δ_C 65.4) were suggested to be attached to a nitrogen atom. Furthermore, the six sp² carbon signals mentioned above, the ¹H NMR doublet peaks at δ_H 7.05 and 6.87 with coupling constant 7.6 Hz, and the characteristic IR absorptions at 1639 and 1563 cm⁻¹ implied the presence of a 1,2,3,4-tetrasubstituted benzene ring moiety in the structure. Apart from three degrees of unsaturation occupied by three double bonds of the benzene ring, the remaining six degrees of unsaturation indicated that **1** should possess a hexacyclic system.

The ¹H–¹H COSY revealed that **1** possessed three fragments, **a** (C-2 to C-4, C-2 to C-18, and C-18 to C-19 and C-20), **b** (C-6 to C-7 and C-12, C-10 to C-12 and C-17, and C-16 to C-17), and **c** (C-13 to C-14), as shown in Figure 1. Further detailed HMBC studies established the connections among the three fragments, the quaternary carbons, and a nitrogen atom. HMBC correlations of H-4 to C-19 (δ_C 50.1), H-19 to C-7 (δ_C 58.1), and H-7 to C-4 (δ_C 65.5) confirmed that C-4, C-7, and C-19 were linked at the nitrogen atom. Connectivity of C-21 to C-4, C-6, and C-8 through C-5 was indicated by HMBC correlations of H-4 to C-8 and C-21, H-21 to C-5, C-6, and C-8. The presence of a tetrasubstituted benzene ring, composed of C-1, C-8, C-9, C-13, C-14, and C-15, was suggested by HMBC cross-peaks for H-13 to C-8, C-9 (J^4) and C-15, and H-14 to C-1. Furthermore, HMBC correlations of H-17 to C-9 and H-16 to C-14 indicated the connectivity of fragment **b** and the benzene ring through bonds C-15/C-16 and C-9/C-10. Meanwhile, the fragment **a** and the benzene ring were connected through bond C-1/C-2, which was supported by the HMBC correlations of H-3 to C-1. Thus, the gross structure of daphenylline was assigned as **1** with an unusual fused-

(5) (a) Tan, C.-J.; Di, Y.-T.; Wang, Y.-H.; Wang, Y.; Mu, S.-Z.; Gao, S.; Zhang, Y.; Kong, N.-C.; He, H.-P.; Zhang, J.-X.; Fang, X.; Li, C.-S.; Lu, Y.; Hao, X.-J. *Tetrahedron Lett.* **2008**, *49*, 3376–3379. (b) Di, Y.-T.; He, H.-P.; Wang, Y.-S.; Li, L.-B.; Lu, Y.; Gong, J.-B.; Fang, X.; Kong, N.-C.; Li, S.-L.; Zhu, H.-J.; Hao, X.-J. *Org. Lett.* **2007**, *9*, 1355–1358. (c) Li, C.-S.; Di, Y.-T.; He, H.-P.; Gao, S.; Wang, Y.-H.; Lu, Y.; Zhong, J.-L.; Hao, X.-J. *Org. Lett.* **2007**, *9*, 2509–2512. (d) Li, C.-S.; He, H.-P.; Di, Y.-T.; Wang, Y.-H.; Mu, S.-Z.; Li, S.-L.; Gao, S.; Gao, Z.-L.; Hao, X.-J. *Tetrahedron Lett.* **2007**, *48*, 2737–2740. (e) Zhang, Y.; He, H.-P.; Di, Y.-T.; Mu, S.-Z.; Wang, Y.-H.; Wang, J.-S.; Li, C.-S.; Kong, N.-C.; Gao, S.; Hao, X.-J. *Tetrahedron Lett.* **2007**, *48*, 9104–9107. (f) Di, Y.-T.; He, H.-P.; Liu, H.-Y.; Du, Z.-Z.; Tian, J.-M.; Yang, X.-W.; Wang, Y.-H.; Hao, X.-J. *Tetrahedron Lett.* **2006**, *47*, 5329–5331. (g) Li, L.; He, H.-P.; Di, Y.-T.; Gao, S.; Hao, X.-J. *Tetrahedron Lett.* **2006**, *47*, 6259–6262.

(6) Previous phytochemical research on *D. longracemosum*: (a) Li, C.-S.; Di, Y.-T.; Mu, S.-Z.; He, H.-P.; Zhang, Q.; Fang, X.; Zhang, Y.; Li, S.-L.; Lu, Y.; Gong, Y.-Q.; Hao, X.-J. *J. Nat. Prod.* **2008**, *71*, 1202–1206. (b) Li, L.; He, H.-P.; Di, Y.-T.; Tian, J.-M.; Hao, X.-J. *Helv. Chim. Acta* **2007**, *89*, 1457–1462. (c) Yang, S.-P.; Zhang, H.; Zhang, C.-R.; Cheng, H.-D.; Yue, J.-M. *J. Nat. Prod.* **2006**, *69*, 79–82. (d) Yang, S.-P.; Yue, J.-M. *Helv. Chim. Acta* **2006**, *89*, 2783–2788. (e) Di, Y.-T.; He, H.-P.; Lu, Y.; Yi, P.; Li, L.; Wu, L.; Hao, X.-J. *J. Nat. Prod.* **2006**, *69*, 1074–1076, and 5c, 5d, 5g.

(7) **Daphenylline**: colorless oiliness; $[\alpha]_D^{16} = -45.6$ (c 0.19, MeOH); UV (MeOH) λ_{max} nm 204.4; IR (KBr) ν_{max} : 2922, 2848, 2388, 2283, 1639, 1563, 1458, 1412, 1383, 1292, 1035, 846 and 799 cm⁻¹; ¹H and ¹³C NMR data (Table 1); FABMS and ESIMS m/z 294 $[M + H]^+$; HRESIMS m/z 294.2219 (calcd for $[M + H]^+$ 294.2221).




Figure 1. Selected two-dimensional NMR correlations of daphenylline (**1**).

hexacyclic ring system (two five-, three six-, and one seven-membered rings).


The relative stereochemistry of **1** was fixed by ROESY as shown in Figure 1. The ROESY interaction pairs of H₃-21/H-4, H₃-21/H-6, and H₃-21/H-10 indicated their β -oriented configuration. Interaction pairs of H-2/H₃-20, H₃-20/H α -3, and H β -3/H₃-21 suggested both H-2 and H₃C-20 were β -oriental.

To assign the absolute configuration, the optical rotation (OR) values of **1** and its enantiomer were calculated by the density functional theory (DFT) methods⁸ in the Gaussian 03 program package.⁹ The minimum geometries of **1** and its enantiomer were optimized by B3LYP/6-311G(d, p). The OR values were calculated by B3LYP/6-311G+(d, p) under the Self-Consistent Reaction Field model of solvation: for **1** it was -64.3 , and for its enantiomer was $+64.3$. The former was close to the experimental value of -45.6 , which indicated the absolute configuration of daphenylline as given in structure **1**. This conclusion was also in accord with the previously hypothetical

stereochemistry of the calyciphylline A-type alkaloids based on their biogenetic pathways.¹⁰

A plausible biogenetic pathway for **1** was proposed as shown in Scheme 1. The biogenetic origin of **1** should be calyciphylline

Scheme 1. Plausible Biosynthetic Pathway of **1**



A-type alkaloids such as **2** which was also acquired from the title plant in quantity. Alkaloid **2** might be reduced and dehydrated to form a ring expanded intermediate **4** via the Wagner–Meerwein rearrangement. Then the intermediate **4** could be involved in the elimination of the carboxyl group at C-21 by decarboxylase to generate **5** or **6** with a double bond between C-13 and C-14 or between C-14 and C-15, followed by one or two steps of syn-[1,3] sigmatropic rearrangement to yield **1**.

Daphenylline (**1**) was evaluated for cytotoxic activities against the human tumor cell lines (HL-60, SMMC-7721, A-549, SK-BR-3). The results indicated that **1** was inactive against the above cancer cells ($IC_{50} > 40 \mu M$).¹¹

Acknowledgment. The authors thank Prof. Xun Gong, Kunming Institute of Botany, Chinese Academy of Sciences (CAS), for the identification and collection of the plant material from Hekou area of Yunnan Province PRC in October 2005. A specimen was deposited at the State Key Laboratory of Phytochemistry and Plant Resources in West China, CAS. This work was financially supported by grants from the Ministry of Science and Technology (2009CB522303), National Science Foundation (1130830114 and 3930830114), and West Doctor Foundation of Chinese Academy of Sciences (2908065112w1), PRC.

Supporting Information Available: Experimental section, optical rotation calculation, one- and two-dimensional NMR spectra, and mass spectrum for daphenylline **1**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL9007958

(10) Morita, H.; Kobayashi, J. *Org. Lett.* **2003**, *5*, 2895–2898.

(11) DDP was used as the positive control compound, and the IC_{50} values against the cell lines (HL-60, SMMC-7721, A-549, and SK-BR-3) were 3.99, 11.65, 25.67, and 13.70 $\mu g/mL$, respectively.

(8) (a) Zhang, H. W.; Huang, W. Y.; Chen, J. R.; Yan, W. Z.; Xie, D. Q.; Tan, R. X. *Chem.–Eur. J.* **2008**, *14*, 10670–10674. (b) Ge, H. M.; Zhu, C. H.; Shi, D. H.; Zhang, L. D.; Xie, D. Q.; Yang, J.; Ng, S. W.; Tan, R. X. *Chem.–Eur. J.* **2008**, *14*, 376–381. (c) Ruud, K.; Zanasi, R. *Angew. Chem., Int. Ed.* **2005**, *44*, 3594–3596. (d) Stephens, P. J.; Devlin, F. J.; Cheeseman, J. R.; Frisch, M. J.; Rosini, C. *Org. Lett.* **2002**, *4*, 4595–4598. (g) Stephens, P. J.; Devlin, F. J.; Cheeseman, J. R.; Frisch, M. J. *J. Phys. Chem. A* **2001**, *105*, 5356–5371. (h) Cheeseman, J. R.; Frisch, M. J.; Delvin, G. J.; Stephens, P. J. *J. Phys. Chem. A* **2000**, *104*, 1039–1046.

(9) Frisch, M. J. *Gaussian 03*, revision D.01; Gaussian, Inc.: Wallingford, CT, 2005 (see Supporting Information for complete reference).