

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8491569>

Chamuvarinin, an Acetogenin Bearing a Tetrahydropyran Ring from the Roots of *Uvaria chamae* 1

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · JULY 2004

Impact Factor: 3.8 · DOI: 10.1021/np030521a · Source: PubMed

CITATIONS

16

READS

64

5 AUTHORS, INCLUDING:



Djibril Fall

Cheikh Anta Diop University, Dakar

22 PUBLICATIONS 141 CITATIONS

SEE PROFILE



Romain Duval

Institute of Research for Development

15 PUBLICATIONS 184 CITATIONS

SEE PROFILE

Chamuvarinin, an Acetogenin Bearing a Tetrahydropyran Ring from the Roots of *Uvaria chamae*¹

Djibril Fall, Romain A. Duval, Christophe Gleye, Alain Laurens,* and Reynald Hocquemiller

Laboratoire de Pharmacognosie, UMR 8076 CNRS (BioCIS), Faculté de Pharmacie, Université Paris XI, 92296 Châtenay-Malabry Cedex, France

Received December 4, 2003

A new cytotoxic acetogenin, chamuvarinin (**1**), containing a tetrahydropyran ring with an adjacent bis-tetrahydrofuran ring, which corresponds to a novel carbon skeleton in this series, was isolated from the roots of *Uvaria chamae*, together with the previously reported acetogenins squamocin (**2**), desacetylavaricin (**3**), and neoannonin (**4**). The structure determination of chamuvarinin (**1**) was based on extensive NMR studies and high-resolution mass spectral measurements. This new compound shows significant cytotoxicity toward the KB 3-1 cell line ($IC_{50} = 8 \times 10^{-10}$ M). In addition, a biosynthetic relationship between **1** and **2** is briefly discussed.

Annonaceous acetogenins are a relatively new class of natural polyketides that have promising anticancer, anti-parasitic, and pesticidal properties.² Structurally, most of these long-chain fatty acid derivatives may be classified into three major groups: mono-tetrahydrofuran (THF), adjacent bis-THF, and nonadjacent bis-THF classes. Among the 400 or so annonaceous acetogenins previously reported, only seven possessed a tetrahydropyran (THP) ring. The THP ring of these molecules is linked in a nonadjacent manner to the mono-THF ring. These compounds have been recently isolated from *Rollinia mucosa*,^{3–5} *Goniothalamus giganteus*,⁶ and *Annona montana*.⁷


A cyclohexane extract of the roots of *Uvaria chamae* P. Beauv. (Annonaceae) was cytotoxic in vitro against human tumor cell lines in our preliminary screening tests. From this plant, we previously isolated mono-THF annonaceous acetogenins, namely, *cis*-bullatencin and seven known acetogenins (bullatencin, annotemoyin-1, uvariamicin-I, -II, and -III, *cis*-reticulatacin, and *cis*-uvariamicin-I).⁸ A close study of the cyclohexane extract of *U. chamae* led to the isolation of chamuvarinin (**1**), which is the first acetogenin with a THP ring adjacent to a bis-THF ring, in addition to three known compounds (squamocin (**2**), desacetylavaricin (**3**), and neoannonin (**4**)). The cytotoxic activity of this compound was evaluated on the KB 3-1 cell line.

The cyclohexane extract of *U. chamae* roots was subjected to repeated column chromatography and preparative HPLC to yield four compounds; **1** was found to be new, and it was given the trivial name chamuvarinin (Figure 1).

Compound **1** has a molecular mass 604, determined by HRESIMS $[M + Na]^+$ at m/z 627.460160 (calc 627.460059), corresponding to the molecular formula $C_{37}H_{64}O_6Na$.

The presence of an α,β -unsaturated γ -methyl- γ -lactone moiety, the common feature of acetogenins of type 1a, was suggested by a strong IR absorption peak (carbonyl) at 1756 cm^{-1} and a weak UV λ_{max} at 219.1 nm. The ¹H NMR spectrum indicated seven protons at δ 6.98 (CH-35), 4.98 (CH-36), 2.26 (CH₂-3), and 1.42 (CH₃-37), and the ¹³C NMR spectrum showed six carbon at δ 173.8 (C-1), 148.8 (C-35), 134.3 (C-2), 77.4 (C-36), 25.1 (C-3), and 19.2 (C-37), corresponding to the spectroscopic features of the lactone ring (Table 1).

The ¹H NMR spectrum of compound **1** exhibited six well-defined signals in the range δ_H 3.0–4.0. Those at δ 3.37



(*) Absolute configurations may be inverted

Figure 1. Chamuvarinin.

Table 1. NMR Spectroscopic Data of Chamuvarinin (**1**) in $CDCl_3$ ^a

position	¹ H	¹³ C
1		173.8
2		134.3
3	2.26 t (7.4)	25.1
4	1.55 m	26.9
5–13	1.25–1.29	25.5–32.4
14	1.38 m	25.7
15	3.37 m	74.1
16	3.82 m	83.0
17a, 18a	1.65 m	25.5–32.4
17b, 18b	1.97 m	25.5–32.4
19	3.93 m	81.4*
20	3.93 m	81.9*
21a, 22a	1.71 m	25.5–32.4
21b, 22b	1.92 m	25.5–32.4
23	3.88 m	82.0
24	3.28 m	79.9
25a	1.26 m	25.5–32.4
25b	1.48 m	25.5–32.4
26	1.82 m	23.4
27a	1.12 m	25.5–32.4
27b	1.33 m	25.5–32.4
28	3.23 m	77.8
29	1.52 m	25.5–32.4
30, 31	1.25–1.29	25.5–32.4
32	1.25 m	31.8
33	1.25 m	22.6
34	0.87 t (7.3)	14.1
35	6.98 d (1.5)	148.8
36	4.98 dq (6.7; 1.5)	77.4
37	1.42 d (6.7)	19.2

^a Chemical shifts (δ) are in ppm relative to TMS, observed splittings J are in Hz, superscript * corresponds to interchangeable attributions.

(H-15), 3.82 (H-16), 3.88 (H-23), and 3.93 (H-19, H-20) were assigned to a bis-THF ring with a flanking hydroxyl group.² The remaining signals at δ 3.23 (H-28) and 3.28 (H-24) were consistent with the presence of a THP ring in the molecule.^{3–7} The HOHAHA correlation found between

* To whom correspondence should be addressed. Phone: 01 46835639. Fax: 01 46835399. E-mail: alain.laurens@cep.u-psud.fr.




Figure 2. Structural subunit, assigned on the basis of ^1H - ^1H HOHAHA NMR.




Figure 3. EIMS fragmentations of chamuvarinin (**1**).

H-28/H-27_a, H-28/H-27_b, H-28/H-26, H-28/H-25_a, H-28/H-25_b and H-24/H-25_a, H-24/H-25_b, H-24/H-26, H-24/H-27_a, H-24/H-27_b, H-24/H-23 confirmed the presence of a THP moiety adjacent to the bis-THF ring (Figure 2).

The disposition of the adjacent THF-THP unit on the aliphatic chain was determined by the analysis of the fragmentation pattern displayed by **1** in EIMS (Figure 3). Thus, the fragment ion peaks at m/z 291, 295, 309 (cleavage at C-15/C-16), m/z 221, 239, 347 (cleavage at C-19/C-20), and m/z 169, 399 (cleavage at C-23/C-24) permitted location of the THF ring with a flanking hydroxyl between C-15 and C-23 and of the THP ring between C-24 and C-28.

The relative configuration at C-15/C-16 was assigned according to Born's rule.¹² Thus, the chemical shift values of both C-15 (δ 74.1) and H-15 (δ 3.37) indicated a *threo* relationship. On the other hand, the relative *trans* configuration of the chiral carbon centers of the first bis-THF moiety is demonstrated by a relatively large δ difference between the C-17 or C-18 methylene protons.¹³ The absolute configuration of the C-36 stereocenter was established as *S* by Latypov's method using Pirkle reagent.¹

Compounds **2**–**4** showed the characteristic NMR signals of bis-THF α,α' -dihydroxy acetogenins² and were identified by comparison of the physicochemical data of the previously isolated acetogenins squamocin⁹ (**2**), desacetylavaricin¹⁰ (**3**), and neoannonin¹¹ (**4**, a C₃₅ acetogenin with the configuration of desacetylavaricin).

Chamuvarinin could biogenetically be derived from squamocin via desacetylavaricin (**3**). The triepoxidation of a triene such as $\Delta^{15,19,23}$ -chatenaytrienin-4, previously isolated from roots of *Annona nutans*,¹⁴ leads to triepoxyrollin, isolated from the seeds of *Rollinia membranacea*.¹⁵ This triepoxide in turn converts to the bis-THF acetogenin desacetylavaricin (**3**), which is presumably C-28 hydroxylated to squamocin (**2**). Therefore, a regioselective intramolecular nucleophilic substitution may lead to the THP ring of chamuvarinin (**1**).

Cytotoxic activity of chamuvarinin (**1**) was evaluated relative to squamocin (**2**) on KB 3-1 cells. With an IC_{50} of 8×10^{-10} M, chamuvarinin (**1**) appears to be less cytotoxic than squamocin (**2**) ($\text{IC}_{50} = 3 \times 10^{-12}$ M).

These results may rely on an insufficient amphiphilicity of chamuvarinin (**1**), in accordance with numerous examples.^{15–17} On the other hand, it has to be noted that the C-24/C-28 absolute configurations of the THF nucleus




Figure 4. Biogenetic pathway proposed for chamuvarinin (**1**). (*) Absolute configurations may be inverted.

of chamuvarinin remain unknown and that strong differences of cytotoxicities are frequently observed for diastereoisomeric acetogenins.^{18–23}

Experimental Section

General Experimental Procedures. UV spectra were determined in MeOH on a Philips PU 8720 spectrophotometer. IR spectra were recorded on a Bruker Vector 22 spectrophotometer. The ^1H NMR spectra were obtained with a Bruker AC-400 (at 400 MHz) and AC-200 (at 200 MHz). The ^{13}C NMR spectra were obtained with a Bruker AC-200 at 50 MHz. EIMS (70 eV) were recorded with an Automass multi spectrometer R10-10C, and HRESIMS were registered with a navigator spectrometer (ThermoFinnigan, France). HPLC was performed with a pump (Waters 590), UV detector (Waters 84), and injector (Waters SSV).

Plant Material. Roots of *Uvaria chamae* P. Beauv. were collected in Casamance (Senegal) in August 1999. A voucher specimen (DF126) has been deposited at the Faculty of Medicine and Pharmacy of Dakar.

Extraction and Isolation. The dried and powdered roots (1.7 kg) were extracted by percolation (10 L of cyclohexane) during 48 h and evaporated to give a brown extract (66 g). Thirty grams of this extract was subjected to silica gel column chromatography (silica gel Merck 60 70–230 mesh) eluted with *n*-hexane/EtOAc in a gradient from 90:10 to 10:90. The eluate was collected in 16 fractions controlled by TLC (silica gel Merck 60 F 254). The solvent of the fraction 7 was evaporated under reduced pressure. The resulting residue (2.1 g) was subjected to silica gel column chromatography (silica gel Merck 60 H 230–400 mesh) eluted with *n*-hexane/ CH_2Cl_2 /EtOH, 70:30:2 v/v/v. The fraction corresponding to **1** was chromatographed on the same stationary phase, eluted with *n*-hexane/ CH_2Cl_2 /EtOH, 70:30:1 v/v/v. A final purification by preparative HPLC using a reversed-phase Waters μ Bondapak C₁₈ column (10 μm [250 \times 20 mm] cartridge column, flow rate 9 mL/min, 20 mg/injection, and eluant $\text{CH}_3\text{OH}/\text{H}_2\text{O}/\text{THF}$, 90:10:5 v/v/v) led to 14 mg of chamuvarinin (**1**) ($t_R = 22.4$ min), 73 mg of squamocin (**2**) ($t_R = 38.0$ min), 40 mg of desacetylavaricin (**3**) ($t_R = 31.2$ min), and 12 mg of neoannonin (**4**) ($t_R = 21.6$ min).

Chamuvarinin (1): oil (14 mg); $[\alpha]_D^{25} +27.0^\circ$ (c 0.026, CHCl_3); UV (MeOH) λ_{max} (log ϵ) 219.1 (3.46) nm; IR ν_{max} 3474, 2924, 2854, 1756, 1464, 1373, 1318, 1199, 1074, 1028 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz) and ^{13}C NMR (CDCl_3 , 50 MHz), see Table 1; HRESIMS m/z 627.460160 (calc 627.460059 for $\text{C}_{37}\text{H}_{64}\text{O}_6\text{Na}$); EIMS, see Figure 3.

Determination of the Absolute C-36 Configuration. About 1 mg of acetogenin is dissolved in 1 mL of CDCl_3 and divided exactly in two tubes containing separately 15 equiv of the *R* and *S* alcohol of the Pirkle reagent. The ^1H NMR (400 MHz) is performed at low temperature (213–223 K). The difference of the resonance of H-36 ($\delta_R - \delta_S$) between the two complexes formed gives the absolute configuration of C-36. We assign the *S* absolute configuration when $(\delta_R - \delta_S) > 0$ and *R* when $(\delta_R - \delta_S) < 0$. With δ_R 5.021 and δ_S 4.955, the stereo-center was assigned as *S*.

Cytotoxic Assay. Cytotoxicities were colorimetrically evaluated through a 96-well plate assay after 72 h cell exposure to the acetogenins on the KB 3-1 cell line by the method of Fleury et al.²⁴

Acknowledgment. This research was sponsored by the “Direction de la Recherche et des Études Doctorales (DRED)”, through a biennial contract with the “Réseau de Recherche Pharmacochimie” and EGIDE (Centre Français pour l'accueil et les échanges internationaux). The authors are grateful to J. Mahuteau and J. C. Jullian for NMR measurements and O. Laprévotte (ICSN) for MS.

References and Notes

- (1) Part 91 in the series Acetogenins of Annonaceae. For part 90 see: Latypov, Sh.; Jullian, J. C.; Franck, X.; Hocquemiller, R.; Figadère, B. *Chem. Eur. J.* **2002**, *8*, 5662–5666.
- (2) Cavé, A.; Figadère, B.; Laurens, A.; Cortes, D. In *Progress in the Chemistry of Organic Natural Products*; Herz, W., Kirby, G. W., Moore, R. E., Steglich, W., Tamm, Ch., Eds.; Springer-Verlag: Vienna, 1997; Vol. 70, pp 81–288.
- (3) Shi, G.; Alfonso, D.; Fatope, M. O.; Zeng, L.; Gu, Z.-M.; Zhao, G.-X.; He, K.; MacDougall, J. M.; McLaughlin, J. L. *J. Am. Chem. Soc.* **1995**, *117*, 10409–10410.
- (4) Shi, G.; Kozłowski, J. F.; Schwedler, J. T.; Wood, K. V.; MacDougall, J. M.; McLaughlin, J. L. *J. Org. Chem.* **1996**, *61*, 7988–7989.
- (5) Chavez, D.; Acevedo, L. A.; Mata, R. *J. Nat. Prod.* **1998**, *61*, 419–421.
- (6) Alali, F. Q.; Rogers, L.; Zhang, Y.; McLaughlin, J. L. *Tetrahedron* **1998**, *54*, 5833–5844.
- (7) Wang, L.-Q.; Nakamura, N.; Meselhy, M. R.; Hattori, M.; Zhao, W.-M.; Cheng, K.-F.; Yang, R.-Z.; Qin, G.-W. *Chem. Pharm. Bull.* **2000**, *48*, 1109–1113.
- (8) Fall, D.; Gleye, C.; Franck, X.; Laurens, A.; Hocquemiller, R. *Nat. Prod. Lett.* **2002**, *16*, 315–321.
- (9) Fujimoto, Y.; Eguchi, T.; Kakinuma, K.; Ikekawa, N.; Sahai, M.; Gupta, Y. K. *Chem. Pharm. Bull.* **1988**, *36*, 4802–4806.
- (10) Jolad, S. D.; Hoffmann, J. J.; Cole, J. R.; Barry, C. E.; Bates, R. B.; Linz, G. S.; König, W. A. *J. Nat. Prod.* **1985**, *48*, 644–645.
- (11) Kawazu, K.; Alcantara, J. P.; Kobayashi, A. *Agric. Biol. Chem.* **1989**, *53*, 2719–2722.
- (12) Born, L.; Lieb, F.; Lorentzen, J. P.; Moeschler, H.; Nonfon, M.; Söllner, R.; Wendisch, D. *Planta Med.* **1990**, *56*, 312–316.
- (13) Fujimoto, Y.; Murasaki, C.; Shimada, H.; Nishioka, S.; Kakinuma, K.; Singh, S.; Singh, M.; Gupta, Y. K.; Sahai, M. *Chem. Pharm. Bull.* **1994**, *42*, 1175–1184.
- (14) Gleye, C.; Raynaud, S.; Hocquemiller, R.; Laurens, A.; Fourneau, C.; Serani, L.; Laprévotte, O.; Roblot, F.; Lebœuf, M.; Fournet, A.; Rojas de Arias, A.; Figadère, B.; Cavé, A. *Phytochemistry* **1998**, *47*, 749–754.
- (15) Sahpaz, S.; Figadère, B.; Saez, J.; Hocquemiller, R.; Cavé, A.; Cortes, D. *Nat. Prod. Lett.* **1993**, *2*, 301–308.
- (16) Rupprecht, J. K.; Hui, Y.-H.; McLaughlin, J. L. *J. Nat. Prod.* **1990**, *53*, 237–278.
- (17) Landolt, J. L.; Ahammadsahib, K. I.; Hollingworth, R. M.; Barr, R.; Crane, F. L.; Buerck, N. L.; McGrabe, G. P.; McLaughlin, J. L. *Chem. Biol. Interact.* **1995**, *98*, 1–13.
- (18) Li, X.-H.; Hui, Y.-H.; Rupprecht, J. K.; Liu, Y.-M.; Wood, K. V.; Smith, D. L.; Chang, C.-J.; McLaughlin, J. L. *J. Nat. Prod.* **1990**, *53*, 81–86.
- (19) Fang, X.-P.; Rupprecht, J. K.; Alkofahi, A.; Hui, Y.-H.; Liu, Y.-M.; Smith, D. L.; Wood, K. V.; McLaughlin, J. L. *Heterocycles* **1991**, *32*, 11–17.
- (20) Zhao, G.; Hui, Y.; Rupprecht, J. K.; McLaughlin, J. L.; Wood, K. V. *J. Nat. Prod.* **1992**, *55*, 347–356.
- (21) Rieser, M. J.; Fang, X.-P.; Anderson, J. E.; Miesbauer, L. R.; Smith, D. L.; McLaughlin, J. L. *Helv. Chim. Acta* **1993**, *76*, 2433–2444.
- (22) Gu, Z.-M.; Zhao, G.-X.; Oberlies, N. H.; Zeng, L.; McLaughlin, J. L. In *Recent Advances in Phytochemistry*; Arnason, J. T., Mata, R., Romeo, J. T., Eds.; Plenum Press: New York, 1995; Vol. 29, pp 249–310.
- (23) Shi, G.; Zeng, L.; Gu, Z.-M.; MacDougall, J. M.; McLaughlin, J. L. *Heterocycles* **1995**, *41*, 1785–1796.
- (24) Fleury, C.; Cotte, J.; Quéro, A.-M. *Path. Biol.* **1984**, *32*, 628–630.

NP030521A