

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/248738429>

14,15- Dihydroxygermacranolides and Other Constituents of Mikania minima

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · MAY 1990

Impact Factor: 3.8 · DOI: 10.1021/np50069a024

CITATIONS

5

READS

3

3 AUTHORS, INCLUDING:



Cesar Catalan

National University of Tucum...

215 PUBLICATIONS 2,475

CITATIONS

SEE PROFILE

14,15-DIHYDROXYGERMACRANOLIDES AND OTHER CONSTITUENTS
OF *MIKANIA MINIMA*

MARÍA DEL R. CUENCA, CÉSAR A. N. CATALÁN,*

Instituto de Química Orgánica, Facultad de Bioquímica, Química y Farmacia, Universidad Nacional de Tucumán,
Ayacucho 491, 4000 S.M. de Tucumán, Argentina

and W. C. M. C. KOKKE

Department of Synthetic Chemistry, SmithKline Beecham Pharmaceuticals, King of Prussia, Pennsylvania 19406

ABSTRACT.—The aerial parts of *Mikania minima* yielded a large amount of 14-acetoxyartemisiifolin-6 α -O-acetate [**1**], a known lactone, along with two closely related new analogues, 14-hydroxyartemisiifolin-6 α -O-acetate [**3**] and 11 β H-11,13-dihydro-14-hydroxyartemisiifolin-6 α -O-acetate [**5**]. Several common triterpenes and sterols were also identified. The structures were determined by spectroscopic methods and chemical transformations.

Mikania minima (Bak.) Robinson (Compositae, tribe Eupatorieae) is a rare vine found only in Tucumán province, Argentina. With its small flower heads, hexagonal branches, and typical location of the bractlet, *M. minima* is one of the best marked species of the perplexing *Mikania scandens* complex (1). Sesquiterpene dilactones of the miscandenin and mikanolide group are frequently found in the genus (2–7), although other sesquiterpene lactone types (2,3,6–10), *ent*-kaurene and pimarene diterpenes (3,11,12), and geranylnerol derivatives (2–4) are also relatively common. Recently, bisabolone derivatives were isolated from a newly described *Mikania* species (13). Continuing our work on this genus (5) we report here the chemical composition of *M. minima*.

The main constituent of *M. minima* was a crystalline lactone, mp 110°, identified as 14-acetoxyartemisiifolin-6 α -O-acetate [**1**]. This lactone has previously been isolated as a gum from two subspecies of *Dicoma anomala* (14). Even though the ¹H-nmr spectrum of **1** was reported to give broad signals at room temperature (14), our crystalline sample exhibited quite well-defined signals in CDCl₃, which were assigned by spin decoupling and COSY experiments. The signals at higher fields were better resolved in C₆D₆ solution (Table 1). The absence of nOe's between H-1 or H-5 and any of the two proton AB systems of H-14 and H-15 confirmed that the stereochemistry of the 1,10 and 4,5 double bonds was *Z* in both cases. Oxidation of **1** with MnO₂ afforded **4**, the ¹H-nmr spectrum of which, in accordance with Herz's rule (15,16), showed the aldehyde proton at 9.98 ppm as a doublet allylically coupled to H-5 at 6.00 ppm (Table 1). The acetate **2** prepared from **1** showed the expected downfield shift for the H-15 protons (Table 1). A




TABLE 1. ^1H -nmr Data of Compounds 1-4.^a

Proton	Compound			
	1 ^{b,c} C ₆ D ₆	2 ^c CDCl ₃	3 ^c CDCl ₃	4 ^d CDCl ₃
H-1	4.71 br dd	5.26 br dd	5.34 br dd	5.26 br dd
H-2 α	2.14 dddd	2.3-2.45 m	2.1-2.6 m	1.9-3.1 m
H-2 β	1.88 m			
H-3 α	1.70 ddd			
H-3 β	2.37 ddd	2.15 m		
H-5	4.43 d	2.56 ddd		
H-6	4.43 d	5.04 d	4.87 br d	6.00 dd
H-7	4.86 dd	4.98 dd	5.14 dd	5.61 dd
H-8	2.55 dddd	3.06 dddd	3.07 dddd	3.15 dddd
H-9	4.98 br dd	5.06 br dd	5.10 br dd	5.05 ddd
H-9 α	2.04 dd	2.47 dd	2.48 dd	2.66 dd
H-9 β	2.72 br d	2.71 br d	2.66 br d	2.39 dd
H-13a	6.37 dd	6.38 dd	6.36 dd	6.42 dd
H-13b	5.53 dd	5.83 dd	5.87 dd	5.88 dd
H-14a	4.55 br d	4.57 br d	4.24 br d	4.35 d
H-14b	4.33 br d	4.40 br d	3.87 br d	4.24 d
H-15a	3.88 d	4.61 d	4.34 d	9.98 d
H-15b	3.78 d	4.55 d	3.98 d	—
Ac ₁ ^e	1.85 s	2.13 s	2.09 s	2.07 s
Ac ₂ ^e	1.59 s	2.11 s	—	2.07 s
Ac ₃ ^e	—	2.08 s	—	—

^aThe spectrum of 4 was recorded at 80 MHz; all other data were obtained at 400 MHz. The numbering is the same in all skeletons.

^bSee Bohlmann *et al.* (14) for CDCl₃ data.

^cCouplings (Hz) 1-3: $J_{1,2\alpha} = 4$, $J_{1,2\beta} = 12$, $J_{2\alpha,2\beta} = 13$, $J_{2\beta,3\alpha} = 10$, $J_{2\beta,3\beta} = 5$, $J_{2\alpha,3\alpha} = 5$, $J_{2\alpha,3\beta} = 2$, $J_{3\alpha,3\beta} = 12$, $J_{5,6} = 10$, $J_{6,7} = 7.5$, $J_{7,13a} = 3.5$, $J_{7,13b} = 3.1$, $J_{7,8} = 7$, $J_{8,9\alpha} = 9$, $J_{8,9\beta} = 0$, $J_{9\alpha,9\beta} = 13$, $J_{13a,13b} = 0.8$, $J_{14a,14b} = 12.5$, $J_{15a,15b} = 13.5$.

^dCouplings (Hz) 4: $J_{1,2\alpha} = 6.5$, $J_{1,2\beta} = 9$, $J_{5,6} = 10.5$, $J_{6,7} = 7$, $J_{7,8} = 7$, $J_{8,9\alpha} = 10$, $J_{8,9\beta} = 3$, $J_{9\alpha,9\beta} = 13$, $J_{7,13a} = 3.5$, $J_{7,13b} = 3.1$, $J_{13a,13b} = 0.7$, $J_{14a,14b} = 12.5$.

^eIntensity 3H.

triacetate reported to have an identical stereoformula (17) was actually a melampolide as noted in a later publication (14).

The previously unreported ^{13}C -nmr data of 1 are listed in Table 2. The proton-bearing carbons were assigned by ^1H - ^{13}C heteronuclear correlation. The assignment of the signals of the quaternary carbons in the germacradiene ring of 1 was made possible by the regioselective and stereoselective conversion of 1 into the corresponding 4 α ,5 β -epoxide 6 with *m*-chloroperbenzoic acid. NOESY data of 6 (in DMSO-*d*₆) were consistent with the 4 α ,5 β stereochemistry of the epoxide ring and a conformation with H-6, H-8, H-14, and H-15 above and H-1, H-5, and H-7 below the plane of the ten-membered ring. The dihedral angles measured on a Dreiding model of 6 in this conformation are in excellent agreement with the coupling constants observed (Table 2). The value of $J_{5,6}$ of the epoxide ring (9.5 Hz) of 6 is consistent with literature data for the coupling H-5 α -H-6 β (9-10 Hz) in structurally related 5 α ,6 β epoxides (18,19). Smaller coupling constants ($J_{5\beta,6\beta} = 3-4$ Hz) have been observed in 4 β ,5 α isomers (19,20). The structure of 6 indicates that, under conditions employed for the selective epoxidation of 1, lactone 1 exists in the 1D^{14} , 15D_5 conformation (21).


Rearrangement of 1 in boiling toluene cleanly produced a 3:7 equilibrium mixture of 1 with the Cope product 7 which is an attractive starting material for the partial syn-

TABLE 2. ^1H nmr Data of Compounds 5–7.^a


Proton	Compound				
	5 CDCl_3	5 ^b $\text{C}_5\text{D}_5\text{N}$	6 ^c $\text{DMSO}-d_6$	6 ^d $\text{C}_5\text{D}_5\text{N}$	7 ^{e,f} CDCl_3
H-1	5.12 br dd	5.12 br dd	5.80 br dd	5.83	5.70 dd
H-2 α	2.30 m	2.14 m	2.18 br dd	2.25	5.07 d
H-2 β	2.54 m	2.61 dddd	2.54 dddd	2.81	5.16 d
H-3 α	~2.6	2.72 m	1.03 ddd	1.29	5.02 br s
H-3 β	2.09 m	~2.0	2.38 br dd	2.74	5.48 br s
H-5	4.83 d	4.92 d	3.02 d	3.16	2.67 d
H-6	5.03 dd	5.41 dd	4.35 dd	5.08	4.43 dd
H-7	2.21 dddd	2.40 dddd	3.59 dddd	3.67	2.85 dddd
H-8	5.32 ddd	5.86 ddd	4.52 ddd	5.01	5.25 ddd
H-9 α	2.47 dd	2.49 dd	2.55 dd	2.70	1.61 dd
H-9 β	2.58 br d	3.11 dd	2.45 br d	2.81	2.35 dd
H-11	2.75 dq	2.75 dq	—	—	—
H-13 α	} 1.43 ^g d	} 1.50 ^g d	6.19 d	6.48	6.16 d
H-13 β			5.77 d	5.81	5.59 d
H-14 α	4.29 d	4.42 d	4.78 d	5.06	4.32 d
H-14 β	3.91 d	4.39 d	4.54 d	4.91	4.12 d
H-15 α	4.34 d	4.36 d	3.56 dd	4.14	4.10 d
H-15 β	3.98 d	4.35 d	3.21 br dd	3.90	4.02 d
Ac ₁ ^h	2.13 s	2.03 s	2.01 s	1.98	2.12 s
Ac ₂ ^h	—	—	1.99 s	1.92	2.11 s

^aAll data were obtained at 400 MHz. The numbering is the same in all skeletons.^bCouplings (Hz) 5 in $\text{C}_5\text{D}_5\text{N}$: $J_{1,2\alpha} = 4.7$, $J_{1,2\beta} = 12.5$, $J_{5,6} = 10$, $J_{6,7} = 9.5$, $J_{7,8} = 8.5$, $J_{7,11} = 11.5$, $J_{8,9\alpha} = 10$, $J_{8,9\beta} = 1.5$, $J_{9\alpha,9\beta} = 12$, $J_{11,13} = 7$, $J_{14\alpha,14\beta} = 12.5$, $J_{15\alpha,15\beta} = 14$.^cCouplings (Hz) 6 in $\text{DMSO}-d_6$: $J_{1,2\alpha} = 3.6$, $J_{1,2\beta} = 12.4$, $J_{2\alpha,2\beta} = 12$, $J_{2\beta,3\alpha} = 12.5$, $J_{2\beta,3\beta} = 4.6$, $J_{2\alpha,3\alpha} = 6.5$, $J_{2\alpha,3\beta} = 0.7$, $J_{3\alpha,3\beta} = 12.5$, $J_{5,6} = 9.5$, $J_{6,7} = 6.7$, $J_{7,8} = 4.3$, $J_{7,13\alpha} = 3.4$, $J_{7,13\beta} = 3$, $J_{8,9\alpha} = 11.5$, $J_{8,9\beta} = 1.5$, $J_{9\alpha,9\beta} = 12.3$, $J_{14\alpha,14\beta} = 12.1$, $J_{15\alpha,15\beta} = 12.5$, $J_{15\alpha,\text{OH}} = 5.4$, $J_{15\beta,\text{OH}} = 3.5$; OH δ 5.05 dd.^dMultiplicities of all signals are the same as in CDCl_3 .^eRead 2 α , 2 β , 3 α , 3 β instead of 2 α , 2 β , 3 α , 3 β .^fCouplings (Hz) 7: $J_{1,2 \text{ cis}} = 11$, $J_{1,2 \text{ trans}} = 17.7$, $J_{5,6} = 12$, $J_{6,7} = 12$, $J_{7,8} = 11$, $J_{7,13\alpha} = 3$, $J_{7,13\beta} = 3$, $J_{8,9\alpha} = 11$, $J_{8,9\beta} = 4.3$, $J_{9\alpha,9\beta} = 13.5$, $J_{14\alpha,14\beta} = 12$, $J_{15\alpha,15\beta} = 14$.^gIntensity 2H.^hIntensity 3H.

thesis of the bioactive elemanolides vernomenin and vernolepin (22,23). Dihedral angles measured in a Dreiding model of 7 agree with all observed splitting constants, as does comparison with ^1H -nmr data of similar elemanolides in the literature (23–26). If the Cope product had the opposite stereochemistry at C-5 and C-10, H-5 and H-6



6



7

would be *cis* and the expected maximum value for $J_{5,6}$ would be 7.5 Hz. The observed value is 12 Hz (Table 2).

Two minor lactones of *M. minima* were the new analogues **3** and **5** whose structures were deduced by ms, extensive ^1H -nmr studies to verify coupling constants, ^{13}C nmr, and comparison with the spectral data of **1** (Tables 1–3). Compound **3** was obtained only as a 1:2 mixture with **5**. The chemical shifts and coupling constants showed that both **3** and **5** possessed identical stereochemistry around the 1(10) and 4,5 double bonds and a lactone ring trans fused as in **1**. The C-13 methyl group of **5** had to be α -oriented because of the value of $J_{7,11}$ (11.5 Hz).

Other substances identified in the extract were lupeol, α - and β -amyrin, stigmasterol, sitosterol, and isofucosterol.

TABLE 3. ^{13}C -nmr Data of Compounds 1–7.^a

Carbon	Compound						
	1 ^b CDCl ₃	2 CDCl ₃	3 ^c MeOH- <i>d</i> ₄	4 CDCl ₃	5 MeOH- <i>d</i> ₄	6 DMSO- <i>d</i> ₆	7 CDCl ₃
C-1	135.99 d	134.90 d	134.29 d	135.98 d	134.96 d	133.96 d	141.64 d
C-2	26.07 t	25.99 t	26.63 t	25.62 t	26.85 t	26.25 t	114.86 t ^e
C-3	34.23 t	34.53 t	35.01 t	29.93 t	35.24 t	33.08 t	115.77 t ^e
C-4	143.92 s	138.99 s	145.21 s	142.20 s	144.13 s	63.18 s	142.94 s
C-5	129.00 d	130.07 d	130.26 d	145.93 d	130.68 d	64.41 d	50.61 d ^f
C-6	77.02 d	77.05 d	79.23 d	74.63 d	78.08 d	78.83 d	78.36 d
C-7	52.82 d	52.81 d	53.86 d	51.89 d	59.06 d	47.65 d	51.68 d ^f
C-8	72.61 d	72.56 d	74.64 d	72.01 d	74.84 d	72.78 d	69.11 d
C-9	44.97 t	45.24 t	45.58 t	^d	46.66 t	44.77 t	40.71 t
C-10	129.98 s	131.38 s	137.38 s	131.06 s	136.05 s	128.25 s	44.33 s
C-11	135.26 s	135.94 s	135.37 s	133.65 s	41.42 d	134.09 s	136.53 s
C-12	169.80 s	169.52 s	171.93 s	168.61 s	180.89 s	170.05 s	169.15 s ^g
C-13	125.41 t	125.57 t	125.11 t	125.98 t	17.22 q	124.80 t	120.23 t
C-14	62.04 t	62.19 t	60.25 t ^e	61.89 t	60.08 t ^e	60.16 t ^e	67.23 t ^h
C-15	61.14 t	61.90 t	61.02 t ^e	188.45 d	61.03 t ^e	60.04 t ^e	66.32 t ^h
C-1'	171.06 s	170.86 s	172.05 s	170.22 s	171.72 s	170.24 s	170.03 s ^g
C-2'	21.10 q	21.07 q	21.17 q	20.76 q	21.10 q	20.85 q	20.95 q
C-1''	169.60 s	170.54 s	—	169.44 s	—	168.97 s	170.42 s ^g
C-2''	20.95 q	20.29 q	—	20.65 q	—	20.58 q	21.00 q
C-1'''	—	169.44 s	—	—	—	—	—
C-2'''	—	20.85 q	—	—	—	—	—

^aThe spectrum of **4** was recorded at 20 MHz. All other ^{13}C data were obtained at 100.61 MHz. Multiplicities were determined by DEPT experiments. Carbons 1'–2'' are acetate carbons. The numbering is the same in all skeletons.

^bResults of a ^1H - ^{13}C heteronuclear correlation facilitated the assignment.

^cFrom a mixture with **5**.

^dSignal not observed.

^{e-h}Assignments with the same superscript in a column are interchangeable.

EXPERIMENTAL

GENERAL EXPERIMENTAL PROCEDURES.—For separation of mixtures we used a Konik-500-A liquid chromatograph with RI detector, a Rheodyne injector (2-ml loop), and, unless stated otherwise, an Alltech RSil C18 column (10 mm i.d. \times 50 cm, 10 μ).

PLANT MATERIAL.—Aerial parts of *M. minima* were collected in Rio Vipos, Tucuman Province, Argentina, in May 1988. A voucher specimen (CANC #38) has been deposited at the Instituto Miguel Lillo, S.M. de Tucumán.

EXTRACTION OF *M. MINIMA*.—Flower heads and leaves (84 g) were extracted with CHCl₃ (2 \times 600 ml) at room temperature for 7 days to give 11.1 g of crude extract, which was suspended in EtOH (100 ml) at 50–55°, diluted with H₂O (75 ml), and extracted successively with hexane (3 \times 100 ml) and CHCl₃ (3 \times 100 ml). Evaporation of the hexane extract gave 5.2 g of residue, which was chromatographed over Si gel using hexane and increasing amounts of Et₂O (0–33%), yielding crude triterpenes (234 mg) and crude sterols (81 mg). The triterpene sample was saponified with dilute KOH, and the neutral unsaponifiables were chromatographed over Si gel with hexane-Et₂O (4:1) and gave purified triterpenes (96 mg). Reversed-

phase hplc of part (32 mg) of the purified triterpenes (MeOH, flow rate 3 ml/min) gave crystalline lupeol (3 mg), crystalline β -amyrin (14 mg), and 5.2 mg of α -amyrin contaminated with an unidentified triterpene. The crude sterol sample, processed in the same way as the crude triterpenes, yielded stigmasterol (14 mg), β -sitosterol (13 mg), and isofucosterol (1.2 mg).

The residue of the CHCl_3 extract (4.3 g) was chromatographed over Si gel using C_6H_6 with increasing amounts of EtOAc (20–33%). The separation was monitored by Si gel tlc using C_6H_6 -EtOAc mixtures (2:1, 1:1, and 2:3) as developers, and 60 fractions were collected. Fractions 11–33 afforded crystalline **1** [1.48 g, mp 109.5–110.5°, from heptane/EtOAc, R_f 0.50, C_6H_6 -EtOAc (1:1)]. Fractions 48–53 were combined (residue 193 mg) and rechromatographed over Si gel C_6H_6 -EtOAc (5:3). A 1:2 mixture (50 mg) of **3** and **5** was obtained [unresolved spot on Si gel, R_f 0.28, C_6H_6 -EtOAc (2:3)]. Separation by reversed-phase hplc [C8 column (Phenomenex, Palos Verdes, California), 10 mm i.d. \times 50 cm, 10 μ , MeOH- H_2O (3:2), flow 2 ml/min] gave pure **5**, Rt 5 min, as a gum. Lactone **3** partially decomposed during workup of the column effluent.

IDENTIFICATION OF STEROLS AND TRITERPENES.—All compounds were first tentatively identified on the basis of their relative retention times in gc and hplc. ^1H - and ^{13}C -nmr spectra (run at 400 and 100.61 MHz, respectively) confirmed these assignments. The ^{13}C -nmr spectra showed sitosterol and stigmasterol to be sterically pure (27). This is of interest because it is known, mainly through the work of Akihisa and co-workers (28,29), that 24-alkyl sterols of higher plants are usually mixtures of epimers at C-24.

14-Acetoxyartemisiifolin-6 α -O-acetate [1].—Mp 109.5–110.5° from heptane-EtOAc (9:1); ir (KBr) ν max cm^{-1} 3445 (OH), 1770 (shoulder) (γ -lactone), 1734 (OAc), 1650, 1233, 1222, 1035, 999; cims (reagent gas CH_4) m/z (rel. int.) [$\text{C}_{19}\text{H}_{24}\text{O}_7 + \text{H}$] $^+$ 365 (44), 347 (38), 333 (42), 323 (58), 305 (15), 245 (100), 227 (99).

14-Acetoxy-4 α ,5 β -epoxyartemisiifolin-6 α -O-acetate [6].—To **1** (182 mg) in CH_2Cl_2 (3 ml) and 0.5 M NaHCO_3 (1.25 ml) cooled in ice was added with magnetic stirring *m*-chloroperbenzoic acid (135 mg) in small portions. The progress of the reaction was followed by tlc. The reaction was complete in 5 h. The organic layer was diluted with CH_2Cl_2 (15 ml) and extracted with 10% aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (2 \times 2 ml), 1 M NaOH (3 \times 2 ml), and H_2O (2 \times 2 ml). After drying and solvent evaporation the residue **6** was shown to be impure by hplc and tlc. Recrystallization from heptane-EtOAc (1:4) gave pure **6** (98 mg); mp 207–209°; ir (KBr) ν max cm^{-1} 3445, 1745, 1736, 1650, 1382, 1290, 1269, 1244, 1226, 1161, 1042, 1024, 1008, 968, 952, 923; cims (reagent gas CH_4) m/z (rel. int.) [$\text{C}_{19}\text{H}_{24}\text{O}_8 + \text{H}$] $^+$ 381 (61), 363 (39), 349 (51), 339 (75), 321 (20), 261 (90), 243 (100).

14-Acetoxyartemisiifolin-6 α ,15-di-O-acetate [2].— Ac_2O (0.30 ml) was added to **1** (90 mg) in pyridine (3 ml). After the usual workup the product was purified by cc over Si gel to give **2** (49 mg) as a gum; ir (KBr) ν max cm^{-1} 1766, 1742, 1651, 1374, 1239, 1145, 1023, 961. The ms sample of **2** decomposed prior to analysis.

6 α ,14-Diacetoxy-15-oxo-(Z)1(10),(Z)4-germacradien-8 α ,12-olide [4].—To **1** (50 mg) in CHCl_3 (50 ml) was added active MnO_2 (300 mg) at room temperature and with magnetic stirring. The progress of the reaction was monitored by tlc. It was complete after 2 h. Filtration and solvent evaporation yielded **4** (42 mg) as a gum. This aldehyde was very unstable, and most of it had decomposed after one day at room temperature.

6 α ,14-Diacetoxy-15-hydroxyeleman-8 α ,12-olide [7].—Compound **1** (30 mg) was refluxed in toluene (6 ml) under N_2 . The progress of the reaction was monitored by tlc. Only one product could be detected. The equilibrium was reached in about 5 h. Preparative Si gel tlc [EtOAc- C_6H_6 (4:3), two developments] gave **7** (gum, 18.2 mg) and **1** (8.3 mg). Compound **7**: ir (film) ν max cm^{-1} 3570 (OH), 1770 (γ -lactone), 1740 (OAc), 1674, 1642, 1372, 1245, 1143, 1041, 992, 974, 924, 757.

14-Hydroxyartemisiifolin-6 α -O-acetate [3].—Compound **3** was obtained as the minor component of a 1:2 mixture with **5**. It partially decomposed during workup after hplc separation from **5**. The ^1H spectrum of this material indicated that one of the components was 13-methoxy-11,13-dihydro-14-hydroxyartemisiifolin-6 α -O-acetate, probably produced by 1,4-addition of MeOH to the α,β -unsaturated γ -lactone **3** catalyzed by traces of acid present in the CHCl_3 used for extraction. Mixture of **3** and **5**: ir (KBr) ν max cm^{-1} 3480, 3430, 1750, 1710, 1652; cims (reagent gas CH_4) m/z (rel. int.) [$\text{C}_{17}\text{H}_{24}\text{O}_6 + \text{H}$] $^+$ 325 (30), [$\text{C}_{17}\text{H}_{22}\text{O}_6 + \text{H}$] $^+$ 323 (11). Compound **5** could be isolated pure (see below) and was properly identified. Subtraction of the corresponding signals in the high field ^{13}C and ^1H spectra permitted assignment of all signals corresponding to **3**.

11 β H-11,13-Dihydro-14-hydroxyartemisiifolin-6 α -O-acetate [5].—Gum, ir (KBr) ν max cm^{-1} 3430, 1760, 1720, 1655, 1414, 1380, 1260, 1220, 1035, 957; cims (reagent gas CH_4) m/z (rel. int.) [$\text{C}_{17}\text{H}_{24}\text{O}_6 + \text{H}$] $^+$ 325 (36), 307 (77), 283 (21), 265 (19), 205 (61), 187 (100).

ACKNOWLEDGMENTS

We thank the Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina and the Consejo de Investigaciones de la Universidad Nacional de Tucumán for financial support.

LITERATURE CITED

1. B.L. Robinson, *Contrib. Gray Herb., Harv. Univ.*, **104**, 60 (1934).
2. A.B. Gutierrez, J.C. Oberti, P. Kulanthaivel, and W. Herz, *Phytochemistry*, **24**, 2967 (1985).
3. W. Herz and P. Kulanthaivel, *Phytochemistry*, **24**, 1761 (1985), and references cited therein.
4. A.B. Gutierrez, J.C. Oberti, and W. Herz, *Phytochemistry*, **27**, 938 (1988).
5. M. del R. Cuenca, A. Bardón, C.A.N. Catalán, and W.C.M.C. Kokke, *J. Nat. Prod.*, **51**, 625 (1988).
6. R. Boeker, J. Jakupovic, F. Bohlmann, and G. Schmeda-Hirschmann, *Planta Med.*, 105 (1987).
7. W. Herz, *Stud. Org. Chem. (Amsterdam)*, **26**, 143 (1986).
8. V. Castro, J. Jakupovic, and F. Bohlmann, *Phytochemistry*, **25**, 1750 (1986).
9. A.B. Gutierrez, J.C. Oberti, V.E. Sosa, and W. Herz, *Phytochemistry*, **26**, 2315 (1987).
10. V. Castro, J. Jakupovic, and F. Bohlmann, *Phytochemistry*, **28**, 527 (1989).
11. F.S. Knudsen, W. Vilegas, F. Oliveira, and N.F. Roque, *Phytochemistry*, **25**, 1240 (1986).
12. V. Castro and J. Jakupovic, *Phytochemistry*, **24**, 2450 (1985).
13. A. Gutierrez and W. Herz, *Phytochemistry*, **27**, 3871 (1988).
14. F. Bohlmann, P. Singh, and J. Jakupovic, *Phytochemistry*, **21**, 2029 (1982).
15. W. Herz and P.S. Kalyanaraman, *J. Org. Chem.*, **40**, 3486 (1975).
16. N.H. Fischer, E.J. Olivier, and H.D. Fischer, *Prog. Chem. Organ. Nat. Prod.*, **38**, 47 (1979).
17. F. Bohlmann and N.L. Van, *Phytochemistry*, **17**, 570 (1978).
18. F. Bohlmann, A. Adler, J. Jakupovic, R.M. King, and H. Robinson, *Phytochemistry*, **21**, 1349 (1982), compounds **16** and **20**.
19. E. Maldonado, A.M. Flores, and A. Ortega, *Phytochemistry*, **27**, 861 (1988), compounds **5-7** and discussion therein.
20. F. Bohlmann, J. Jakupovic, M. Ahmed, and A. Schuster, *Phytochemistry*, **22**, 1623 (1983), compound **103**.
21. Z. Samek and J. Harmatha, *Collect. Czech. Chem. Commun.*, **43**, 2779 (1978).
22. S.M. Kupchan, R.J. Hemingway, D. Werner, A. Karim, A.T. McPhail, and G.A. Sim, *J. Am. Chem. Soc.*, **90**, 3596 (1968).
23. S.M. Kupchan, R.J. Hemingway, D. Werner, and A. Karim, *J. Org. Chem.*, **34**, 3903 (1969).
24. M.L. Cardona, B. Garcia, J.R. Pedro, and J.F. Sinisterra, *Phytochemistry*, **28**, 1264 (1989).
25. F. Bohlmann, C. Zdero, M.R. King, and H. Robinson, *Liebigs Ann. Chem.*, 240 (1984).
26. F. Bohlmann, J. Jakupovic, and A. Schuster, *Phytochemistry*, **22**, 1637 (1983).
27. J.L.C. Wright, A.G. McInnes, S. Shimizu, D.G. Smith, J.A. Walter, D. Idler, and W. Khalil, *Can. J. Chem.*, **56**, 1889 (1978).
28. T. Akihisa, W.C.M.C. Kokke, T. Tamura, and T. Matsumoto, *Lipids*, accepted for publication, and references therein.
29. T. Akihisa, T. Tamua, T. Matsumoto, W.C.M.C. Kokke, P. Ghosh, and S. Thakur, *J. Chem. Soc., Perkin Trans. 1*, (in press).

Received 29 December 1989