

Note: Inverted time-ordering in two-dimensional-Raman-terahertz spectroscopy of water

Peter Hamm, Janne Savolainen, Junichi Ono, and Yoshitaka Tanimura

Citation: *The Journal of Chemical Physics* **136**, 236101 (2012); doi: 10.1063/1.4729945

View online: <http://dx.doi.org/10.1063/1.4729945>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/136/23?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

Assessment of mode-mixing and Herzberg-Teller effects on two-photon absorption and resonance hyper-Raman spectra from a time-dependent approach

J. Chem. Phys. **140**, 094107 (2014); 10.1063/1.4867273

Two-dimensional-Raman-terahertz spectroscopy of water: Theory

J. Chem. Phys. **136**, 094516 (2012); 10.1063/1.3691601

Polarization of molecular targets using infrared stimulated Raman adiabatic passage

J. Chem. Phys. **133**, 094301 (2010); 10.1063/1.3475523

Electronic polarization effect on low-frequency infrared and Raman spectra of aprotic solvent: Molecular dynamics simulation study with charge response kernel by second order Møller–Plesset perturbation method

J. Chem. Phys. **127**, 244502 (2007); 10.1063/1.2813421

The formation of ClOO following the photoexcitation of aqueous OCIO studied by two-color, time-resolved resonance Raman spectroscopy

J. Chem. Phys. **112**, 505 (2000); 10.1063/1.480540



AIP | Journal of
Applied Physics

Journal of Applied Physics is pleased to
announce **André Anders** as its new Editor-in-Chief

Note: Inverted time-ordering in two-dimensional-Raman-terahertz spectroscopy of water

Peter Hamm,^{1,a)} Janne Savolainen,¹ Junichi Ono,² and Yoshitaka Tanimura²

¹Physikalisch-Chemisches Institut, Universität Zürich, Switzerland

²Department of Chemistry, Kyoto University, Japan

(Received 25 April 2012; accepted 5 June 2012; published online 15 June 2012)

[<http://dx.doi.org/10.1063/1.4729945>]

In a previous paper,¹ we have studied the 2D-Raman-THz response of liquid water, based on an all-atom molecular dynamics (MD) simulation. This novel two-dimensional spectroscopy is designed to directly work in the low-frequency spectral range of the *intermolecular* degrees of freedom of water. Its information content is similar to 2D-Raman^{2–9} or 2D-THz spectroscopy,¹⁰ but the experimental implementation should be easier. In 2D-Raman-THz spectroscopy, a near-infrared laser pulse excites a coherence of an intermolecular vibrational mode through a Raman process (see Fig. 1(b)). After time t_1 , the THz pulse interrogates the molecular system again and switches it into another coherence, which is finally read out at time $t_1 + t_2$ by the emission of a THz field. It is this switching between coherences which ultimately allows one to measure couplings and correlations between the various degrees of freedom of the system under study.

In the present note, we explore the inverted time-ordered pulse sequence, in which the first coherence is generated by a direct THz excitation, while the switching of coherences is achieved by a Raman interaction (Fig. 1(a)). The dependence of the polarizability on the nuclear coordinates is expected to be more nonlinear compared to that of the dipole moment. Both pulse sequences involve a two-quantum transition for the switching of coherences, which requires such an electrical anharmonicity. Since the second interaction in the inverted pulse sequence is the one with the larger nonlinearity, one might expect that it reveals more of the couplings and correlations between the various degrees of freedom. The second motivation for looking at the inverted time-ordered response function lies in the fact that in an experimental realization of 2D-Raman-THz spectroscopy, where the laser pulses will not be infinitely short, one will convolute over both response functions during pulse overlap, so both will be needed for a modeling of an experimental result.

The response function of the original pulse sequence is¹

$$R^{(I)}(t_2, t_1) \propto \text{Tr}\{\mu(t_2 + t_1)[\Pi(t_1), [\Pi(0), \rho_{\text{eq}}]]\}, \quad (1)$$

where [..., ...] is a commutator, $\Pi(t)$ and $\mu(t)$ are polarizability and dipole operators, respectively, and ρ_{eq} the equilibrium density matrix. For the inverted time-ordered response function, this is changed to

$$R^{(II)}(t_2, t_1) \propto \text{Tr}\{\mu(t_2 + t_1)[\Pi(t_1), [\mu(0), \rho_{\text{eq}}]]\}. \quad (2)$$

The hybrid equilibrium-non-equilibrium approach introduced by Tanimura and co-workers⁸ is used to compute this

response function from an all-atom MD simulation:

$$R^{(II)}(t_2, t_1) \propto \langle (\mu_+(t_2) - \mu_-(t_2))\dot{\mu}(-t_1) \rangle_{\text{nonequ}}. \quad (3)$$

Here, the average is taken over an ensemble of non-equilibrium trajectories, and $\mu_+(t_2)$ and $\mu_-(t_2)$ refer to the dipole moments at time t_2 obtained after perturbing the momenta of the individual atoms by $\pm \Delta p_i$, respectively, at time $t = 0$. This perturbation now requires the calculation of the forces on the individual water molecules upon acting of a short electric field pulse on the polarizability tensor (rather than the dipole moment). All other simulation details, in particular the MD protocol, the parametrization for the calculation of the dipole moment and the polarizability tensor of the simulation box and the electric field strength are the same as in Ref. 1.

Fig. 1(c) shows the response function $R^{(II)}$ calculated in this way and combines it with $R^{(I)}$ in Fig. 1(d). The time-axis t_1 has been inverted for $R^{(II)}$ to emphasize the continuity of the overall response function with $R^{(I)}(t_1 = 0, t_2) = R^{(II)}(t_1 = 0, t_2)$ for all t_2 . This continuity is, of course, expected as both response functions are supposed to be the same when the THz and the Raman-pulses coincide with $t_1 = 0$. However, given the fact that Eq. (3) and its counterpart for $R^{(I)}$ (Ref. 1) require the evaluations of different derivatives, the continuity serves as a good consistency check of the numerical procedures.

Most interesting is the signature of an echo along the diagonal in the inverted THz-Raman-THz sequence along the diagonal for $t_1 = t_2$ (Fig. 1(c), label *). An echo arises when a coherence dephases very quickly, and then rephases upon a second perturbation. Traces of a photon echo appear in simulations of 2D-Raman spectroscopy, but only when approximating the signal with instantaneous or quenched normal modes.^{3,6} The photon echo gets completely blurred in 2D-Raman spectroscopy when all the dynamic effects of a fully anharmonic MD force field are included. To our knowledge, the echo signal in Fig. 1(c) is the clearest signature of that kind observed so far for a fully anharmonic MD simulation, highlighting the potential that lies in the THz-Raman-THz pulse sequence. The reason for the larger sensitivity to an echo might be the following: Assuming that the dependence of the polarizability Π on nuclear coordinates q is more nonlinear than that of the dipole moment μ , the THz-Raman-THz sequence is dominated by the term $\text{Tr}\{q(t_2 + t_1)[q^2(t_1), [q(0), \rho_{\text{eq}}]]\}$, where the $q^2(t_1)$ -term is responsible for a two-quantum transition. Indeed, it has been




FIG. 1. (a) The inverted pulse sequences and (b) the original pulse sequence of Ref. 1. A red pulse depicts a THz field, a pair of blue pulses depicts the two fields necessary to achieve a Raman transition. Panels (c) and (d) show the response functions according to Eqs. (1) and (2), respectively, where red and blue colors depict response of opposite sign. The response function $R^{(l)}$ in panel (d) is adapted from Ref. 1. The time-axis t_1 has been inverted in panel (c) to emphasize the continuity of the overall response function at $t_1 = 0$. The feature labeled with an asterisk (*) indicates an echo.

shown that this term dephases as a function of $(t_1 - t_2)$, i.e., forms an echo at $t_1 = t_2$ (Ref. 4) (the mechanism for the formation of an echo is different from resonant 2D spectroscopy, where two phase-conjugate pathways exist). On the other hand, the term $\text{Tr}\{q^2(t_2 + t_1)[q(t_1), [q(0), \rho_{\text{eq}}]]\}$ contributes equally strongly in the Raman-Raman-Raman sequence of 2D-Raman spectroscopy. The latter does not switch coherences after time t_1 , but is superimposed and hence might blur the echo. Even worse, in our original Raman-THz-THz sequence $R^{(l)}$, this second term dominates. Another possible explanation for the different responses might be connected to non-coincidence effect we have identified for the librational mode,¹ in which case Raman and THz transitions are mutually exclusive to a certain extent. Calculating the various signals based on a normal mode analysis will clarify this effect.

We conclude that indeed the inverted pulse sequence more sensitively measures the coupling between the various degrees of freedom of water. Simulations of that kind, on the one hand, will guide the design of an experimental implementation of 2D-Raman-THz spectroscopy, and on the other

hand, will contribute to validating the water model on which the simulation is based, once experimental data are available. Should one observe an echo, it will give unprecedented insights into water structure and dynamics.

The work has been supported by the Swiss National Science Foundation (SNF) through the National Center of Competence and Research (NCCR) MUST.

^{a)} Author to whom correspondence should be addressed. Electronic mail: phamm@pci.uzh.ch.

¹P. Hamm and J. Savolainen, *J. Chem. Phys.* **136**, 094516 (2012).

²Y. Tanimura and S. Mukamel, *J. Chem. Phys.* **99**, 9496 (1993).

³S. Saito and I. Ohmine, *J. Chem. Phys.* **108**, 240 (1998).

⁴J. T. Fourkas, *Adv. Chem. Phys.* **117**, 235 (2001).

⁵L. J. Kaufman, J. Heo, L. D. Ziegler, and G. R. Fleming, *Phys. Rev. Lett.* **88**, 207402 (2002).

⁶S. Saito and I. Ohmine, *J. Chem. Phys.* **119**, 9073 (2003).

⁷K. J. Kubarych, C. J. Milne, and R. J. D. Miller, *Int. Rev. Phys. Chem.* **22**, 497 (2003).

⁸T. Hasegawa and Y. Tanimura, *J. Chem. Phys.* **125**, 074512 (2006).

⁹Y. L. Li, L. Huang, R. J. D. Miller, T. Hasegawa, and Y. Tanimura, *J. Chem. Phys.* **128**, 234507 (2008).

¹⁰W. Kuehn, K. Reimann, M. Woerner, T. Elsaesser, and R. Hey, *J. Phys. Chem. B* **115**, 5448 (2011).