

Erratum: Monte Carlo studies on the hydrophobic hydration in dilute aqueous solutions of nonpolar molecules [J. Chem. Phys. 7 1, 2421 (1979)]

Susumu Okazaki, Koichiro Nakanishi, Hidekazu Touhara, and Yoshinori Adachi

Citation: *The Journal of Chemical Physics* **72**, 4253 (1980); doi: 10.1063/1.439831

View online: <http://dx.doi.org/10.1063/1.439831>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/72/7?ver=pdfcov>

Published by the AIP Publishing

Articles you may be interested in

Erratum: Trajectory studies of the molecular dynamics of ethyl radical decomposition [J. Chem. Phys. 7 1, 2911 (1979)]

J. Chem. Phys. **76**, 2771 (1982); 10.1063/1.443587

Erratum: Repulsive intermolecular potential between two H₂ molecules [J. Chem. Phys. 7 1, 5362 (1979)]

J. Chem. Phys. **73**, 4712 (1980); 10.1063/1.440675

Erratum: Elliptic phase boundaries between smectic and nematic phases [J. Chem. Phys. 7 1, 1874 (1979)]

J. Chem. Phys. **73**, 4149 (1980); 10.1063/1.440651

Erratum: Vibrational relaxation in jet-cooled alkyl benzenes [J. Chem. Phys. 7 1, 3886 (1979)]

J. Chem. Phys. **72**, 2905 (1980); 10.1063/1.439828

Monte Carlo studies on the hydrophobic hydration in dilute aqueous solutions of nonpolar molecules

J. Chem. Phys. **71**, 2421 (1979); 10.1063/1.438647



NEW Special Topic Sections

NOW ONLINE
Lithium Niobate Properties and Applications:
Reviews of Emerging Trends

AIP Applied Physics Reviews

trusive, convenient, and accurate method for determining D_m . This is even true for unliganded Hb, which must be considered a worst case because of its large molar absorbance. In contrast to this, the classical diaphragm method, while deceptively simple to analyze, is beset with difficulties when applied to proteins.

We wish to thank Professor G. D. J. Phillies for helpful comments. This material is based upon work supported by the National Science Foundation under Grant No. CHE 77-23564. Also, the facilities of the Departmental Laser Laboratory, which was funded by the National Science Foundation (CHE 77-14547), were used.

¹K. H. Keller, E. R. Canales, and S. I. Yum, *J. Phys. Chem.* **75**, 379 (1971).

²L. Onsager and R. M. Fuoss, *J. Phys. Chem.* **36**, 2689 (1932).

³J. G. Kirkwood, R. L. Baldwin, P. J. Dunlop, L. J. Gosting, and G. Kegeles, *J. Chem. Phys.* **33**, 1505 (1960).

⁴G. D. J. Phillies, *J. Chem. Phys.* **60**, 976, 983 (1974).

⁵J. L. Anderson and C. C. Reed, *J. Chem. Phys.* **64**, 3240, 4336 (1976).

⁶R. S. Hall, Y. Oh, and C. S. Johnson, Jr., *J. Chem. Phys.* (in press).

⁷C. R. Jones and C. S. Johnson, Jr., *Biopolymers* **17**, 1581 (1978).

⁸S. S. Alpert and G. Banks, *Biophys. Chem.* **4**, 287 (1976).

⁹W. B. Veldkamp and J. R. Votano, *J. Phys. Chem.* **80**, 2794 (1976).

¹⁰A. P. Minton and P. D. Ross, *J. Phys. Chem.* **82**, 1934 (1978).

¹¹R. G. Kitchen, B. N. Preston, and J. D. Wells, *J. Polymer Sci. Symposium No. 55*, 39 (1976).

¹²B. D. Fair, D. Y. Chao, and A. M. Jamieson, *J. Colloid Interface Sci.* **66**, 32 (1978).

¹³L. J. Gosting, *Adv. Protein Chem.* **11**, 429 (1956).

¹⁴R. F. Probststein, W.-F. Leung, and Y. Alliance, *J. Phys. Chem.* **83**, 1228 (1979).

¹⁵G. Gros, *Biophys. J.* **22**, 453 (1978).

¹⁶V. Riveros-Moreno and J. B. Wittenberg, *J. Biol. Chem.* **247**, 895 (1972).

ERRATA

Erratum: Monte Carlo studies on the hydrophobic hydration in dilute aqueous solutions of nonpolar molecules [*J. Chem. Phys.* **71**, 2421(1979)]

Susumu Okazaki, Koichiro Nakanishi, and Hidekazu Touhara

Department of Industrial Chemistry, Kyoto University, Kyoto 606, Japan

Yoshinori Adachi

Nagoya Institute of Technology, Nagoya 466, Japan

Some of internal energy values given in Table III are incorrectly calculated. Tables III and IV should be corrected to read as shown below.

TABLE III.

	Pure water	Aqueous methane solution	Aqueous isobutane solution
U_N	-35.1 ± 0.2	-36.1 ± 0.2	-35.0 ± 0.7
U_{vw}		-36.4 ± 0.6	-35.0 ± 0.5
U_{dw}		-36.8 ± 0.4	-35.7 ± 0.8
U_s		-0.5 ± 0.2	-6.1 ± 1.2
$U_{vw} - U_{dw}$		0.3 ± 0.9	0.8 ± 0.5
z	-0.10	0.95	1.28

TABLE IV. Thermodynamic functions of dissolution of methane into water at 295.15 K in kJ mol^{-1} .

$A - A_0$	$U - U_0$	$T(S - S_0)$
0.8	-1.0	-1.8

Some numerical values cited in the text should also be corrected according to the above figures. Revised discussion will appear in the forthcoming article.