See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/234845997

Erratum: The structure of liquid 1,2 dichloroethane by neutron diffraction. I. Molecular structure, full pair correlation function, and temperature effects [J. Chem. Phys. 87, 7171...

ARTICLE in THE JOURNAL OF CHEMICAL PHYSICS · DECEMBER 1987

Impact Factor: 2.95 · DOI: 10.1063/1.453360

CITATIONS

7

READS

11

6 AUTHORS, INCLUDING:



Francico J Bermejo

Spanish National Research Council

365 PUBLICATIONS **3,683** CITATIONS

SEE PROFILE



Noé G. Almarza

Spanish National Research Council

95 PUBLICATIONS 1,073 CITATIONS

SEE PROFILE



Jorge Santoro

Spanish National Research Council

123 PUBLICATIONS 3,245 CITATIONS

SEE PROFILE



Erratum: The structure of liquid 1,2 dichloroethane by neutron diffraction. I. Molecular structure, full pair correlation function, and temperature effects [J. Chem. Phys. 87, 7171 (1987)]

F. J. Bermejo, E. Enciso, J. C. Dore, P. Chieux, N. Garcia et al.

Citation: J. Chem. Phys. 88, 6680 (1988); doi: 10.1063/1.454760

View online: http://dx.doi.org/10.1063/1.454760

View Table of Contents: http://jcp.aip.org/resource/1/JCPSA6/v88/i10

Published by the AIP Publishing LLC.

Additional information on J. Chem. Phys.

Journal Homepage: http://jcp.aip.org/

Journal Information: http://jcp.aip.org/about/about_the_journal Top downloads: http://jcp.aip.org/features/most_downloaded

Information for Authors: http://jcp.aip.org/authors

ADVERTISEMENT



Erratum: The structure of liquid 1,2 dichloroethane by neutron diffraction. I. Molecular structure, full pair correlation function, and temperature effects [J. Chem. Phys. 87, 7171 (1987)]

F. J. Bermejo

Instituto de Estructura de la Materia, C.S.I.C., Serrano 119, 28006 Madrid, Spain

E. Enciso

Departamento de Química Física, Facultad de Ciencias Químicas, Universidad Complutense, 28040 Madrid, Spain

J. C. Dore

Physics Laboratory, University of Kent, Canterbury, CT2 7NR, Great Britain

P Chieux

Institut Laue-Langevin, rue des Martyrs, Grenoble, France

N. Garcia

Departamento di Química Física, Facultad de Ciencias Químicas, Universidad Complutense, 28040 Madrid, Spain

J. Santoro

Instituto de Estructura de la Materia, C.S.I.C., Serrano 119, 28006 Madrid, Spain

Due to an error in the data corrections and subsequent extrapolation to $Q \rightarrow 0$ of the measured differential cross section $(0.3 \leqslant Q \leqslant 23)$, the (extrapolated) small-angle region was incorrectly treated. The low-Q data region was recorrected using a cadmium-rod dataset taken for the same geometrical arrangement as well as an experimental value for the isothermal compressibility (at 298 K) of 1.119×10^{-19} m² N⁻¹.

A reanalysis of the structure factors affected by this correction was performed and the corrected estimates [total Fourier transforms and full pair correlation functions g(r)] are shown in Figs. 1 and 2.

The abovementioned errors introduced a phase distortion in Figs. 7 of the paper which should be replaced by the two figures given here.

The predictions for the intramolecular structure and temperature-difference effects are not affected by the error in the small-angle data.

Most of the conclusions of the paper pertaining to intermolecular structure are still valid since they were taken on qualitative grounds and compared with a previously reported Monte Carlo simulation. The main effect of the mentioned correction is to shift the most prominent g(r) peak from 8 to \sim 6 Å, with a second maximum \sim 10 Å.




FIG. 1. Total Fourier transforms of the $S_m(Q)$ structure factors at two extreme temperatures.




FIG. 2. Full pair correlation functions for -20 °C (full curve) and +70 °C (dashed curve).

J. Chem. Phys. 88 (10), 15 May 1988

0021-9606/88/106680-02\$02.10

© 1988 American Institute of Physics

Further work on decomposition of the total g(r) into its pair contributions using chlorine isotopes is, at present, in progress as well as the analysis of the measured correlation

function by means of new simulation data.

¹D. M. Newit and K. W. Weale, J. Chem. Soc. 1951, 3092.

Erratum: Accurate theoretical modeling of NaCl [J. Chem. Phys. 84, 867 (1986)]

E. Clementi, G. Corongiu, and P. K. Swaminathan IBM Corporation, Department 48B/Mail Station 428, Kingston, New York 12401

Misprints have been uncovered in recent papers¹⁻³ reporting the interaction potentials between halogen anions and alkaline cations. Concerning Ref. 1, Table IV has a misprint: the F coefficient for the + interaction has a wrong sign and should read + 3.994 29. Concerning Ref. 2, the potential function, Eq. (10) on page 500, should read

$$V(r) = Z_i Z_i e^2 / (4\pi\epsilon_0 r) + (A_{ii} r^{B_{ij}} + C_{ii} r) e^{-D_{ij} r} + E_{ii} r^{-F_{ij}}.$$

Concerning Ref. 3 the correct analytical expression for the ++ interaction is the one given above. Finally, the potential for the F^-F^- interaction is also incorrectly reported in Ref. 3. Using the analytical expression given above, the following parameters (in a.u.) should be used:

$$A = -28.01022$$
, $B = -4.51542$, $C = -0.05446$, $D = 0.63766$, $E = 23.87259$, $F = 5.55292$.

An alternative set is A = 1.03733, B = 0.01024, C = 0.32410, D = 0.79816, E = 0.0, and F = 0.0.

We thank Dr. F. Jenc (Philipps-Marburg University) and Dr. V. K. W. Cheng (University of Hong Kong) for having called our attention to problems in Refs. 1, 2, and 3.

¹P. K. Swaminathan, A. Laaksonen, G. Corongiu, and E. Clementi, J. Chem. Phys. **84**, 495 (1986).

²A. Laaksonen and E. Clementi, Mol. Phys. **56**, 495 (1986).

³P. K. Swaminathan and E. Clementi, J. Phys. Chem. 91, 1020 (1987).