

Published in final edited form as:

ACS Chem Neurosci. 2015 July 15; 6(7): 1040-1054. doi:10.1021/acschemneuro.5b00007.

# Top-down control of serotonin systems by the prefrontal cortex: a path towards restored socioemotional function in depression

Collin Challis<sup>1,2</sup> and Olivier Berton<sup>1,2</sup>

<sup>1</sup>Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104

<sup>2</sup>Neuroscience Graduate Group, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104

#### **Abstract**

Social withdrawal, increased threat perception and exaggerated reassurance seeking behaviors are prominent interpersonal symptoms in major depressive disorder (MDD). Altered serotonin (5-HT) systems and corticolimbic dysconnectivity have long been suspected to contribute to these symptomatic facets, however, the underlying circuits and intrinsic cellular mechanisms that control 5-HT output during socioemotional interactions remain poorly understood. We review literature that implicates a direct pathway between the ventromedial prefrontal cortex (vmPFC) and dorsal raphe nucleus (DRN) in the adaptive and pathological control of social approachavoidance behaviors. Imaging and neuromodulation during approach-avoidance tasks in humans point to the cortical control of brainstem circuits as an essential regulator of socioemotional decisions and actions. Parallel rodent studies using viral-based connectomics and optogenetics are beginning to provide a cellular blueprint of the underlying circuitry. In these studies, manipulations of vmPFC synaptic inputs to the DRN have revealed bidirectional influences on socioaffective behaviors via direct monosynaptic excitation and indirect disynaptic inhibition of 5-HT neurons. Additionally, adverse social experiences that result in permanent avoidance biases, such as social defeat, drive long-lasting plasticity in this microcircuit, potentiating the indirect inhibition of 5-HT output. Conversely, neuromodulation of the vmPFC via deep brain stimulation (DBS) attenuates avoidance biases by restoring the direct excitatory drive of 5-HT neurons and strengthening a key subset of forebrain 5-HT projections. Better understanding the cellular organization of the vmPFC-DRN pathway and identifying molecular determinants of its neuroplasticity can open fundamentally novel avenues for the treatment of affective disorders.

#### **Keywords**

dorsal raphe; ventromedial prefrontal cortex; serotonin; optogenetics; electrophysiology
depression; mood disorders; social perception; social defeat; social avoidance

<sup>\*</sup>Correspondence should be addressed to Olivier Berton (bertonol@mail.med.upenn.edu).

## Why study socioemotional deficits in MDD?

Humans are social by nature. Almost all of their actions are directed towards or generated in response to other individuals. The ability to regulate one's social behavior in order to optimize social relationships and capitalize on opportunities in the social environment is referred to as social competence, which involves a multitude of cognitive and affective abilities. <sup>1, 2</sup> For example, achieving emotional attunement and tempering avoidant and aggressive impulses are necessary skills to establish novel relationships and sustain those relationships through adversity. Impairments in social competence are one of the most prominent and disabling characteristics of psychiatric illnesses that affect a third of the world's population at some point in their life.<sup>3</sup> Socioemotional deficits figure among the core symptoms of several psychiatric disorders such as autism, schizophrenia, and social anxiety disorder, but are not traditionally viewed as a cardinal characteristic of major depressive disorder (MDD). As a result, socioemotional behaviors are rarely used as endpoints in antidepressant clinical trials (Table 1).<sup>4, 5</sup> Yet consistent evidence shows that humans suffering from MDD withdraw socially and engage in behaviors that trigger hostility and rejection from interaction partners. These patients often rate their interpersonal difficulties as the most disabling aspect of the disease, which results in the erosion of social support and eventually isolation.<sup>6</sup>

Clinical studies that did include interpersonal functioning as a measurable outcome found that patients with remitted mood-related symptoms, but residual socioemotional deficits had a greater risk of relapse.<sup>7, 8</sup> In contrast, the ability to maintain social engagement or affiliation in the immediate aftermath of adverse life events was a robust early predictor of resilience and stable recovery from depressive episodes.<sup>9</sup> Furthermore, evidence that interpersonal psychotherapies specifically targeting social function are among the most efficacious forms of behavioral treatment for MDD points to socioemotional and interpersonal deficits as an essential pathogenic dimension of the disorder.<sup>10, 11</sup> Thus, a greater emphasis on the social facets of MDD symptomatology in neurobiological and preclinical studies is essential for successfully moving towards novel and more effective treatments.

# Human studies of socioemotional processing

Substantial progress has been made during the last decade in identifying circuits that mediate our ability to detect, appraise and respond to the emotional state of others using non-verbal cues. These emotion-relevant cues are delivered during human social interaction via multiple sensory modalities. Visually detected facial expressions are the most salient of these cues and the neural mechanisms underlying this form of emotional recognition have been the focus of most clinical investigations, <sup>12</sup> however other sensory modalities (e.g. vocal prosody, olfaction and biological motion) have also been emphasized in related research. <sup>13–17</sup> Though a thorough description of the circuits involved in each of these specific forms of sensory processing and their multimodal integration is beyond the scope of this paper, this topic has been the object of several recent reviews. <sup>18, 19</sup>

A widely used standardized laboratory task to evaluate socioaffective processing consists of exposing subjects to sets of composite images representing calibrated facial emotions (i.e. sad, happy, angry).<sup>20</sup> The subjective evaluation of the valence of these images is commonly assessed through the use of rating scales (Figure 1). Typical questions include "how many feet would you stay away from this person" or "how trustworthy is this person."<sup>21</sup> Other experiments have relied on implicit measures of emotion processing, such as reaction time in emotional Stroop tasks.<sup>22</sup> These studies have consistently shown that presentation durations as low as 50 milliseconds are sufficient for humans to extract basic emotional content from visual facial stimuli.<sup>21</sup> This affective information is processed unconsciously and takes precedence over other aspects of visual processing that allow individual recognition. This is revealed by the fact that subjects are often able to emit consistent valence judgments or inferences about complex traits of a facial stimulus (e.g. trustworthiness, dominance) while lacking the ability to describe the physical features of the face.<sup>21, 23</sup>

Socioemotional cues processed either unconsciously or explicitly have been shown to prompt valence-specific social tendencies in the perceiver, particularly the facilitation of approach or avoidance behavioral responses. These effects are evident in tasks where subjects are instructed to produce motor responses upon presentation of valenced facial stimuli.<sup>24</sup> For instance, in a recently developed approach-avoidance laboratory task, subjects must move towards or away from a visual social target presented on a screen using push-pull movements of a joystick (Figure 1).<sup>25</sup> Convergent results during this task show that healthy subjects avoid threatening stimuli faster than neutral pictures, even if these stimuli are not consciously detected. Functional imaging and neurophysiological methods such as EEG have been used in combination with these laboratory tasks to identify regions and neural networks implicated in the automatic and explicit processing of socioemotuional stimuli as well as during the execution of rapid approach-avoidance responses. Additional studies (mentioned in table 2) have utilized pharmacology or non-invasive neuromodulation methods such as transcranial magnetic stimulation (TMS) and direct current stimulation to provide causal evidence for the role of specific biological mechanisms, brain regions or circuits in healthy subjects and patients with affective disorders. <sup>25, 26</sup> Although early views posited a predominant role of subcortical alarm systems in the rapid processing of salient biologically relevant affective signals, current neurobiological models have evolved to emphasize the interactions between cortical and subcortical structures through multiple parallel bottom-up and top-down modulations.<sup>27</sup>

# Biased socioemotional processing and social avoidance in depression

Interpersonal deficits of MDD patients may arise in part from neurobiological abnormalities that disrupt the normal processing of socioaffective stimuli. 13, 28 This concept is based on the observation that subjects suffering from MDD and anxiety tend to preferentially attend to, process and remember negatively valenced information in their environment, whereas a bias towards positive information is generally observed in healthy subjects. 10 In social contexts, this cognitive bias leads to neutral or ambiguous social stimuli that are normally neglected by healthy subjects to break into consciousness as threats for MDD patients. 29–31 Indeed, during explicit rating tasks, MDD patients consistently rated ambiguous or negative

social stimuli more negatively than a healthy individual would. Administration of the Approach-Avoidance Task to social anxiety<sup>20</sup> or MDD<sup>31</sup> patients also revealed increased attention towards negative or threatening social stimuli (i.e. angry, sad) and a tendency to initiate withdrawal responses faster regardless of the emotional valence of the stimuli,<sup>23, 32–34</sup>. In real world interactions, this negatively biased socioemotional processing might lead MDD patients to misattribute negative intentions and motives to others and anticipate negative judgment or rejection with exaggerated defensiveness and withdrawal.<sup>32, 35</sup>

# Circuits underlying biased socioaffective processing in MDD: Key role of the ventromedial prefrontal cortex

In the last 10 years, consistent imaging and neurophysiological data have revealed the aberrant morphology and reactivity of the prefrontal cortex in MDD patients<sup>36–43</sup> and its normalization after successful chronic antidepressant treatment. 32, 44-50 In socioemotional processing studies, classical antidepressants rapidly remediate negative biases in MDD patients. This rapid action contrasts with the slow onset of the therapeutic effects on mood. It has been proposed that restored emotional processing may slowly foster improvement in mood by providing a more positive social environment in which the patient can relearn positive emotional associations.<sup>51</sup> Neural signatures of this neuroplastic process have been repeatedly identified in the ventromedial prefrontal cortex (vmPFC), a cluster of contiguous and functionally related areas in the primate frontal lobe comprised of Brodmann areas 10. 11, 12, 13, 14, 25 and 32.<sup>52–54</sup> It is important to note that the rationale for considering the vmPFC a single functionally homogenous structure remains controversial, <sup>55</sup> as several areas within the vmPFC cluster have distinct cytorachitecture and a unique pattern of input-output connections. <sup>56, 57</sup> Nevertheless, the vmPFC has been the focus of numerous studies in humans and animals leading to its classification as a control center for executive function, emotional valuation, autonomic control and social cognition. 58, 59

Functional abnormalities in the vmPFC may contribute in particular to the socioemotional facets of MDD symptoms, as this collection of regions shows striking overlap with the networks engaged when individuals make social approach or avoidance decisions. 60 Seminal observations that connected the vmPFC to social function were made by Damasio and colleagues who characterized the deficits of patients suffering from frontal lesions.<sup>61</sup> These individuals had abnormal autonomic responses to visual social stimuli and social decision-making deficits. A more recent eye tracking study in a cohort with similar lesions revealed a decreased capacity of these patients to attend to the specific features of the face that carry high emotional value such as the eyes or mouth. 62 These observations are in line with results in healthy individuals that show that the vmPFC is engaged during social attention and the processing of social sensory information. <sup>63, 64</sup> The essential role of the vmPFC in the effortful control of avoidance responses during social interactions was recently revealed by a study that used continuous theta bursts with TMS to non-invasively inhibit the activity of the vmPFC in subjects during an approach-avoidance task.<sup>25</sup> They found that subjects whose vmPFC was selectively inhibited committed more errors when they had to override automatic avoidance tendencies in order to apply rule-based responses.

The errors coincided with an upregulation in the activity of the insula and amygdala, two regions downstream of the vmPFC implicated in automatic processing.

It is likely that the vmPFC executes adaptive social behavioral decisions by computing valence and personal relevance of social stimuli and integrating this high-level multisensory information with the activity of subcortical limbic and brainstem structures, which control other elementary aspects of emotions such as autonomic and motor responses. Among the downstream regions controlled by the vmPFC are the nucleus accumbens and amygdala, as well mesencephalic and pontine monoaminergic nuclei containing dopamine and serotonin producing neurons. The following sections of this review focus on the circuitry underlying the reciprocal interaction of serotonin systems and the vmPFC during social interaction.

#### Serotonin influences on social behaviors

Serotonin (5-HT) is a phylogenetically ancient neuromodulator that exerts a wide range of influences on physiology and behavior through its action on 15 different receptors.<sup>67</sup> One of the most consistent roles attributed to 5-HT based on human and animal behavioral data relates to its influence on the expression of behaviors along the affiliative-aggressive axis. In humans, depleting 5-HT by reducing the dietary source of its precursor L-tryptophan, or boosting 5-HT via administration of selective serotonin reuptake inhibitors has respectively been shown to induce negative and positive shifts in social perception (Table 2).<sup>68–73</sup> Variants of genes that control 5-HT synaptic availability, such as the serotonin transporter (5-HTT)<sup>74–76</sup> and the 5-HT synthesizing enzyme tryptophan hydroxylase (TPH) <sup>77, 78</sup> have also been associated with alterations in socioemotional behaviors, such as differential sensitivity to the rewarding properties of social cues or reactivity to unfairness.<sup>79</sup> Differential corticolimbic responses to these cues have also been reported in carriers of these polymorphisms. <sup>76–78</sup> The effect of 5-HT on social behavior is likely to be mediated in part by its effects on neurons from the frontal cortex, as this area is one of the most enriched in serotonergic axons and 5-HT receptors. 5-HT modulates the excitability of cortical neurons and their discharge rate through the activation of several receptor subtypes, of which the 5-HT1<sub>A</sub>, 5-HT1<sub>B</sub>, 5-HT2<sub>A</sub>, and 5-HT3 play major roles. <sup>80–82</sup>

The primary serotonergic nuclei that innervate the prefrontal cortex in mammals are the median and dorsal raphe nucleus (MRN, DRN), two brainstem structures that provide the bulk of serotonergic afferents to the forebrain. Though the DRN and MRN are the two largest serotonergic cell groups, imaging them *in vivo* in humans remains a challenge due to their relatively small size and deep location in the brainstem.<sup>83, 84</sup> Only with the most recent advances in imaging methods has it become possible to visualize individual raphe nuclei in awake subjects and conduct probabilistic tractographic analyses of major fiber tracts to these brainstem nuclei.<sup>85</sup> This is an important issue as tractography indicates that brainstem nuclei are among key downstream regions likely to mediate the effects of vmPFC neurostimulation for the treatment of MDD.<sup>86</sup> Imaging studies together with postmortem investigations indicate that the distribution of raphe nuclei cell clusters and the architecture of the frontopontine fiber tract are well conserved across mammals.<sup>87, 88</sup> A small number of studies have begun identifying volumetric alterations and structural abnormalities in

brainstem fiber tracts that connect the raphe with the frontolimbic systems in MDD patients, although abnormalities in functional connectivity have yet to be reported.<sup>89, 90</sup> Nevertheless, combined use of PET and fMRI in healthy subjects have provided evidence for negative correlations between the density of 5-HT<sub>1A</sub> receptors in the DRN and amygdala reactivity to threats, suggesting that the capacity for DRN neuron autoinhibition predicts cotricolimbic reactivity. 91 Histological studies of the DRN in post-mortem tissues of MDD and suicide victims have revealed differences in levels of certain 5-HT-specific markers such as 5-HT<sub>1A</sub> and 5-HT<sub>2A</sub> receptors, <sup>92, 93</sup> TPH<sup>94–96</sup> and certain transcription factors controlling gene expression in 5-HT neurons. 97-100 However, evidence directly linking alterations of serotonergic systems with symptoms of mood disorders remains limited and confirmation will require replications in larger cohorts. <sup>86, 90</sup> The fact that over half of the DRN cell population is comprised of non-serotonergic cells<sup>101, 102</sup> including glutamatergic, <sup>103</sup> GABAergic, <sup>104</sup> and dopaminergic neurons <sup>105</sup> that in some cases exert opposite effects on behavior and connectivity complicates the interpretation of human data derived from imaging or tissue homogenates. 106, 107 Because studies in many different organisms ranging from insects and fishes <sup>108–111</sup> to rodents and primates have shown that altering levels of 5-HT influences the expression of affiliative-aggressive behaviors, <sup>112</sup> researchers have turned to these simpler models to more finely dissect the DRN microcircuit and its interaction with corticolimbic systems during socioaffective behaviors.

### Dissecting the cellular architecture of the vmPFC-DRN pathways in animals

There is strong evidence dating from the 1990's that the vmPFC is connected with the DRN via reciprocal monosynaptic projections, however the methods used to characterize these circuits have left areas of uncertainty regarding the cellular architecture of the pathway. Within the last few years, recent advances in mouse transgenics, viral tracing and connectomics methods have allowed for exponential progress in the characterization of this pathway. The first efforts to trace the vmPFC-DRN pathway relied on conventional retrograde and anterograde tracers such as biotinylated dextran amine and horseradish peroxidase. 87, 113-120 Most of these studies were conducted in the rat or non-human primates and in a few instances, combined tracing with ultrastructural methods or in vivo electrophysiology to test functional connectivity. These studies showed that the vmPFC sends robust direct afferents to the DRN, however the majority of axons anterogradely traced in the DRN represented fibers of passage and only 6% of traced axons formed synaptic contacts. 115, 121 Electron microscopy analysis of dual labeling in the DRN for mPFC afferents and Tph2 (5-HT neurons) or GABA revealed a greater frequency of mPFC terminals synapsing on GABA-labeled dendrites versus Tph2-labeled dendrites. 115 This suggested the existence of a disynaptic microcircuit mediating the inhibitory influence of the cortex on 5-HT neurons (Figure 2). Electrophysiological evidence supported this hypothetical organization when electrical stimulation of the rat mPFC caused a predominant reduction in the firing rates of DRN 5-HT neurons in vivo that was blocked by GABA antagonists. 114, 122 However, these earlier studies provided little information about the actual topographical organization of these cellular ensembles.

To overcome these limitations, our group recently generated two lines of transgenic mice and with fluorescently labeled GABA (*GAD2-tdTomato*) or 5-HT (*Pet1-tdTomato*)

neurons. 123 Using these mice, we mapped the relative distribution of GABA versus 5-HT cell types in the DRN and defined for the first time the morphological and electrophysiological signatures of genetically identified GABA neurons in the DRN. 123 We found little overlap between the areas occupied by 5-HT and GABA neurons in the DRN (Figure 2). Serotonergic neurons sit in the midline portion of the nucleus, whereas GABAergic neurons were found primarily at paramedial locations such as the lateral wings and ventrolateral periaqueductal grey (PAG). We also determined that GABA neurons are of smaller size than serotonergic neurons, have a greater excitability, and are often spontaneously active in slice preparations, while 5-HT neurons are silent <sup>104, 124</sup> Using optogenetics in these mice we demonstrated that DRN GABA neurons inhibit 5-HT neurons concentrated in the midline in the same slice. To examine the relative distribution of vmPFC afferents with regard to GABA and 5-HT cells, we developed a viral vector system to selectively express a fluorescent Synaptophysin-GFP (SynP-GFP) fusion protein in excitatory neurons of the vmPFC. 125 Because this protein is targeted to presynaptic densities, it acts as an anterograde tracer that labels synaptic contacts while excluding fibers of passage. Using this vector we found that synapses formed by vmPFC axons in the DRN are topographically distributed and follow a rostrocaudal gradient (Figure 2). In the rostral DRN, most vmPFC synapses were found in the lateral wings area overlapping with GABA neurons clusters. In contrast, vmPFC synapses were concentrated in the midline in the caudal DRN and overlapped with 5-HT neurons. This gradient suggests that vmPFC inputs are likely to disynaptically inhibit serotonergic output from the rostral DRN, but monosynaptically stimulate serotonergic output from the caudal DRN. 125 Because rostral and caudal DRN 5-HT neurons project to distinct forebrain targets, this hypothetical microcircuit places the DRN GABA population in a critical position to gate top-down influence from vmPFC on a key subset of mood-regulating rostral 5-HT neurons. 126-128

Our results showing regional overlap between SynP-GFP-labeled vmPFC axonal boutons and GABA or 5-HT neurons suggest the existence of synaptic contacts, but do not provide functional evidence for this connectivity. A number of approaches have been used to address this question. One takes advantage of the retrograde transsynaptic properties of genetically modified rabies viruses to map synaptic connections between identified cell-types in a network. Weissbourd et al. and Pollak Dorocic et al. recently used this type of rabies virusbased approach in combination with in situ hybridization and brain slice electrophysiology to comprehensively map and quantitate the origin of synaptic inputs to DRN serotonergic and GABAergic neurons. 127, 128 They utilized GAD2-Cre and SERT-Cre transgenic drivers to target GABA and 5-HT neurons respectively and employed two different forms of an avian virus receptor (TVA) to optimize their anatomical tracing of either long-range or local intra-DRN connectivity. Their results show that prefrontocortical inputs amount to roughly 15% of the total synaptic input received by DRN neurons. This suggests that the functional influence of cortical inputs on the DRN is likely to be weaker than certain subcortical inputs such as the hypothalamus (30% of input). Surprisingly, retrograde labeling of cortical neurons was denser when starter cells were DRN 5-HT neurons rather than GABA neurons, indicating, in contrast with our data, that the PFC preferentially innervates DRN 5-HT cells (10%) over GABA neurons (5%). In situ hybridization results showed that virtually all of these cortical inputs were glutamatergic, in agreement with previous tracing studies showing

that most DRN-projecting cortical neurons are localized in layer V, <sup>117, 118</sup> a typical location for subcortically projecting pyramidal neurons. <sup>57, 129, 130</sup> Approximately 2% of DRN inputs identified by Weissbourd *et al.* originated from the prelimbic cortex (PL) and less than 0.5% were from the infralimbic cortex (IL), two regions whose combination corresponds to the human vmPFC. These results are in line with earlier tracing studies showing that the PL is the predominant source of vmPFC inputs to the DRN. <sup>113, 116, 118</sup> Unexpectedly though, among all prefrontocortical areas, the insular cortex (3%) proved to be the major source of inputs to the DRN.

To test the functional architecture of the PFC-DRN pathway, several groups including our own have conducted *Channelrhopsin2*-assisted circuit mapping (CRACM). This analysis uses optogenetically evoked EPSCs as an index of functional connectivity between 2 regions. Our group was the first to report this type of analysis between the vmPFC and DRN. We expressed *Channelrhodopsin2* (*ChR2*) in vmPFC pyramidal neurons using an AAV virus driven by the *CaMKIIA* promoter and recorded from identified serotonergic and GABAergic neurons in the DRN of *tdTomato* reporter lines. Optogenetically activating vmPFC terminals reliably triggered *cFos* expression and laser-locked EPSCs in GABAergic neurons of the lateral wings, but not in midline 5-HT cells, suggesting that the former are the preferential functional targets of vmPFC projections. In contrast, using similar approaches, Weissbourd *et al.* and Pollak Dorocic *et al.* 127 found that cortical axon photoactivation was twice as efficient at triggering EPSCs in 5-HT neurons than in GABA neurons.

Given the gradient described above, inconsistencies between the results from ours and other's CRACM experiments investigating the PFC-DRN pathway might reflect distinctions in the rostrocaudal localization of the DRN cells selected for recording, as well as in the cortical areas targeted by viral injections (vmPFC in our study versus anterior cingulate in Weissbourd *et al.* versus anterior PFC in Pollak Dorocic *et al.*). In addition, differences may also be explained by technical factors such as differences in the viral vectors used to express *ChR2* (i.e. AAV serotypes and promoters) or strategies used to identify cells for recording (e.g. viral versus transgenic expression of *tdTomato*; *Pet1-Cre* versus *SERT-Cre* to identify 5-HT neurons). Although our understanding of the architecture of the vmPFC-DRN could still be improved by an even greater refinement in the definition of various cellular subsets in the DRN, the recent studies described above provide detailed groundwork to start examining the function and plasticity of these circuits *in vivo* and test their influence on socioaffective behaviors.

# Chronic Social Defeat Stress (CSDS): an animal model of socioemotional bias?

Modeling psychiatric disorders in animals poses major challenges,<sup>131</sup> however a number of animal procedures offer opportunities to isolate specific dimensions or endophenotypes of diseases in order to probe their underlying neurobiology. Chronic social defeat stress (CSDS) takes advantage of ethologically relevant threats in mice (i.e. dyadic aggressive interactions) to precipitate a multidimensional stress-induced syndrome that resembles certain features of clinical depression and comorbid anxiety disorders, including circadian

disruption, HPA axis dysregulation, anhedonia, and socioaffective deficits, each of which occur with distinct temporal dynamics. 132-136 The primary behavioral endpoint of the CSDS model is altered social approach-avoidance behavior. This variable is tested in a social interaction task where experimental mice must confront an unfamiliar social target contained in a wire mesh box and are evaluated for their decisions to approach or avoid the target (Figure 1). In many ways this task provides an animal parallel to the approach-avoidance laboratory task used in humans and described above. To induce avoidance, mice in the "defeat" experimental group are subjected to brief experiences of physical aggression from trained CD1 aggressors followed by continuous protected sensory contact with the aggressor (Figure 3). In contrast, control mice are housed peacefully among dyads of the same strain and live under sensory contact conditions replicating those of the defeated mice. For naïve mice, unfamiliar social targets appear positively valenced as virtually all undefeated control mice make the rapid choice to approach the unfamiliar target during the social interaction task and remain engaged in social interaction during the entire duration of the test. In contrast, most defeated mice (roughly 70%, termed vulnerable) develop a social aversion, characterized by a delayed latency to start investigating the target followed by vigorous retreat and persistent immobility in the corners of the arena (Figure 1).

Our experiments have determined that the avoidance response is a "learned behavior" and its encoding depends critically on the daily sensory contact period that follows each episode of physical defeat. 123 We determined that 20 minutes of post-defeat sensory exposure each day is necessary and sufficient to trigger a maximal avoidance response. This is suggestive of an associative process that occurs during the early phase of the post defeat sensory exposure and leads to the consolidation of the social avoidance response. During the social interaction test, the expression of the avoidance behavior is triggered by social cues emitted by the behaviorally active social target. This was deduced from the observation that defeated mice do not avoid an inanimate novel object or an anesthetized social target. 137 However the avoidance response is not conditioned to cues that are specific of the aggressors, as avoidance generalizes to any unfamiliar social target, including mice from a different strain and devoid of territorial aggression. 137 The exact nature of these sensory cues is currently unknown, but could involve a combination of visual, olfactory and ultrasonic cues. 138 The avoidance response is long lasting and does not extinguish upon repeated testing. 137, 139, 140 CSDS-induced avoidance can thus be conceptualized as a learned aversive response to socioemotional cues that is overgeneralized and overconsolidated. The reliance of CSDS on this learned socioemotional response instead of a response to an innately aversive stimulus (i.e. shock or forced water immersion used in other antidepressant screens) is a major distinguishing feature of the model. This difference implies that the brain systems and plasticity mechanisms engaged by CSDS are likely to differ considerably from those at play in other models of depression, such as the learned helplessness paradigm, the tail suspension test or the forced swim test. 141, 142 The CSDS model does not respond to acute administration of benzodiazepine anxiolytics, but is responsive to SSRIs, tricyclic antidepressants, ketamine and DBS with clinically relevant time-courses. 137, 143

The primary sensory modalities humans and rodents rely on to execute rapid approachavoidance decisions are very likely to differ. Vision is predominant in humans while olfaction is likely to be the primary sensory process at play in the murine test. However, it is

reasonable to draw a translational parallel regarding the executive processes at play. A subpopulation in each cohort of defeated mice (roughly 30%, termed resilient) will maintain social approach despite experiencing similar level of aggression. Like in the human approach-avoidance task, the sustained approach behavior of these resilient mice can be interpreted as an ability to actively overcome automatic avoidance responses triggered by the ambiguous social target. The comparison of cohorts stratified into resilient and susceptible subpopulations (Figure 1) allows the dissection of the neurobiological mechanisms underlying active adaptation versus vulnerability to the effects of psychosocial stress.

Since our original description of the CSDS model, the work from several groups applying innovative approaches such as optogenetics and multiregion in vivo recording has contributed to an increased molecular and circuit-level understanding of the mechanisms underlying resilient and vulnerable phenotypes (Table 2). In parallel with the human literature on the role of stress hormones in the development of socioaffective bias, <sup>144</sup> glucocorticoids have been shown to mediate the development of social avoidance, in part by acting on adult-born neurons 145 as well as dopaminergic and serotonergic systems. 146-148 Tph2 knockin mice with a mutation analogous to the rare human R441H variant of the TPH2 gene, which leads to an 80% reduction in 5-HT levels, have corticolimbic hypersynchrony and increased vulnerability to various stressors including social defeat. 149 This genetic effect can be mimicked by pharmacological depletion of 5-HT, <sup>150, 151</sup> an effect that resembles the influence of tryptophan depletion in approach-avoidance tests and socioemotional processing in human subjects. <sup>22, 68, 152</sup> High frequency electrical stimulation of the vmPFC, a treatment that enhances 5-HT release in the forebrain, <sup>153</sup> has been shown to restore social approach behavior in defeated mice. 154 This effect replicates the action of vmPFC DBS on negative bias and mood related symptoms in MDD patients. 155, 156

In all, these results underlie the striking cross-species parallels that exist regarding the roles of the vmPFC and serotonergic systems in socioemotional responses. This supports the hypothesis that the plasticity of the vmPFC-DRN pathway that links these two systems may contribute to the encoding or expression of social avoidance across species.

# Social-experience-induced structural and functional plasticity of the vmPFC-DRN microcircuits

We have reported that exposure to CSDS followed by protected sensory contact with an aggressor induces the accumulation of *cFos* and *FosB* in the DRN.<sup>123, 157</sup> These markers of activity-dependent neuroplasticity were expressed specifically by a cluster of GABA cells in the lateral wings of the DRN, the area that receives the densest axonal input from the vmPFC. This suggests that neuroplastic changes occur in DRN GABA neurons might be driven by vmPFC inputs and facilitate in encoding a shift in social valence after social defeat. To test this hypothesis we examined whether the electrical and morphological properties of serotonergic and GABAergic neurons were altered in the DRN after CSDS in a manner that correlates with the expression of social avoidance. After stratifying mice as resilient or susceptible after 10 days of CSDS, we performed whole-cell recordings on fluorescently tagged 5-HT or GABA neurons from transgenic mice to unambiguously

identify the cellular populations. In susceptible mice that developed social avoidance, we found that DRN GABAergic neurons in the lateral wings were hyperexcitable and had increased glutamatergic input compared to controls, whereas serotonergic neurons showed opposite changes, namely, a reduced intrinsic excitability and increased GABAergic input (Figure 4). 123 Furthermore, morphological analysis of filled 5-HT neurons revealed an increase in size and complexity of the proximal dendrites in vulnerable mice. 146 Lastly, we also observed a drastic reduction in the number and density of *SynP-GFP* labeled varicosities along serotonergic axons, a sensitive index of presynaptic plasticity and release efficacy. 154 None of these electrophysiological or structural changes were observed in resilient mice.

Given our previous observations that DRN GABA neurons receive synaptic contacts from the vmPFC and inhibit nearby 5-HT neurons, the results summarized above suggest that CSDS may lead to maladaptive sensitization of the disynaptic inhibition of 5-HT cells while simultaneously depressing the intrinsic excitability of 5-HT neurons. <sup>123</sup> In line with this prediction, microdialysis experiments ongoing in our lab indicate that CSDS-resilient mice respond to sensory re-exposure to an aggressor with a dramatic increase in 5-HT release in the vmPFC whereas vulnerable mice do not show this response. This suggests that activation of serotonergic neurons and 5-HT release in the PFC and possibly other regions may contribute to the ability of resilient mice to maintain social approach after CSDS (Figure 5). This mechanism would be in line with recent evidence that the firing of certain subsets of serotonergic neurons tonically increases in vivo during expectation of a reward 158, 159 and mediates social reward signals through release of 5-HT and activation of 5-HT1<sub>B</sub> receptors on corticostriatal axon terminals in the nucleus accumbens (NAc). 160 Several recent studies have also implicated subsets of ventral tegmental area (VTA) projecting DRN serotonergic and glutamatergic neurons in the encoding of reward. <sup>161</sup> These findings provide mechanistic bases to understand the reduced social motivation and negative shifts in social perception observed after depleting 5-HT (Table 2).<sup>22, 73, 95</sup>

To determine whether vmPFC-driven disynaptic inhibition of 5-HT neurons was sufficient to promote social avoidance we performed in vivo photoactivation of glutamatergic vmPFC terminals in the DRN of mice that were exposed only to sensory contact with a CD1 aggressor through a plexiglass partition. Even in the absence of physical aggression, photoactivation resulted in the development of subsequent avoidance responses that recapitulated the behavior of defeated mice. 125 To test whether this circuit mechanism was necessary for the development of social avoidance, we photoinhibited either the vmPFC inputs to the DRN or DRN GABA neurons during the daily post-defeat sensory contact period. Under these conditions we found that mice exposed to CSDS did not become avoidant, confirming the causal role of the vmPFC-DRN disynaptic circuit in the encoding of social avoidance. <sup>123, 125</sup> Interestingly, in contrast with this key role in the associative process leading to avoidance, altering the top-down inhibitory control of raphe circuits by the vmPFC did not appear to affect the expression of a previously learned avoidance response. Indeed, acute optogenetic inhibition of vmPFC inputs to the DRN or silencing DRN GABAergic neurons during the social interaction task did not inhibit avoidance of mice previously defeated without stimulation. 123, 125 On the other hand, chronic treatment

with vmPFC DBS, when applied for 5 hours daily over a week, did restore approach behaviors of mice previously characterized as vulnerable. This effect was obtained using parameters that replicated clinically used stimulation conditions and was associated with a desensitization of DRN GABA neurons and a restoration of the intrinsic excitability and morphology of 5-HT neurons. <sup>154</sup> Furthermore, DBS potentiated the direct glutamatergic drive of 5-HT cells, as revealed by an enhanced frequency of mEPSCs and increased density of PSD95 puncta on 5-HT dendrites above baseline. Surprisingly, DBS also restored the density of 5-HT axonal boutons in a projection-specific manner, leading to a greater serotonin release capacity in the vmPFC. The fact that high frequency stimulation of the vmPFC by DBS can restore social approach once avoidance has already been stabilized, but not local bidirectional optogenetic modulation of vmPFC axons in the DRN, suggests that this therapeutic effect may require the combined activation of several downstream targets of the vmPFC, which in turn may converge on the DRN.

Altogether our results thus confirm the hypothesis that CSDS sensitizes vmPFC driven disynaptic inhibition of 5-HT neurons, thereby reducing 5-HT output in innervated regions to alter the affective impact of an adverse social experience and promoting generalized social aversion in vulnerable mice (Figure 4). Although the mechanism whereby resilient mice are protected from the effects of CSDS and evade sensitization of vmPFC-DRN pathway remains unclear, it is possible to formulate hypotheses based on recent studies using the same model. The Dzirasa group recently combined optogenetics and in vivo multiregion recordings in mice to show that stimulation of the vmPFC resulted in evoked potentials in numerous subcortical and limbic structures, including the amygdala and NAc as well as in aminergic neuromodulatory centers like the DRN and VTA. 162 Surprisingly, vmPFC evoked responses occurred significantly faster in the DRN that in limbic regions despite this structure being more anatomically distal from the vnPFC. This faster transmission velocity in the vmPFC-DRN pathway places the DRN in a position to modulate threat sensitivity by acting as a closed-loop regulator that can synchronize or desynchronize corticolimbic networks by independently controlling excitability of the hub (i.e. vmPFC) and its distributed downstream targets. Further work by the Dzirasa laboratory has also shown that naturally occurring differences in corticolimbic synchrony correlates with trait vulnerability to CSDS, with greater phase locking between the PFC and amygdala predicting increased vulnerability. 163 Interestingly, as mentioned above, animal and human studies have shown that 5-HT is a major regulator of vmPFC-amygdala coupling and reducing 5-HT using genetic mouse models of 5-HT deficiency or pharmacological treatments has been shown to favor cortico-amygdala hypersynchrony and enhance vulnerability to social defeat (Figure 5).<sup>164</sup>

## **Concluding remarks**

Evidence grows to suggest that biased socioemotional processing is a core cognitive component of MDD that contributes to interpersonal deficits in this disorder. Socioemotional processing is under key control by the prefrontal cortex and serotonergic systems in part via a direct vmPFC-DRN circuit whose plasticity is targeted by long-term antidepressant treatment and DBS. Relying on translationally meaningful models for the recapitulation of socioaffective symptoms of MDD, studies from our lab and others

summarized in this review have started establishing a cellular blueprint for the organization of this pathway and identifying key neuroplastic events that mediate normal and pathological regulation of socioaffective function. An important next step for future studies will be to define how vmPFC-driven neuroplastic adaptations of specific subsets of serotonergic neurons translate into modifications of corticolimbic circuit dynamics. A further goal will be to delve deeper into the molecular mechanisms that underlie the synaptic adaptations of DRN neurons. Our hope is that these approaches will offer new insights into more efficacious pharmacological or circuit-based therapeutics that target the socioemotional aspects of depressive symptomology.

#### References

- 1. Oliveira RF. Social behavior in context: Hormonal modulation of behavioral plasticity and social competence. Integr Comp Biol. 2009; 49:423–440. [PubMed: 21665831]
- 2. Taborsky B, Oliveira RF. Social competence: an evolutionary approach. Trends Ecol Evol. 2012; 27:679–688. [PubMed: 23040461]
- 3. Whitton SW, Larson JJ, Hauser ST. Depressive symptoms and bias in perceived social competence among young adults. J Clin Psychol. 2008; 64:791–805. [PubMed: 18442044]
- 4. Bech P. Social functioning: should it become an endpoint in trials of antidepressants? CNS Drugs. 2005; 19:313–324. [PubMed: 15813645]
- 5. Billeke P, Boardman S, Doraiswamy PM. Social cognition in major depressive disorder: A new paradigm? Translational Neuroscience. 2013; 4:437–447.
- Hames JL, Hagan CR, Joiner TE. Interpersonal processes in depression. Annu Rev Clin Psychol. 2013; 9:355–377. [PubMed: 23297787]
- 7. Paykel ES. Partial remission, residual symptoms, and relapse in depression. Dialogues Clin Neurosci. 2008; 10:431–437. [PubMed: 19170400]
- 8. Hirschfeld RMA, Dunner DL, Keitner G, Klein DN, Koran LM, Kornstein SG, Markowitz JC, Miller I, Nemeroff CB, Ninan PT, Rush AJ, Schatzberg AF, Thase ME, Trivedi MH, Borian FE, Crits-Christoph P, Keller MB. Does psychosocial functioning improve independent of depressive symptoms? a comparison of nefazodone, psychotherapy, and their combination. Biological Psychiatry. 2002; 51:123–133. [PubMed: 11822991]
- Mathews A, MacLeod C. Cognitive vulnerability to emotional disorders. Annu Rev Clin Psychol. 2005; 1:167–195. [PubMed: 17716086]
- 10. Beck AT. The evolution of the cognitive model of depression and its neurobiological correlates. Am J Psychiatry. 2008; 165:969–977. [PubMed: 18628348]
- 11. Markowitz JC, Weissman MM. Interpersonal psychotherapy: principles and applications. World Psychiatry. 2004; 3:136–139. [PubMed: 16633477]
- 12. Brecht M, Freiwald WA. The many facets of facial interactions in mammals. Curr Opin Neurobiol. 2012; 22:259–266. [PubMed: 22209040]
- 13. Chang SW, Gariepy JF, Platt ML. Neuronal reference frames for social decisions in primate frontal cortex. Nat Neurosci. 2013; 16:243–250. [PubMed: 23263442]
- 14. Leitman DI, Wolf DH, Ragland JD, Laukka P, Loughead J, Valdez JN, Javitt DC, Turetsky BI, Gur RC. "It's Not What You Say, But How You Say it": A Reciprocal Temporo-frontal Network for Affective Prosody. Front Hum Neurosci. 2010; 4:19. [PubMed: 20204074]
- 15. Oosterhof NN, Todorov A. Shared perceptual basis of emotional expressions and trustworthiness impressions from faces. Emotion. 2009; 9:128–133. [PubMed: 19186926]
- Allison T, Puce A, McCarthy G. Social perception from visual cues: role of the STS region. Trends Cogn Sci. 2000; 4:267–278. [PubMed: 10859571]
- 17. Brancucci A, Lucci G, Mazzatenta A, Tommasi L. Asymmetries of the human social brain in the visual, auditory and chemical modalities. Philos Trans R Soc Lond B Biol Sci. 2009; 364:895–914. [PubMed: 19064350]

 Fossati P. Neural correlates of emotion processing: from emotional to social brain. Eur Neuropsychopharmacol. 2012; 22(Suppl 3):S487–491. [PubMed: 22959113]

- 19. Pessoa L, Adolphs R. Emotion processing and the amygdala: from a 'low road' to 'many roads' of evaluating biological significance. Nat Rev Neurosci. 2010; 11:773–783. [PubMed: 20959860]
- 20. Heuer K, Rinck M, Becker ES. Avoidance of emotional facial expressions in social anxiety: The Approach-Avoidance Task. Behav Res Ther. 2007; 45:2990–3001. [PubMed: 17889827]
- 21. Todorov A. Evaluating faces on trustworthiness: an extension of systems for recognition of emotions signaling approach/avoidance behaviors. Ann N Y Acad Sci. 2008; 1124:208–224. [PubMed: 18400932]
- 22. Munafo MR, Hayward G, Harmer C. Selective processing of social threat cues following acute tryptophan depletion. J Psychopharmacol. 2006; 20:33–39. [PubMed: 16204324]
- 23. Morris JS, Ohman A, Dolan RJ. Conscious and unconscious emotional learning in the human amygdala. Nature. 1998; 393:467–470. [PubMed: 9624001]
- 24. Bargh JA, Chen M, Burrows L. Automaticity of social behavior: direct effects of trait construct and stereotype-activation on action. J Pers Soc Psychol. 1996; 71:230–244. [PubMed: 8765481]
- 25. Volman I, Roelofs K, Koch S, Verhagen L, Toni I. Anterior prefrontal cortex inhibition impairs control over social emotional actions. Curr Biol. 2011; 21:1766–1770. [PubMed: 22000109]
- Novakovic V, Sher L, Lapidus KA, Mindes J, JAG, Yehuda R. Brain stimulation in posttraumatic stress disorder. Eur J Psychotraumatol. 2011; 2
- Pantazatos SP, Talati A, Pavlidis P, Hirsch J. Cortical functional connectivity decodes subconscious, task-irrelevant threat-related emotion processing. Neuroimage. 2012; 61:1355– 1363. [PubMed: 22484206]
- 28. Cahn DD, Frey LR. Listeners' perceived verbal and nonverbal behaviors associated with communicators' perceived understanding and misunderstanding. Percept Mot Skills. 1992; 74:1059–1064. [PubMed: 1501968]
- 29. Davidson RJ, Ekman P, Saron CD, Senulis JA, Friesen WV. Approach-withdrawal and cerebral asymmetry: emotional expression and brain physiology. I. J Pers Soc Psychol. 1990; 58:330–341. [PubMed: 2319445]
- 30. Lang PJ, Bradley MM, Cuthbert BN. Emotion, attention, and the startle reflex. Psychol Rev. 1990; 97:377–395. [PubMed: 2200076]
- Seidel EM, Habel U, Finkelmeyer A, Schneider F, Gur RC, Derntl B. Implicit and explicit behavioral tendencies in male and female depression. Psychiatry Res. 2010; 177:124–130. [PubMed: 20199811]
- 32. Derntl B, Seidel EM, Eickhoff SB, Kellermann T, Gur RC, Schneider F, Habel U. Neural correlates of social approach and withdrawal in patients with major depression. Soc Neurosci. 2011; 6:482–501. [PubMed: 21777105]
- 33. Godard J, Grondin S, Baruch P, Lafleur MF. Psychosocial and neurocognitive profiles in depressed patients with major depressive disorder and bipolar disorder. Psychiatry Res. 2011; 190:244–252. [PubMed: 21764461]
- 34. Radke S, Guths F, Andre JA, Muller BW, de Bruijn ER. In action or inaction? Social approach-avoidance tendencies in major depression. Psychiatry Res. 2014; 219:513–517. [PubMed: 25060832]
- 35. Fava M, Kendler KS. Major depressive disorder. Neuron. 2000; 28:335–341. [PubMed: 11144343]
- 36. Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood disorders: implications for neurocircuitry models of depression. Brain Struct Funct. 2008; 213:93–118. [PubMed: 18704495]
- 37. Hamilton JP, Etkin A, Furman DJ, Lemus MG, Johnson RF, Gotlib IH. Functional neuroimaging of major depressive disorder: a meta-analysis and new integration of base line activation and neural response data. Am J Psychiatry. 2012; 169:693–703. [PubMed: 22535198]
- 38. Liao C, Feng Z, Zhou D, Dai Q, Xie B, Ji B, Wang X, Wang X. Dysfunction of fronto-limbic brain circuitry in depression. Neuroscience. 2012; 201:231–238. [PubMed: 22119640]
- Mayberg HS. Limbic-cortical dysregulation: a proposed model of depression. J Neuropsychiatry Clin Neurosci. 1997; 9:471–481. [PubMed: 9276848]

40. Mayberg HS. Modulating dysfunctional limbic-cortical circuits in depression: towards development of brain-based algorithms for diagnosis and optimised treatment. Br Med Bull. 2003; 65:193–207. [PubMed: 12697626]

- 41. de Kwaasteniet B, Ruhe E, Caan M, Rive M, Olabarriaga S, Groefsema M, Heesink L, van Wingen G, Denys D. Relation between structural and functional connectivity in major depressive disorder. Biol Psychiatry. 2013; 74:40–47. [PubMed: 23399372]
- 42. Johnstone T, van Reekum CM, Urry HL, Kalin NH, Davidson RJ. Failure to regulate: counterproductive recruitment of top-down prefrontal-subcortical circuitry in major depression. J Neurosci. 2007; 27:8877–8884. [PubMed: 17699669]
- 43. Murray EA, Wise SP, Drevets WC. Localization of dysfunction in major depressive disorder: prefrontal cortex and amygdala. Biol Psychiatry. 2011; 69:e43–54. [PubMed: 21111403]
- 44. Smith R, Chen K, Baxter L, Fort C, Lane RD. Antidepressant effects of sertraline associated with volume increases in dorsolateral prefrontal cortex. J Affect Disord. 2013; 146:414–419. [PubMed: 23017544]
- 45. Vasic N, Walter H, Hose A, Wolf RC. Gray matter reduction associated with psychopathology and cognitive dysfunction in unipolar depression: a voxel-based morphometry study. J Affect Disord. 2008; 109:107–116. [PubMed: 18191459]
- 46. Bellani M, Dusi N, Yeh PH, Soares JC, Brambilla P. The effects of antidepressants on human brain as detected by imaging studies. Focus on major depression. Prog Neuropsychopharmacol Biol Psychiatry. 2011; 35:1544–1552. [PubMed: 21138750]
- 47. Delaveau P, Jabourian M, Lemogne C, Guionnet S, Bergouignan L, Fossati P. Brain effects of antidepressants in major depression: a meta-analysis of emotional processing studies. J Affect Disord. 2011; 130:66–74. [PubMed: 21030092]
- 48. Drevets WC, Bogers W, Raichle ME. Functional anatomical correlates of antidepressant drug treatment assessed using PET measures of regional glucose metabolism. Eur Neuropsychopharmacol. 2002; 12:527–544. [PubMed: 12468016]
- Romera I, Perez V, Menchon JM, Delgado-Cohen H, Polavieja P, Gilaberte I. Social and occupational functioning impairment in patients in partial versus complete remission of a major depressive disorder episode. A six-month prospective epidemiological study. Eur Psychiatry. 2010; 25:58–65. [PubMed: 19553092]
- Harmer CJ, Bhagwagar Z, Perrett DI, Vollm BA, Cowen PJ, Goodwin GM. Acute SSRI administration affects the processing of social cues in healthy volunteers. Neuropsychopharmacology. 2003; 28:148–152. [PubMed: 12496951]
- 51. Pringle A, Browning M, Cowen PJ, Harmer CJ. A cognitive neuropsychological model of antidepressant drug action. Prog Neuropsychopharmacol Biol Psychiatry. 2011; 35:1586–1592. [PubMed: 20673783]
- 52. Der-Avakian A, Markou A. The neurobiology of anhedonia and other reward-related deficits. Trends Neurosci. 2012; 35:68–77. [PubMed: 22177980]
- 53. Finger EC, Marsh AA, Mitchell DG, Reid ME, Sims C, Budhani S, Kosson DS, Chen G, Towbin KE, Leibenluft E, Pine DS, Blair JR. Abnormal ventromedial prefrontal cortex function in children with psychopathic traits during reversal learning. Arch Gen Psychiatry. 2008; 65:586–594. [PubMed: 18458210]
- 54. Phan KL, Wager T, Taylor SF, Liberzon I. Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. Neuroimage. 2002; 16:331–348. [PubMed: 12030820]
- Haber SN, Knutson B. The reward circuit: linking primate anatomy and human imaging. Neuropsychopharmacology. 2010; 35:4–26. [PubMed: 19812543]
- 56. Heidbreder CA, Groenewegen HJ. The medial prefrontal cortex in the rat: evidence for a dorsoventral distinction based upon functional and anatomical characteristics. Neurosci Biobehav Rev. 2003; 27:555–579. [PubMed: 14599436]
- 57. Shepherd GM. Corticostriatal connectivity and its role in disease. Nat Rev Neurosci. 2013; 14:278–291. [PubMed: 23511908]
- 58. Fuster JM. Executive frontal functions. Exp Brain Res. 2000; 133:66-70. [PubMed: 10933211]

59. Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. Annu Rev Neurosci. 2001; 24:167–202. [PubMed: 11283309]

- 60. Barbey AK, Colom R, Paul EJ, Chau A, Solomon J, Grafman JH. Lesion mapping of social problem solving. Brain. 2014; 137:2823–2833. [PubMed: 25070511]
- 61. Damasio AR, Tranel D, Damasio H. Individuals with sociopathic behavior caused by frontal damage fail to respond autonomically to social stimuli. Behav Brain Res. 1990; 41:81–94. [PubMed: 2288668]
- Wolf RC, Philippi CL, Motzkin JC, Baskaya MK, Koenigs M. Ventromedial prefrontal cortex mediates visual attention during facial emotion recognition. Brain. 2014; 137:1772–1780. [PubMed: 24691392]
- 63. Adolphs R. Social attention and the ventromedial prefrontal cortex. Brain. 2014; 137:1572–1574. [PubMed: 24776972]
- 64. Winecoff A, Clithero JA, Carter RM, Bergman SR, Wang L, Huettel SA. Ventromedial prefrontal cortex encodes emotional value. J Neurosci. 2013; 33:11032–11039. [PubMed: 23825408]
- 65. Boksem MA, Smolders R, De Cremer D. Social power and approach-related neural activity. Soc Cogn Affect Neurosci. 2012; 7:516–520. [PubMed: 19304842]
- Quirin M, Meyer F, Heise N, Kuhl J, Kustermann E, Struber D, Cacioppo JT. Neural correlates of social motivation: an fMRI study on power versus affiliation. Int J Psychophysiol. 2013; 88:289– 295. [PubMed: 22841755]
- 67. Graeff FG, Guimaraes FS, De Andrade TG, Deakin JF. Role of 5-HT in stress, anxiety, and depression. Pharmacol Biochem Behav. 1996; 54:129–141. [PubMed: 8728550]
- 68. Bilderbeck AC, McCabe C, Wakeley J, McGlone F, Harris T, Cowen PJ, Rogers RD. Serotonergic activity influences the cognitive appraisal of close intimate relationships in healthy adults. Biol Psychiatry. 2011; 69:720–725. [PubMed: 21396628]
- Crockett MJ, Apergis-Schoute A, Herrmann B, Lieberman MD, Muller U, Robbins TW, Clark L. Serotonin modulates striatal responses to fairness and retaliation in humans. J Neurosci. 2013; 33:3505–3513. [PubMed: 23426678]
- Crockett MJ, Clark L, Apergis-Schoute AM, Morein-Zamir S, Robbins TW. Serotonin modulates the effects of Pavlovian aversive predictions on response vigor. Neuropsychopharmacology. 2012; 37:2244–2252. [PubMed: 22643930]
- 71. Harmer CJ, Shelley NC, Cowen PJ, Goodwin GM. Increased positive versus negative affective perception and memory in healthy volunteers following selective serotonin and norepinephrine reuptake inhibition. Am J Psychiatry. 2004; 161:1256–1263. [PubMed: 15229059]
- 72. Robinson OJ, Overstreet C, Allen PS, Letkiewicz A, Vytal K, Pine DS, Grillon C. The role of serotonin in the neurocircuitry of negative affective bias: serotonergic modulation of the dorsal medial prefrontal-amygdala 'aversive amplification' circuit. Neuroimage. 2013; 78:217–223. [PubMed: 23583742]
- 73. Young SN. The effect of raising and lowering tryptophan levels on human mood and social behaviour. Philos Trans R Soc Lond B Biol Sci. 2013; 368:20110375. [PubMed: 23440461]
- Miller JM, Hesselgrave N, Ogden RT, Sullivan GM, Oquendo MA, Mann JJ, Parsey RV. Positron emission tomography quantification of serotonin transporter in suicide attempters with major depressive disorder. Biol Psychiatry. 2013; 74:287–295. [PubMed: 23453288]
- 75. Verhagen M, van der Meij A, Janzing JG, Arias-Vasquez A, Buitelaar JK, Franke B. Effect of the 5-HTTLPR polymorphism in the serotonin transporter gene on major depressive disorder and related comorbid disorders. Psychiatr Genet. 2009; 19:39–44. [PubMed: 19125107]
- Volman I, Verhagen L, den Ouden HE, Fernandez G, Rijpkema M, Franke B, Toni I, Roelofs K. Reduced serotonin transporter availability decreases prefrontal control of the amygdala. J Neurosci. 2013; 33:8974–8979. [PubMed: 23699508]
- 77. Brown SM, Peet E, Manuck SB, Williamson DE, Dahl RE, Ferrell RE, Hariri AR. A regulatory variant of the human tryptophan hydroxylase-2 gene biases amygdala reactivity. Mol Psychiatry. 2005; 10:884–888. 805. [PubMed: 16044172]
- 78. Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR. Serotonin transporter genetic variation and the response of the human amygdala. Science. 2002; 297:400–403. [PubMed: 12130784]

79. Fisher PM, Hariri AR. Identifying serotonergic mechanisms underlying the corticolimbic response to threat in humans. Philos Trans R Soc Lond B Biol Sci. 2013; 368:20120192. [PubMed: 23440464]

- 80. Cervantes MC, Delville Y. Serotonin 5-HT1A and 5-HT3 receptors in an impulsive-aggressive phenotype. Behav Neurosci. 2009; 123:589–598. [PubMed: 19485565]
- 81. Clark MS, Neumaier JF. The 5-HT1B receptor: behavioral implications. Psychopharmacol Bull. 2001; 35:170–185. [PubMed: 12397864]
- 82. Fisher PM, Price JC, Meltzer CC, Moses-Kolko EL, Becker C, Berga SL, Hariri AR. Medial prefrontal cortex serotonin 1A and 2A receptor binding interacts to predict threat-related amygdala reactivity. Biol Mood Anxiety Disord. 2011; 1:2. [PubMed: 22738071]
- 83. Kranz GS, Hahn A, Savli M, Lanzenberger R. Challenges in the differentiation of midbrain raphe nuclei in neuroimaging research. Proc Natl Acad Sci U S A. 2012; 109:E2000. [PubMed: 22711836]
- 84. Son YD, Cho ZH, Choi EJ, Kim JH, Kim HK, Lee SY, Chi JG, Park CW, Kim JH, Kim YB. Individually differentiated serotonergic raphe nuclei measured with brain PET/MR imaging. Radiology. 2014; 272:541–548. [PubMed: 24654972]
- 85. Ford AA, Colon-Perez L, Triplett WT, Gullett JM, Mareci TH, Fitzgerald DB. Imaging white matter in human brainstem. Front Hum Neurosci. 2013; 7:400. [PubMed: 23898254]
- 86. Lehman JF, Greenberg BD, McIntyre CC, Rasmussen SA, Haber SN. Rules ventral prefrontal cortical axons use to reach their targets: implications for diffusion tensor imaging tractography and deep brain stimulation for psychiatric illness. J Neurosci. 2011; 31:10392–10402. [PubMed: 21753016]
- 87. Freedman LJ, Insel TR, Smith Y. Subcortical projections of area 25 (subgenual cortex) of the macaque monkey. J Comp Neurol. 2000; 421:172–188. [PubMed: 10813780]
- 88. Jbabdi S, Lehman JF, Haber SN, Behrens TE. Human and monkey ventral prefrontal fibers use the same organizational principles to reach their targets: tracing versus tractography. J Neurosci. 2013; 33:3190–3201. [PubMed: 23407972]
- 89. Song YJ, Korgaonkar MS, Armstrong LV, Eagles S, Williams LM, Grieve SM. Tractography of the brainstem in major depressive disorder using diffusion tensor imaging. PLoS One. 2014; 9:e84825. [PubMed: 24465436]
- 90. Gutman DA, Holtzheimer PE, Behrens TE, Johansen-Berg H, Mayberg HS. A tractography analysis of two deep brain stimulation white matter targets for depression. Biol Psychiatry. 2009; 65:276–282. [PubMed: 19013554]
- 91. Fisher PM, Meltzer CC, Ziolko SK, Price JC, Moses-Kolko EL, Berga SL, Hariri AR. Capacity for 5-HT1A-mediated autoregulation predicts amygdala reactivity. Nat Neurosci. 2006; 9:1362–1363. [PubMed: 17013380]
- 92. Lee HY, Tae WS, Yoon HK, Lee BT, Paik JW, Son KR, Oh YW, Lee MS, Ham BJ. Demonstration of decreased gray matter concentration in the midbrain encompassing the dorsal raphe nucleus and the limbic subcortical regions in major depressive disorder: an optimized voxelbased morphometry study. J Affect Disord. 2011; 133:128–136. [PubMed: 21546094]
- Stockmeier CA, Dilley GE, Shapiro LA, Overholser JC, Thompson PA, Meltzer HY. Serotonin receptors in suicide victims with major depression. Neuropsychopharmacology. 1997; 16:162– 173. [PubMed: 9015799]
- 94. Boldrini M, Underwood MD, Mann JJ, Arango V. More tryptophan hydroxylase in the brainstem dorsal raphe nucleus in depressed suicides. Brain Res. 2005; 1041:19–28. [PubMed: 15804496]
- 95. Passamonti L, Crockett MJ, Apergis-Schoute AM, Clark L, Rowe JB, Calder AJ, Robbins TW. Effects of acute tryptophan depletion on prefrontal-amygdala connectivity while viewing facial signals of aggression. Biol Psychiatry. 2012; 71:36–43. [PubMed: 21920502]
- 96. Rogers RD. The roles of dopamine and serotonin in decision making: evidence from pharmacological experiments in humans. Neuropsychopharmacology. 2011; 36:114–132. [PubMed: 20881944]
- 97. Albert PR. Transcriptional regulation of the 5-HT1A receptor: implications for mental illness. Philos Trans R Soc Lond B Biol Sci. 2012; 367:2402–2415. [PubMed: 22826341]

98. Homberg JR, Molteni R, Calabrese F, Riva MA. The serotonin-BDNF duo: developmental implications for the vulnerability to psychopathology. Neurosci Biobehav Rev. 2014; 43:35–47. [PubMed: 24704572]

- 99. Wellman CL, Camp M, Jones VM, MacPherson KP, Ihne J, Fitzgerald P, Maroun M, Drabant E, Bogdan R, Hariri AR, Holmes A. Convergent effects of mouse Pet-1 deletion and human PET-1 variation on amygdala fear and threat processing. Exp Neurol. 2013; 250:260–269. [PubMed: 24100022]
- 100. Kerman IA, Bernard R, Bunney WE, Jones EG, Schatzberg AF, Myers RM, Barchas JD, Akil H, Watson SJ, Thompson RC. Evidence for transcriptional factor dysregulation in the dorsal raphe nucleus of patients with major depressive disorder. Front Neurosci. 2012; 6:135. [PubMed: 23087602]
- 101. Fu W, Le Maitre E, Fabre V, Bernard JF, David Xu ZQ, Hokfelt T. Chemical neuroanatomy of the dorsal raphe nucleus and adjacent structures of the mouse brain. J Comp Neurol. 2010; 518:3464–3494. [PubMed: 20589909]
- 102. Vasudeva RK, Lin RC, Simpson KL, Waterhouse BD. Functional organization of the dorsal raphe efferent system with special consideration of nitrergic cell groups. J Chem Neuroanat. 2011; 41:281–293. [PubMed: 21640185]
- 103. Xie H, Han F, Shi X. Single-prolonged stress induce changes of CaM/CaMKIIalpha in the rats of dorsal raphe nucleus. Neurochem Res. 2012; 37:1043–1049. [PubMed: 22294157]
- 104. Gocho Y, Sakai A, Yanagawa Y, Suzuki H, Saitow F. Electrophysiological and pharmacological properties of GABAergic cells in the dorsal raphe nucleus. J Physiol Sci. 2013; 63:147–154. [PubMed: 23275149]
- 105. Dougalis AG, Matthews GA, Bishop MW, Brischoux F, Kobayashi K, Ungless MA. Functional properties of dopamine neurons and co-expression of vasoactive intestinal polypeptide in the dorsal raphe nucleus and ventro-lateral periaqueductal grey. Eur J Neurosci. 2012; 36:3322–3332. [PubMed: 22925150]
- 106. Calizo LH, Akanwa A, Ma X, Pan YZ, Lemos JC, Craige C, Heemstra LA, Beck SG. Raphe serotonin neurons are not homogenous: electrophysiological, morphological and neurochemical evidence. Neuropharmacology. 2011; 61:524–543. [PubMed: 21530552]
- 107. Crawford LK, Craige CP, Beck SG. Increased intrinsic excitability of lateral wing serotonin neurons of the dorsal raphe: a mechanism for selective activation in stress circuits. J Neurophysiol. 2010; 103:2652–2663. [PubMed: 20237311]
- 108. Alekseyenko OV, Lee C, Kravitz EA. Targeted manipulation of serotonergic neurotransmission affects the escalation of aggression in adult male Drosophila melanogaster. PLoS One. 2010; 5:e10806. [PubMed: 20520823]
- 109. Kravitz EA. Serotonin and aggression: insights gained from a lobster model system and speculations on the role of amine neurons in a complex behavior. J Comp Physiol A. 2000; 186:221–238. [PubMed: 10757238]
- 110. Stevenson PA, Schildberger K. Mechanisms of experience dependent control of aggression in crickets. Curr Opin Neurobiol. 2013; 23:318–323. [PubMed: 23537901]
- 111. Teles MC, Dahlbom SJ, Winberg S, Oliveira RF. Social modulation of brain monoamine levels in zebrafish. Behav Brain Res. 2013; 253:17–24. [PubMed: 23850359]
- 112. Young SN, Leyton M. The role of serotonin in human mood and social interaction. Insight from altered tryptophan levels. Pharmacol Biochem Behav. 2002; 71:857–865. [PubMed: 11888576]
- 113. Chiba T, Kayahara T, Nakano K. Efferent projections of infralimbic and prelimbic areas of the medial prefrontal cortex in the Japanese monkey, Macaca fuscata. Brain Res. 2001; 888:83–101. [PubMed: 11146055]
- 114. Hajos M, Richards CD, Szekely AD, Sharp T. An electrophysiological and neuroanatomical study of the medial prefrontal cortical projection to the midbrain raphe nuclei in the rat. Neuroscience. 1998; 87:95–108. [PubMed: 9722144]
- 115. Jankowski MP, Sesack SR. Prefrontal cortical projections to the rat dorsal raphe nucleus: ultrastructural features and associations with serotonin and gamma-aminobutyric acid neurons. J Comp Neurol. 2004; 468:518–529. [PubMed: 14689484]

116. Vertes RP. Differential projections of the infralimbic and prelimbic cortex in the rat. Synapse. 2004; 51:32–58. [PubMed: 14579424]

- 117. Aghajanian GK, Wang RY. Habenular and other midbrain raphe afferents demonstrated by a modified retrograde tracing technique. Brain Res. 1977; 122:229–242. [PubMed: 837230]
- 118. Gabbott PL, Warner TA, Jays PR, Salway P, Busby SJ. Prefrontal cortex in the rat: projections to subcortical autonomic, motor, and limbic centers. J Comp Neurol. 2005; 492:145–177. [PubMed: 16196030]
- 119. Goncalves L, Nogueira MI, Shammah-Lagnado SJ, Metzger M. Prefrontal afferents to the dorsal raphe nucleus in the rat. Brain Res Bull. 2009; 78:240–247. [PubMed: 19103268]
- 120. Vazquez-Borsetti P, Cortes R, Artigas F. Pyramidal neurons in rat prefrontal cortex projecting to ventral tegmental area and dorsal raphe nucleus express 5-HT2A receptors. Cereb Cortex. 2009; 19:1678–1686. [PubMed: 19029064]
- 121. Varga V, Szekely AD, Csillag A, Sharp T, Hajos M. Evidence for a role of GABA interneurones in the cortical modulation of midbrain 5-hydroxytryptamine neurones. Neuroscience. 2001; 106:783–792. [PubMed: 11682163]
- 122. Celada P, Puig MV, Casanovas JM, Guillazo G, Artigas F. Control of dorsal raphe serotonergic neurons by the medial prefrontal cortex: Involvement of serotonin-1A, GABA(A), and glutamate receptors. J Neurosci. 2001; 21:9917–9929. [PubMed: 11739599]
- 123. Challis C, Boulden J, Veerakumar A, Espallergues J, Vassoler FM, Pierce RC, Beck SG, Berton O. Raphe GABAergic neurons mediate the acquisition of avoidance after social defeat. J Neurosci. 2013; 33:13978–13988. [PubMed: 23986235]
- 124. Shikanai H, Yoshida T, Konno K, Yamasaki M, Izumi T, Ohmura Y, Watanabe M, Yoshioka M. Distinct neurochemical and functional properties of GAD67-containing 5-HT neurons in the rat dorsal raphe nucleus. J Neurosci. 2012; 32:14415–14426. [PubMed: 23055511]
- 125. Challis C, Beck SG, Berton O. Optogenetic modulation of descending prefrontocortical inputs to the dorsal raphe bidirectionally bias socioaffective choices after social defeat. Front Behav Neurosci. 2014; 8:43. [PubMed: 24596546]
- 126. Amemori K, Graybiel AM. Localized microstimulation of primate pregenual cingulate cortex induces negative decision-making. Nat Neurosci. 2012; 15:776–785. [PubMed: 22484571]
- 127. Pollak Dorocic I, Furth D, Xuan Y, Johansson Y, Pozzi L, Silberberg G, Carlen M, Meletis K. A whole-brain atlas of inputs to serotonergic neurons of the dorsal and median raphe nuclei. Neuron. 2014; 83:663–678. [PubMed: 25102561]
- 128. Weissbourd B, Ren J, DeLoach KE, Guenthner CJ, Miyamichi K, Luo L. Presynaptic partners of dorsal raphe serotonergic and GABAergic neurons. Neuron. 2014; 83:645–662. [PubMed: 25102560]
- 129. Dembrow N, Johnston D. Subcircuit-specific neuromodulation in the prefrontal cortex. Front Neural Circuits. 2014; 8:54. [PubMed: 24926234]
- Hattox AM, Nelson SB. Layer V neurons in mouse cortex projecting to different targets have distinct physiological properties. J Neurophysiol. 2007; 98:3330–3340. [PubMed: 17898147]
- 131. Berton O, Nestler EJ. New approaches to antidepressant drug discovery: beyond monoamines. Nat Rev Neurosci. 2006; 7:137–151. [PubMed: 16429123]
- 132. Bjorkqvist K. Social defeat as a stressor in humans. Physiol Behav. 2001; 73:435–442. [PubMed: 11438372]
- 133. Kudryavtseva NN, Bakshtanovskaya IV, Koryakina LA. Social model of depression in mice of C57BL/6J strain. Pharmacol Biochem Behav. 1991; 38:315–320. [PubMed: 2057501]
- 134. Koolhaas JM, Meerlo P, de Boer SF, Strubbe JH, Bohus B. The temporal dynamics of the stress response. Neuroscience & Biobehavioral Reviews. 1997; 21:775–782. [PubMed: 9415902]
- 135. Miczek KA, Yap JJ, Covington HE 3rd. Social stress, therapeutics and drug abuse: preclinical models of escalated and depressed intake. Pharmacol Ther. 2008; 120:102–128. [PubMed: 18789966]
- 136. Chaouloff F. Social stress models in depression research: what do they tell us? Cell Tissue Res. 2013; 354:179–190. [PubMed: 23532563]
- 137. Berton O, McClung CA, Dileone RJ, Krishnan V, Renthal W, Russo SJ, Graham D, Tsankova NM, Bolanos CA, Rios M, Monteggia LM, Self DW, Nestler EJ. Essential role of BDNF in the

- mesolimbic dopamine pathway in social defeat stress. Science. 2006; 311:864–868. [PubMed: 16469931]
- 138. Lehmann ML, Geddes CE, Lee JL, Herkenham M. Urine scent marking (USM): a novel test for depressive-like behavior and a predictor of stress resiliency in mice. PLoS One. 2013; 8:e69822. [PubMed: 23875001]
- 139. Cooper MA, Huhman KL. Corticotropin-releasing factor receptors in the dorsal raphe nucleus modulate social behavior in Syrian hamsters. Psychopharmacology (Berl). 2007; 194:297–307. [PubMed: 17581742]
- 140. Kollack-Walker S, Watson SJ, Akil H. Social stress in hamsters: defeat activates specific neurocircuits within the brain. J Neurosci. 1997; 17:8842–8855. [PubMed: 9348352]
- 141. Golden SA, Covington HE 3rd, Berton O, Russo SJ. A standardized protocol for repeated social defeat stress in mice. Nat Protoc. 2011; 6:1183–1191. [PubMed: 21799487]
- 142. Krishnan V, Berton O, Nestler E. The use of animal models in psychiatric research and treatment. Am J Psychiatry. 2008; 165:1109. [PubMed: 18765492]
- 143. Donahue RJ, Muschamp JW, Russo SJ, Nestler EJ, Carlezon WA Jr. Effects of striatal DeltaFosB overexpression and ketamine on social defeat stress-induced anhedonia in mice. Biol Psychiatry. 2014; 76:550–558. [PubMed: 24495460]
- 144. van Peer JM, Spinhoven P, Roelofs K. Psychophysiological evidence for cortisol-induced reduction in early bias for implicit social threat in social phobia. Psychoneuroendocrinology. 2010; 35:21–32. [PubMed: 19836898]
- 145. Vincent MY, Jacobson L. Glucocorticoid receptor deletion from the dorsal raphe nucleus of mice reduces dysphoria-like behavior and impairs hypothalamic-pituitary-adrenocortical axis feedback inhibition. Eur J Neurosci. 2014; 39:1671–1681. [PubMed: 24684372]
- 146. Espallergues J, Teegarden SL, Veerakumar A, Boulden J, Challis C, Jochems J, Chan M, Petersen T, Deneris E, Matthias P, Hahn CG, Lucki I, Beck SG, Berton O. HDAC6 regulates glucocorticoid receptor signaling in serotonin pathways with critical impact on stress resilience. J Neurosci. 2012; 32:4400–4416. [PubMed: 22457490]
- 147. Haller J, Halasz J, Mikics E, Kruk MR. Chronic glucocorticoid deficiency-induced abnormal aggression, autonomic hypoarousal, and social deficit in rats. J Neuroendocrinol. 2004; 16:550– 557. [PubMed: 15189330]
- 148. Lehmann ML, Brachman RA, Martinowich K, Schloesser RJ, Herkenham M. Glucocorticoids orchestrate divergent effects on mood through adult neurogenesis. J Neurosci. 2013; 33:2961–2972. [PubMed: 23407954]
- 149. Sachs BD, Rodriguiz RM, Tran HL, Iyer A, Wetsel WC, Caron MG. Serotonin deficiency alters susceptibility to the long-term consequences of adverse early life experience. Psychoneuroendocrinology. 2015; 53C:69–81. [PubMed: 25602134]
- 150. Temel Y, Helmy A, Pinnock S, Herbert J. Effect of serotonin depletion on the neuronal, endocrine and behavioural responses to corticotropin-releasing factor in the rat. Neurosci Lett. 2003; 338:139–142. [PubMed: 12566172]
- 151. Zhang WQ, Smolik CM, Barba-Escobedo PA, Gamez M, Sanchez JJ, Javors MA, Daws LC, Gould GG. Acute dietary tryptophan manipulation differentially alters social behavior, brain serotonin and plasma corticosterone in three inbred mouse strains. Neuropharmacology. 2014; 90C:1–8. [PubMed: 25445490]
- 152. Beacher FD, Gray MA, Minati L, Whale R, Harrison NA, Critchley HD. Acute tryptophan depletion attenuates conscious appraisal of social emotional signals in healthy female volunteers. Psychopharmacology (Berl). 2011; 213:603–613. [PubMed: 20596858]
- 153. Hamani C, Diwan M, Macedo CE, Brandao ML, Shumake J, Gonzalez-Lima F, Raymond R, Lozano AM, Fletcher PJ, Nobrega JN. Antidepressant-like effects of medial prefrontal cortex deep brain stimulation in rats. Biol Psychiatry. 2010; 67:117–124. [PubMed: 19819426]
- 154. Veerakumar A, Challis C, Gupta P, Da J, Upadhyay A, Beck SG, Berton O. Antidepressant-like effects of cortical deep brain stimulation coincide with pro-neuroplastic adaptations of serotonin systems. Biol Psychiatry. 2014; 76:203–212. [PubMed: 24503468]
- 155. Holtzheimer PE, Kelley ME, Gross RE, Filkowski MM, Garlow SJ, Barrocas A, Wint D, Craighead MC, Kozarsky J, Chismar R, Moreines JL, Mewes K, Posse PR, Gutman DA,

- Mayberg HS. Subcallosal cingulate deep brain stimulation for treatment-resistant unipolar and bipolar depression. Arch Gen Psychiatry. 2012; 69:150–158. [PubMed: 22213770]
- 156. Lozano AM, Mayberg HS, Giacobbe P, Hamani C, Craddock RC, Kennedy SH. Subcallosal cingulate gyrus deep brain stimulation for treatment-resistant depression. Biol Psychiatry. 2008; 64:461–467. [PubMed: 18639234]
- 157. Berton O, Covington HE 3rd, Ebner K, Tsankova NM, Carle TL, Ulery P, Bhonsle A, Barrot M, Krishnan V, Singewald GM, Singewald N, Birnbaum S, Neve RL, Nestler EJ. Induction of deltaFosB in the periaqueductal gray by stress promotes active coping responses. Neuron. 2007; 55:289–300. [PubMed: 17640529]
- 158. Liu Z, Zhou J, Li Y, Hu F, Lu Y, Ma M, Feng Q, Zhang JE, Wang D, Zeng J, Bao J, Kim JY, Chen ZF, El Mestikawy S, Luo M. Dorsal raphe neurons signal reward through 5-HT and glutamate. Neuron. 2014; 81:1360–1374. [PubMed: 24656254]
- 159. Miyazaki KW, Miyazaki K, Tanaka KF, Yamanaka A, Takahashi A, Tabuchi S, Doya K. Optogenetic activation of dorsal raphe serotonin neurons enhances patience for future rewards. Curr Biol. 2014; 24:2033–2040. [PubMed: 25155504]
- 160. Dolen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin. Nature. 2013; 501:179–184. [PubMed: 24025838]
- 161. Qi J, Zhang S, Wang HL, Wang H, de Jesus Aceves Buendia J, Hoffman AF, Lupica CR, Seal RP, Morales M. A glutamatergic reward input from the dorsal raphe to ventral tegmental area dopamine neurons. Nat Commun. 2014; 5:5390. [PubMed: 25388237]
- 162. Kumar S, Black SJ, Hultman R, Szabo ST, DeMaio KD, Du J, Katz BM, Feng G, Covington HE 3rd, Dzirasa K. Cortical control of affective networks. J Neurosci. 2013; 33:1116–1129. [PubMed: 23325249]
- 163. Kumar S, Hultman R, Hughes D, Michel N, Katz BM, Dzirasa K. Prefrontal cortex reactivity underlies trait vulnerability to chronic social defeat stress. Nat Commun. 2014; 5:4537. [PubMed: 25072279]
- 164. Dzirasa K, Kumar S, Sachs BD, Caron MG, Nicolelis MA. Cortical-amygdalar circuit dysfunction in a genetic mouse model of serotonin deficiency. J Neurosci. 2013; 33:4505–4513. [PubMed: 23467366]
- 165. Amat J, Aleksejev RM, Paul E, Watkins LR, Maier SF. Behavioral control over shock blocks behavioral and neurochemical effects of later social defeat. Neuroscience. 2010; 165:1031–1038. [PubMed: 19909791]
- 166. Stoltenberg SF, Christ CC, Carlo G. Afraid to help: social anxiety partially mediates the association between 5-HTTLPR triallelic genotype and prosocial behavior. Soc Neurosci. 2013; 8:400–406. [PubMed: 23789884]
- 167. Carlson JM, Mujica-Parodi LR, Harmon-Jones E, Hajcak G. The orienting of spatial attention to backward masked fearful faces is associated with variation in the serotonin transporter gene. Emotion. 2012; 12:203–207. [PubMed: 21875187]
- 168. Perez-Edgar K, Bar-Haim Y, McDermott JM, Gorodetsky E, Hodgkinson CA, Goldman D, Ernst M, Pine DS, Fox NA. Variations in the serotonin-transporter gene are associated with attention bias patterns to positive and negative emotion faces. Biol Psychol. 2010; 83:269–271. [PubMed: 19723555]
- 169. Pergamin-Hight L, Bakermans-Kranenburg MJ, van Ijzendoorn MH, Bar-Haim Y. Variations in the promoter region of the serotonin transporter gene and biased attention for emotional information: a meta-analysis. Biol Psychiatry. 2012; 71:373–379. [PubMed: 22138391]
- 170. Watson KK, Ghodasra JH, Platt ML. Serotonin transporter genotype modulates social reward and punishment in rhesus macaques. PLoS One. 2009; 4:e4156. [PubMed: 19142220]
- 171. Bartolomucci A, Carola V, Pascucci T, Puglisi-Allegra S, Cabib S, Lesch KP, Parmigiani S, Palanza P, Gross C. Increased vulnerability to psychosocial stress in heterozygous serotonin transporter knockout mice. Dis Model Mech. 2010; 3:459–470. [PubMed: 20371729]
- 172. Holmes A, Murphy DL, Crawley JN. Reduced aggression in mice lacking the serotonin transporter. Psychopharmacology (Berl). 2002; 161:160–167. [PubMed: 11981596]

173. Lewejohann L, Kloke V, Heiming RS, Jansen F, Kaiser S, Schmitt A, Lesch KP, Sachser N. Social status and day-to-day behaviour of male serotonin transporter knockout mice. Behav Brain Res. 2010; 211:220–228. [PubMed: 20347882]

- 174. Kane MJ, Angoa-Perez M, Briggs DI, Sykes CE, Francescutti DM, Rosenberg DR, Kuhn DM. Mice genetically depleted of brain serotonin display social impairments, communication deficits and repetitive behaviors: possible relevance to autism. PLoS One. 2012; 7:e48975. [PubMed: 23139830]
- 175. Murphy SE, Longhitano C, Ayres RE, Cowen PJ, Harmer CJ. Tryptophan supplementation induces a positive bias in the processing of emotional material in healthy female volunteers. Psychopharmacology (Berl). 2006; 187:121–130. [PubMed: 16767422]
- 176. Moskowitz DS, Pinard G, Zuroff DC, Annable L, Young SN. The effect of tryptophan on social interaction in everyday life: a placebo-controlled study. Neuropsychopharmacology. 2001; 25:277–289. [PubMed: 11425511]
- 177. Browning M, Reid C, Cowen PJ, Goodwin GM, Harmer CJ. A single dose of citalopram increases fear recognition in healthy subjects. J Psychopharmacol. 2007; 21:684–690. [PubMed: 17259206]
- 178. Hysek CM, Schmid Y, Simmler LD, Domes G, Heinrichs M, Eisenegger C, Preller KH, Quednow BB, Liechti ME. MDMA enhances emotional empathy and prosocial behavior. Soc Cogn Affect Neurosci. 2014; 9:1645–1652. [PubMed: 24097374]
- 179. Majumder I, White JM, Irvine RJ. Antidepressant-like effects of ecstasy in subjects with a predisposition to depression. Addict Behav. 2012; 37:1189–1192. [PubMed: 22704044]
- 180. Jochems J, Boulden J, Lee BG, Blendy JA, Jarpe M, Mazitschek R, Van Duzer JH, Jones S, Berton O. Antidepressant-like properties of novel HDAC6-selective inhibitors with improved brain bioavailability. Neuropsychopharmacology. 2014; 39:389–400. [PubMed: 23954848]
- 181. Morley KC, Arnold JC, McGregor IS. Serotonin (1A) receptor involvement in acute 3,4-methylenedioxymethamphetamine (MDMA) facilitation of social interaction in the rat. Prog Neuropsychopharmacol Biol Psychiatry. 2005; 29:648–657. [PubMed: 15908091]
- 182. Morley KC, McGregor IS. (±)-3,4-Methylenedioxymethamphetamine (MDMA, 'Ecstasy') increases social interaction in rats. European Journal of Pharmacology. 2000; 408:41–49. [PubMed: 11070182]
- 183. Hilimire MR, Mayberg HS, Holtzheimer PE, Broadway JM, Parks NA, DeVylder JE, Corballis PM. Effects of Subcallosal Cingulate Deep Brain Stimulation on Negative Self-bias in Patients With Treatment-resistant Depression. Brain Stimul. 2014
- 184. Covington HE 3rd, Lobo MK, Maze I, Vialou V, Hyman JM, Zaman S, LaPlant Q, Mouzon E, Ghose S, Tamminga CA, Neve RL, Deisseroth K, Nestler EJ. Antidepressant effect of optogenetic stimulation of the medial prefrontal cortex. J Neurosci. 2010; 30:16082–16090. [PubMed: 21123555]
- 185. Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, Sohal VS, Goshen I, Finkelstein J, Paz JT, Stehfest K, Fudim R, Ramakrishnan C, Huguenard JR, Hegemann P, Deisseroth K. Neocortical excitation/inhibition balance in information processing and social dysfunction. Nature. 2011; 477:171–178. [PubMed: 21796121]


# Approach Avoidance Task for human studies

#### **EXPLICIT RATING TASK**




"How many steps would you go towards or away from this person?"

#### **IMPLICIT TASK**



Push when presented with a red frame Pull when presented with a blue frame

## Social Interaction Test for rodent studies



in interaction zone (sec) Cumulative time spent 20 50 100 150 Elapsed time (sec)

Figure 1. Behavioral tasks used to probe social approach/avoidance behaviors

In the Approach Avoidance task for humans, approach or avoidant behaviors are quantified using two tasks. In the explicit task, individuals are asked to define how close they would approach the subject on the picture or rate properties associated with its social valence, such as safety or trustworthiness. In the implicit task, automatic processing and tendencies are assessed. Pictures of faces with different emotions (i.e. happy, neutral, angry) are presented for short durations (50ms). A colored frame surrounds these pictures and subjects are instructed to pull or push a joystick according to the color of the frame presented. Even in the absence of conscious perception, subjects are faster at avoiding threatening faces. In the social interaction test for mice, experimental subjects are placed in the corner of an open field arena with a caged off unfamiliar social target on the opposite side. Mice are free to explore the arena while movement is recorded using video tracking software. Undefeated control mice unconditionally approach the target within seconds. One subset of defeated mice display strong avoidance tendencies (vulnerable) while the other will maintain social approach and interaction (resilient).




Figure 2. Overview of vmPFC-DRN circuitry

Based on data from previous tracing studies, glutamatergic layer V vmPFC neurons send afferents to the DRN that preferentially synapse on GABA interneurons, which then locally inhibit 5-HT neurons and gate serotonergic output. Representative confocal images depict typical localization of GABAergic (*GAD2-tdTomato*), serotonergic (*Pet1-tdTomato*) and vmPFC terminals (*CaMKIIA*-driven *SynP-GFP*) in the rostral and caudal DRN. Analysis reveals higher correlation of vmPFC terminals with GABA neurons in the rostral DRN and with 5-HT neurons in the caudal DRN.

# Social Defeat Training (10 days) Physical aggression Sensory contact Social Interaction Test Social interaction zone

Figure 3. Chronic social defeat stress paradigm

In CSDS, mice are first exposed to social defeat training, which involves physical aggression from a trained CD-1 mouse followed by sensory contact protected by a plexiglass divider. After 10 days of defeat with exposure to novel aggressor each day, social behaviors are characterized in the social interaction test (described in Figure 1). Depicted here is photomanipulation of the subject mouse during the sensory contact period.





Figure 4. Hypothesized adaptations in the vmPFC-DRN microcircuit associated with defeat-induced avoidance and its normalization by vmPFC DBS  $\,$ 

Schematic summarizing the differential neuroplastic adaptations in the vmPFC-DRN microcircuit and hypothesized effect on 5-HT output. Depicted are changes in intrinsic cellular excitability, synaptic activity (EPSC, IPSC) and corresponding social approach/avoidance behavior profiles (shown as video tracking heat maps). Compared to controls, in susceptible mice we observed hyperexcitable GABA neurons that had increased glutamatergic input and hypoexcitable 5-HT neurons that had increased GABAergic input. After chronic vmPFC DBS, there was a neuroplastic remodeling of DRN microcircuitry that normalized GABA-mediated hyperinhibition of 5-HT neurons and facilitated a direct excitatory synaptic drive of 5-HT neurons. Though the direct source of synaptic adaptations was not causally determined in this work, our mapping studies provide strong evidence for this circuit and adaptive mechanisms.



**Figure 5. Neuronal targets downstream of the vmPFC-DRN pathway in socioaffective behaviors** Schematic depicts vmPFC control of the DRN and the structures likely to be affected by modulated DRN serotonergic projections in socioaffective behavior. Reciprocal projections to the vmPFC have been implicated in impulsivity and negative affective bias, 5-HT activity in the amygdala has been linked to fear and threat processing, and projections to the pathway between the VTA and NAc have been shown to affect reward behaviors.

Table 1

#### Symptoms associated with MDD.

Psychological symptoms*	Physical symptoms*	Social symptoms
Guilt, self-doubt or loss of self-esteem     Lack of interest, pleasure or enthusiasm     Concentration or attention span is reduced     Recurring thoughts of death or suicide	Disturbances in sleep – insomnia or hypersomnia     Abnormal psychomotor activity     Fatigue or loss of energy     Impaired appetite resulting in weight loss or gain	Poor performance at work or school     Interpersonal difficulties at home or with family     Social withdrawal     Neglecting hobbies or interests

In addition to depressed mood and sadness, listed are the core symptoms associated with MDD. It is important to note that only symptoms in the categories marked with an asterisk are used to clinically diagnose MDD.

 Table 2

 Parallel human and animal studies on socioaffective behaviors

	Human studies	Animal studies
Stress hormones	Cortisol administration increased attention to social threat especially in social anxiety patients <sup>144</sup>	CORT administration facilitated social avoidance after CSDS <sup>146</sup> ; blocking CORT release via adrenalectomy prevented avoidance <sup>147,148</sup>
Mood disorders	Patients clinically diagnosed with MDD reacted differently to emotional faces, rated ambiguous social expression more negatively and displayed stronger avoidance tendencies in the explicit condition of the AAT, whereas social withdrawal was less pronounced in the implicit condition 31, 32	CSDS <sup>125</sup> and LH <sup>165</sup> models of depression result in long lasting social withdrawal and avoidant behavioral tendencies
Genetic models	Subjects with the short allele of 5-HTTLPR polymorphism displayed stronger social avoidance <sup>166</sup> and increased attention to negative social stimuli <sup>167–169</sup>	In macaques, pictures of high status male were rewarding and induced risk seeking behaviors in 5-HTTLPR long allele carriers, while the opposite was observed in short allele carriers <sup>170</sup>
		SERT -/- mice displayed increased vulnerability to social stress <sup>171</sup> and decreased social dominance <sup>172,173</sup>
		Humanized TPH2 (R439H) knock-in ( <i>Tph2KI</i> ) mice displayed increased social avoidance after social defeat <sup>174</sup>
Decreased serotonin	ATD decreased positive appraisal of happy faces 152 or evaluation of social relationships of intimate couples 68 and increased attention towards threatening cues 22	TRP depletion <sup>151</sup> or 5,7-DHT administration <sup>150</sup> decreased sociability in mice
Increased serotonin	<ul> <li>TRP supplementation increased recognition of happiness<sup>175</sup> and decreased quarrelsome behaviors<sup>176</sup></li> <li>Citalopram administration increased recognition of happy<sup>50</sup> or fearful<sup>177</sup> faces and increased rating of ambiguous faces as positively valenced<sup>71</sup></li> <li>MDMA administration enhanced prosocial behaviors<sup>178</sup> and decreased depressive tendencies in social gatherings<sup>179</sup></li> </ul>	TRP administration improved sociability <sup>151</sup> Citalopram <sup>180</sup> or fluoxetine <sup>137</sup> administration following CSDS restored social approach  MDMA administration decreased aggression and increased social interaction duration <sup>181,182</sup>
Corticolimbic synchrony	Enhanced prefrontal-amygdala coupling was associated with negatively biased evaluation of socioemotional cues in healthy subjects and avoidance tendencies in depressed patients; coupling was reduced by increased serotonin <sup>72</sup>	Enhanced cortico-amygdala synchrony predicted vulnerability to CSDS-induced avoidance <sup>163</sup> and is increased in TPH2 -/ - mice <sup>164</sup>
vmPFC manipulation	<ul> <li>cTBS of the aPFC increased emotional dysfunction during social interaction<sup>76</sup></li> <li>DBS of the SCG normalized negative self bias<sup>183</sup> and resulted in social reintegration and lowered social dysfunction<sup>155,156</sup></li> </ul>	High frequency (>100 Hz) electrical <sup>154</sup> or optogenetic <sup>184</sup> stimulation of the vmPFC reverses avoidance after social defeat  Elevating excitatory/inhibitory balance via 80Hz optogenetic activation induced social deficits <sup>185</sup>

Abbreviations: 5-HTTLPR – serotonin transporter linked polymorphic region, AAT – approach avoidance task, aPFC – anterior prefrontal cortex, ATD – acute tryptophan depletion, CORT – corticosterone, CSDS – chronic social defeat stress, cTBS – chronic theta burst stimulation, DBS – deep brain stimulation, LH – learned helplessness, MDD – major depressive disorder, MDMA – methylenedioxymethamphetamine, SCG – subcallosal cingulate, SERT – serotonin transporter, TRP – tryptophan, TPH – tryptophan hydroxylase, vmPFC – ventromedial prefrontal cortex