

Biochemistry. Author manuscript; available in PMC 2008 November 10.

Published in final edited form as:

Biochemistry. 2007 July 10; 46(27): 8146-8155. doi:10.1021/bi700686p.

Tropomyosin Regulates Elongation by Formin at the Fast-Growing End of the Actin Filament[†]

Barbara Wawro^{‡, ⊥}, Norma J. Greenfield[‡], Martin A. Wear^{§,#}, John A. Cooper[§], Henry N. Higgs^{||}, and Sarah E. Hitchcock-DeGregori^{*,‡}

‡Department of Neuroscience and Cell Biology, Robert Wood Johnson Medical School, 675 Hoes Lane, Piscataway, New Jersey 08854

§Department of Cell Biology, Washington UniVersity, 660 South Euclid AVenue, St. Louis, Missouri 63110 ||Department of Biochemistry, Dartmouth Medical School, HanoVer, New Hampshire 03755-3844

Abstract

The balance between dynamic and stable actin filaments is essential for the regulation of cellular functions including the determination of cell shape and polarity, cell migration, and cytokinesis. Proteins that regulate polymerization at the filament ends and filament stability confer specificity to actin filament structure and cellular function. The dynamics of the barbed, fast-growing end of the filament are controlled in space and time by both positive and negative regulators of actin polymerization. Capping proteins inhibit the addition and loss of subunits, whereas other proteins, including formins, bind at the barbed end and allow filament growth. In this work, we show that tropomyosin regulates dynamics at the barbed end. Tropomyosin binds to constructs of FRL1 and mDia2 that contain the FH2 domain and modulates formin-dependent capping of the barbed end by relieving inhibition of elongation by FRL1- FH1FH2, mDia1-FH2, and mDia2-FH2 in an isoformdependent fashion. In this role, tropomyosin functions as an activator of formin. Tropomyosin also inhibits the binding of FRL1-FH1FH2 to the sides of actin filaments independent of the isoform. In contrast, tropomyosin does not affect the ability of capping protein to block the barbed end. We suggest that tropomyosin and formin act together to ensure the formation of unbranched actin filaments, protected from severing, that could be capped in stable cellular structures. This role, in addition to its cooperative control of myosin function, establishes tropomyosin as a universal regulator of the multifaceted actin cytoskeleton.

> Reversible polymerization of actin is essential for many cellular functions including determination of cell shape and polarity, cell movement, cytokinesis, and intracellular transport. The ends of actin filaments are structurally distinct and differ in the rates of incorporation and dissociation of actin subunits. Elongation at the fast growing, barbed end predominates in cells and is precisely controlled in space and time by positive and negative regulators of actin polymerization. Capping proteins bind and prevent addition and loss of subunits, whereas other proteins, including formins, bind at the barbed end and allow filament growth and depolymerization [reviewed in (1,2)].

[†]This work was supported by NIH Grants GM36326 and GM63257 to S.E.H.D., GM38542 to J.A.C., and GM69818 to H.N.H.

^{*}To whom correspondence should be addressed. Tel: 732-235-5236. Fax: 732-235-4029. E-mail: hitchcoc@umdnj.edu. _Present address: Gwiazdzista 21/91, 01-651 Warsaw, Poland.

^{*}Present address: The Center for Translational and Chemical Biology, Institute of Structural and Molecular Biology, University of Edinburgh, EH9 3JR, U.K.

Formins are a widespread family of barbed-end binding proteins that are implicated in the assembly of unbranched actin-containing structures including microvilli and filopodia, stress fibers, and contractile rings. They are required for the formation of actin cables in yeast. Formins prevent the binding of capping protein and modulate the rate of monomer addition to the barbed end [reviewed in (3,4)].

Formins are classified on the basis of the presence of formin homology domains, FH1¹ and FH2, which are responsible for most of its effects on actin [reviewed in (5)]. Fragments containing the FH2 domain or FH1FH2 domains form dimers and nucleate *de novo* filament assembly of monomeric actin. Formin remains associated with the elongating barbed end of the filament, leading to its description as a processive, leaky capper (6-8). Although formins share common mechanisms, isoforms differ in the ability to nucleate polymerization, inhibit elongation, bind to the sides of filaments, and bundle and sever filaments, and in their processivity (9).

The capping protein family of proteins, expressed in virtually all eucaryotic cells, are antagonistic to formins. Capping protein is an $\alpha\beta$ heterodimer localized at the ends of filaments where unbranched actin filaments end in stable structures such as Z-discs in muscle and adhering junctions, and at filament ends in lamellipodia [(10); reviewed in (11)]. Capping protein is also associated with actin patches and endosomal vesicles in yeast (12-14). In vertebrates, there are two isoforms of each subunit with tissue-specific expression patterns and functions. Negative regulators of capping protein include the phosphoinositide, PIP2 (15), and proteins that specifically complex with capping proteins such as CARMIL and myotrophin/V-1 (16,17). No positive regulators have been reported.

Formins and capping protein are often associated with the formation and maintenance of long, unbranched actin filaments [reviewed in (3,18)] that contain tropomyosin [reviewed in (19)]. Tropomyosins are coiled-coil proteins that associate N-terminus to C-terminus on the actin filament to form a continuous strand that follows the filament helix, one strand on each side of the actin filament [reviewed in (20,21)]. A large protein family, tropomyosins have cell-and tissue-specific expression patterns and functions [reviewed in (19)]. It is best known for its classic role in the regulation of striated muscle contraction together with troponin and myosin. In addition, tropomyosin physically stabilizes actin filaments and protects them against the action of depolymerizing and severing proteins, such as DNase I, cofilin, and gelsolin as well as the nucleation of branches by the Arp2/3 complex (22-30). Tropomyosin, together with tropomodulin, caps the pointed ends of actin filaments in the sarcomeres of striated muscles and in the erythrocyte membrane cytoskeleton [reviewed in (31)], and can itself depress depolymerization from the pointed end (32-34).

A role for tropomyosin in regulating the barbed end of the filament is less well established. Direct interaction with an actin-binding domain of gelsolin, a barbed end-capping protein, is well documented, as is the ability of tropomyosin to dissociate gelsolin and promote annealing of short actin filaments (35-38). There is circumstantial evidence for an effect of tropomyosin on formin function from studies in *Saccharomyces cerevisiae*. Actin cables, which are polarized arrays of tropomyosin-containing filaments that are needed for cellular polarization, require both formin (Bni1p) and tropomyosin to form (39-41).

Here, we show that tropomyosin regulates filament dynamics at the barbed end, as it does the pointed end. Because tropomyosin is often present in formin-nucleated filaments in cells, we postulated that it may regulate formin. Tropomyosin binds directly to FRL1 and mDia2 constructs that contain the FH2 domain and relieves the inhibition of barbed-end elongation

¹Abbreviations: FH, formin homology domain; TM, tropomyosin; CP, capping protein.

by formin in an isoform-specific manner. Binding of tropomyosin to actin filaments inhibits the association of FRL1-FH1FH2 with the filament sides. In contrast, the tropomyosin isoforms we studied do not relieve blocking of the barbed end by capping protein. Tropomyosin's roles in stabilizing the filament from severing, and in regulating the dynamics of the filament ends, alone and in concert with other regulatory proteins, positions it as a universal regulator of the multifaceted actin cytoskeleton.

EXPERIMENTAL PROCEDURES

Protein Purification

Chicken skeletal muscle actin was extracted from acetone powder according to the method of Spudich and Watt (42) and further purified by gel filtration on Sephacryl S-100 HR. Immediately after purification, G-actin (in 2 mM Tris-HCl at pH 8.0, 0.2 mM ATP, 0.2 mM CaCl₂, 0.5 mM DTT, and 0.01% NaN₃) was quickly frozen in liquid nitrogen at -80 °C or stored at 4 °C and used within 2 weeks. Actin labeled with *N*-(1-pyrenyl)-iodoacetamide was prepared as described in ref 43, followed by gel filtration on Sephacryl S-100 HR.

Recombinant rat α-tropomyosins, TM5a, TM2, and unacetylated stTM (stTM-rec), were expressed in *Escherichia coli* BL21(DE3) cells and purified (44). N-acetylated striated muscle α-TM (stTM) was purified from chicken pectoral muscle (45). Recombinant FRL1-FH1FH2 (a.a. res. 449–1094), mDia1-FH2 (a.a. res. 748–1175), and mDia2-FH2 (a.a. res. 612–1034) were expressed in *E. coli* and purified (46). Recombinant capping proteins (CP) were expressed and purified (47): chicken muscle $\text{CP}\alpha1\beta1$, mouse $\text{CP}\alpha1\beta2$, and mouse $\text{CP}\alpha2\beta2$.

Protein concentrations were determined spectrophotometrically, using absorption coefficients of 22,100 M⁻¹ cm⁻¹, at 280 nM for mDia2-FH2; 25,700 at 280 nm for FRL1-FH1FH2; and 20,160 M⁻¹ cm⁻¹ at 280 nM for mDia1-FH2. Tropomyosin and actin concentrations were calculated by measuring the difference spectrum between pH 12.5 and pH 6 in 6 M guanidine-HCl, using a molar absorption coefficient for tyrosine of 2480 (48).

Elongation Assays

Elongation of 0.5 μ M Mg-G-actin (10% pyrenyl-labeled, in F-buffer: 10 mM Tris at pH 7.4, 0.2 mM ATP, 0.2 mM DTT, 0.01% NaN₃, 0.2 mM EGTA, 100 mM NaCl, 2 mM MgCl₂) on 1–2 μ M F-actin seeds, in the presence or absence of CP, formin, and tropomyosin, was monitored by an increase in pyrene fluorescence, at 25 °C. Mg-G actin was prepared by treating Ca-G-actin with 0.2 mM EGTA and 0.05 mM MgCl₂ for 3 min immediately prior to the experiment. For the seeds, 20–28 μ M Mg-G-actin was polymerized overnight in 100 mM NaCl and 2 mM MgCl₂ to allow spontaneous annealing of filaments before the addition of tropomyosin. Elongation was initiated by the addition of F-buffer (10×) and concentrated (10–14×) F-actin seeds to the cuvettes containing Mg-G-actin. The F-actin seeds were added using a Microman pipet and gently mixed with a plastic stirrer to minimize shearing. The figure legends include specific experimental details.

Time courses of pyrene fluorescence were measured on a PTI fluorimeter (Lawrenceville, NJ) with the excitation and emission wavelengths of 365 and 386 nm, respectively. Each data set was normalized by setting the final fluorescence increase at steady state, when available, or the value reported following curve fitting, as 100%. Normalized data were fit to the single- or double-exponential growth model, and initial rates of the reaction were calculated as first derivatives at t = 0 (GraphPad Prizm 4).

Actin Filament Sedimentation

A stock solution of $10{\text -}30\,\mu\text{M}$ Mg-G-actin was polymerized overnight with $100\,\text{mM}$ NaCl and $2\,\text{mM}$ MgCl $_2$. F-Actin was diluted with polymerization buffer for a final concentration of 5 μM prior to addition of phalloidin (5 μM), tropomyosin, and/or FRL1-FH1FH2. After 0.5-1 h, the samples (150 μL) were centrifuged for 20 min, at 60,000 rpm, in a TLA-100 rotor (Beckman) at 20 °C. One hundred microliters of supernatant were removed and dried on SpeedVac Plus (Savant). Pellets and supernatants were analyzed using SDS-PAGE (9% gels) according to Laemmli (49) and quantified on a Molecular Dynamics model 300A densitometer.

Circular Dichroism Spectroscopy

Circular dichroism experiments were performed on an Aviv model 215 spectrometer equipped with a 5 sample changer. Spectra of the tropomyosins, TM5a and TM2, and the formins, FRL1-FH1FH2 and mDia2-FH2, and their mixtures were obtained at 0 °C in 100 mM NaCl, 10 mM Tris-HCl, 0.2 mM EGTA, 0.1 mM MgCl₂, 0.01% NaN₂, and 0.5 mM dithiothreitol at pH 7.4. Data were collected at 0.25 to 0.5 nm intervals, corrected for baselines and smoothed with a polynomial order of 3 and a window of 21 to 11 points, respectively. The concentrations of the proteins ranged from 0.1 to 2 μ M. Data were obtained in 0.1 and 1 cm cells. Thermal denaturation curves of 1–2 μ M samples were obtained between 0 and 60–70 °C at 0.2 degree intervals with 5 s time averaging. The data were not smoothed. The first derivatives of the denaturation curves were obtained using data smoothed with a polynomial order of 3 and a window of 21 points and fitting the data to a third order polynomial. The helical content of FRL-FH1FH2 was estimated from its CD spectra by fitting it to standards for α -helical, β -pleated sheets, β -turns, and random structures using non-constrained least-squares analysis (50).

RESULTS

Tropomyosin Regulates the Dynamics of the Actin Filament's Barbed End

We evaluated the influence of muscle and non-muscle tropomyosins on actin filament barbedend elongation alone, in the presence of three formins that contain the FH2 actin binding domain, FRL1-FH1FH2, mDia1-FH2, and mDia2-FH2 (46,51), and with three capping protein isoforms, $CP\alpha1\beta1$, $CP\alpha1\beta2$, and $CP\alpha2\beta2$ (47). We compared three tropomyosins, all products of the α -TM gene [TPM1, (19)]. TM5a is a recombinant short, non-muscle isoform; TM2 is a recombinant long, non-muscle isoform; and stTM is a long tropomyosin isolated from skeletal muscle. TM2 and TM5a have the same C-terminal sequence, the end that is oriented toward the barbed end of the actin filament (52). The two long tropomyosins have the same N-terminal sequence that differs from TM5a, but stTM is N-acetylated (19).

We measured the polymerization of the pyrene-actin monomer (10% labeled) on actin filaments polymerized overnight to maximize filament length prior to the addition of tropomyosin because annealing by muscle tropomyosin is inversely related to filament length (38,53). The initial velocity of elongation (V_0) in the presence of regulatory proteins was calculated and compared to that of F-actin (taken as 100%). Tropomyosin influenced the rate of elongation in an isoform-specific manner. Figure 1 illustrates data from a representative experiment, along with the values relative to actin alone averaged from several experiments. Actin filaments saturated with TM5a and TM2 elongated at a faster rate than actin alone (137 \pm 7%, n = 4; 127 \pm 14%, n = 4, respectively). The increased rates suggest a specific barbedend effect that is not the result of new barbed ends formed by severing because TM5a and TM2 had no effect on the rate of elongation of actin filaments in the presence of capping protein (Figure 7) or on gelsolin-capped filaments in which TM5a partially blocked the pointed end (34). Striated muscle TM (stTM) slowed the overall rate of elongation (62 \pm 8%, n = 5). In our earlier work using a microscopic assay, we found that rabbit $\alpha\alpha$ -stTM (vs chicken $\alpha\alpha$ -stTM

used here) had no significant effect on the rate of barbed-end elongation, but rabbit $\alpha\beta$ -stTM reduced the rate (54). The effects of tropomyosin on elongation rate were independent of the time of incubation with F-actin (minutes to 8 h).

Tropomyosin Regulates Capping of the Actin Filament by Formin

The three formins inhibited barbed-end elongation to different extents, as previously reported (5,46). mDia2-FH2 was the most effective ($13 \pm 1\%$, relative to actin alone, n = 4) (Figure 2C and D). The kinetics of the fluorescence increase in the presence of FRL1-FH1FH2 (Figure 2A) and mDia1-FH2 (Figure 2E) closely followed a single-exponential reaction model. However, the best or only fits for the curves in the presence of mDia2-FH2 (Figure 2C) were obtained using a two-phase exponential model that describes two independent, simultaneous reactions. The result suggests the presence of two populations of filament ends elongating at different rates.

Tropomyosin relieved the inhibition by formins in an isoform-specific manner. The inhibition of barbed-end elongation by FRL1-FH1FH2 was almost completely relieved by TM5a, whereas TM2 and stTM had little effect (Figure 2A and B). In contrast, inhibition by mDia2-FH2 was partially relieved by TM2 but was little affected by TM5a or stTM (Figure 2C and D). The inhibition by mDia1-FH2 was less, but it was relieved by TM5a and to a lesser extent by TM2 (Figure 2E and F). In no case did tropomyosin relieve inhibition by formin to the level of actin with tropomyosin alone (compare with Figure 1). The results were independent of the time of incubation or order of addition of formin and tropomyosin to the pre-assembled filaments.

FRL1-FH1FH2 Inhibits ActiVation of Barbed-End Elongation by TM5a

We measured the rate of barbed-end elongation as a function of tropomyosin concentration to understand the relationship between the relief of inhibition by formins and assembly of a tropomyosin-actin filament. The ability of TM2 to relieve inhibition by mDia2-FH2 and the activation by TM2 alone corresponded to saturation of the actin filament (results not shown.).

In contrast, the dependence of activation and relief of inhibition by FRL1-FH1FH2 on TM5a concentration were more complicated (Figure 3). At saturation, TM5a activated the rate of barbed-end elongation ~1.4-fold compared to actin alone (Figure 1). The unanticipated finding was that activation by TM5a was greatest at substoichiometric levels and reached a minimum at saturation, as illustrated in a representative experiment in Figures 3. The activation at 0.05 μ M was 540 \pm 76% (680%, 530%, and 420% in three experiments), and at 0.1 μ M TM5a, the activation was 370 \pm 160% (530% and 200% in two experiments) when F-actin was less than 10% saturated with TM5a (i.e., almost undetectable; Figure 3D) compared to 137 \pm 7% (n = 3) at saturation. The variability in the magnitude of activation may be because the number of free barbed ends in the filaments used to nucleate polymerization differs from one experiment to the next.

The presence of FRL1-FH1FH2 inhibited the activation of barbed-end elongation at the lowest TM5a concentration, whereas TM5a relieved the capping activity of the formin. In the presence of FRL1-FH1FH2, the rate of barbed-end elongation was similar at all TM5a concentrations. The amount of FRL1-FH1FH2 in the pellet was independent of tropomyosin (Figure 3D), suggesting that formin remains on the barbed end in the presence of tropomyosin.

The FRL1 concentration and ionic strength in these experiments (0.05 μ M FRL1-FH1FH2 and 100 mM NaCl) minimized the formation of filament bundles (51), verified by no observed change in the actin content in the supernatant following low-speed centrifugation (results not shown.). We conclude that TM5a specifically influences the barbed end and that activation

may be inhibited by interaction with FRL1-FH1FH2, independent of stoichiometric TM5a binding to actin. The mutual neutralization may take place via direct complex formation because the FRL1-FH1FH2 concentration used (0.05 μ M) was sufficient to inhibit activation by 0.05 μ M TM5a.

Formins Containing the FH2 Domain Bind Tropomyosin

We measured binding between formin and tropomyosin using circular dichroism, a method we have used for the analysis of binding between tropomyosin and other proteins (Figure 4) (34, 55-57). When TM5a and FRL1-FH1FH2 were combined in an ~1:1 ratio, the negative ellipticity at 222 nm of the mixture of the proteins increased $9 \pm 1\%$ (n = 5) compared to the sum of the curves, evidence for complex formation (Figure 4A). The percent increase in ellipticity was independent of tropomyosin concentration over the range of 0.01 to 0.2 μ M, indicating that the binding is relatively tight ($K_d < 1 \mu M$). Because TM5a is ~100% helical, we infer that binding of TM5a to FRL1 induces the formation of the α-helix in FRL1-FH1FH2. Although the X-ray structures of FH2 domains are highly helical, the CD spectrum of FRL1-FH1FH2 is at most ~50% helical in the absence of a binding partner. The increase in helical content was lost upon heat denaturation (Figure 4B). TM5a has multiple unfolding transitions that overlap to give one broad unfolding transition at ~41 °C, close to that of FRL1-FH1FH2 (~39.5 °C, Figure 4C). There was no shift in the overall $T_{\rm M}$ of unfolding of the FRL1-FH1FH2-TM5a mixture, compared to that of the unmixed components. However, the unfolding of the complex was more cooperative, as evidenced by the greater peak height of the first derivative of the unfolding curve.

Thermal denaturation experiments show that mDia2-FH2 also binds to TM5a and TM2 and indicate the involvement of the FH2 domain (Figures 4D and E). The mDia2-FH2 domain is more helical and more stable than FRL1-FH1FH2 and unfolded with a $T_{\rm M}$ of ~46 °C. The C-terminal region of TM5a and TM2, encoded by exon 9d, is relatively unstable, and it unfolds during the first major transition of TM2 at ~40 °C (58). When mDia2-FH2 was bound to TM5a (Figure 3D) or TM2 (Figure 3E), the $T_{\rm M}$ of unfolding of the first transitions near 40 °C increased by ~1 to 2 °C, suggesting that the FH2 domain binds to the C-terminal region of tropomyosin.

Even though the CD binding studies indicate that the formin FH2 domain binds the C-terminal domain of tropomyosin, the end oriented toward the barbed end of the filament (52), relief of formin-dependent capping is isoform-specific and is influenced by the rest of the tropomyosin molecule, not just the C-terminus. TM5a relieves FRL1-FH1FH2 inhibition of barbed-end elongation, whereas TM2 affects mDia2, even though the two tropomyosins contain the same C-terminal sequence.

Tropomyosin Dissociates FRL1-FH1FH2 from the Sides of Filaments

Constructs containing the FH2 domain of some formins can bind to the sides of actin filaments and bundle them (51). In the ionic conditions optimal for tropomyosin binding (100 mM NaCl and 2 mM MgCl₂), FRL1-FH1FH2 bound poorly to actin (Figure 5). This is consistent with the observation of Harris et al. (59) and others (60) who found that side binding of formins to filamentous actin is sensitive to ionic strength. However, 85–88% of the FRL1-FH1FH2 bound to phalloidin-stabilized actin filaments (Figure 5).

Tropomyosin inhibited the side binding of FRL1-FH1FH2 to phalloidin-stabilized actin filaments, independent of the tropomyosin isoform (Figure 6). Filaments saturated with tropomyosin reduced formin binding by about 50%. Recombinant striated muscle TM, a form that lacks the N-terminal acetyl group required for high affinity binding (61,62), did not inhibit FRL1-FH1FH2 binding and serves as a control to show that inhibition of side binding of FRL1-FH1FH2 requires tropomyosin binding to the actin filament. The lack of isoform specificity

suggests that tropomyosin's inhibition of side binding is via a general mechanism, comparable to its inhibition of cross-linking and severing by cofilin, gelsolin, and related proteins (24-26,28) and the inhibition of nucleation of branches of filaments by the Arp2/3 complex (63).

Tropomyosin Does Not Unblock Filaments with Capping Protein at the Barbed End

Stable actin filaments with capping protein at the barbed end, such as those in adhering junctions or at the Z-disc or dense bodies in muscle, contain tropomyosin. We tested the ability of tropomyosin to regulate barbed-end capping by three recombinant CapZ isoforms: $\alpha 1\beta 1$ ($\beta 1$ is muscle-specific; CapZ localized at the Z-line), $\alpha 1\beta 2$ and $\alpha 2\beta 2$ ($\beta 2$ is widely expressed) (47,64,65). Actin polymerization was nucleated by actin filaments with bound capping protein and tropomyosin. We selected conditions in which capping in the absence of tropomyosin was incomplete. Tropomyosin neither relieved the inhibition of elongation by capping protein as it did formin-capped filaments (above) nor improved capping activity (Figure 7). In all cases, the amount of polymer formed was well below that of the actin and similar to that with capping protein alone, indicating no significant effect of tropomyosin on capping protein function. The results of binding measurements using circular dichroism were negative (results not shown).

DISCUSSION

We show here for the first time that tropomyosin regulates the dynamics of the barbed, plus end of the actin filament, alone, by uncoupling the capping function of formins and by inhibiting the binding of FRL1 to the sides of filaments. Tropomyosin's role as a universal regulator of the actin filament now extends to both ends of the filament, in addition to overall filament stiffening and stabilization and the allosteric regulation of the actin filament in contractile function via its cooperative regulation of myosin and with troponin [reviewed in (19-21)].

Formins are leaky, processive cappers [reviewed in (3,4)]. The binding of the mDia1-FH2 dimer to the barbed end makes the actin filament more flexible over a long range and the entire filament more dynamic (60,66), just as binding of gelsolin at the filament end cooperatively influences filament structure and dynamics (67). Assuming that the FH2 domains of other formins have similar allosteric effects, the stabilization of the filament by tropomyosin may antagonize the effect of formin and reduce its ability to block the barbed end, thereby making it a leakier but still processive capper.

An effect of tropomyosin itself on the barbed end is illustrated by the marked stimulation of elongation at substoichiometric TM5a and its neutralization by FRL1-FH1FH2. The results infer that TM5a has a specific influence on the barbed end where the first molecules may bind the filament. As tropomyosin cooperatively saturates the filament and stabilizes the structure, stimulation of barbed-end elongation is dampened. The binding of FRL1-FH1FH2 at the barbed end also depresses the effect of substoichiometric TM5a, presumably because the cap formed by the FH2 domain with the two terminal actin subunits (68) impedes the addition of actin subunits. Tropomyosin, by binding to the FH2 domain, relieves capping by formin. In this scenario, tropomyosin is an *activator*. Although there is no specific structural information available, we postulate that the flexible hinge near the C-terminus of tropomyosin (69-71) allows interaction of the C-terminal chains with helical regions of the FH2 domain, possibly forming a coiled coil as they do with the N-terminus of tropomyosin and troponin T (69,71).

Tropomyosin has no observed effect on capping protein function, and we have no evidence for direct interaction between the proteins. Formins compete with CapZ proteins for the barbed end, but they should operate by distinct mechanisms because they have different structures and different proposed interactions with the filament end (68,72-74). Tropomyosin binds to the G2 domain of gelsolin (37), but it has no marked effect on gelsolin capping of long filaments

(34). The cartoons in Figure 8 illustrate the different relationships of tropomyosin to the formin FH2 domain and capping protein at the barbed end of the actin filament (based on models in refs 68 and 74). Tropomyosin binds along the sides of the actin filament and, at the barbed end, binds to the formin FH2 domain but not capping protein.

The influence of tropomyosin on the side binding of FRL1-FH1FH2 (and presumably other formins) occurs by a general mechanism, without tropomyosin isoform specificity. In this regard, it is similar to the effect of tropomyosin on the side binding, severing, and bundling by other actin binding proteins, including DNase I, gelsolin, the Arp2/3 complex, and cofilin (23-26,28). Tropomyosin and formin acting together would ensure the formation of a stable, straight filament, protected from severing that could eventually be capped by a CapZ family protein in a stable cellular structure such as an adhering junction or a Z-disk in striated muscle. We suggest that tropomyosin may have a similar role in modulating the formation of stable actin filaments in concert with Ena/VASP family proteins (75-77).

Acknowledgements

We thank Elizabeth Harris for the preparation of the formin constructs and Dr. Nandini Bhattacharya for the capping proteins.

References

- dos Remedios CG, Chhabra D, Kekic M, Dedova IV, Tsubakihara M, Berry DA, Nosworthy NJ. Actin binding proteins: regulation of cytoskeletal microfilaments. Physiol ReV 2003;83:433–473. [PubMed: 12663865]
- 2. Nicholson-Dykstra S, Higgs HN, Harris ES. Actin dynamics: growth from dendritic branches. Curr Biol 2005;15:R346–357. [PubMed: 15886095]
- 3. Faix J, Grosse R. Staying in shape with formins. DeV Cell 2006;10:693-706. [PubMed: 16740473]
- 4. Kovar DR. Cell polarity: formin on the move. Curr Biol 2006;16:R535–R538. [PubMed: 16860728]
- 5. Higgs HN. Formin proteins: a domain-based approach. Trends Biochem Sci 2005;30:342–353. [PubMed: 15950879]
- Kovar DR, Pollard TD. Insertional assembly of actin filament barbed ends in association with formins produces piconewton forces. Proc Natl Acad Sci U S A 2004;101:14725–14730. [PubMed: 15377785]
- Romero S, Le Clainche C, Didry D, Egile C, Pantaloni D, Carlier MF. Formin is a processive motor that requires profilin to accelerate actin assembly and associated ATP hydrolysis. Cell 2004;119:419– 429. [PubMed: 15507212]
- 8. Watanabe N, Higashida C. Formins: processive cappers of growing actin filaments. Exp Cell Res 2004;301:16–22. [PubMed: 15501440]
- Michelot A, Derivery E, Paterski-Boujemaa R, Guerin C, Huang S, Parcy F, Staiger CJ, Blanchoin L. A novel mechanism for the formation of actin-filament bundles by a nonprocessive formin. Curr Biol 2006;16:1924–1930. [PubMed: 17027489]
- 10. Mejillano MR, Kojima S, Applewhite DA, Gertler FB, Svitkina TM, Borisy GG. Lamellipodial versus filopodial mode of the actin nanomachinery: pivotal role of the filament barbed end. Cell 2004;118:363–373. [PubMed: 15294161]
- 11. Wear MA, Cooper JA. Capping protein: new insights into mechanism and regulation. Trends Biochem Sci 2004;29:418–428. [PubMed: 15362226]
- Waddle JA, Karpova TS, Waterston RH, Cooper JA. Movement of cortical actin patches in yeast. J Cell Biol 1996;132:861–870. [PubMed: 8603918]
- 13. Kim K, Yamashita A, Wear MA, Maeda Y, Cooper JA. Capping protein binding to actin in yeast: biochemical mechanism and physiological relevance. J Cell Biol 2004;164:567–580. [PubMed: 14769858]
- 14. Engqvist-Goldstein AE, Drubin DG. Actin assembly and endocytosis: from yeast to mammals. Annu ReV Cell DeV Biol 2003;19:287–332. [PubMed: 14570572]

15. Schafer DA, Jennings PB, Cooper JA. Dynamics of capping protein and actin assembly in vitro: uncapping barbed ends by polyphosphoinositides. J Cell Biol 1996;135:169–179. [PubMed: 8858171]

- 16. Uruno T, Remmert K, Hammer JA III. CARMIL is a potent capping protein antagonist: identification of a conserved CARMIL domain that inhibits the activity of capping protein and uncaps capped actin filaments. J Biol Chem 2006;281:10635–10650. [PubMed: 16434392]
- 17. Bhattacharya N, Ghosh S, Sept D, Cooper JA. Binding of myotrophin/V-1 to actin-capping protein: implications for how capping protein binds to the filament barbed end. J Biol Chem 2006;281:31021–31030. [PubMed: 16895918]
- Kovar DR, Harris ES, Mahaffy R, Higgs HN, Pollard TD. Control of the assembly of ATP- and ADPactin by formins and profilin. Cell 2006;124:423–435. [PubMed: 16439214]
- 19. Gunning PW, Schevzov G, Kee AJ, Hardeman EC. Tropomyosin isoforms: divining rods for actin cytoskeleton function. Trends Cell Biol 2005;15:333–341. [PubMed: 15953552]
- 20. Perry SV. Vertebrate tropomyosin: distribution, properties and function. J Muscle Res Cell Motil 2001;22:5–49. [PubMed: 11563548]
- 21. Brown JH, Cohen C. Regulation of muscle contraction by tropomyosin and troponin: how structure illuminates function. Adv Protein Chem 2005;71:121–159. [PubMed: 16230111]
- 22. Ishiwata S, Fujime S. Effect of calcium ions on the flexibility of reconstituted thin filaments of muscle studied by quasielastic scattering of laser light. J Mol Biol 1972;68:511–522. [PubMed: 4672238]
- 23. Hitchcock SE, Carisson L, Lindberg U. Depolymerization of F-actin by deoxyribonuclease I. Cell 1976;7:531–542. [PubMed: 133761]
- 24. Bernstein BW, Bamburg JR. Tropomyosin binding to F-actin protects the F-actin from disassembly by brain actin-depolymerizing factor (ADF). Cell Motil 1982;2:1–8. [PubMed: 6890875]
- Kobayashi R, Nonomura Y, Okano A, Tashima Y. Purification and some properties of porcine kidney tropomyosin. J Biochem (Tokyo) 1983;94:171–179. [PubMed: 6619108]
- 26. Pruliere G, d'Albis A, der Terrossian E. Effect of tropomyosin on the interactions of actin with actin-binding proteins isolated from pig platelets. Eur J Biochem 1986;159:535–547. [PubMed: 3019696]
- 27. Ludescher RD, Liu Z. Characterization of skeletal muscle actin labeled with the triplet probe erythrosin-5-iodoacetamide. Photochem Photobiol 1993;58:858–866. [PubMed: 8310009]
- 28. DesMarais V, Ichetovkin I, Condeelis J, Hitchcock-DeGregori SE. Spatial regulation of actin dynamics: a tropomyosin-free, actin-rich compartment at the leading edge. J Cell Sci 2002;115:4649–4660. [PubMed: 12415009]
- 29. Ono S, Ono K. Tropomyosin inhibits ADF/cofilindependent actin filament dynamics. J Cell Biol 2002;156:1065–1076. [PubMed: 11901171]
- 30. Yu R, Ono S. Dual roles of tropomyosin as an F-actin stabilizer and a regulator of muscle contraction in *Caenorhabditis elegans* body wall muscle. Cell Motil Cytoskeleton 2006;63:659–672. [PubMed: 16937397]
- 31. Fischer RS, Fowler VM. Tropomodulins: life at the slow end. Trends Cell Biol 2003;13:593–601. [PubMed: 14573353]
- 32. Broschat KO, Weber A, Burgess DR. Tropomyosin stabilizes the pointed end of actin filaments by slowing depolymerization. Biochemistry 1989;28:8501–8506. [PubMed: 2605200]
- 33. Broschat KO. Tropomyosin prevents depolymerization of actin filaments from the pointed end. J Biol Chem 1990;265:21323–21329. [PubMed: 2250026]
- 34. Kostyukova AS, Hitchcock-DeGregori SE. Effect of the structure of the N terminus of tropomyosin on tropomodulin function. J Biol Chem 2004;279:5066–5071. [PubMed: 14660556]
- 35. Ishikawa R, Yamashiro S, Matsumura F. Annealing of gelsolin-severed actin fragments by tropomyosin in the presence of Ca2+. Potentiation of the annealing process by caldesmon. J Biol Chem 1989;264:16764–16770. [PubMed: 2550459]
- 36. Koepf EK, Burtnick LD. Interaction of plasma gelsolin with tropomyosin. FEBS Lett 1992;309:56–58. [PubMed: 1324850]
- 37. Maciver SK, Ternent D, McLaughlin PJ. Domain 2 of gelsolin binds directly to tropomyosin. FEBS Lett 2000;473:71–75. [PubMed: 10802062]

38. Nyakern-Meazza M, Narayan K, Schutt CE, Lindberg U. Tropomyosin and gelsolin cooperate in controlling the microfilament system. J Biol Chem 2002;277:28774–28779. [PubMed: 12048198]

- 39. Liu HP, Bretscher A. Disruption of the single tropomyosin gene in yeast results in the disappearance of actin cables from the cytoskeleton. Cell 1989;57:233–242. [PubMed: 2649250]
- 40. Evangelista M, Blundell K, Longtine MS, Chow CJ, Adames N, Pringle JR, Peter M, Boone C. Bni1p, a yeast formin linking cdc42p and the actin cytoskeleton during polarized morphogenesis. Science 1997;276:118–122. [PubMed: 9082982]
- 41. Sagot I, Klee SK, Pellman D. Yeast formins regulate cell polarity by controlling the assembly of actin cables. Nat Cell Biol 2002;4:42–50. [PubMed: 11740491]
- 42. Spudich JA, Watt S. The regulation of rabbit skeletal muscle contraction. I. Biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. J Biol Chem 1971;246:4866–4871. [PubMed: 4254541]
- 43. Kouyama T, Mihashi K. Fluorimetry study of N-(1-pyrenyl)iodoacetamide-labelled F-actin. Local structural change of actin protomer both on polymerization and on binding of heavy meromyosin. Eur J Biochem 1981;114:33–38. [PubMed: 7011802]
- 44. Moraczewska J, Nicholson-Flynn K, Hitchcock-DeGregori SE. The ends of tropomyosin are major determinants of actin affinity and myosin subfragment 1-induced binding to F-actin in the open state. Biochemistry 1999;38:15885–15892. [PubMed: 10625454]
- 45. Hitchcock-DeGregori SE, Lewis SF, Chou TM. Tropomyosin lysine reactivities and relationship to coiled-coil structure. Biochemistry 1985;24:3305–3314. [PubMed: 3927977]
- 46. Harris ES, Li F, Higgs HN. The mouse formin, FRLalpha, slows actin filament barbed end elongation, competes with capping protein, accelerates polymerization from monomers, and severs filaments. J Biol Chem 2004;279:20076–20087. [PubMed: 14990563]
- 47. Schafer DA, Korshunova YO, Schroer TA, Cooper JA. Differential localization and sequence analysis of capping protein beta-subunit isoforms of vertebrates. J Cell Biol 1994;127:453–465. [PubMed: 7929588]
- 48. Edelhoch H. Spectroscopic determination of tryptophan and tyrosine in proteins. Biochemistry 1967;6:1948–1954. [PubMed: 6049437]
- 49. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 1970;227:680–685. [PubMed: 5432063]
- 50. Brahms S, Brahms J. Determination of protein secondary structure in solution by vacuum ultraviolet circular dichroism. J Mol Biol 1980;138:149–178. [PubMed: 7411608]
- 51. Harris ES, Rouiller I, Hanein D, Higgs HN. Mechanistic differences in actin bundling activity of two mammalian formins, FRL1 and mDia2. J Biol Chem 2006;281:14383–14392. [PubMed: 16556604]
- 52. Ohtsuki I. Molecular arrangement of troponin-T in the thin filament. J Biochem (Tokyo) 1979;86:491–497. [PubMed: 479140]
- 53. Teubner A, Wegner A. The rate of annealing of actin tropomyosin filaments depends strongly on the length of the filaments. Biochim Biophys Acta 1996;1297:214–218. [PubMed: 8917624]
- 54. Hitchcock-DeGregori SE, Sampath P, Pollard TD. Tropomyosin inhibits the rate of actin polymerization by stabilizing actin filaments. Biochemistry 1988;27:9182–9185. [PubMed: 3242622]
- 55. Greenfield NJ. Analysis of circular dichroism data. Methods Enzymol 2004;383:282–317. [PubMed: 15063655]
- 56. Greenfield NJ, Kostyukova AS, Hitchcock-DeGregori SE. Structure and tropomyosin binding properties of the N-terminal capping domain of tropomodulin 1. Biophys J 2005;88:372–383. [PubMed: 15475586]
- 57. Kostyukova AS, Rapp BA, Choy A, Greenfield NJ, Hitchcock-DeGregori SE. Structural requirements of tropomodulin for tropomyosin binding and actin filament capping. Biochemistry 2005;44:4905–4910. [PubMed: 15779917]
- 58. Hammell RL, Hitchcock-DeGregori SE. Mapping the functional domains within the carboxyl terminus of alpha-tropomyosin encoded by the alternatively spliced ninth exon. J Biol Chem 1996;271:4236–4242. [PubMed: 8626768]
- 59. Harris ES, Higgs HN. Biochemical analysis of mammalian formin effects on actin dynamics. Methods Enzymol 2006;406:190–214. [PubMed: 16472659]

60. Bugyi B, Papp G, Hild G, Lorinczy D, Nevalainen EM, Lappalainen P, Somogyi B, Nyitrai M. Formins regulate actin filament flexibility through long range allosteric interactions. J Biol Chem 2006;281:10727–10736. [PubMed: 16490788]

- 61. Heald RW, Hitchcock-DeGregori SE. The structure of the amino terminus of tropomyosin is critical for binding to actin in the absence and presence of troponin. J Biol Chem 1988;263:5254–5259. [PubMed: 2965699]
- 62. Urbancikova M, Hitchcock-DeGregori SE. Requirement of amino-terminal modification for striated muscle alpha-tropomyosin function. J Biol Chem 1994;269:24310–24315. [PubMed: 7929088]
- 63. Blanchoin L, Pollard TD, Hitchcock-DeGregori SE. Inhibition of the Arp2/3 complex-nucleated actin polymerization and branch formation by tropomyosin. Curr Biol 2001;11:1300–1304. [PubMed: 11525747]
- 64. Hart MC, Korshunova YO, Cooper JA. Vertebrates have conserved capping protein alpha isoforms with specific expression patterns. Cell Motil Cytoskeleton 1997;38:120–132. [PubMed: 9331217]
- 65. Hart MC, Cooper JA. Vertebrate isoforms of actin capping protein beta have distinct functions in vivo. J Cell Biol 1999;147:1287–1298. [PubMed: 10601341]
- 66. Papp G, Bugyi B, Ujfalusi Z, Barko S, Hild G, Somogyi B, Nyitrai M. Conformational changes in actin filaments induced by formin binding to the barbed end. Biophys J 2006;91:2564–2572. [PubMed: 16829561]
- 67. Prochniewicz E, Zhang Q, Janmey PA, Thomas DD. Cooperativity in F-actin: binding of gelsolin at the barbed end affects structure and dynamics of the whole filament. J Mol Biol 1996;260:756–766. [PubMed: 8709153]
- Otomo T, Tomchick DR, Otomo C, Panchal SC, Machius M, Rosen MK. Structural basis of actin filament nucleation and processive capping by a formin homology 2 domain. Nature 2005;433:488– 494. [PubMed: 15635372]
- 69. Greenfield NJ, Palm T, Hitchcock-DeGregori SE. Structure and interactions of the carboxyl terminus of striated muscle alpha-tropomyosin: it is important to be flexible. Biophys J 2002;83:2754–2766. [PubMed: 12414708]
- 70. Li Y, Mui S, Brown JH, Strand J, Reshetnikova L, Tobacman LS, Cohen C. The crystal structure of the C-terminal fragment of striated-muscle alpha-tropomyosin reveals a key troponin T recognition site. Proc Natl Acad Sci U S A 2002;99:7378–7383. [PubMed: 12032291]
- 71. Greenfield NJ, Huang YJ, Swapna GV, Bhattacharya A, Rapp B, Singh A, Montelione GT, Hitchcock-Degregori SE. Solution NMR structure of the junction between tropomyosin molecules: implications for actin binding and regulation. J Mol Biol 2006;364:80–96. [PubMed: 16999976]
- 72. Yamashita A, Maeda K, Maeda Y. Crystal structure of CapZ: structural basis for actin filament barbed end capping. EMBO J 2003;22:1529–1538. [PubMed: 12660160]
- 73. Xu Y, Moseley JB, Sagot I, Poy F, Pellman D, Goode BL, Eck MJ. Crystal structures of a formin homology-2 domain reveal a tethered dimer architecture. Cell 2004;116:711–723. [PubMed: 15006353]
- 74. Narita A, Takeda S, Yamashita A, Maeda Y. Structural basis of actin filament capping at the barbedend: a cryo-electron microscopy study. EMBO J 2006;25:5626–5633. [PubMed: 17110933]
- 75. Krause M, Dent EW, Bear JE, Loureiro JJ, Gertler FB. Ena/VASP proteins: regulators of the actin cytoskeleton and cell migration. Annu ReV Cell DeV Biol 2003;19:541–564. [PubMed: 14570581]
- 76. Barzik M, Kotova TI, Higgs HN, Hazelwood L, Hanein D, Gertler FB, Schafer DA. Ena/VASP proteins enhance actin polymerization in the presence of barbed end capping proteins. J Biol Chem 2005;280:28653–286562. [PubMed: 15939738]
- Scott JA, Shewan AM, den Elzen NR, Loureiro JJ, Gertler FB, Yap AS. Ena/VASP proteins can regulate distinct modes of actin organization at cadherin-adhesive contacts. Mol Biol Cell 2006;17:1085–1095. [PubMed: 16371509]
- 78. Casella JF, Maack DJ, Lin S. Purification and initial characterization of a protein from skeletal muscle that caps the barbed ends of actin filaments. J Biol Chem 1986;261:10915–10921. [PubMed: 3733738]




Figure 1. Effect of tropomyosin on elongation at the barbed end of actin filaments. (A) A representative time course of elongation of $0.5\,\mu\mathrm{M}$ Mg-G-actin (10% pyrene labeled) from $2\,\mu\mathrm{M}$ F-actin seeds (1) or F-actin in the presence of saturating amounts of tropomyosin (44): $0.5\,\mu\mathrm{M}$ TM5a (2), 2 $\mu\mathrm{M}$ TM2 (3), and 1 $\mu\mathrm{M}$ stTM (4) in the absence of formin or capping protein, or G-actin alone without F-actin seeds (5) in 100 mM NaCl, 2 mM MgCl₂, 10 mM TrisHCl at pH 7.4, 0.2 mM ATP, 0.2 mM EGTA, and 0.2 mM DTT. (B) The initial velocities (V_0) relative to F-actin alone, taken as 100%, with standard error (n = 4–5). The gray curves are the fits to the data.




Figure 2. Effect of tropomyosin on elongation at the barbed end of actin filaments in the presence of formin. (A, C, E) Representative time courses. (B, D, F) Relative initial rates of elongation. (A, B) 25 nM FRL1-FH1FH2, 1 μ M F-actin seeds; n=4. (C, D) 50 nM mDia2-FH2, 2 μ M F-actin seeds; no TM, n=4; TM5a, n=4; TM2, n=2; stTM, n=3. (E, F) 25 nM mDia1-FH2, 2 μ M F-actin seeds; noTM, n=3; TM5a, n=1; TM2, n=2; stTM, n=2. The tropomyosin concentrations and ionic conditions are the same as those in Figure 1. Curves: 1, F-actin alone (no formin or tropomyosin); 2, F-actin with formin, no tropomyosin; 3, formin and TM5a; 4, formin and TM2; 5, formin and stTM; 6, G-actin alone, no F-actin seeds; and 7, G-actin with formin, no F-actin seeds. FRL1-FH1FH2 had no effect with G-actin alone and is not shown. The gray curves are the fits to the data. Ac: F-actin, without tropomyosin or formin (from curve 1).




Figure 3. Relief of FRL1-FH1FH2 inhibition of barbed-end elongation by TM5a does not require saturation of the actin filaments with tropomyosin. The results from a representative experiment are shown; 50 nM FRL1-FH1FH2, 2 μ M F-actin seeds, TM5a, varied. (A) The V_0 of barbed-end elongation as a function of TM5a concentration in the absence of FRL1-FH1FH2 (\bullet) and in the presence of 50 nM FRL1-FH1FH2 (\bullet). The panel shows the data with FRL1-FH1FH2 on an expanded scale. (B) The time course of increase in pyrene fluorescence in the absence of FRL1-FH1FH2 in the absence of TM5a (1), a subsaturating concentration, 0.05 μ M TM5a (2), and a saturating concentration, 1 μ M TM5a (3). (C) The time course of increase in pyrene fluorescence in the presence of 50 nM FRL1-FH1FH2 in the absence of

TM5a (1), a subsaturating concentration, 0.05 μ M TM5a (2), and a saturating concentration, 1 μ M TM5a (3). The gray curves are the fits to the data. (D) SDS–PAGE gels of the pellets following centrifugation at the end of the polymerization experiment. Even numbers, with FRL1-FH1FH2; odd numbers, no FRL1. TM5a concentration: 1 and 2, no TM5a; 3 and 4, 0.05 μ M; 5 and 6, 0.1 μ M; 7 and 8, 0.3 μ M; 9 and 10, 0.5 μ M; 11 and 12, 0.7 μ M; 13 and 14, 1.0 μ M. The ionic conditions are the same as those in Figure 1.




Figure 4. Circular dichroism measurements of the interactions of formin domains with tropomyosin. (A) Spectra of FRL1-FH1FH2 (—); TM5a (····); a 1:1 mixture of FRL1-FH1FH2 and TM5a (- - -); and addition of the curves of unmixed FRL and TM5a (-··-). (B) The change in ellipticity at 222 nm as a function of temperature of the samples in panel A. (C) The first derivatives of the curves in panel B. (D) The first derivative of the unfolding curves of mDia2-FH2 (—); TM5a (····); a 1:1 mixture of mDia2-FH2 and TM5a (- - -); and addition of the curves of the unmixed components (-··-). (E) The first derivatives of the change in ellipticity at 222 nm as a function of temperature of mDia2-FH2 (—); TM2 (····); a 1:1 mixture of mDia2- FH2 and TM2 (- - -); and addition of the curves of the unmixed components (-··-). The $T_{\rm M}$ of the unfolding transition of TM5a increases ~1 °C when bound to mDia2-FH2 (D), and the $T_{\rm M}$ of the first unfolding transition of TM2 is increased by ~2 °C (E). Conditions: 0 °C in 100 mM NaCl, 10 mM Tris-HCl, 0.2 mM EGTA, 0.1 mM MgCl₂, 0.01% NaN₂, and 0.5 mM dithiothreitol at pH 7.4. The formin and tropomyosin concentrations were 1 μ M.




Figure 5. Stabilization of F-actin by phalloidin allows side binding of FRL1-FH1FH2 at near physiological ionic strength. F-actin (5 μ M), in the absence or presence of 5 μ M phalloidin, was cosedimented with 0.5 μ M FRL1-FH1FH2 in 100 mM NaCl, 2 mM MgCl₂, 10 mM TrisHCl at pH 7.4, 0.2 mM ATP, 0.2 Mm EGTA, and 0.2 mM DTT. The actin and the formin were quantified in SDS gels of the pellets and supernatants. The percent actin (gray bars) and FRL1-FH1FH2 (black bars) in the pellets are illustrated with standard errors, n=3.




Figure 6. Tropomyosin binding to F-actin partially dissociates FRL1-FH1FH2 from the filament side independent of the isoform. Phalloidin-stabilized F-actin (5 μ M) was cosedimented with 0.5 μ M FRL1-FH1FH2 and 1.3 μ M TM5a, TM2, stTM, or recombinant, unacetylated stTM expressed in *E. coli*, stTM(rec). The proteins in the pellets were quantified following SDS–PAGE. (A) A gel from a representative experiment. (B) FRL1-FH1FH2/actin ratio in the pellet; the value in the absence of TM was set at 100%. (C) TM/actin ratio in the pellet with arbitrary units. A value of 0.35–0.4 represents saturation. stTM(rec) has low affinity for actin and is used here as a control to show that dissociation from the sides of the filaments requires binding of tropomyosin. The bar graphs show standard error, n = 3.




Figure 7. Tropomyosin does not unblock filaments with capping protein at the barbed end. (A) Spectrinactin seeds were used to nucleate polymerization of pyrene-labeled monomeric actin (5% labeled). The seeds were prepared as previously described (15,78), by incubating spectrin-actin with F-actin overnight to give F-actin seeds ~200 actins long, capped at their pointed ends by spectrin (1.7 nM spectrin-actin seeds and 0.5 μ M F-actin). stTM was added to the F-actin to 2 μ M before addition of the spectrin-actin and incubated overnight. The capping protein (α 1 β 2) was added to the filaments just prior to the initiation of polymerization. The F-actin seeds were diluted 1:10 into Mg-G-actin (2 μ M, 5% pyrene labeled) to seed polymerization in 100 mM KCl, 2 mM MgCl₂, 10 Mm TrisHCl at pH 7.5, 0.2 mM ATP, 0.5 mM DTT, 0.2 mM CaCl₂,

and 1 mM EGTA. When present, tropomyosin concentration was 2 μ M to ensure saturation of any polymer formed. Curves: 1, F-actin seeds; 2, F-actin seeds and 1.5 nM CP; 3, F-actin seeds, stTM, and 1.5 nM CP; 4, F-actin seeds and 4 nM CP; 5, F-actin seeds, stTM, and 4 nM CP; 6, G-actin alone. (B, C) Elongation experiments were carried out as described in Figures 1 and 2, and Experimental Procedures, at a final concentration of 1.5 nM CP. (B) Data from a representative data set with CP α 2 β 2: 1, actin alone; 2, CP and no tropomyosin; 3, CP + TM5a; 4, CP + TM2; 5, CP + stTM. (C) F_{max} (at 60 min) relative to the initial fluorescence (F_{o}) for actin alone, and the three capping proteins with the three tropomyosin isoforms, mean with standard error for 3 experiments with CP alone and TM5a, two to three experiments with TM2, and one to two experiments with stTM. The numbering corresponds to that in B.




Figure 8.

Regulatory proteins at the barbed end of the actin filament. A cartoon of the barbed end of the actin filament shows the formin FH2 domain based on ref 68 or the capping protein based on ref 74. Tropomyosin binds along the actin filament (not precisely positioned in the cartoon; the tropomyosin on only one side of the filament is shown) such that the C-terminal end interacts with formin-FH2, regulating its assembly function but with no effect on the capping protein.