

Med Chem. Author manuscript; available in PMC 2012 January 13.

Published in final edited form as:

J Med Chem. 2011 January 13; 54(1): 342–353. doi:10.1021/jm1012165.

Design and Synthesis of an Orally Active Metabotropic Glutamate Receptor Subtype-2 (mGluR2) Positive Allosteric Modulator (PAM) that Decreases Cocaine Self-administration in Rats

Raveendra-Panickar Dhanya^{1,†}, Shyama Sidique^{1,†}, Douglas J. Sheffler², Hilary Highfield Nickols³, Ananda Herath¹, Li Yang¹, Russell Dahl¹, Robert Ardecky¹, Svetlana Semenova⁴, Athina Markou⁴, P. Jeffrey Conn², and Nicholas D. P. Cosford^{1,*}

- ¹ Apoptosis and Cell Death Research Program and Conrad Prebys Center for Chemical Genomics, Sanford-Burnham Medical Research Institute, 10901 N. Torrey Pines Rd., La Jolla, CA 92037
- ² Department of Pharmacology, Vanderbilt University Medical Center, Nashville, TN 37232
- ³ Department of Pathology, Vanderbilt University Medical Center, Nashville, TN 37232
- ⁴ Department of Psychiatry, School of Medicine, University of California San Diego, La Jolla, CA 92093

Abstract

The modification of 3'-((2-cyclopentyl-6,7-dimethyl-1-oxo-2,3-dihydro-1*H*-inden-5-yloxy)methyl)biphenyl-4-carboxylic acid (BINA, 1) by incorporating heteroatoms into the structure and replacing the cyclopentyl moiety led to the development of new mGluR2 positive allosteric modulators (PAMs) with optimized potency and superior drug-like properties. These analogues are more potent than 1 *in vitro*, and are highly selective for mGluR2 *vs.* other mGluR subtypes. They have significantly improved pharmacokinetic (PK) properties, with excellent oral bioavailability and brain penetration. The benzisothiazol-3-one derivative 14 decreased cocaine self-administration in rats, providing proof-of-concept for the use of mGluR2 PAMs for the treatment of cocaine dependence.

Keywords

Metabotropic glutamate receptors; agonist; positive allosteric modulators; BINA; cocaine self-administration

Cocaine addiction is a chronic relapsing disorder affecting more than 1.6 million Americans.

Relapse rates among cocaine users is in the range of 94–99%, the highest among all commonly abused drugs.
Chronic cocaine abuse can lead to severe neurotoxicity, psychosis, lethargy, depression, or potentially death through a fatal overdose. Currently there is no effective treatment for cocaine dependence, and therefore cocaine addiction constitutes a major public health problem. Consequently, there is a significant need to identify new therapeutic agents for the treatment of cocaine and other psychomotor stimulant addictions. Recent findings suggest that neuroadaptations in glutamatergic

^{*}Corresponding Author: To whom correspondence should be addressed. Tel: 858-646-3100. Fax: 858-795-5225. pcosford@sanfordburnham.org.

[†]These authors contributed equally to this work.

transmission produced by repeated exposure to cocaine or other drugs of abuse are likely to contribute to the maintenance of addictive behaviors including drug use, craving and relapse to drug taking in humans.³ Specifically, it has been shown that repeated cocaine exposure alters the function of Group II metabotropic glutamate receptors (mGluRs).^{4a} The Group II mGluRs include the mGluR2 and mGluR3 subtypes, which couple to G_{i/o} proteins to negatively regulate the activity of adenylyl cyclase.^{4b} Brain regions implicated in different aspects of drug abuse and drug dependence, including the cerebral cortex, hippocampus, striatum, amygdala, frontal cortex and nucleus accumbens display high levels of mGluR2 and mGluR3 binding,⁵ suggesting a role for the mGluR2/3 subtypes in the development of cocaine dependence and as potential targets for therapeutic agents.^{3a,3e,3f,6}

Orthosteric (glutamate site) mGluR2/3 agonists such as LY379268^{6g} are constrained amino acid analogues that do not exhibit selectivity for mGluR2 versus mGluR3, presumably because of the high degree of sequence homology at the glutamate binding site for these two receptors. 6h LY379268 has been shown to decrease glutamate levels 7 and to attenuate cocaine self-administration both in rats^{6c,8} and in squirrel monkeys.^{6a} However, LY379268 also inhibits responding for food and food-seeking behavior^{6c,6e,9}, suggesting that mGluR2/3 agonists exhibit non-selective actions on responding for drug and non-drug reinforcers. We recently initiated investigations to determine the effect of selectively activating mGluR2 on cocaine dependence. 8 Localization studies suggest that mGluR2 acts predominantly as a presynaptic autoreceptor to modulate the release of glutamate into the synaptic cleft. ¹⁰ In recent years, there have been several reports on the use of highthroughput screening (HTS) for the identification of small molecule mGluR2 positive allosteric modulators (PAMs). 11 The activity of these compounds is of interest because of their subtype-selectivity, ability to overcome receptor desensitization and potential for improved drug-like properties. 12 The use of a selective mGluR2 PAM in an appropriate in vivo model therefore represents an attractive approach to elucidate the role of mGluR2 in cocaine dependence. We previously disclosed the structure activity relationship (SAR) around a series of pyrimidine derivatives with activity as mGluR2 PAMs. 11b The synthesis and characterization of 3'-((2-cyclopentyl-6,7-dimethyl-1-oxo-2,3-dihydro-1H-inden-5yloxy)methyl)biphenyl-4-carboxylic acid (BINA, 1) has been previously reported 13 and we have shown that this compound possesses antipsychotic and anxiolytic effects in mice. 14 Most relevant to the present report, we recently showed that compound 1, unlike mGluR2/3 agonists, decreased cocaine self-administration in rats at doses that did not affect responding for food. 8 A recent report by the Merck group on a series of oxazolobenzimidazole-based mGluR2 PAMs that are orally active in an in vivo model of schizophrenia prompts us to disclose our own efforts on the development of orally active mGluR2 PAMs. 15b

While compound 1 is brain penetrant and selective, it lacks potency for mGluR2 both *in vitro* and *in vivo* and has sub-optimal pharmacokinetic (PK) properties. We therefore set out to design and synthesize new selective mGluR2 PAMs based on the compound 1 scaffold in order to improve both potency and PK properties. These new PAMs are suitable for *in vivo* proof-of-concept (POC) studies to evaluate the potential of mGluR2 as a therapeutic target for the treatment of cocaine dependence. We hypothesized that modifying the indanone ring of 1 by incorporating heteroatoms may lead to analogues with improved properties (Figure 1). Interestingly, to date there have been no reports on the synthesis and evaluation of analogues of compound 1 in which the indanone ring has been systematically modified in this way. In addition, as far as we are aware there have been no studies documenting the effects of modifying the cyclopentyl ring in this scaffold. Furthermore, despite several recent reports disclosing mGluR2 PAMs, ^{11,15} none have been evaluated in an *in vivo* model of drug dependence.

Initially, we synthesized and tested compound $\mathbf{2}$, an analogue of compound $\mathbf{1}$ previously reported to be roughly 4-fold more potent than $\mathbf{1}$ at mGluR2 (Figure 1)¹³ The *in vitro* potency of the compounds was assessed using a thallium flux assay performed in HEK-GIRK cells expressing rat mGluR2.¹⁶ In this assay compound $\mathbf{1}$ exhibited an EC₅₀ value of 380 nM, while compound $\mathbf{2}$ had an EC₅₀ of 180 nM, confirming an improved potency of $\mathbf{2}$ compared with $\mathbf{1}$ (Table 1). We therefore incorporated the 4-chlorobiphenyl-3-carboxylic acid moiety from $\mathbf{2}$ into the design of the new analogues.

We next focused on the synthesis of isoindolinone analogues (8a-8k). As depicted in Scheme 1, the preparation of 8a-8k commenced with the conversion of commercially available 4-hydroxyphthalic acid 3 into the corresponding phthalic anhydride derivative 4 under thermal conditions, which was further converted to the phthalimide derivatives 5a-5k by treatment with the appropriate amine derivative. Regioselective reduction of the 3-carbonyl group with zinc in acetic acid afforded the desired 5-hydroxyisoindolinone derivatives 6a-6k. To verify the regioselectivity of the reduction, intermediate 6a (R¹ = cyclopentyl) was also synthesized *via* an alternate procedure to confirm the structure unambiguously. Completion of the synthesis of compounds 8a-8k was achieved in two steps using an O-alkylation reaction with methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate to give the ester derivative 7a-7k which was saponified to the corresponding acid derivative 8a-8k using LiOH in THF.

The synthesis of an analogue of compound 1 based on the benzisothiazol-3-one scaffold is illustrated in Scheme 2. Methyl 4-methoxy thiosalicylate 9¹⁸ was converted to the corresponding amide 10 by treatment with cyclopentylamine in the presence of trimethylaluminum. The cyclization to access the benzisothiazol-3-one derivative 11 was accomplished by a PIFA-meditated *in situ* formation of an *N*-acylnitrenium ion, and its intramolecular trapping by the thiol moiety. Demethylation of 11 was achieved with BBr₃ in hot benzene, followed by O-alkylation with methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate to provide 13. Hydroxide-mediated demethylation led to the formation of byproducts, however clean saponification of 13 was accomplished with lithium iodide to complete the synthesis of 14.

We also designed and synthesized an analogue of compound 1 based on the isoxazol-3-one scaffold (21). Our approach to the synthesis of 21 began with the preparation of *N*-cyclopentylbenzohydroxamic acid derivative 18 as shown in Scheme 3. The synthesis began with 4-methoxysalicylic acid 15, which was transformed to 18 using a previously described procedure. ²⁰ Thus, 4-methoxysalicylic acid 15 was acetylated at the phenolic hydroxyl group prior to conversion to the acid chloride derivative 16. *N*-cyclopentylhydroxylamine 17 was reacted with 16 to provide hydroxamic acid 18 as a colorless solid. Cyclization of 18 under Mitsunobu conditions afforded *N*-cyclopentylbenzisoxazolinone 19, which was converted to the desired analogue 21 *via* intermediate 20 using the same conditions as those described for the preparation of compounds 8 and 14.

In addition, we designed and synthesized an analogue of compound **1** (**28**) based on the tetrahydroisoquinolinone scaffold. Our approach to the synthesis of **28** is outlined in Scheme 4. Thus, commercially available 3-methoxyphenethylamine was alkylated with iodocyclopentane to provide *N*-(3-methoxyphenethyl)cyclopentanamine (**23**). Treatment of **23** with methylchloroformate in the presence of triethylamine gave methyl cyclopentyl(3-methoxyphenethyl)carbamate **24**. Carbamate **24** was subjected to the Bischler-Napieralski cyclization conditions of Wang *et al* ²¹ (POCl₃.P₂O₅) to afford the corresponding tetrahydroisoquinolone derivative **25**. Compound **25** was then converted to the target BINA analogue **28** in three steps as described for the other analogues.

The new analogues 8a-k, 14, 21 and 28 were tested, along with 1 and 2, to determine in vitro activity and the data are summarized in Table 1. For these in vitro experiments, a thallium flux assay was performed in HEK-GIRK cells¹⁶ expressing rat mGluR2. The concentration-response relationship that potentiates the effect of an EC₂₀ concentration of glutamate was determined for each of the mGluR2 PAMs and this potency is expressed as an EC₅₀ value. The efficacy for potentiation of an EC₂₀ concentration of glutamate is presented as a percentage of the maximal glutamate response. Compounds 8a-j, 14, 21 and **28** were all more potent than **1** with compound **8a** being especially potent (EC₅₀ = 50 nM). Increase (8h) or decrease in ring size (8d, 8f) or the presence of bulky aryl substituents (8g, 8i) led to a decrease in potency compared with the N-cyclopentyl derivative 8a. On the other hand, branching (8b) or lengthening of the branched chain (8e, 8j) led to compounds with potency similar to that of 8a. Interestingly simple straight chain aliphatic substitution, as in the N-propyl derivative (8k), led to complete loss of activity. Enlargement of the lactam ring in the core scaffold from 5- (8a) to 6-membered (28) led to a 6-fold decrease in EC₅₀. In addition, the new analogues are highly selective for mGluR2 vs. other mGluR subtypes with no activity observed up to 10 µM for 8a and 14 (data not shown). We also determined the membrane permeability, in vitro plasma stability, and microsomal stability of the analogues to predict their in vivo pharmacological profile. As shown in Table 1, several of the new analogues displayed superior drug-like properties compared with 1, suggesting that these improvements may translate to an in vivo test.

Based on their overall *in vitro* profile, an isoindolinone derivative (**8a**) and a benzothiazolone derivative (**14**) were therefore selected for an *in vivo* assessment of their PK properties in rats. The rat PK data for compounds **1**, **8a** and **14** are shown in Table 2. Gratifyingly, both **8a** and **14** exhibited vastly superior PK profiles compared with **1**, with high values for C_{max} , AUC, and $t_{1/2}$. Incorporation of sulfur into the isoindolinone ring (**14**) did lead to a three fold decrease in potency, but showed superior PK properties (Table 2) compared to **8a**. Importantly, both compounds had excellent oral bioavailability and were highly brain penetrant. In particular, compound **14** showed exceptional oral bioavailability (86%) and brain levels (4.8 μ M) following a 10 mg/kg oral dose, with a brain:plasma ratio of 4.8. In contrast, the brain levels for **1** were below the limit of quantitation (LOQ) following oral dosing (10 mg/kg). Based on these data compound **14** was selected for evaluation in the cocaine self-administration procedure in rats.

The *in vivo* model was designed to investigate the effects of acute systemic (oral) administration of the mGluR2 PAM **14** on intravenous cocaine self-administration in rats that had 1 h access to cocaine (see Experimental for details). Additionally, we examined the effects of compound **14** on food-maintained responding to assess the selectivity of an orally active mGluR2 PAM on responding for cocaine compared to responding for the natural reinforcer food.

The results of the *in vivo* behavioral studies are shown in Figure 2. Oral administration of compound **14** significantly decreased cocaine self-administration in rats (Figure 2). Similarly, our previous work demonstrated that systemic (i.p.) administration of the mGluR2/3 agonist LY379268 significantly decreased cocaine self-administration in rats. However, compound **14** had a much less pronounced effect on food responding compared with the effects of LY379268. The effects of compound **14** on cocaine self-administration and the small effect of this compound on responding for food only at the highest dose tested (40 mg/kg) are consistent with our previous findings with compound **1**. These data indicate that a selective, orally active mGluR2 PAM is as effective as an mGluR2/3 agonist at reversing the effects of cocaine self administration in rats, but with a diminished or no effect on food-maintained responding. This selectivity of the effect may be due to the selectivity of **14** for mGluR2 compared with the dual acting mGluR2/3 agonist, and/or because of the

modulatory properties of a mGluR2 PAM compared to the direct actions of an agonist on the receptor(s).

In conclusion, using the mGluR2 PAM **1** as a template, we have designed and synthesized a series of new, highly potent mGluR2 PAMs. These compounds are highly selective for mGluR2 *vs.* other mGluR subtypes and have significantly improved *in vitro* ADME and *in vivo* PK properties, providing compounds with excellent oral bioavailability and brain penetration. On the basis of these data compound **14** was selected for *in vivo* studies in a model of cocaine dependence in rats, providing proof-of-concept for the use of an orally active mGluR2 PAM for the treatment of cocaine addiction.

Experimental

General Chemistry

All reactions were performed in oven-dried glassware under an atmosphere of argon with magnetic stirring. All solvents and chemicals used were purchased from Sigma-Aldrich or Acros, and were used as received without further purification. Purity and characterization of compounds were established by a combination of liquid chromatography-mass spectroscopy (LC-MS) and NMR analytical techniques and was >95% for all tested compounds. Silica gel column chromatography was carried out using prepacked silica cartridges from RediSep (ISCO Ltd.) and eluted using an Isco Companion system. Melting points were reordered on a MEL-TEMP® apparatus and are uncorrected. ¹H- and ¹³C-NMR spectra were obtained using a Jeol 400 spectrometer at 400 MHz and 100 MHz, respectively, unless otherwise mentioned, in which case a Varian 300 MHz spectrometer was used. Chemical shifts are reported in δ (ppm) relative to residual solvent peaks or TMS as internal standards. Coupling constants are reported in Hz. High-resolution ESI-TOF mass spectra were acquired at the Center for Mass Spectrometry at The Scripps Research Institute (La Jolla, CA). HPLC-MS analyses were performed on a Shimadzu 2010EV LCMS using the following conditions: Kromisil C18 column (reverse phase, 4.6 mm × 50 mm); a linear gradient from 10% acetonitrile and 90% water to 95% acetonitrile and 5% water over 4.5 min; flow rate of 1 mL/min; UV photodiode array detection from 200 to 300 nm.

5-Hydroxyisobenzofuran-1,3-dione (4).²³—4-Hydroxyphthalic acid (10 g, 54.9 mmol) was heated at 200 °C overnight in a round bottomed flask in the absence of solvent. The resulting material was cooled and washed with ethyl acetate to obtain 5-hydroxyisobenzofuran-1,3-dione 4 as a white solid in quantitative yield.

General procedure for the synthesis of thalimide derivatives (5a-5k)—4-Hydroxyphthalic anhydride **4** (5.4 g, 32.9 mmol), commercial amine (49.6 mmol) and 4Å molecular sieves in toluene (136 mL) were stirred at reflux overnight. The reaction mixture was filtered and the solvent was evaporated *in vacuo* to obtain **5a-5k** as yellow solids in quantitative yield.

2-Cyclopentyl-5-hydroxyisoindoline-1,3-dione (5a)²⁴—¹H NMR (DMSO-d₆): δ 10.8 (s,1H), 7.60 (d, J = 7.8 Hz, 1H), 7.05-7.01 (m, 2H), 4.50-4.40 (m, 1H), 1.97-1.52 (m, 8H). ¹³C NMR (DMSO-d₆): δ 167.7, 163.1, 134.2, 124.9, 121.0, 120.2, 109.5, 50.0, 29.2, 24.7.

5-Hydroxy-2-isopropylisoindoline-1,3-dione (5b)— 1 H NMR (300 MHz, DMSO-d6): δ 7.65 (d, J = 8.7 Hz, 1H), 7.22-7.09 (m, 2H), 4.38-4.29 (m, 1H), 1.37 (d, J = 6.9 Hz, 6H).

2-(Cyclopropylmethyl)-5-hydroxyisoindoline-1,3-dione (5c)— 1 H NMR (300 MHz, DMSO-d6): δ 7.70 (d, J = 8.1 Hz, 1H), 7.16-7.11 (m, 2H), 3.39 (d, J = 7.5 Hz, 2H),

1.11-1.06 (m, 1H), 0.47-0.39 (m, 2H), 0.32-0.28 (m, 2H). ^{13}C NMR (75 MHz, DMSO-d6): δ 168.5, 163.9, 135.1, 125.9, 122.4, 121.0, 110.5, 42.4, 11.0, 4.2.

- **2-Cyclobutyl-5-hydroxyisoindoline-1,3-dione (5d)**— 1 H NMR (300 MHz, DMSO-d6): δ 7.61 (d, J = 8.4 Hz, 1H), 7.06-7.04 (m, 2H), 4.59-4.50 (m, 1H), 2.79-2.68 (m, 2H), 2.16-2.10 (m, 2H), 1.78-1.66 (m, 2H).
- **5-Hydroxy-2-isobutylisoindoline-1,3-dione (5e)**—¹H NMR (300 MHz, DMSO-d6): δ 7.65 (d, J = 8.4 Hz, 1H), 7.10-7.06 (m, 2H), 3.32 (d, J = 7.2 Hz, 2H), 2.00-1.91 (m, 1H), 0.84 (d, J = 6.3 Hz, 6H). ¹³C NMR (75 MHz, DMSO-d6): δ 168.7, 134.9, 125.9, 121.1, 110.6, 45.3, 28.1, 20.6.
- **2-Cyclopropyl-5-hydroxyisoindoline-1,3-dione (5f)**— 1 H NMR (300 MHz, DMSO-d6): δ 7.70 (+d, J = 8.4 Hz, 1H), 7.20 7.10 (m, 2H), 2.70-2.60 (m, 1H), 0.91-0.82 (m, 4H).
- **5-Hydroxy-2-phenylisoindoline-1,3-dione (5g)**— 1 H NMR (DMSO-d₆): δ 11.03 (s, 1H), 7.78 (d, J = 8.2 Hz, 1H), 7.52-7.16 (m, 7H). 13 C NMR (DMSO-d₆): δ 166.7, 163.5, 134.3, 132.1, 128.8, 127.8, 127.3, 125.6, 120.8, 109.9.
- **2-Cyclohexyl-5-hydroxyisoindoline-1,3-dione (5h)**— 1 H NMR (300 MHz, DMSO-d6): δ 7.48 (d, J = 7.8 Hz, 1H), 6.86-6.79 (m, 2H), 3.93-3.87 (m, 1H), 1.82-1.09 (m, 10H).
- **5-Hydroxy-2-(4-(trifluoromethoxy)benzyl)isoindoline-1,3-dione (5i)**— 1 H NMR (300 MHz, DMSO-d6): δ 7.68-7.03 (m, 7H), 4.75 (s, 2H). 13 C NMR (75 MHz, DMSO-d6): δ 168.5, 166.4, 137.2, 135.2, 130.1, 130.0, 126.2, 121.9, 121.6, 111.1, 44.7.
- **5-Hydroxy-2-isopentylisoindoline-1,3-dione (5j)**— 1 H NMR (300 MHz, DMSO-d6): δ 7.62 (d, J = 7.8 Hz, 1H), 7.07-7.02 (m, 2H), 3.54-3.49 (m, 2H), 1.47-1.40 (m, 3H), 0.89 (d, J = 6.9 Hz, 6H).
- **5-Hydroxy-2-propylisoindoline-1,3-dione (5k)**—¹H NMR (DMSO-d₆): δ 7.74 (d, J = 8.2 Hz, 1H), 7.19-7.16 (m, 2H), 3.56-3.52 (m, 2H), 1.66-1.61 (m, 2H), 0.92-0.89 (m, 3H). ¹³C NMR (DMSO-d₆): δ 167.8, 163.2, 134.3, 125.1, 121.7, 120.2, 109.7, 21.4, 11.2.
- **General procedure for the synthesis of isoindolinones (6a-6k)**—To a solution of 1,3-dione **5a-5k** (18.0 mmol), in CH₃CO₂H (129 mL) was added Zn dust (11.6 g, 180 mmol) and the mixture was heated at reflux for 5 min. The volatiles were removed *in vacuo* and the residue was dissolved in water and extracted with ethyl acetate. The crude residue was purified using automated preparative HPLC to yield the desired compounds (**6a-6k**) as amorphous solids.
- **2-Cyclopentyl-5-hydroxyisoindolin-1-one (6a)**— $(1.7~{\rm g}, 44\%)$ ¹H NMR (DMSO-d₆): δ 10.10 (bs, 1H), 7.44 (d, J = 8.5 Hz, 1H), 6.89 (s, 1H), 6.83 (d, J = 7.9 Hz, 1H), 4.52 4.48 (m, 1H), 4.31 (s, 2H), 1.83–1.57 (m, 8H). ¹³C NMR (DMSO-d₆): δ 167.2, 160.4, 144.1, 123.9, 123.7, 115.4, 109.6, 51.8, 45.4, 29.6, 23.8. LCMS calcd. for $C_{13}H_{15}NO_2$ [M+H]⁺: 218; found: 218.
- **5-Hydroxy-2-isopropylisoindolin-1-one (6b)**—(23 mg, 22%) 1 H NMR (DMSO-d₆): δ 10.04(s, 1H), 7.43 (d, J = 8.2 Hz, 1H), 6.89-6.81 (m, 2H), 4.38-4.31 (m, 1H), 4.28 (s, 2H), 1.16 (d, J = 6.4 Hz, 6H). 13 C NMR (DMSO-d₆): δ 166.7, 160.4, 144.1, 123.9, 115.4, 109.6, 44.4, 41.9, 20.4. LRMS (ESI) calcd. for $C_{11}H_{13}NO_{2}$ [M+H]⁺: 192; found 191.95.

2-(Cyclopropylmethyl)-5-hydroxyisoindolin-1-one (6c)—(60 mg, 63%). LRMS (ESI) calcd. for $C_{12}H_{13}NO_2[M+H]^+$: 204; found 204.

- **2-Cyclobutyl-5-hydroxyisoindolin-1-one (6d)**—(108 mg, 45%). 1 H NMR (DMSO-d₆): δ 7.43, (d, J = 8.2 Hz, 1H), 6.89-6.80 (m, 2H), 4.71-4.62 (m, 1H), 4.42 (s, 2H), 2.31-2.06 (m, 4H), 1.72-1.62 (m, 2H). 13 C NMR (DMSO-d₆): δ 166.7, 160.6, 144.1, 124.2, 123.6, 115.5, 109.6, 45.8, 45.6, 28.1, 14.5. LRMS (ESI) calcd. for $C_{12}H_{13}NO_{2}[M+H]^{+}$: 204; found 203.95.
- **5-Hydroxy-2-isobutylisoindolin-1-one (6e)**—(86 mg, 34%). 1 H NMR (DMSO-d₆): δ 10.07 (s, 1H), 7.45 (d, J = 8.2 Hz, 1H), 6.88- 6.81 (m, 2H), 4.32 (s, 2H), 3.23 (d, J = 7.8 Hz, 2H), 2.00-1.90 (m, 1H), 0.83 (d, J = 6.8 Hz, 6H). 13 C NMR (DMSO-d₆): δ 167.6, 160.5, 144.1, 124.1, 123.5, 115.4, 109.5, 49.6, 49.1, 27.1, 20.0. LRMS (ESI) calcd. for $C_{12}H_{15}NO_2[M+H]^+$: 206; found 205.90.
- **2-Cyclopropyl-5-hydroxyisoindolin-1-one (6f)**—(53 mg, 23%). 1 H NMR (DMSO-d₆):8 7.36 (d, J = 8.2 Hz, 1H), 6.84-6.79 (m, 2H), 4.36 (s, 2H), 2.84-2.80 (m, 1H), 0.77-0.71 (m, 4H). 13 C NMR (DMSO-d₆): 8 168.9, 143.9, 123.8, 116.1, 109.7, 49.3, 24.9, 24.5, 5.1. LRMS (ESI) calcd. for $C_{11}H_{11}NO_{2}[M+H]^{+}$: 190; found 190.00.
- **5-Hydroxy-2-phenylisoindolin-1-one (6g)**—(14 mg, 19%). 1 H NMR (DMSO-d₆): δ 7.83-7.81(m, 2H), 7.55-7.53 (m, 1H), 7.38-7.34 (m, 2H), 7.10-7.06 (m, 1H), 6.92-6.91 (m, 1H), 6.87-6.84 (m, 1H). LRMS (ESI) calcd. for $C_{14}H_{11}NO_{2}[M+H]^{+}$: 226; found 226.
- **2-Cyclohexyl-5-hydroxyisoindolin-1-one (6h)**—(120 mg, 64%). 1 H NMR (300 MHz, DMSOd6): δ 10.08 (s, 1H), 7.47 (d, J = 8.7 Hz, 1H), 6.92-6.84 (m, 2H), 4.33 (s, 2H), 3.99-3.93 (m, 1H), 1.82-1.31 (m, 10H). 13 C NMR (75 MHz, DMSO-d6): δ 167.4, 161.1, 144.9, 124.7, 116.1, 110.3, 50.7, 46.1, 31.4, 25.9, 25.8. LRMS (ESI) calcd. for $C_{14}H_{17}NO_{2}[M+H]^{+}$: 232; found 232.0.
- **5-Hydroxy-2-(4-(trifluoromethoxy)benzyl)isoindolin-1-one (6i)—**(19 mg, 20%). LRMS (ESI) calcd. for $C_{16}H_{12}$ $F_3NO_3[M+H]^+$: 324; found 323.95.
- **5-Hydroxy-2-isopentylisoindolin-1-one (6j)**—(213 mg, 32%). 1 H NMR (300 MHz, DMSO-d6): δ 7.46 (d, J = 8.1 Hz, 1H), 6.91 (s, 1H), 6.84 (d, J = 8.4 Hz, 1H), 4.34 (s, 2H), 1.49-1.42 (m, 3H), 0.91 (d, J = 5.4 Hz, 6H). LRMS (ESI) calcd. for $C_{13}H_{17}NO_{2}[M+H]^{+}$: 220; found 219.85 (M+1)⁺.
- **5-Hydroxy-2-propylisoindolin-1-one (6k)**—(12 mg, 22%). LRMS (ESI) calcd. for $C_{11}H_{13}NO_2[M+H]^+$: 192; found 191.95.
- **Methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate**— $Pd(OAc)_2$ (0.393 g, 1.75 mmol) was added to a solution of 2-chloro-5-iodobenzoic acid (9.9 g, 35 mmol), mtolylboronic acid (5.71 g, 42 mmol) and Na_2CO_3 (11.45 g, 105 mmol) in water (45 mL) under a nitrogen atmosphere. The resulting mixture was stirred at 50 °C for 2 h. The reaction mixture was cooled to room temperature and filtered. The filtrate was acidified with 1M HCl. The precipitated 4-chloro-3'-methylbiphenyl-3-carboxylic acid was filtered, washed with water and dried to provide a tan solid that was used without further purification in the next step (quantitative yield). 1 H NMR (DMSO-d₆, 300 MHz): δ 8.03 (s, 1H), 7.83 (d, J = 8.1 Hz, 1H), 7.63 (d, J = 7.8 Hz, 1H), 7.55-7.50 (m, 2H), 7.40 (t, J = 7.8 Hz, 1H), 7.25 (d, J = 6.6 Hz, 1H), 2.41 (s, 3H). 13 C NMR (DMSO-d₆, 75 MHz): δ 167.4, 139.9, 139.1, 138.6, 132.7, 131.9, 131.2, 129.7, 129.5, 129.3, 128.0, 124.5, 21.8.

To a solution of 4-chloro-3'-methylbiphenyl-3-carboxylic acid (7 g, 28.5 mmol) in acetone was added K₂CO₃ (27.5 g, 199 mmol) and MeI (12.5 mL, 28 mmol). The resulting solution was heated at 50 °C for 1 h. After cooling the excess solvent was removed *in vacuo*. The crude material was dissolved in CH₂Cl₂ and washed with water. The organic layer was dried over Na₂SO₄, and the solvent was evaporated to afford methyl 4-chloro-3'-methylbiphenyl-3-carboxylate as a reddish yellow viscous liquid (7.3 g, 98%) which was used in the next step without further purification. ¹H NMR (DMSO-d₆): δ 8.04 (s, 1H), 7.85 (d, J = 7.8 Hz, 1H), 7.64 (d, J = 8.1 Hz, 1H), 7.52-7.47 (m, 2H), 7.38 (t, J = 7.8 Hz, 1H), 7.23 (d, J = 7.5 Hz, 1H), 3.90 (s, 3H), 2.39 (s, 3H). ¹³C NMR (DMSO-d₆, 75 MHz): δ 166.2, 139.9, 139.1, 138.4, 131.9, 131.7, 131.4, 131.3, 129.7, 129.6, 129.5, 128.0, 124.5, 53.3, 21.7.

A catalytic amount of AIBN was added to a solution of methyl 4-chloro-3'-methylbiphenyl-3- carboxylate (7.3 g, 28 mmol) and NBS (5.5 g, 30.8 mmol) in CCl₄. The reaction mixture was heated under reflux for 12 h. After cooling, the precipitated succinimide was removed by filtration. The filtrate was concentrated *in vacuo* to afford the crude product. Silica gel column chromatography using 4:1 hexanes/ethyl acetate afforded methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate (8.8 g, 91%) as a colorless solid. 1 H NMR (CDCl₃): δ 8.07(s, 1H), 7.68-7.45 (m, 6H), 4.58 (s, 2H), 4.00 (s, 3H). 13 C NMR (CDCl₃): δ 166.1, 139.5, 139.2, 138.7, 133.0, 131.6, 131.0, 130.4, 129.9, 129.6, 128.7, 127.7, 52.6, 33.2.

4-Chloro-3'-((2-cyclopentyl-1-oxoisoindolin-5-yloxy)-methyl)biphenyl-3-carboxylic acid (8a)—To a solution of 2-cyclopentyl-5-hydroxyisoindolin-1-one (50 mg, 0.23 mmol) in acetone (15 mL), K₂CO₃ (181 mg, 1.31mmol) was added methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate (92 mg, 0.27 mmol) and the mixture was heated at reflux for 1 h. The reaction mixture was filtered and the solvent was evaporated *in vacuo* to obtain methyl 4-chloro-3'-((2- cyclopentyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylate **7a** as a yellow oil in quantitative yield.

Crude methyl 4-chloro-3'-((2-cyclopentyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylate **7a** (109 mg, 0.23 mmol) was dissolved in tetrahydrofuran (17 mL), 2M LiOH (0.57 mL, 1.15 mmol) was added and the mixture was heated at reflux for 3 h. The solvent was evaporated and the residue was dissolved in water and neutralized using 2M HCl. The aqueous layer was extracted with ethyl acetate and the organic layer was dried over anhydrous Na₂SO₄. The solvent was evaporated *in vacuo* to obtain the crude acid as a yellow solid. The crude residue was purified using automated preparative HPLC to yield the desired compound **8a** (48 mg, 46%) as an amorphous white solid. ¹H NMR (CDCl₃): δ 8.18 (s, 1H), 7.78 (d, J = 8.5 Hz, 1H), 7.65-7.41 (m, 6H), 7.05 (d, J = 8.5 Hz, 1H) 6.99 (s, 1H), 5.16 (s, 2H), 4.74 (m, 1H), 4.30 (s, 2H), 1.97 (m, 2H), 1.76 – 1.59 (m, 6H). ¹³C NMR (CDCl₃): δ 168.7, 161.5, 143.3, 139.3, 137.2, 133.5, 131.7, 131.4, 130.7, 129.7, 129.4, 127.0, 126.8, 126.0, 125.9, 125.1, 115.4, 108.7, 70.1, 52.7, 46.1, 30.1, 24.1. HRMS: calcd. for $C_{27}H_{24}ClNO_4$ [M+H]⁺: 462.1467; found: 462.1468.

Following the above mentioned procedure **8b-8k** were synthesized using the appropriate starting materials.

4-Chloro-3'-((2-isopropyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8b)—Amorphous white solid (14 mg, 32%). 1 H NMR (300 MHz, DMSO-d₆): δ 8.06 (s, 1H), 7.85-7.82 (m, 2H), 7.68-7.51 (m, 5H), 7.24 (s, 1H), 7.13 (d, J = 8.4 Hz, 1H), 5.26 (s, 2H), 4.45-4.40 (m, 1H), 4.34 (s, 2H), 1.19 (d, J = 6.9 Hz, 6H). 13 C NMR (75 MHz, DMSO-d₆): δ 166.8, 166.6, 161.1, 144.2, 138.8, 138.3, 137.7, 132.1, 131.4, 131.0, 130.6, 129.5, 128.8, 127.6, 126.4, 126.2, 125.7, 124.1, 115.4, 109.2, 69.5, 44.8, 42.3,

20.5. LRMS (ESI): 435.95 (M+1)^+ . HRMS: calcd. for $C_{25}H_{22}CINO_4$ 436.131; found 436.1309.

4-Chloro-3'-((2-(cyclopropylmethyl)-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3- carboxylic acid (8c)—Amorphous white solid (49 mg, 39%). 1 H NMR (300 MHz, DMSO-d₆): δ 8.07 (s, 1H), 7.88 – 7.84 (m, 2H), 7.71 – 7.52 (m, 5H), 7.28 (s, 1H), 7.14 (d, J = 8.4 Hz, 1H), 5.28 (s, 2H), 4.50 (s, 2H), 3.33 (d, J = 7.5 Hz, 2H), 1.06 – 0.98 (m, 1H), 0.51 – 0.48 (m, 2H), 0.31 – 0.28 (m, 2H). 13 C NMR (75 MHz, DMSO-d₆): δ 167.5, 167.3, 161.8, 144.8, 139.4, 138.9, 138.4, 132.8, 131.9, 131.6, 131.2, 130.1, 129.4, 128.3, 127.1, 126.9, 126.0, 124.7, 116.0, 109.8, 70.2, 50.0, 46.8, 10.6, 4.1. LRMS (ESI): 448.00 (M+1)⁺. HRMS: calcd. for C₂₆H₂₂ClNO₄ 448.131; found 448.1313.

- **4-Chloro-3'-((2-cyclobutyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8d)**—Amorphous white solid (183 mg, 41%). 1 H NMR (300 MHz, DMSO-d₆): δ 8.05 (s, 1H), 7.85-7.83 (m, 2H), 7.68-7.51 (m, 5H), 7.25 (s, 1H), 7.12 (d, J = 8.7 Hz, 1H), 5.27 (s, 2H), 4.78-4.62 (m, 1H), 4.49 (s, 2H), 2.38- 2.22 (m, 2H), 2.15-2.09 (m, 2H), 1.78-1.66 (m, 2H). 13 C NMR (75 MHz, DMSO-d₆): δ 166.6, 166.4, 161.1, 144.0, 138.7, 138.2, 137.6, 132.0, 131.2, 130.9, 130.5, 129.4, 128.7, 127.5, 126.3, 126.2, 125.4, 124.1, 115.4, 109.1, 69.4, 46.0, 45.7, 28.1, 14.5. LRMS (ESI): 448.00 (M+1)⁺. HRMS: calcd. for $C_{26}H_{22}$ ClNO₄ 448.131; found 448.131.
- **4-Chloro-3'-((2-isobutyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8e)**—Amorphous white solid (75 mg, 34%). 1 H NMR (CDCl₃): δ 8.17 (s, 1H), 7.78 (d, J = 8.5 Hz, 1H), 7.66 7.62 (m, 2H), 7.55 7.41 (m, 4H), 7.07 7.05 (m, 1H), 7.00 (s, 1H), 5.17 (s, 2H), 4.33 (s, 2H), 3.39 (d, J = 7.3 Hz, 2H), 2.07 1.96 (m, 1H), 0.92 (d, J = 6.7 Hz, 6H). 13 C NMR (CDCl₃): δ 169.1, 168.2, 161.6, 143.4, 139.4, 137.2, 133.6, 131.7, 131.5, 130.7, 129.4, 127.1, 126.8, 125.9, 125.6, 125.3, 115.4, 108.7, 70.1, 50.6, 50.2, 27.8, 20.1. LRMS (ESI): 450.00 (M+1)⁺. HRMS: calcd. for C₂₆H₂₄ClNO₄ 450.1467; found 450.1472.
- **4-Chloro-3'-((2-cyclopropyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8f)**—Amorphous white solid (78 mg, 23%). 1 H NMR (300 MHz, DMSO-d₆): δ 8.05 (s, 1H), 7.86-7.81 (m, 2H), 7.67-7.51 (m, 5H), 7.21 (s, 1H), 7.12 (d, J=7.8 Hz, 1H), 5.26 (s, 2H), 4.32 (s, 2H), 2.92-2.86 (m, 1H), 0.80-0.76 (m, 4H). 13 C NMR (75 MHz, DMSO-d₆): δ 168.3, 166.8, 161.2, 144.1, 138.8, 138.3, 137.7, 132.1, 131.4, 130.9, 130.6, 129.5, 128.7, 127.6, 126.4, 126.2, 125.7, 124.0, 115.5, 108.9, 69.5, 49.5, 25.1, 5.11. LRMS (ESI): 434.05 (M+1)⁺. HRMS: calcd. for C₂₅H₂₀ClNO₄ 434.1154; found 434.1150.
- **4-Chloro-3'-((1-oxo-2-phenylisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8g)**—Amorphous white solid (28 mg, 36%). 1 H NMR (CDCl₃): δ 8.20 (s, 1H), 7.86 7.36 (m, 11H), 7.15 7.05 (m, 3H), 5.19 (s, 2H), 4.77 (s, 2H). 13 C NMR (CDCl₃): δ 169.2, 167.4, 162.2, 142.4, 139.5, 139.4, 139.3, 137.1, 133.7, 131.9, 131.7, 130.8, 129.5, 129.1, 127.1, 126.9, 126.1, 126.0, 125.8, 124.3, 119.4, 115.9, 108.4, 70.2, 50.6. LRMS (ESI): 470. (M+1)⁺. HRMS: calcd. for $C_{28}H_{20}CINO_4$: 470.1154; found 470.1157.
- **4-Chloro-3'-((2-cyclohexyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8h)**—Amorphous white solid (87 mg, 36%). 1 H NMR (300 MHz, DMSO-d₆): δ 8.05 (s, 1H), 7.83 (m, 2H), 7.68 7.45 (m, 5H), 7.24 (s, 1H), 7.12 (d, J = 8.4 Hz, 1H), 5.27 (s, 2H), 4.36 (s, 2H), 3.99 3.92 (m, 1H), 1.79 1.14 (m, 10H). 13 C NMR (75 MHz, DMSO-d₆): δ 167.3, 167.0, 161.7, 144.8, 139.4, 138.9, 138.4, 132.8, 131.9, 131.6, 131.2, 130.1, 129.4, 128.2, 127.0, 126.8, 126.3, 124.6, 116.0, 109.8, 70.1, 50.8, 46.3, 31.3, 25.9, 25.7. LRMS (ESI): 476.05 (M+1)⁺. HRMScalcd. for $C_{28}H_{26}CINO_4$: 476.1623; found 476.1621.

4-Chloro-3'-((1-oxo-2-(4-(trifluoromethoxy)benzyl)isoindolin-5-yloxy)methyl)biphenyl-3- carboxylic acid (8i)—Amorphous white solid (12 mg, 30%). 1 H NMR (CDCl₃): δ 8.13 (s, 1H), 7.80 (d, J = 8.5 Hz, 1H), 7.62 – 7.28 (m, 8H),7.14 (d, J = 7.9 Hz, 2H), 7.07 – 7.05 (m, 1H), 6.93 (s, 1H), 5.14 (s, 2H), 4.74 (s, 2H), 4.20 (s, 2H). 13 C NMR (CDCl₃): δ 168.5, 161.8, 148.7, 143.4, 139.9, 139.4, 137.1, 135.8, 131.7, 131.5, 130.7, 129.5, 127.1, 126.8, 125.9, 125.4, 125.3, 121.3, 115.5, 108.9, 70.2, 49.3, 45.7. LRMS (ESI): 567.95 (M+1) $^+$. HRMS: calcd. for C₃₀H₂₁ClF₃NO₅ 568.1133; found 568.1136.

4-Chloro-3'-((2-isopentyl-1-oxoisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8j)—Amorphous white solid (82 mg, 41%). 1 H NMR (CDCl₃): δ 8.18 (s, 1H), 7.78 (d, J = 8.5 Hz, 1H), 7.66 – 7.62 (m, 2H), 7.55 – 7.42 (m, 4H), 7.06 (dd, J = 8.5, 1.8 Hz, 1H), 6.99 (s, 1H), 5.17 (s, 2H), 4.31 (s, 2H), 3.62 – 3.58 (m, 2H), 1.63- 1.49 (m, 3H), 0.93 (d, J = 6.1 Hz, 6H). 13 C NMR (CDCl₃): δ 168.6, 161.5, 143.3, 139.4, 137.2, 133.6, 131.8, 131.5, 130.7, 129.4, 127.1, 126.8, 125.9, 125.2, 115.3, 108.7, 70.1, 49.8, 40.8, 37.2, 25.8, 22.5. LRMS (ESI): 463.95 (M+1)⁺. HRMS: calcd. for C₂₇H₂₆ClNO₄ 464.1623; found 464.1626.

4-Chloro-3'-((1-oxo-2-propylisoindolin-5-yloxy)methyl)biphenyl-3-carboxylic acid (8k)—Amorphous white solid (30 mg, 34%). 1 H NMR (CDCl₃): δ 8.18 (s, 1H), 7.78 (d, J = 8.5 Hz, 1H), 7.65 – 7.62 (m, 2H), 7.53 – 7.43 (m, 4H), 7.06 (dd, J = 8.5 Hz, J = 2.4 Hz, 1H), 6.99 (s, 1H), 5.17 (s, 2H), 4.31 (s, 2H), 3.54 (t, J = 7.3 Hz, 2H), 1.70 – 1.61 (m, 2H), 0.93 (t, J = 7.3 Hz, 3H). 13 C NMR (CDCl₃): δ 168.7, 168.4, 161.5, 143.3, 139.4, 137.2, 133.5, 131.7, 131.4, 130.7, 129.5, 129.4, 127.0, 126.8, 125.9, 125.9, 125.2, 115.3, 108.7, 70.1, 49.9, 44.1, 21.7, 11.3. LRMS (ESI): 436.00 (M+1)⁺. HRMS: calcd. for C₂₅H₂₂ClNO₄ 436.131; found 436.1315.

N-Cyclopentyl-2-mercapto-4-methoxybenzamide (10)—The title compound was prepared using the literature procedure as follows. AlMe₃ (10 mmol, 2.0 M in toluene) was added dropwise to a cooled (0 °C) suspension of cyclopentylamine (0.85 g, 10 mmol) in CH₂Cl₂ (30 mL). When the addition was complete, the reaction mixture was allowed to warm to room temperature and stirring was continued for 30 min until gas evolution had ceased. A solution of methyl 2-mercapto-4-methoxybenzoate 9 (0.990 g, 5 mmol) was then added. After stirring at 60 °C for 12 h, the mixture was cooled, and carefully quenched with 5% aq HCl (20 mL). The organic layer was separated and the aqueous layer was extracted with CH₂Cl₂ (3 × 15 mL). The combined organic extracts were washed with a saturated aqueous solution of NaHCO₃ (15 mL) and brine. The organic phase was dried using MgSO₄ and evaporated, filtered and the solvent was evaporated in vacuo. The resulting residue was purified by crystallization from Et₂O to afford amide **10** as a tan solid (1.18 g, 94%). ¹H NMR (300 MHz, DMSO- d_6): δ 8.26 (s, 1H), 7.51 (d, J = 8.7 Hz, 1H), 7.01 (s, 1H), 6.72 (d, J = 8.1 Hz, 1H), 5.62 (s, 1H), 4.16-4.14 (m, 1H), 3.78 (s, 3H), 1.99-1.80 (m, 2H), 1.79-1.68 (m, 2H), 1.66-1.53 (m, 4H). 13 C NMR: δ 164.6, 162.7, 142.7, 127.1, 118.4, 115.2, 105.0, 56.3, 54.8, 32.0, 23.9. LC-MS (ESI) calcd. for C₁₃H₁₇NO₂S [M+H]⁺: 252.09; found: 251.95.

2-Cyclopentyl-6-methoxybenzo[d]isothiazol-3(2H)-one (11)—A solution of PIFA (645 mg, 1.5 mmol) in CH₂Cl₂ (20 mL) was added at 0 °C to a solution of *N*-cyclopentyl-2-mercapto-4-methoxybenzamide 10 (251mg, 1 mmol) and TFA (0.23 mL, 3 mmol) in CH₂Cl₂ (5 mL). The resulting mixture was gradually warmed to room temperature. After 1 h, the solvent was evaporated under reduced pressure and the resulting residue was purified by column chromatography using CH₂Cl₂ to afford the 2-cyclopentyl-6-methoxybenzo[d]isothiazol-3(2H)-one **11** as a white amorphous solid (84%, 0.210 g). 1 H NMR (300 MHz, CDCl₃): δ 7.87 (d, J = 8.5 Hz, 1H), 6.93 (s, 1H), 6.92 (d, J = 8.2 Hz, 1H),

5.50-5.01 (m, 1H), 3.86 (s, 1H), 2.13-2.12 (m, 2H), 1.83-1.67 (m, 6H). 13 C NMR (75 MHz, CDCl₃): δ 165.3, 162.7, 142.2, 127.6, 118.8, 114.6, 103.1, 55.8, 55.0, 32.2, 24.4. LC-MS (ESI) calcd. for $C_{13}H_{15}NO_2S$ [M+H]⁺: 250.08; found: 249.95.

2-Cyclopentyl-6-hydroxybenzo[d]isothiazol-3(2H)-one (12)—A solution of 2-cyclopentyl-6-methoxybenzo[d]isothiazol-3(2H)-one 11 (0.250 g, 1 mmol) in anhydrous C_6H_6 (20 mL) was cooled 0 °C before BBr₃ (0.54 g, 2 mmol) was added dropwise. The mixture was gradually warmed to room temperature and then heated at 80 °C for 30 min. After that, the reaction mixture was cooled to room temperature and carefully quenched with H_2O (20 mL). The resulting mixture was then heated at reflux for 1 h and filtered; the precipitate was washed with H_2O and dried *in vacuo* to give the crude product. This material was purified by flash chromatography (silica gel, CHCl₃:MeOH, 9:1) to afford 2-cyclopentyl-6-hydroxybenzo[d]isothiazol-3(2H)-one **12** (0.200 g, 85%) as a colorless amorphous solid. 1H NMR (300 MHz, DMSO-d₆): δ 10.36 (s, 1H), 7.60 (d, J = 8.5 Hz, 1H), 7.20 (s, 1H), 6.80 (d, J = 8.6 Hz, 1H), 4.84-4.81 (m, 1H), 2.12-1.98 (m, 2H), 1.75-1.60 (m, 6H). ^{13}C NMR (75 MHz, DMSO-d₆): δ 164.8, 161.3, 142.5, 127.4, 117.2, 115.6, 106.9, 54.6, 32.1, 24.4. LC-MS (ESI) calcd. for $C_{12}H_{13}NO_{2}S$ [M+H]+: 236.0667; found: 235.95.

3-Chloro-3'-((2-cyclopentyl-3-oxo-2,3-dihydrobenzo[d]isothiazol-6-yloxy)methyl)biphenyl-4-carboxylic acid (14)—Potassium carbonate (0.052 g, 0.38 mmol) was added to a solution of 2-cyclopentyl-6- hydroxybenzo[d]isothiazol-3(2H)-one 12 (0.075g, 0.32 mmol) and methyl 3'-(bromomethyl)-3-chlorobiphenyl-4-carboxylate (0.129 g, 0.38 mmol) in CH₃CN (5 mL). After stirring for 12 h at 80 °C, the organic phase was evaporated under reduced pressure and the crude material was partioned between water and CH₂Cl₂. The aqueous layer was extracted with CH₂Cl₂ (3 × 15 mL). The organic layer was dried using Na₂SO₄ and evaporated to give methyl 3-chloro-3'-((2-cyclopentyl-3-oxo-2,3-dihydrobenzo[d]isothiazol-6-yloxy)methyl)biphenyl-4-carboxylate **13** (0.158 g, quantitative yield). The crude product was used in the next step without further purification. LC-MS (ESI) calcd. for C₂₇H₂₄NO₂S [M+H]⁺: 494.11; found: 494.00.

LiI (0.042 g, 0.32 mmol) was added to a solution of the crude product (0.158 g, 0.32 mmol) in pyridine (5 mL). The reaction mixture was heated under reflux for 12 h and then cooled to room temperature. The excess solvent was removed *in vacuo*. The product was purified by preparative HPLC using isopropanol:water as the solvent system to afford the title compound **14** as a colorless solid (70 mg) in 46% yield. mp: 189-191 °C. 1 H NMR (DMSO-d₆): δ 8.01 (s, 1H), 7.82 (d, J = 8.5 Hz, 1H), 7.80 (d, J = 8.6 Hz, 1H), 7.71-7.76 (m, 4H), 7.49-7.47 (m, 2H), 7.05 (d, J = 8.5 Hz, 1H.), 5.23 (s, 2H), 4.83-4.81 (m, 1H), 2.02-1.99 (m, 2H), 1.62-1.60 (m, 6H). 13 C NMR (DMSO-d₆): δ 167.2, 164.0, 140.2, 138.7, 137.8, 131.7, 130.6, 131.0, 129.9, 129.2, 128.2, 127.2, 127.0, 126.9, 119.0, 115.73, 106.6, 70.2, 54.8, 32.2, 24.4. LC-MS (ESI) calcd. for $C_{26}H_{22}$ CINO₄S [M+H]+: 480.09; found: 479.95. HRMS (ESI) calcd. for $C_{26}H_{22}$ CINO₄S [M+H]+: 480.1031; found: 480.1031.

N-Cyclopentyl-2-dihydroxy-4-methoxybenzamide (18)—To a solution of 4-methoxysalicylic acid (0.84 g, 5 mmol) in CH_2Cl_2 was added pyridine (2.05 mL, 25 mmol) and acetyl chloride (0.89 mL, 12.5 mmol) at 0 °C. The reaction mixture was gradually warmed to room temperature over 30 min and then poured into water. The resulting biphasic mixture was stirred for 1h and then acidified with 1 M HCl and extracted with EtOAc. The crude acetoxy derivative was then converted to the corresponding acid choride by heating with oxalyl chloride (1.3 mL, 15 mmol) in CH_2Cl_2 for 1h at reflux. After removal of the volatile materials, the acid chloride (1.05 g, 4.38 mmol) in CH_2Cl_2 was added to a stirred mixture of sodium carbonate (1M, 17.5 mL) and C_5H_9NH . OH (1.32 g, 13.2 mmol) in THF at 0 °C. The resulting mixture was stirred at room temperature for 1 h, acidified with 0.5 M HCl and then extracted with EtOAc. The crude product was purified by column

chromatography on silica gel using hexanes:ethyl acetate afforded hydroxamic acid **18** (1 g, 79%) as a colorless solid. 1H NMR (DMSO- d_6): δ 7.65 (d, J=9.1 Hz, 1H), 6.37-6.36 (overlapping doublet and singlet, 2 H), 4.10-3.98 (m, 1H), 3.70 (s, 1H), 1.69-1.63 (m, 2H), 1.55-1.52 (m, 6H). ^{13}C NMR (DMSO- d_6): δ 167.5, 162.5, 160.0, 131.4, 113.3, 105.5, 101.6, 56.6, 55.7, 29.5, 25.5. LC-MS (ESI) calcd. for $C_{13}H_{17}NO_4$ [M+H]+: 251.12; found: 252.00.

2-Cyclopentyl-6-methoxybenzo[d]isoxazol-3(2H)-one (19)—

Diethylazodicarboxylate was added dropwise to a cooled solution of hydroxamic acid 18 (0.71 g, 2.8 mmol) and triphenyl phosphine (0.89 g, 3.4 mmol) in THF. The reaction mixture was warmed to room temperature and quenched with 1:1 MeOH:AcOH (0.2 mL). Concentration of the solvent followed by chromatographic separation on silica gel afforded **19** as a pale yellow solid (0.3 g, 46%). ¹H NMR (CDCl₃): δ 7.67 (d, J = 8.7 Hz, 1H), 6.78 (d, J = 8.6 Hz, 1H), 6.32 (s, 1H), 4.35-4.33 (m, 1H), 3.84 (s, 3H), 1.89-1.68 (m, 8H). ¹³C NMR (CDCl₃): δ 164.5, 162.8, 162.3, 124.9, 112.9, 109.8, 103.1, 55.8, 55.0, 32.2, 24.4. LC-MS (ESI) calcd. for C₁₃H₁₅NO₃ [M+H]⁺: 233.10; found: 232.90.

4-Chloro-3'-((2-cyclopentyl-3-oxo-2,3-dihydrobenzo[d]isoxazol-6yloxy)methyl)biphenyl- 3-carboxylic acid (21)—BBr₃ (0.54 g, 2 mmol) was added dropwise to a solution of 19 (0.232 g, 1 mmol) in anhydrous benzene (20 mL) at 0 °C. The mixture was gradually warmed to room temperature and then heated at 80 °C for 30 min. After that the reaction mixture was cooled to room temperature and carefully quenched with H₂O (20 mL). The resulting mixture was then heated at reflux for 1 h and then cooled. Extraction with EtOAc and removal of the solvent in vacuo afforded the crude demethylated product (0.170 g, 78%) which was used in the next step without further purification. A solution of 2-cyclopentyl-6-hydroxyybenzo[d]isoxazol-3(2H)-one (0.109 g, 0.5 mmol) was treated with methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate under the same conditions as described for compound 14 to give the corresponding ester in 76% yield (0.180 g). LC-MS (ESI) calcd. for $C_{27}H_{24}ClNO_5 [M+H]^+$: 477.13. Found: 478.05. The crude product obtained above was treated with LiI in pyridine to give the final compound 21 (0.063 g, after HPLC purification) as a colorless solid in 36% yield. mp: 185–187 °C. ¹H NMR (DMSO- d_6): δ 8.21 (s, 1H), 7.70-7.56 (m, 7H), 6.92 (d, J = 6.1 Hz, 1H), 5.18 (s, 2H), 4.98-4.95 (m, 1H), 2.06-2.02 (m, 2H), 1.92-1.88 (m, 4H), 1.64-1.63 (m, 2H). ¹³C NMR (DMSO- d_6): δ 164.0, 163.4, 162.7, 139.4, 137.6, 131.7, 130.9, 129.9, 129.2, 127.1, 126.9, 125.2, 114.3, 109.4, 70.5, 95.5, 56.6, 30.2, 25.4. LC-MS (ESI) calcd. for C₂₆H₂₂ClNO₅ [M +H]⁺: 464.13; found: 463.95. HRMS (ESI) calcd. for $C_{26}H_{22}CINO_5$ [M+H]⁺: 464.1259; found: 464.1261.

N-(3-Methoxyphenethyl)cyclopentanamine (23)—3-Methoxyphenethylamine (4.5 mL, 30 mmol), and iodocyclopentane (1.15 mL, 10 mmol) in CH₃CN were irradiated for 10 min in a microwave reactor at 110 °C. After cooling to room temperature the mixture was diluted with CH₂Cl₂ and washed with NaHCO₃ solution. The organic layer was dried over anhydrous Na₂SO₄ and evaporated *in vacuo*. The crude product was separated by silica gel chromatography using hexane as eluent and the pure product *N*-(3-methoxyphenethyl)cyclopentanamine (1.45 g, 66%) was obtained as a colorless liquid which solidified upon standing. ¹H NMR (CDCl₃): δ 8.48 (s, 1H), 7.20 (t, J = 8.0 Hz, 1H), 6.79-6.73 (overlapping doublets and singlet, 3H), 3.74 (s, 3H), 3.44-3.39 (m, 1H), 3.38-3.11 (m, 4H), 1.84-1.83 (m, 2H), 1.82-1.81 (m, 4H), 1.66-1.56 (m, 2H). ¹³C NMR (CDCl₃): δ 159.63, 140.93, 129.38, 121.07, 114.49, 111.45, 59.26, 55.18, 48.08, 32.46, 29.65, 23.84. LC-MS (ESI) Calcd for: C₁₄H₂₁NO [M+H]⁺: 220.16. Found: 220.00.

Methyl cyclopentyl(3-methoxyphenethyl)carbamate (24)—To a solution of *N*-(3-methoxyphenethyl)cyclopentanamine **23** (1.31 g, 6 mmol) in diethyl ether was added triethylamine (1.3 mL, 9 mmol) and methyl chloroformate (0.7 mL, 9 mmol) at 0 °C. The reaction mixture was gradually warmed to room temperature and stirred for 30 min. Filtration and concentration of the solvent under reduced pressure preceded silica gel column chromatography using hexanes/ethyl acetate to afford methyl cyclopentyl(3-methoxyphenethyl)carbamate as a pale yellow liquid (1.41g, 84%). ¹H NMR (CDCl₃): δ 7.19 (t, J = 7.6 Hz, 1H), 6.78-6.73 (overlapping doublets and singlet, 3H), 4.27-4.24 (m, 1H), 3.78 (s, 3H), 3.70 (s, 3H), 3.28 (t, J = 7.9 Hz, 2H), 2.80 (t, J = 7.9 Hz, 2H), 1.68-1.67 (m, 2H), 1.53-1.50 (m, 6H). ¹³C NMR (CDCl₃): δ 159.63, 156.74, 140.93, 129.38, 121.07, 114.49, 111.45, 57.94, 55.09, 52.30, 45.75, 36.50, 29.37, 23.50. LC-MS (ESI) Calcd for C₁₆H₂₃NO₃ [M+H]⁺: 278.17. Found: 278.00.

2-Cyclopentyl-6-methoxy-3,4-dihydroisoquinolin-1(2H)-one (25)—To a solution of methyl cyclopentyl(3-methoxyphenethyl)carbamate **24** (1.39 g, 5 mmol) in POCl₃ (20 mL) was added P₂O₅ (1.4 g, 10 mmol). The reaction mixture was heated at reflux for 2 h. Excess POCl₃ was removed under reduced pressure and quenched with iced water. The mixture was neutralized with Na₂CO₃, extracted with EtOAc, and the organic phase was dried over Na₂SO₄. Concentration of the solvent followed by silica gel column chromatography using hexanes/ethyl acetate afforded 2-cyclopentyl-6-methoxy-3,4-dihydroisoquinolin-1(2H)-one (1.30 g, 94%). ¹H NMR (CDCl₃): δ 8.00 (d, J = 8.6 Hz, 1H), 6.81 (d, J = 8.6 Hz, 1H), 6.63 (s, 1H), 5.18-5.14 (m, 1H), 3.85 (t, J = 6.7 Hz, 1H), 2.88 (t, J = 6.7 Hz, 1H), 1.88-1.70 (m, 2H), 1.64-1.63 (m, 4H), 1.62-1.61 (m, 2H). ¹³C NMR (CDCl₃): δ 164.44, 162.08, 139.97, 130.56, 123.05, 112.43, 111.79, 55.43, 53.70, 40.32, 28.96, 28.81, 24.57. LC-MS (ESI) Calcd for C₁₅H₁₉NO₂ [M+H]⁺: 246.14. Found: 246.00.

2-Cyclopentyl-6-hydroxy-3,4-dihydroisoquinolin-1(2H)-one (26)—BBr₃ (3.7 g, 14.7 mmol) was added dropwise to a solution of 2-cyclopentyl-7-methoxy-3,4-dihydroisoquinolin-1(2H)-one **25** (1.2 g, 4.9 mmol) in benzene at 0 °C. The mixture was gradually warmed to room temperature and then heated at 80 °C for 30 min. The reaction mixture was then cooled to room temperature and carefully quenched with H₂O. The resulting mixture was then heated at reflux for 1 h and then cooled. Extraction with EtOAc and removal of the solvent *in vacuo*, followed by chromatographic separation using MeOH-CHCl₃ provided 2-cyclopentyl-6-hydroxy-3,4-dihydroisoquinolin-1(2H)-one **26** as a colorless solid (0.735 g, 65%). ¹H NMR (DMSO-D₆): δ 9.96 (s, 1H), 7.65 (d, J = 8.6 Hz, 1H), 6.63 (d, J = 8.5 Hz, 1H), 6.56 (s, 1H), 4.95-4.93 (m, 1H), 3.32 (t, 2H, J = 6.1 Hz), 2.73 (t, 2 H, J = 6.7 Hz), 1.73-1.64 (m, 4H), 1.53-1.45 (m, 4H). ¹³C NMR (DMSO-d₆): 163.82, 160.69, 141.02, 130.29, 121.51, 114.28, 113.62, 53.44, 42.20, 28.79, 28.57, 24.52. LC-MS (ESI) Calcd for C₁₄H₁₇NO₂ [M+H]⁺: 232.13. Found: 232.00.

4-Chloro-3'-((2-cyclopentyl-1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yloxy)methyl)biphenyl-3- carboxylic acid (28)—2-Cyclopentyl-6-hydroxy-3,4-dihydroisoquinolin-1(2H)-one **26** (0.106 g, 0.5 mmol) and methyl 3'-(bromomethyl)-4-chlorobiphenyl-3-carboxylate (0.208 g, 0.6 mmol) were processed according to the general procedure to provide methyl 4-chloro-3'-((2-cyclopentyl-1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yloxy)methyl)biphenyl-3-carboxylate **27**. NaOH (50% aqueous) was added to a solution of the crude product (0.150 g, 0.3 mmol) in dioxane (5 mL). The reaction mixture was heated at reflux for 1 h, and then cooled to room temperature. Excess solvent was removed *in vacuo*, the residue was ddiluted with water and then acidified using 1M HCl. Extraction using EtOAc, followed by purification using preparative HPLC afforded **28** (0.162 g, 69% over 2 steps) as a colorless solid. mp: 147–149 °C. ¹H NMR (DMSO-d₆): δ 8.00 (s, 1H), 7.87-7.75 (m, 3H), 7.65-7.57 (m, 2H), 7.49-7.41 (m, 2H),

6.95-6.90 (overlapping doublets and singlet, 2H), 5.20 (s, 2H), 4.97-4.93 (m, 1H), 3.34 (t, J=6.1 Hz, 2H), 2.84 (t, J=6.1 Hz, 2H), 1.69-1.46 (m, 8 H). 13 C NMR (DMSO-d₆): 163.32, 161.79, 141.60, 138.58, 137.55, 131.75, 130.99, 129.84, 129.16, 128.86, 126.45,113.78, 112.80, 69.70, 53.54, 42.20, 28.81, 28.63, 23.83. LC-MS (ESI) Calcd for C₂₈H₂₆ClNO₄ [M+H]⁺: 476.16. Found: 476.00. HRMS (ESI) calcd for C₂₈H₂₆ClNO₄ [M+H]⁺: 476.1623. Found: 476.1623.

mGluR2 in vitro assay

Human Embryonic Kidney (HEK-293) cell lines co-expressing rat mGluR 2, 3, 4, 6, 7 or 8 and G protein inwardly rectifying potassium (GIRK) channels 16 were grown in Growth Media containing 45% DMEM, 45% F-12, 10% FBS, 20 mM HEPES, 2 mM L-glutamine, antibiotic/antimycotic, non-essential amino acids, 700 µg/ml G418, and 0.6 µg/ml puromycin at 37°C in the presence of 5% CO₂. Rat mGluR1 and 5 cells were cultured as described in Hemstapat et al.²⁵ All cell culture reagents were purchased from Invitrogen Corp. (Carlsbad, CA) unless otherwise noted. Calcium assays were used to assess activity of compounds at mGluRs 1 and 5 as previously described in Engers et al. ²⁶ Compound activity at the rat group II (mGluRs 2 and 3) and group III (mGluRs 4, 6, 7, and 8) mGluRs was assessed using thallium flux through GIRK channels, a method that has been described in detail. 16 Briefly, cells were plated into 384-well, black-walled, clear-bottomed poly-Dlysine-coated plates at a density of 15,000 cells/20 µl/well in DMEM containing 10% dialyzed FBS, 20 mM HEPES, and 100 units/ml penicillin/streptomycin (assay media). Plated cells were incubated overnight at 37 °C in the presence of 5% CO₂. The following day, the medium was removed from the cells, and 20 μL/well of 1.7 μM BTC-AM, an indicator dye (Invitrogen; prepared as a stock in DMSO and mixed in a 1:1 ratio with Pluronic acid F-127) in assay buffer [Hanks' balanced salt solution (Invitrogen) containing 20 mM HEPES, pH 7.3] was added to the plated cells. Cells were incubated for 1 h at room temperature, and the dye was replaced with 20 µl assay buffer. Test compounds were diluted to 2 times their final desired concentration in assay buffer. Agonists were diluted in thallium buffer [125 mM sodium bicarbonate (added fresh the morning of the experiment), 1 mM magnesium sulfate, 1.8 mM calcium sulfate, 5 mM glucose, 12 mM thallium sulfate, and 10 mM HEPES, pH 7.3] at 5 times the final concentration to be assayed. Cell plates and compound plates were loaded onto a kinetic imaging plate reader (FDSS 6000; Hamamatsu Corporation, Bridgewater, NJ). Appropriate baseline readings were taken (10 images at 1 Hz; excitation, 470 ± 20 nm; emission, 540 ± 30 nm) and test compounds were added in a 20 µl volume and incubated for 2.5 min before the addition of 10 µl of thallium buffer with or without agonist. After the addition of agonist, data were collected for an additional 2.5 min. Data were analyzed using Excel (Microsoft Corp, Redmond, WA). The slope of the fluorescence increase beginning 5 s after thallium/agonist addition and ending 15 s after thallium/agonist addition was calculated, corrected to vehicle and maximal agonist control slope values, and plotted in Prism software (GraphPad Software, San Diego, CA) to generate concentration-response curves. Potencies were calculated from fits using a fourpoint parameter logistic equation. For concentration-response curve experiments, compounds were serially diluted 1:3 into 10 point concentration response curves and were transferred to daughter plates using an Echo acoustic plate reformatter (Labcyte, Sunnyvale, CA). Test compounds were applied and followed by EC20 concentrations of glutamate. For selectivity experiments, full concentration-response curves of glutamate or L-AP4 (for mGluR7) were performed in the presence of a 10 µM concentration of compound and compounds that affected the concentration-response by less than 2 fold in terms of potency or efficacy were designated as inactive.

Microsomal stability in vitro assay

Pooled rat liver microsomes (BD Biosciences, # 452701) were preincubated with test compounds at 37.5 °C for 5 min in the absence of NADPH. The reaction was initiated by addition of NADPH and incubated under the same conditions. The final incubation concentrations were 4 μM test compound, 2 mM NADPH, and 1 mg/mL (total protein) liver microsomes in phosphate-buffered saline (PBS) at pH 7.4. One aliquot (100 μL) of the incubation mixture was withdrawn at 15 min time points and combined immediately with 100 μL of ACN/MeOH. After mixing, the sample was centrifuged at approximately 13000 rpm for 12 min. The supernatant was filtered and transferred into an autosampler vial and the amount of test compound was quantified using a Shimadzu LCMS 2010EV mass spectrometer. The change of the AUC (area under the curve) of the parent compound as a function of time was used as a measure of microsomal stability. Test compounds were run in duplicate with a positive control.

Plasma stability in vitro assay

A 20 μ L aliquot of a 10 mM solution in DMSO of the test compound was added to 2.0 mL of heparinized rat plasma (Lampire, P1-150N) to obtain a 100 μ M final solution. The mixture was incubated for 1 h at 37.5 °C. Aliquots of 100 μ L were taken at 15 min intervals and diluted with 100 μ L of MeOH/ACN. After mixing, the sample was centrifuged at approximately 13000 rpm for 12 min. The supernatant was filtered and transferred into an autosampler vial and the amount of test compound was quantified using the Shimadzu LCMS-2010EV system. The change of the AUC of the parent compound in function of time was used as a measure of plasma stability.

Parallel artificial membrane permeation assay (PAMPA)

A 96-well microtiter plate (Millipore, # MSSACCEPTOR) was completely filled with aqueous buffer solution (pH 7.4) and covered with a microtiter filterplate (Millipore, # MAPBMN310) to create a sort of sandwich construction. The hydrophobic filter material was impregnated with a 10% solution of polar brain lipid extract in chloroform (Avanti) as the artificial membrane, and the organic solvent was allowed to completely evaporate. Permeation studies were started by the transfer of 200 μ L of a 100 μ M test compound solution on top of the filter plate. In general phosphate pH 7.2 buffer was used. The maximum DMSO content of the stock solutions was <1.5%. In parallel, an equilibrium solution lacking a membrane was prepared using the exact concentrations and specifications but lacking the membrane. The concentrations of the acceptor and equilibrium solutions were determined using the Shimadzu LCMS-2010EV and AUC methods. The Acceptor plate and equilibrium plate concentrations were used to calculate the permeability rate (Log P_e) of the compounds. The log P_e values were calculated using the following equation:

$$\begin{split} \operatorname{Log} Pe = & log\{C \bullet - ln(1 - [drug]_{\scriptscriptstyle Acceptor} / [drug]_{\scriptscriptstyle equilibrium})\} \\ & C = & (V_{\scriptscriptstyle D} \bullet V_{\scriptscriptstyle A}) / ((V_{\scriptscriptstyle D} + V_{\scriptscriptstyle A}) Area \bullet time)) \end{split}$$

Behavioral procedures

Apparati—Intravenous cocaine self-administration and food-maintained responding took place in 24 Plexiglas experimental chambers $(25 \times 31 \times 24 \text{ cm}; \text{MED Associates}, \text{St.}$ Albans, VT), each housed in a sound-attenuating box (San Diego Instruments, San Diego, CA). One wall of the chamber contained two levers, measuring approximately 3 cm in width and located approximately 3 cm above the metal grid floor of the chamber. There was also an opaque disk (2.5 cm diameter) mounted 10 cm above each lever which could be illuminated by a 2.5 W, 24 V white light bulb and served as a cue light paired with cocaine

or food delivery. All data collection and test session functions were controlled by computers and MED-PC IV software (MED Associates).

Food training and food-maintained responding—Methodological details of food training and food-maintained responding procedures have been described previously. ^{27–30}, Rats were placed under food restriction (20 g/day/rat) and trained to lever press for 45 mg food pellets (Research Diets, New Brunswick, NJ). Training started on a fixed-ratio 1 time-out 1 s (FR1 TO1 s) schedule of reinforcement, which then changed to FR1 TO 20 s. During food training, only the active lever was used and there was no cue light presentation. Successful acquisition of the food response was defined as earning 100 pellets during a 60-min session. The training period lasted approximately 5 days. After successful acquisition of food-maintained responding, these rats were allowed to continue responding for food on an FR1 TO20 s schedule during 1-h test sessions 5–7 days per week. During these sessions, two levers (active and inactive) were present in the box. Responses on an active lever resulted in a food pellet delivery paired with a 20 s cue light presentation. Responses on inactive lever had no programmed consequences. An identical training procedure was used for both the subjects that responded for food and the rats that later were allowed to selfadminister cocaine.

Cocaine self-administration procedure—Methodological details of catheter construction and surgery and self-administration procedure have been described previously. ^{26–29} Briefly, rats were prepared with intravenous catheters inserted into the right jugular vein under isoflurane anesthesia (1–1.5% isoflurane/oxygen mixture). Cocaine self-administration training began after 7 days of recovery from surgery during daily 1-h self-administration testing sessions. These sessions were initiated by extension of both the active and inactive levers. Responses at the active lever were reinforced under an FR1 TO 20 s schedule of reinforcement by an intravenous injection of cocaine (0.5 mg/kg/infusion salt) delivered over a 2-s period in a volume of 0.05 ml. Upon completion of the operant response for cocaine delivery, a cue light above the lever was turned on for 20 s and signaled the time-out period, during which responses had no consequences. Responses at the inactive lever were recorded but had no scheduled consequences. All rats were trained to selfadminister cocaine for 21 days. The criteria for exhibiting stable performance (more than 10 infusions per session; less than 20% variation over three consecutive sessions) were reached by all rats during this 10-day period.

Experimental design for investigating the effects of the mGluR2 PAM on cocaine self-administration and food-maintained responding—After completion of the cocaine self-administration training, drug treatments were initiated. Compound **14** was administered to rats (n=9) according to a within-subjects Latin-square design. The compound was administered orally at doses of 0, 5, 10 and 20 mg/kg with 60 min pretreatment time. The highest dose (40 mg/kg) was administered to all rats after completion of the Latin-square. At least three days elapsed between drug/vehicle injections to reestablish stable self-administration behavior (less than 20% variation over three days).

To assess possible nonspecific effects, the effects of compound **14** at doses identical to those used for cocaine self-administration were assessed in a separate group of rats (n=8) trained to lever-press for food during 1-h sessions. The FR1 TO20 s schedule of reinforcement used for food-maintained responding was identical in all parameters to the schedule under which cocaine was self-administered.

Statistical analyses—Intravenous self-administration and food-maintained responding data were expressed as a percentage of the baseline number of rewards earned, with baseline defined as the mean number of rewards earned during the three days before each drug

testing session. Percent values are presented to allow for direct comparisons for data derived from the two different reinforcers (i.e., cocaine and food). Acute effects of compound **14** on cocaine self-administration and food-maintained responding were analyzed using two-way repeated-measures ANOVA, with *Dose* as the within-subjects factor, and *Reinforcer* (food or cocaine) as the between-subject factor. *Post hoc* comparisons were conducted with Newman-Keuls and Bonferroni tests. The criterion for significance was set at 0.05.

Results—The number of baseline cocaine infusions was 20 ± 4.0 (mean \pm SEM) and the number of baseline food pellets earned was 100 ± 0.0 (mean \pm SEM). A two-way ANOVA on the number of baseline cocaine/food infusions earned during the three baseline days prior to the drug treatment days indicated no significant differences across days, demonstrating the stability of baseline food and cocaine self-administration in rats during the experiment (data not shown).

Systemic administration of compound **14** significantly decreased both cocaine and food self-administration in rats (Fig. 2 in main text). A two-way ANOVA revealed significant main effects for the factor compound **14** *Dose* ($F_{4,60} = 14,88$, p < 0.0001), the factor *Reinforcer* ($F_{1,60} = 35.51$, p < 0.0001) and their interaction ($F_{4,60} = 8.75$, p < 0.0001). *Post hoc* tests showed a significant decrease in cocaine intake after administration of 20 mg/kg (p<0.01) and 40 mg/kg (p<0.001) of compound **14** compared with the vehicle treatment condition. Compound 14 had no effect on inactive lever presses at any of the doses tested (data not shown). Compound 14 also decreased food-maintained responding and this effect was significant only at the highest dose used (40 mg/kg). Additionally, *post-hoc* tests comparing the effects of the same dose of compound **14** on cocaine- and food-maintained responding indicated that compound **14** had more pronounced effect in decreasing cocaine compared to food self-administration (Figure 2 in main text).

Acknowledgments

This work was supported by National Institutes of Health grant R01 DA023926 to NC. The authors would like to thank Dr. Xinchun Jin for collection of the cocaine and food self-administration data, and Dr. Anthony Pinkerton for helpful comments during the preparation of the manuscript.

Abbreviations

mGluR2 metabotropic glutamate receptor subtype 2

PAM positive allosteric modulator

BINA 3'-((2-cyclopentyl-6,7-dimethyl-1-oxo-2,3-dihydro-1H-inden-5-

yloxy)methyl)biphenyl-4-carboxylic acid

HTS high-throughput screening

PK pharmacokinetic
POC proof-of-concept
AUC area under curve
LOQ limit of quantitation

AIBN 2,2'-azo bisisobutyronitrile

NBS *N*-bromosuccinimide

HEK Human Embryonic Kidney

GIRK G protein inwardly rectifying potassium

PBS phosphate-buffered saline

DMEM Dulbecco's modified eagle medium

HEPES 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

FBS fetal bovine serum

References

 (a) Dackis CA, O'Brien CP. Cocaine dependence: a disease of the brain's reward centers. J Subst Abuse Treat. 2001; 21:111–117. [PubMed: 11728784] (b) Leshner AI. Addiction is a brain disease, and it matters. Science. 1997; 278:45–47. [PubMed: 9311924] (c) Ohishi H, Neki A, Mizuno N. Distribution of a metabotropic glutamate receptor, mGluR2, in the central nervous system of the rat and mouse: an immunohistochemical study with a monoclonal antibody. Neurosci Res. 1998; 30:65–82. [PubMed: 9572581]

- Yuferov V, Butelman ER, Kreek MJ. Biological clock: biological clocks may modulate drug addiction. Eur J Hum Genet. 2005; 13:1101–1103. [PubMed: 16094306]
- (a) Gass JT, Olive MF. Glutamatergic substrates of drug addiction and alcoholism. Biochem Pharmacol. 2008; 75:218–265. [PubMed: 17706608] (b) Kalivas PW. Glutamate systems in cocaine addiction. Curr Opin Pharm. 2004; 4:23–29. (c) Kalivas PW. The glutamate homeostasis hypothesis of addiction. Nat Rev. 2009; 10:561–572. (d) Kalivas PW, Duffy P. Repeated cocaine administration alters extracellular glutamate in the ventral tegmental area. J Neurochem. 1998; 70:1497–1502. [PubMed: 9523566] (e) Kenny PJ, Markou A. The ups and downs of addiction: role of metabotropic glutamate receptors. Trends Pharmacol Sci. 2004; 25(5):265. [PubMed: 15120493] (f) Knackstedt LA, Kalivas PW. Glutamate and reinstatement. Curr Opin Pharm. 2009; 9:59–64. (g) Mansvelder HD, McGehee DS. Long-term potentiation of excitatory inputs to brain reward areas by nicotine. Neuron. 2000; 27:349–357. [PubMed: 10985354] (h) Markou A. Metabotropic glutamate receptor antagonists: Novel therapeutics for nicotine dependence and depression? Biol Psychiatry. 2007; 61:17–22. [PubMed: 16876138] (i) Ungless MA, Whistler JL, Malenka RC, Bonci A. Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. Nature. 2001; 411:583–587. [PubMed: 11385572]
- 4. (a) Moran MM, McFarland K, Melendez RI, Kalivas PW, Seamans JK. Cystine/glutamate exchange regulates metabotropic glutamate receptor presynaptic inhibition of excitatory transmission and vulnerability to cocaine seeking. J Neurosci. 2005; 25:6389–6393. [PubMed: 16000629] (b) Xi ZX, Ramamoorthy S, Baker DA, Shen H, Samuvel DJ, Kalivas PW. Modulation of group II metabotropic glutamate receptor signaling by chronic cocaine. J Pharmacol Exp Ther. 2002; 303:608–615. [PubMed: 12388642]
- (a) Feltenstein MW, See RE. The neurocircuitry of addiction: an overview. Br J Pharmacol. 2008; 154:261–274. [PubMed: 18311189] (b) Koob GF. Neuroadaptive mechanisms of addiction: studies on the extended amygdala. Eur Neuropsychopharmacol. 2003; 13:442–452. [PubMed: 14636960] (c) Schaffhauser H, Rowe BA, Morales S, Chavez-Noriega LE, Yin R, Jachec C. Pharmacological characterization and identification of amino acids involved in the positive modulation of metabotropic glutamate receptor subtype 2. Mol Pharmacol. 2003; 64:798–810. [PubMed: 14500736] (d) Wright RA, Arnold MB, Wheeler WJ, Ornstein PL, Schoepp DD. [3H]LY341495 binding to group II metabotropic glutamate receptors in rat brain. J Pharmacol Exp Ther. 2001; 298:453–460. [PubMed: 11454905]
- 6. (a) Adewale AS, Platt DM, Spealman RD. Pharmacological stimulation of group ii metabotropic glutamate receptors reduces cocaine self-administration and cocaine-induced reinstatement of drug seeking in squirrel monkeys. J Pharmacol Exp Ther. 2006; 318:922–931. [PubMed: 16675638] (b) Aujla H, Martin-Fardon R, Weiss F. Rats with extended access to cocaine exhibit increased stress reactivity and sensitivity to the anxiolytic-like effects of the mGluR 2/3 agonist LY379268 during abstinence. Neuropharmacology. 2008; 33:1818–1826. (c) Baptista MA, Martin-Fardon R, Weiss F. Preferential effects of the metabotropic glutamate 2/3 receptor agonist LY379268 on conditioned reinstatement versus primary reinforcement: comparison between cocaine and a potent conventional reinforcer. J Neurosc. 2004; 24:4723–4727. (d) Lee B, Platt DM, Rowlett JK, Adewale AS,

Spealman RD. Attenuation of behavioral effects of cocaine by the metabotropic glutamate receptor 5 antagonist 2-methyl-6-(phenylethynyl)-pyridine in squirrel monkeys: comparison with dizocilpine. J Pharmacol Exp Ther. 2005; 312:1232–1240. [PubMed: 15550570] (e) Peters J, Kalivas PW. The group II metabotropic glutamate receptor agonist, LY379268, inhibits both cocaine- and food-seeking behavior in rats. Psychopharmacology. 2006; 186:143–149. [PubMed: 16703399] (f) Weiss F. Neurobiology of craving, conditioned reward and relapse. Curr Opin Pharm. 2005; 5:9–19. (g) Monn JA, Valli MJ, Massey SM, Hansen MM, Kress TJ, Wepsiec JP, Harkness AR, Grutsch JL Jr, Wright RA, Johnson BG, Andis SL, Kingston A, Tomlinson R, Lewis R, Griffey KR, Tizzano JP, Schoepp DD. Synthesis, pharmacological characterization, and molecular modeling of heterobicyclic amino acids related to (+)-2-aminobicyclo[3.1.0] hexane-2,6-dicarboxylic acid (LY354740): identification of two new potent, selective, and systemically active agonists for group II metabotropic glutamate receptors. J Med Chem. 1999; 42:1027–1040. [PubMed: 10090786] (h) Conn PJ, Pin JP. Pharmacology and functions of metabotropic glutamate receptors. Annu Rev Pharmacol Toxicol. 1997; 37:205–237. [PubMed: 9131252]

- Xi ZX, Baker DA, Shen H, Carson DS, Kalivas PW. Group II metabotropic glutamate receptors modulate extracellular glutamate in the nucleus accumbens. J Pharmacol Exp Ther. 2002; 300:162– 171. [PubMed: 11752112]
- Jin X, Semenova S, Yang L, Ardecky A, Sheffler DJ, Dahl R, Conn JP, Cosford NDP, Markou M.
 The mGluR2 positive allosteric modulator BINA decreases cocaine self-administration, cue-induced
 cocaine-seeking and counteracts cocaine-induced enhancement of brain reward function in rats.
 Neuropsychopharmacol. 2010; 35:2021–2036.
- Bossert JM, Poles GC, Sheffler-Collins SI, Ghitza UE. The mGluR2/3 agonist LY379268 attenuates context- and discrete cue-induced reinstatement of sucrose seeking but not sucrose selfadministration in rats. Behav Brain Res. 2006; 173:148–152. [PubMed: 16834996]
- Cartmell J, Schoepp DD. Regulation of neurotransmitter release by metabotropic glutamate receptors. J Neurochem. 2000; 75:889–907. [PubMed: 10936169]
- 11. (a) Pinkerton AB, Cube RV, Hutchinson JH, Rowe BA, Schaffhauser H, Zhao X, Daggett LP, Vernier JM. Allosteric potentiators of the metabotropic glutamate receptor 2 (mGlu2). Part 1: Identification and synthesis of phenyl-tetrazolyl acetophenones. Bioorg Med Chem Lett. 2004; 14:5329-5332. [PubMed: 15454221] (b) Hu E, Chua PC, Tehrani L, Nagasawa JY, Pinkerton AB, Rowe BA, Vernier JM, Hutchinson JH, Cosford NDP. Pyrimidine methyl anilines: selective potentiators for the metabotropic glutamate 2 receptor. Bioorg Med Chem Lett. 2004; 14:5071-5074. [PubMed: 15380201] (c) Pinkerton AB, Cube RV, Hutchinson JH, James JK, Gardner MF, Rowe BA, Schaffhauser H, Rodriguez DE, Campbell UC, Daggett LP, Vernier JM. Allosteric potentiators of the metabotropic glutamate receptor 2 (mGlu2). Part 3: Identification and biological activity of indanone containing mGlu2 receptor potentiators. Bioorg Med Chem Lett. 2005; 15:1565–1571. [PubMed: 15745798] (d) Johnson MP, Baez M, Jagdmann GE Jr, Britton TC, Large TH, Callagaro DO, Tizzano JP, Monn JA, Schoepp DD. Discovert of allosteric potentiators for the metabotropic glutamate 2 receptor: Synthesis and subtype selectivity of N-(4-(2-methoxyphenoxy) phenyl)-N-(2,2,2-trifluroethylsulfonyl)pyrd-3-ylmethyl-amine. J Med Chem. 2003; 46:3189-3192. [PubMed: 12852748] (e) Pinkerton AB, Vernier J-M, Schaffhauser H, Rowe A, Campbell UC, Rodriguez DE, Lorrain DS, Baccei Christopher S, Daggett LP, Bristow LJ. Phenyl-tetrazolyl acetophenones: discovery of positive allosteric potentiatiors for the metabotropic glutamate 2 receptor. J Med Chem. 2004; 47:4595-4599. [PubMed: 15317469] (f) Govek SP, Bonnefous C, Hutchinson JH, Kamenecka T, McQuiston J, Pracitto R, Zhao LX, Gardner MF, James JK, Daggett LP, Rowe BA, Schaffhauser H, Bristow LJ, Campbell UC, Rodriguez DE, Vernier JM. Benzazoles as allosteric potentiators of metabotropic glutamate receptor 2 (mGluR2): efficacy in an animal model for schizophrenia. Bioorg Med Chem Lett. 2005; 15:4068-4072. [PubMed: 16005222] (g) Duplantier AJ, Efremov I, Candler J, Doran AC, Ganong AH, Haas JA, Hanks AN, Kraus KG, Lazzaro JT, Lu J, Maklad N, McCarthy SA, O'Sullivan TJ, Rogers BN, Siuciak JA, Spracklin DK, Zhang L. 3-Benzyl-1,3-oxazolidin-2-ones as mGluR2 positive allosteric modulators: Hit-to lead and lead optimization. Bioorg Med Chem Lett. 2009; 19:4068-4072. (h) Cacciola, J.; Empfield, J.; Folmer, J.; Hunter, AM.; Throner, S. Metabotropic glutamate receptor isoxazole ligands and their use as potentiators. US 2009/0306158. p. A1 (i) Tresadern G, Cid JM, Macdonald GJ, Vega JA, de Lucas AI, Garcia A, Matesanz ELM, Oehlrich D, Lavreysen H, Biesmans I, Trabanco A. Scaffold hopping from pyridones to imidazo[1,2-a]pyridines. New

- positive allosteric modulators of metabotropic glutamate 2 receptor. Bioorg Med Chem Lett. 2010; 1:175–179. [PubMed: 19932615]
- Gjoni T, Urwyler S. Receptor activation involving positive allosteric modulation, unlike full agonism, does not result in GABA_B receptor desensitization. Neuropharmacology. 2008; 55:1293–1299. [PubMed: 18775443]
- Bonnefous C, Vernier J-M, Hutchinson JH, Gardner MF, Cramer M, James JK, Rowe BA, Daggett LP, Schaffhauser H, Kamenecka TM. Biphenyl-indanones: allosteric potentiators of the metabotropic glutamate subtype 2 receptors. Bioorg Med Chem Lett. 2005; 15:4354–4358.
 [PubMed: 16046122]
- 14. Galici R, Jones CK, Hemstapat K, Nong Y, Echemndia NG, Williams LC, de Paulis T, Conn JP. Biphenyl-indanone A, a positive allosteric modulator of the metabotropic receptor subtype 2, has antipsychotic-and anxiolytic-like effects in mice. J Pharmacol Exp Ther. 2006; 318:173–185. [PubMed: 16608916]
- 15. (a) D'Alessandro PL, Corti C, Roth A, Ugolini A, Sava A, Montanari D, Bianchi F, Garland SL, Powney B, Koppe EL, Rocheville M, Osborne G, Perez P, de la Fuente J, Frailes MDL, Smith PW, Branch C, Nash D, Watson SP. The identification of structurally novel, selective, orally bioavailable positive modulators of mGluR2. Bioorg Med Chem Lett. 2010; 20:759–762. [PubMed: 20005096] (b) Brnardic EJ, Fraley ME, Garbaccio RM, Layton ME, Sanders JM, Culberson C, Jacobson MA, Magliaro BC, Hutson PH, O'Brien JA, Huszar SL, Uslaner JM, Fillgrove KL, Tang C, Kuo Y, Sur SM, Hartman GD. 3-Aryl-5-phenoxymethyl-1,3-oxazolidin-2-ones as positive allosteric modulators of mGluR2 for the treatment of Schizophrenia: hit-to-lead efforts. Bioorg Med Chem Lett. 2010; 20:3129–3133. [PubMed: 20409708]
- 16. Niswender CM, Johnson KA, Luo Q, Ayala JE, Kim C, Conn PJ, Weaver CD. A novel assay of Gi/o-linked G protein-coupled receptor coupling to potassium channels provides new insights into the pharmacology of the group III metabotropic glutamate receptors. Mol Pharmacol. 2008; 73:1213–1224. [PubMed: 18171729]
- 17. Favor, DA.; Powers, JJ.; Repine, JT.; White, AD. Isoindole derivatives. WO. 2008/020306. Feb 21. 2008
- 18. Hillemann, CL. Herbicidal sulfonamides. US. 1986/4632693. Dec. 1986
- Correa A, Tellitu I, Domínguez E, SanMartin R. Novel alternative for N-S bond formation and its application to the synthesis of benzisothiazol-3-ones. Org Lett. 2006; 8:4811–4813. [PubMed: 17020309]
- 20. Guo-qiang S. The first general synthesis of *N*-substituted 1,2-benzisoxazolin-3-ones. Tetrahedron Lett. 2000; 41:2295–2298.
- Wang XJ, Tan J, Grozinger K. A significantly improved condition for cyclization of phenethylcarbamates to *N*-alkylated 3,4-dihydroisoquinolones. Tetrahedron Lett. 1998; 39:6609– 6612.
- 22. Di L, Kerns EH, Fan K, McConnell OJ, Carter GT. High throughput artificial membrane permeability assay for blood-brain barrier. Eur J Med Chem. 2003; 38:223–232. [PubMed: 12667689]
- 23. Hinde NJ, Hall CD. Kinetics and mechanism of the formation of mono- and di-phthalate esters catalysed by titanium and tin alkoxides. J Chem Soc, Perkin Trans. 1998; 2:1249–1256.
- 24. Wyrick SD, Smith FT, Kemp WE, Grippo AA. Effects of [(N-Alkyl-1,3-dioxo-1~,3H-isoindolin-5-yl)oxy]alkanoic acids, [(N-Alkyl-l-oxo-1H,3H-isoindolin-5-y1)oxy]butanoic acids, and Related Derivatives on Chloride Influx in Primary Astroglial Cultures. J Med Chem. 1987; 30:1798–1806. [PubMed: 3656355]
- 25. Hemstapat K, Da Costa H, Nong Y, Brady AE, Luo Q, Niswender CM, Tamagnan GD, Conn PJ. A novel family of potent negative allosteric modulators of group II metabotropic glutamate receptors. J Pharmacol Exp Ther. 2007; 322:254–264. [PubMed: 17416742]
- 26. Engers DW, Niswender CM, Weaver CD, Jadhav S, Menon UN, Zamorano R, Conn PJ, Lindsley CW, Hopkins CR. Synthesis and Evaluation of a Series of Heterobiaryl Amides that are Centrally Penetrant Metabotropic Glutamate Receptor 4 (mGluR4) Positive Allosteric Modulators (PAMs). J Med Chem. 2009; 52:4115–4118. [PubMed: 19469556]

27. Kenny PJ, Paterson NE, Boutrel B, Semenova S, Harrison AA, Gasparini F, Koob GF, Skoubis PD, Markou A. Metabotropic glutamate 5 receptor antagonist MPEP decreased nicotine and cocaine self-administration but not nicotine and cocaine-induced facilitation of brain reward function in rats. Ann N Y Acad Sci. 2003; 1003:415–418. [PubMed: 14684476]

- Paterson NE, Markou A. Increased motivation for self-administered cocaine after escalated cocaine intake. Neuroreport. 2003; 14:2229–2232. [PubMed: 14625453]
- 29. Paterson NE, Markou A. The metabotropic glutamate receptor 5 antagonist MPEP decreased break points for nicotine, cocaine and food in rats. Psychopharmacol (Berl). 2005; 179:255–261.
- 30. Semenova S, Markou A. Cocaine-seeking behavior after extended cocaine-free periods in rats: role of conditioned stimuli. Psychopharmacology (Berl). 2003; 168:192–200. [PubMed: 12402103]

Figure 1. Incorporation of heteroatoms into BINA (1).




Figure 2. Effects of compound 14 on cocaine self-administration and food-maintained responding in rats. Data are expressed as percent of baseline responding (mean \pm SEM). The number of baseline cocaine infusions was 20 ± 4.0 (mean \pm SEM) and the number of baseline food pellets earned was 100 ± 0.0 (mean \pm SEM). *p < 0.05, **p < 0.01, ***p < 0.001 significant differences from the corresponding vehicle condition (Newman-Keuls post hoc test). ##, p < 0.01 significant difference between cocaine- and food- maintained responding for specific doses of compound 14 (Bonferroni post-hoc test). Cpd = compound.

Scheme 1.

Synthesis of 4-chloro-3'-((2-cyclopentyl-1-oxoisoindolin-5-yloxy)methyl) biphenyl-3-carboxylic acid derivatives $\bf 8a-8k.^a$

 aReagents and conditions: (a) $\Delta;$ (b) $R^1NH_2,$ toluene, 4Å MS, 110 °C, 12 h. (c) Zn/AcOH, 5 min. (d) $K_2CO_3,$ acetone, 60 °C, 2.5 h (e) LiOH, THF, 80 °C, 1 h.

Scheme 2.

Synthesis of 3-chloro-3'-((2-cyclopentyl-3-oxo-2,3-dihydrobenzo[d]isothiazol-6-yloxy)methyl)biphenyl-4-carboxylic acid $\mathbf{14}^a$ aReagents and conditions: (a) AlMe₃, cyclopentylamine, CH₂Cl₂, 0 °C -60 °C, 12 h. (b) PIFA, TFA, 0 °C, CH₂Cl₂, 1 h. (c) i) BBr₃, C₆H₆, 80 °C, 1 h, ii) H₂O, 100 °C, 1 h. (d) ArCH₂Br, K₂CO₃, MeCN, 80 °C, 12 h. (e) LiI, pyridine, 110 °C, 12 h.

Scheme 3.

Synthesis of 4-chloro-3'-((2-cyclopentyl-3-oxo-2,3-dihydrobenzo[d]isoxazol-6-yloxy)methyl)biphenyl-3-carboxylic acid $\bf 21^a$

 $^{a}Reagents \ and \ conditions: (a) \ i) \ AcCl, \ Py, \ CH_{2}Cl_{2}, \ 0 \ ^{\circ}C\text{-rt}, \ 30 \ min. \ ii) \ H_{2}O, \ 1 \ h. \ (b) \ COCl_{2}, \ CH_{2}Cl_{2}, \ 60 \ ^{\circ}C, \ 1 \ h. \ (c) \ 1M \ Na_{2}CO_{3}, \ THF, \ CH_{2}Cl_{2}, \ 0 \ ^{\circ}C\text{-rt}, \ 1 \ h. \ (d) \ TPP, \ DEAD, \ THF, \ 0 \ ^{\circ}C \ -rt, \ 30 \ min, \ MeOH, \ AcOH \ (e) \ i) \ BBr_{3}, \ C_{6}H_{6}, \ 80 \ ^{\circ}C, \ 1 \ h, \ ii) \ H_{2}O, \ 100 \ ^{\circ}C, \ 1 \ h. \ (f) \ ArCH_{2}Br, \ K_{2}CO_{3}, \ MeCN, \ 80 \ ^{\circ}C, \ 12 \ h. \ (g) \ LiI, \ Py, \ 110 \ ^{\circ}C, \ 12 \ h. \$

Scheme 4.

Synthesis of 4-Chloro-3'-((2-cyclopentyl-1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yloxy)methyl)biphenyl-3-carboxylic acid $\bf 28^a$

^aReagents and conditions: (a) iodocyclopentane, μW, AcCN, 110 °C, 10 min. (b) ClCO₂Me, TEA, Et₂O, 0 °C, 30 min., 1 h. (c) i)P₂O₅, POCl₃, 100 °C, 2 h (d) i) BBr₃, C₆H₆, 80 °C, 1 h, ii) H₂O, 100 °C, 1 h. (e) ArCH₂Br, K₂CO₃, MeCN, 80 °C, 12 h. (f) 50% aq. NaOH, dioxane, 80 °C, 1h.

In vitro data for mGluR2 PAMs.

Table 1

Microsomal stability^c Ę 20 42 91 Plasma stability $^{\mathcal{C}}$ 100 Z 99 86 Permeability $(\log \mathrm{P})^b$ -6.2-5.5 -6.1 Ę % Glutamate Max^a $\mathbf{83.2} \pm 10.0$ 77.3 ± 8.0 88.1 ± 13.6 $\mathbf{81.4} \pm 11.7$ 0.38 ± 0.13 0.18 ± 0.05 0.05 ± 0.02 0.07 ± 0.03 $\begin{array}{c} mGluR2 \\ PAM \\ EC50 \\ \mu M^{\it d} \end{array}$ ${f R}^1$ Compound **8**a **8** 7

Microsomal stability ^c	73	76	43	86
Plasma stability ^c	001	000	66	100
Permeability (log P) ^b	-6.5	9.6	-6.3	-5.6
% Glutamate Max ^a	85.7 ± 15.5	90.3 ± 17.2	63.9 ± 7.5	64.9 ± 8.2
mGluR2 PAM EC50 μM^a	0.08 ± 0.04	0.09 ± 0.04	0.10 ± 0.05	0.22 ± 0.18
R ¹	~~~~	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	**	**
Compound	ĕ	88	8e	8f

Dilang	ya et ai.		
Microsomal stability ^C	108		102
Plasma stability ^C	100	66	100
Permeability $(\log \mathrm{P})^b$	9-9-	-6.5	-7.8
% Glutamate Max ^a	86.2 ± 21.3	63.3 ± 3.6	86.7 ± 15.2
$\begin{array}{c} \text{mGluR2} \\ \text{PAM} \\ \text{EC50} \\ \mu \text{M}^{a} \end{array}$	0.26 ± 0.12	0.27 ± 0.13	0.35 ± 0.04
צי			X OCF
Compound	∞ ∞	₩	≅

		mGluR2 PAM EC50				
Compound	\mathbb{R}^1	$\mu \mathrm{M}^a$	% Glutamate Max ^a	Permeability ($\log \mathrm{P})^b$	Plasma stability $^{\mathcal{C}}$	Microsomal stability ^c
8;	•	0.06 ± 0.03	85.3 ± 10.6	-5.8	94	21
	~~					
8k	×	> 10	ND	-4.9	100	96
14		0.17 ± 0.03	63.0 ± 3.3	-5.4	98	45
21		0.31 ± 0.05	64.0 ± 3.4	-6.3	100	35
28		0.30 ± 0.03	66.9 ± 10.1	-6.1	100	36

 a mGluR2 PAM EC50 μ M data and % Glutamate Max data represent the mean \pm SEM for at least three independent experiments performed in triplicate.

^bPermeability is monitored by measuring the amount of compound that can diffuse through a polar brain lipid membrane to predict BBB permeability.²²

 $^{\rm C}$ Percent remaining after incubation for 60 min. at 37 $^{\circ}{\rm C}.$

Table 2

 $\mathit{In\ vivo\ PK}$ data for mGluR2 PAMs in rats following oral administration. a

	oral dose (mg/kg) C _{max} (μM)	Стах (µМ)	T _{max} (min.)	$T_{max} \ (min.) \ \left \ AUC \ (0\text{-}t) \ (\mu M.h) \ \right \ t_{1/2} \ (h)$	t _{1/2} (h)	F (%)	Brain b ($\mu { m M}$)	Plasma b ($\mu { m M}$)	$(\%) \mid \operatorname{Brain}^b(\mu \mathbb{M}) \mid \operatorname{Plasma}^b(\mu \mathbb{M}) \mid \operatorname{Brain:Plasma ratio}$
1	10	0.097 ± 0.032 120	120	0.281 ± 0.097	1.3 ± 0.25 33	33	0.0	0.027 ± 0.008 ND	ND
8a	10	14.4 ± 15.8	08	105.5 ± 69.1	3.4 ± 0.7 40 3.8 ± 1.2	40	3.8 ± 1.2	12.3 ± 5.7	0.3
14	10	4.1 ± 1.0	110	43.3 ± 14.4	2.7 ± 1.1 86 4.8 ± 1.0	98	4.8 ± 1.0	1.0 ± 0.2	4.8

a Cmax: maximum concentration of the compound detected in plasma. Tmax: time at Cmax. AUC: area under the curve. 11/2: terminal half-life. F: oral bioavailability.

 $\ensuremath{^{b}}$ Brains and plasma were harvested at or near the $T_{max}.$