

# Journal of Medicinal Chemistry

© Copyright 2002 by the American Chemical Society

Volume 45, Number 7

March 28, 2002

## Letters

### Design, Synthesis, and Evaluation of Opioid Analogues with Non-Peptidic $\beta$ -Turn Scaffold: Enkephalin and Endomorphin Mimetics

Masakatsu Eguchi,<sup>\*,†,§</sup> Richard Y. W. Shen,<sup>†</sup>  
J. Paul Shea,<sup>†</sup> Min S. Lee,<sup>†</sup> and Michael Kahn<sup>\*,†,§</sup>

*Molecumetics Ltd., 2023 120th Avenue NE, Suite 400,  
Bellevue, Washington 98005, Department of Pathobiology  
University of Washington, Seattle, Washington 98195,  
and Pacific Northwest Research Institute, 720 Broadway,  
Seattle, Washington 98122*


*Received November 20, 2001*

**Abstract:** We have identified a  $\mu$ -selective opioid receptor agonist without a cationic amino group in the molecule from libraries of bicyclic  $\beta$ -turn peptidomimetics. The biologically active conformation of the lead is proposed to mimic an endomorphin type III 4  $\rightarrow$  1  $\beta$ -turn conformation.

The isolation and identification of the endogenous opioid peptides, methionine enkephalin and leucine enkephalin,<sup>1</sup> initiated an intensive effort to discern the biologically active conformation of enkephalin through comparison with the rigid morphine skeleton, which is believed to share a common binding motif.<sup>2</sup> It has been shown that the proper spatial orientation of a cationic amine, a phenol group, and an additional hydrophobic group is necessary to manifest biological activity through interaction with opioid receptors.<sup>2a,c</sup> However, the inherent conformational flexibility of the opioid peptides has hampered numerous attempts to accurately assess the bioactive conformation(s). Significant effort has been devoted to the elucidation of the receptor-bound conformation through systematic studies of conformationally constrained peptides<sup>3</sup> or peptidomimetics<sup>4</sup> or by

conformational analyses of peptides in membrane-mimicking environments.<sup>5</sup> Several studies suggested reverse turn conformations as the potential biologically active conformation of the opioid peptides based upon computational models,<sup>6</sup> X-ray crystallography,<sup>7</sup> and spectroscopic analyses.<sup>5b,8</sup>

We have developed a  $\beta$ -turn peptidomimetic with four




sites of diversity readily accessible through solid-phase synthesis from commercially available diversity components. This privileged template was shown by X-ray crystal structural analysis<sup>9a</sup> and solution-phase 2D NMR spectroscopy<sup>9</sup> to mimic a type I  $\beta$ -turn conformation accurately on a rigid bicyclic structure. This highly constrained 6,6-bicyclic system incorporating functionality at the  $i$  to  $i + 3$  positions affords us an opportunity to probe the biologically active conformation of peptides that potentially adopt a reverse turn conformation. Therefore, we applied this scaffold to the preparation of mimetics of Leu-enkephalin (YGGFL)<sup>1</sup> to identify potent and selective ligand(s) for opioid receptors ( $\mu$ ,  $\delta$ , and  $\kappa$ )<sup>10</sup> and to elucidate their structure–activity relationship (SAR). In addition to the conformational stability induced by this template, its non-peptidic nature can potentially improve the undesirable therapeutic characteristics of peptides, which include poor bioavailability, short duration of action, and lack of oral activity.<sup>11</sup> During the course of our SAR studies, we found a striking similarity between the three-dimensional structure and the receptor selectivity profile of our lead compounds and those of the recently isolated opioid peptides, endomorphin-1 (YPWF) and endomorphin-2 (YPFF).<sup>12</sup> Herein, we report the design, synthesis, and evaluation of opioid peptide analogues utilizing our privileged bicyclic  $\beta$ -turn scaffold to discover potent

\* To whom correspondence should be addressed. Address: Pacific Northwest Research Institute, 720 Broadway, Seattle, WA 98122. Phone: 206-726-1200. Fax: 206-726-1217. E-mail (for M.E.): meguchi@pnri.org. E-mail (for M.K.): mkahn@pnri.org.

<sup>†</sup> Pacific Northwest Research Institute.

<sup>§</sup> University of Washington.


<sup>‡</sup> Molecumetics Ltd.



**Figure 1.** Diversity components for enkephalin and endomorphin mimetics.


$\mu$ -receptor ligands that bear no cationic amine and we evaluate the relevance of the conformational and the functional requirements for biological activity.

Enkephalin can adopt any type of  $\beta$ -turn conformation in solution because of the presence of two glycine residues at the second and third positions. Indeed, various types of  $\beta$ -turn conformations (types I, I', II', and III)<sup>5a-c,7,8a,b</sup> of enkephalin have been observed in various media. By employing our  $\beta$ -turn scaffold, we can investigate the following hypothetical bioactive  $\beta$ -turn conformation models; enkephalin 4  $\rightarrow$  1  $\beta$ -turn model (A), enkephalin 5  $\rightarrow$  2  $\beta$ -turn model (B), and endomorphin 4  $\rightarrow$  1  $\beta$ -turn model (C). To examine these models,



we incorporate the designed diversity elements into the scaffold at the  $i$  to  $i + 3$  positions. Twelve primary amines at the  $i + 3$  position, 13  $\alpha$ -amino acids at the  $i + 2$  position, one  $\beta$ -alanine at the  $i + 1$  position, and a phenol component with three different lengths of spacer at the  $i$  position were selected as shown in Figure 1.


A focused library of opioid peptide mimetics with these components was prepared according to our previously published solid-phase synthetic protocol.<sup>9</sup> Nucleophilic displacement of the bromide, **1**, with a number of primary amines gave the corresponding secondary amine, **2**, which was then coupled with the appropriate Fmoc- $\alpha$ -amino acids with HOAt/DIC in NMP. Treatment of **3** with 25% piperidine in DMF followed by coupling with Fmoc- $\beta$ -alanine afforded **4**. After deprotection of the Fmoc group of **4** with 25% piperidine, the resin was treated with the corresponding alkyl  $p$ -nitrophenyl carbonates in the presence of DIEA to produce **5**. Cleavage from the acetal resin followed by stereoselective tandem acyliminium cyclization was accomplished by treatment with formic acid at room temperature to give the 6,6-bicyclic  $\beta$ -turn mimetic, **6**. All products were purified by preparative TLC (silica gel) to afford the pure product in good overall yields (11–37%; see Supporting Information). The tryptophan derivatives at the  $i + 2$  position and  $p$ -hydroxybenzyl-carbamate derivatives at the  $i$  position failed to afford the desired bicyclic scaffold.<sup>13</sup>



The inhibitory activity of each mimetic at 1  $\mu\text{M}$  and/or 0.1  $\mu\text{M}$  was evaluated through competition of specific binding of radioligand (<sup>3</sup>H]naloxone) for relatively nonselective opioid receptors in rat cerebral cortex.<sup>14</sup> The specific radioligand binding to the receptors is defined as the difference between total binding and nonspecific binding determined in the presence of an excess of unlabeled naloxone (1  $\mu\text{M}$ ). Results are expressed as a percent inhibition of the specific binding obtained in the presence of the test compounds (see Supporting Information for the preliminary results). Among the 12 hydrophobic residues introduced at the  $i + 3$  position, the benzyl and 2-pyridylmethyl groups showed the highest inhibitory activity and the other alkyl groups provided almost the same level of activity when the  $i + 2$  position is benzyl. Diversification of the  $\alpha$ -amino acids at the  $i + 2$  position varies the inhibitory activity from 99% to 11% while holding the functionality at the  $i + 3$  position as  $n$ -butyl. The phenethyl group at this position afforded the best result. The activities decrease in the order of phenethyl >  $p$ -aminobenzyl > benzyl > other alkyls, hydrogen, and phenyl groups >  $p$ -hydroxybenzyl group. Extension of the alkyl chain between the phenol group and the template at the  $i$  position (from ethyl to propyl) causes significant loss of inhibitory activity.

To validate this preliminary SAR result, we resynthesized four of the most active compounds.  $\beta$ -Turn mimetics with a benzyl group or phenethyl group at the  $i + 2$  position and a benzyl group or  $n$ -butyl group at the  $i + 3$  position were prepared, and the IC<sub>50</sub> values of these four compounds versus the relatively nonselective opioid receptors in rat cerebral cortex were determined as shown in Table 1. These compounds represent an enkephalin 5  $\rightarrow$  2  $\beta$ -turn model with one methylene length shorter at the  $i$  position and an endomorphin type III 4  $\rightarrow$  1  $\beta$ -turn model with one methylene length longer at the  $i$  position.<sup>15</sup> The IC<sub>50</sub> value of **6a** was measured as 149 nM. The replacement of the  $n$ -butyl group of **6a** at the  $i + 3$  position with a benzyl group, **6b**, improved the inhibitory activity almost 2-fold to 80 nM, and the replacement of the benzyl group of **6a** at the  $i + 2$  with a phenethyl group, **6c**, improved the inhibitory activity almost 5-fold to 27 nM. The inhibitory activity of **6c** was further enhanced 3-fold to 9 nM by replacing the  $n$ -butyl group at the  $i + 3$  position with a benzyl group, **6d**. All four compounds are lipophilic and bear no cationic

**Table 1.** Effects of the Mimetics on the Specific Radioligand ( $[^3\text{H}]\text{Naloxone}$ ) Binding to the Relatively Nonselective Opioid Receptor

	Compounds			Inhibition (%)		$\text{IC}_{50}$ (nM)
	$\text{R}^1$	$\text{R}^{1+2}$	$\text{R}^{1+3}$	0.1 $\mu\text{M}$	1 $\mu\text{M}$	
<b>6a</b>	<i>p</i> -OH-Phenethyl	Bn	nBu		84	149 $\pm$ 14
<b>6b</b>	<i>p</i> -OH-Phenethyl	Bn	Bn		96	80
<b>6c</b>	<i>p</i> -OH-Phenethyl	Phenethyl	nBu	79	99	27
<b>6d</b>	<i>p</i> -OH-Phenethyl	Phenethyl	Bn	91		9
<b>7</b>					<10	>1000

**Table 2.** Effects of the **6c** and **6d** on the Specific Radioligand Binding to the Human Opioid Receptors and  $\text{IC}_{50}$  Values for the Reference Compounds<sup>a</sup>



receptors	<b>6c</b> 0.1 $\mu\text{M}$	<b>6d</b> 0.1 $\mu\text{M}$	reference compounds $\text{IC}_{50}$ (nM)	
$\delta$ (h)	<10%	<10%	DPDPE	3.5
$\kappa$ (h)	<10%	<10%	U 50488	0.69
$\mu$ (h)	91%	101%	DAMGO	1.6

<sup>a</sup> Radioligands  $[^3\text{H}]\text{DPDPE}$ ,  $[^3\text{H}]\text{U 69593}$ , and  $[^3\text{H}]\text{DAMGO}$  were used for assays for human  $\delta$ -,  $\kappa$ -, and  $\mu$ -opioid receptors, respectively.  $\text{IC}_{50}$  values are within accepted limits of historic averages obtained  $\pm$  0.5 log unit.

amino group in the molecule.<sup>16</sup> Compounds **6c** and **6d** were further studied using cloned human  $\mu$ -,  $\delta$ -, and  $\kappa$ -receptors to examine the receptor selectivity as shown in Table 2. Both of the compounds, **6c** and **6d**, showed excellent  $\mu$ -selectivity for the human opioid receptors.

For comparison, a linear peptide **7** corresponding to **6c** was prepared and tested for its affinity for the rat cerebral cortex relatively nonselective opioid receptors. Although the same functional groups, such as *p*-hydroxyphenethyl, (*S*)-homophenylalanyl, and *n*-butyl groups, are displayed on the surface of the linear amide backbone structure, no binding activity was observed at the 1  $\mu\text{M}$  level. This result suggests that the correct spatial orientations and conformational restrictions of these functional groups by the rigid template are critical for high receptor affinity.

We carried out an *in vivo* evaluation of the analgesic effect of **6c**. Three groups of 10 male CD-1 mice were administered 0.1, 1.0, or 10 mg/kg of **6c** intravenously (iv) at 5.0 mL/kg. One group of 10 male CD-1 mice was administered vehicle (propylene glycol and saline, 1:1) iv at 5.0 mL/kg. Another group of 10 male CD-1 mice received the positive control, morphine sulfate at 10 mg/kg, iv, at 5.0 mL/kg. The mice were sequentially placed on a hot plate analgesia meter (set at 55  $^{\circ}\text{C}$ ) and observed for a reaction to the heat stimulus. The reaction times were obtained at 5, 15, 30, 60, and 120 min following iv administration. The administration of morphine sulfate at 10 mg/kg, iv, produced statistically significant ( $p \leq 0.05$ ) increases in mean reaction times of 279%, 234%, 164%, 113%, and 22% analgesia respectively at 5, 15, 30, 60, and 120 min postadministration. The administration of **6c** at 1.0 mg/kg, iv, produced a statistically significant ( $p \leq 0.05$ ) increase in the mean reaction time of analgesia 104% at 5 min postdose. The administration of **6c** at 10 mg/kg, iv, showed statistically significant ( $p \leq 0.05$ ) increases in the mean reaction times of 337%, 189%, and 75% analgesia respectively at 5, 15, and 30 min postadministration. The results of the intravenous mouse hot plate analgesia assay are

**Figure 2.** Intravenous mouse hot plate analgesia assay of **6c**.**Figure 3.** Overlay of the dominant low-energy conformer of **6d** (azure) and type III  $\beta$ -turn conformation of endomorphin (rose, ref 18).

summarized in Figure 2. It appears that the initial level of analgesic activity of **6c** is the same as the initial activity level of morphine, although the *in vivo* half-life of **6c** was almost 2-fold less than that of morphine.

Our 2D NMR experiments in  $\text{CDCl}_3$  at  $-20^{\circ}\text{C}$  showed that the bicyclic scaffold of **6c** and **6d** adopts the same ring conformation as observed by X-ray crystallography for the corresponding sulfonamide derivative<sup>9a</sup> and that **6c** and **6d** exist as a mixture of two rotamers about the urethane linkage. A Monte Carlo conformational search for **6d** using the MMFF force field in vacuo as implemented in MacroModel 7.1<sup>17</sup> generated the bicyclic ring structure of the lowest energy conformer that matched the core ring structure deduced from the 2D NMR experiments. The overlaid structure of **6d** with a trans-endomorphin-1  $\beta$ -turn type III like structure<sup>18</sup> revealed the structural similarity of these molecules with an rmsd value of 0.39  $\text{\AA}$  at six atom positions<sup>19</sup> as shown in Figure 3. Judging from the similarity of the three-dimensional structure and the  $\mu$ -selectivity profile of **6d**, it would be reasonable to deduce that the opioid analogue **6d** mimics the endomorphin type III 4  $\rightarrow$  1  $\beta$ -turn model.

In conclusion, we have identified a nanomolar affinity  $\mu$ -opioid receptor agonist lacking a positive charge in the molecule by employing a privileged  $\beta$ -turn scaffold. The stereochemistries of **6c** and **6d** in solution were analyzed by 2D NMR in combination with molecular mechanics, which suggest that the biologically active conformation of endomorphin is a type III 4  $\rightarrow$  1  $\beta$ -turn conformation and that the  $\mu$ -active conformation of enkephalin is a 5  $\rightarrow$  2  $\beta$ -turn conformation. Finally, we have demonstrated the application of our  $\beta$ -turn pepti-




domimetic library for the discovery of potent and selective ligands for a member of the G-protein-coupled receptor superfamily.

**Supporting Information Available:** Experimental details and a list of abbreviations. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Hughes, J.; Smith, T. W.; Kosterlitz, H. W.; Fothergill, L. A.; Morgan, B. A.; Morris, H. R. Identification of Two Related Pentapeptides from the Brain with Potent Opiate Agonist Activity. *Nature* **1975**, *258*, 577–580.
- (2) (a) Hansen, P. E.; Morgan, B. A. Structure–Activity Relationships in Enkephalin Peptides. In *The Peptides: Analysis, Synthesis, Biology*; Udenfried, S., Meienhofer, J., Eds.; Academic Press: Orlando, FL, 1990; Vol. 6, pp 269–321. (b) Schiller, P. W. Conformational Analysis of Enkephalin and Conformation–Activity Relationships. In *The Peptides: Analysis, Synthesis, Biology*; Udenfried, S., Meienhofer, J., Eds.; Academic Press: Orlando, FL, 1990; Vol. 6, pp 219–268. (c) Portoghesi, P. S.; Sultana, M.; Takemori, A. E. Design of Peptidomimetic  $\delta$  Opioid Receptor Antagonists Using the Message–Address Concept. *J. Med. Chem.* **1990**, *33*, 1714–1720 and references therein.
- (3) (a) Hrubby, V. J.; Al-Obeidi, F.; Kazmierski, W. Emerging Approaches in the Molecular Design of Receptor-Selective Peptide Ligands: Conformational, Topographical and Dynamic Considerations. *Biochem. J.* **1990**, *268*, 249–262. (b) Schiller, P. W.; Nguyen, T. M.-D.; Weltrowska, G.; Wilkes, B. C.; Mardsen, B. J.; Lemieux, C.; Chung, N. N. Differential Stereochemical Requirements of  $\mu$  vs.  $\delta$  Opioid Receptors for Ligand Binding and Signal Transduction: Development of a Class of Potent and Highly  $\delta$ -Selective Peptide Antagonists. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 11871–11875.
- (4) (a) Bélanger, P. C.; Dufresne, C.; Scheiget, J.; Young, R. N.; Springer, J. P.; Drnitsienko, G. The Design and Synthesis of Nonpeptide Compounds as Mimics of a Conformation of Methionine–Enkephalin. *Can. J. Chem.* **1982**, *60*, 1019–1029. (b) Gardner, B.; Nakanishi, H.; Kahn, M. Conformationally Constrained Nonpeptide  $\beta$ -Turn Mimetics of Enkephalin. *Tetrahedron* **1993**, *49* (17), 3433–3448. (c) Wang, C.; McFadyen, I. J.; Traynor, J. R.; Mosberg, H. I. Design of a High Affinity Peptidomimetics Opioid Agonist from Peptide Pharmacophore Models. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2685–2688. (d) Liao, S.; Alfaro-Lopez, J.; Shenderovich, M. D.; Hosohata, K.; Lin, J.; Li, X.; Stropova, D.; Davis, P.; Jernigan, K. A.; Porreca, F.; Yamamura, H. I.; Hrubby, V. J. De Novo Design, Synthesis, and Biological Activities of High-Affinity and Selective Non-Peptide Agonists of the  $\delta$ -Opioid Receptor. *J. Med. Chem.* **1998**, *41*, 4767–4776.
- (5) (a) Milon, A.; Miyazawa, T.; Higashijima, T. Transferred Nuclear Overhauser Effect Analyses of Membrane-Bound Enkephalin Analogues by  $^1\text{H}$  Nuclear Magnetic Resonance: Correlation between Activities and Membrane-Bound Conformations. *Biochemistry* **1990**, *29*, 65–75. (b) Picone, D.; D'Ursi, A.; Motta, A.; Tancredi, T.; Temussi, P. A. Conformational Preferences of [Leu $^5$ ]–Enkephalin in Biomimetic Media. Investigation by  $^1\text{H}$  NMR. *Eur. J. Biochem.* **1990**, *192*, 433–439. (c) Rudolph-Böhner, S.; Quarzago, D.; Czisch, M.; Ragnarsson, U.; Moroder, L. Conformational Preferences of Leu–Enkephalin in Reverse Micelles as Membrane-Mimicking Environment. *Biopolymers* **1997**, *41*, 591–606. (d) Amodeo, P.; Naider, F.; Picone, D.; Tancredi, T.; Temussi, P. A. Conformational Sampling of Bioactive Conformers: A Low-Temperature NMR Study of  $^{15}\text{N}$ -Leu–Enkephalin. *J. Pept. Sci.* **1998**, *4*, 253–265.
- (6) (a) Hassan, M.; Goodman, M. Computer Simulation of Cyclic Enkephalin Analogues. *Biochemistry* **1986**, *25*, 7596–7606 and references therein. (b) Smith, P. E.; Dang, L. X.; Pettitt, B. M. Simulation of the Structure and Dynamics of the Bis(penicillamine) Enkephalin Zwitterion. *J. Am. Chem. Soc.* **1991**, *113*, 67–73. (c) Chew, C.; Villar, H. O.; Loew, G. H. Theoretical Study of Flexibility and Solution Conformation of the Cyclic Opioid Peptides [D-Pen $^2$ , D-Pen $^5$ ]Enkephalin and [D-Pen $^2$ , L-Pen $^5$ ]–Enkephalin. *Mol. Pharmacol.* **1991**, *39*, 502–510. (d) Pettitt, B. M.; Matsunaga, T.; Al-Obeidi, F.; Gehrig, C.; Hrubby, V. J.; Karplus, M. Dynamic Search for Bis-penicillamine Enkephalin Conformations. *Biophys. J.* **1991**, *60*, 1540–1544.
- (7) (a) Smith, G. D.; Griffin, J. F. Conformation of [Leu $^5$ ]Enkephalin from X-ray Diffraction: Features Important for Recognition at Opiate Receptor. *Science* **1978**, *199*, 1214–1216. (b) Griffin, J. F.; Smith, G. D. X-Ray Diffraction Studies of Enkephalins and Opiates. In *Opioid Peptides: An Update*; Rapaka, R. S., Dhawan, B. N., Eds. NIDA Research Monograph 87; Washington, DC, 1988; pp 41–59. (c) Aubry, A.; Birlirakis, N.; Sakarellos-Daitsiotis, M.; Sakarellos, C.; Marroud, M. A Crystal Conformation of Leucine–Enkephalin Related to the Morphine Molecule. *Biopolymers* **1989**, *28*, 27–40.
- (8) (a) Roques, B. P.; Garbary-Jaureguiberry, C.; Oberlin, R.; Anteunis, M.; Lala, A. K. Conformation of Met $^5$ -Enkephalin Determined by High Field PMR Spectroscopy. *Nature* **1976**, *262*, 778–779. (b) Garbay-Jaureguiberry, C.; Roques, B. P.; Oberlin, R.; Anteunis, M.; Combrisson, S.; Lallemand, J. Y.  $^1\text{H}$  and  $^{13}\text{C}$  NMR Studies of Conformational Behavior of Leu–Enkephalin. *FEBS Lett.* **1977**, *76*, 93–98. (c) Hrubby, V. J.; Kao, L. F.; Pettitt, B. M.; Karplus, M. The Conformational Properties of the Delta Opioid Peptide [D-Pen $^2$ , D-Pen $^5$ ]Enkephalin in Aqueous Solution Determined by NMR and Energy Minimization Calculations. *J. Am. Chem. Soc.* **1988**, *110*, 3351–3359. (d) Mosberg, H. I.; Sobczyk-Kojiro, K.; Subramanian, P.; Crippen, G. M.; Ramalingam, K.; Woodard, R. W. Combined Use of Stereospecific Deuteration, NMR, Distance Geometry, and Energy Minimization for the Conformational Analysis of the Highly  $\delta$  Opioid Receptor Selective Peptide [D-Pen $^2$ , D-Pen $^5$ ]Enkephalin. *J. Am. Chem. Soc.* **1990**, *112*, 822–829.
- (9) (a) Eguchi, M.; Lee, M. S.; Nakanishi, H.; Stasiak, M.; Lovell, S.; Kahn, M. Solid-Phase Synthesis and Structural Analysis of Bicyclic  $\beta$ -Turn Mimetics Incorporating Functionality at the  $i$  to  $i+3$  Positions. *J. Am. Chem. Soc.* **1999**, *121*, 12204–12205. (b) Eguchi, M.; Lee, M. S.; Stasiak, M.; Kahn, M. Solid-Phase Synthesis and Solution Structure of Bicyclic  $\beta$ -Turn Peptidomimetics: Diversity at the  $i$  Position. *Tetrahedron Lett.* **2001**, *42*, 1237–1239.
- (10) Mansour, A.; Khachaturian, H.; Lewis, M. E.; Akil, H.; Watson, S. J. Anatomy of CNS Opioid Receptors. *Trends Neurosci.* **1988**, *11*, 308–314.
- (11) For a recent example: Abe, Y.; Kayakiri, H.; Satoh, S.; Inoue, T.; Sawada, Y.; Imai, K.; Inamura, N.; Asano, M.; Hatori, C.; Katayama, A.; Oku, T.; Tanaka, H. A Novel Class of Orally Active Non-Peptide Bradykinin B2 Receptor Antagonists. I. Construction of the Basic Framework. *J. Med. Chem.* **1998**, *41*, 564–578.
- (12) Zadina, J. E.; Hackler, L.; Ge, L.-I.; Kastin, A. J. A Potent and Selective Endogenous Agonist for the  $\mu$ -Opiate Receptor. *Nature* **1997**, *386*, 499–502.
- (13) Pictet–Spengler cyclization products were isolated in good yield for Trp derivatives.
- (14) Childers, S. R.; Creese, L.; Snowman, A. M.; Snyder, S. H. Opiate Receptor Binding Affected Differentially by Opiates and Opioid Peptides. *Eur. J. Pharmacol.* **1979**, *55*, 11–18.
- (15) Following a referee's suggestion, compound **8** was prepared in 26% overall yield (ref 9b) and evaluated for competition of specific binding of radioligand ( $^3\text{H}$ ]naloxone) to the relatively nonselective opioid receptor. An inhibitory activity of 70% ( $n = 2$ ) at 0.1  $\mu\text{M}$  was observed.
- (16) Schiller, P. W.; Berezowska, I.; Nguyen, T. M.-D.; Schmidt, R.; Lemieux, C.; Chung, N. N.; Falcone-Hindley, M. L.; Yao, W.; Liu, J.; Iwama, S.; Smith, A. B., III; Hirschmann, R. Novel Ligands Lacking a Positive Charge for the  $\delta$ - and  $\mu$ -Opioid Receptors. *J. Med. Chem.* **2000**, *43*, 551–559.
- (17) (a) Mohanmadi, F.; Richards, N. G. J.; Guida, W. C.; Liskamp, R.; Lipton, M.; Caufield, C.; Chang, G.; Hendrickson, T.; Still, W. C. MacroModel—An Integrated Software System for Modeling Organic and Bioorganic Molecules Using Molecular Mechanics. *J. Comput. Chem.* **1990**, *11*, 440. (b) Schrödinger, Inc. (<http://www.schrodinger.com>), Portland, OR.
- (18) Podlogar, B. L.; Paterlini, M. G.; Ferguson, D. M.; Leo, G. C.; Demeter, D. A.; Brown, F. K.; Reitz, A. B. Conformational Analysis of the Endogenous  $\mu$ -Opioid Agonist Endomorphin-1 Using NMR Spectroscopy and Molecular Modeling. *FEBS Lett.* **1998**, *439* (1–2), 13–20.
- (19) C(1), N(2), C(3), C(4), N(5), and C(6) atoms of **6d**, and the corresponding C(Tyr $^1$ ), N(Pro $^2$ ), C $\alpha$ (Trp $^3$ ), C $\beta$ (Trp $^3$ ), N(Phe $^4$ ), and C $\alpha$ (Phe $^4$ ) atoms of endomorphin-2 were compared.



JM0155897