

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/12451479>

Stannyl radical-mediated cleavage of pi-deficient heterocyclic sulfones. Synthesis Of alpha-fluoro esters

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · JULY 2000

Impact Factor: 4.72 · Source: PubMed

CITATIONS

3

READS

10

4 AUTHORS, INCLUDING:



Stanislaw F Wnuk

Florida International University

162 PUBLICATIONS 2,132 CITATIONS

SEE PROFILE

Stannyl Radical-Mediated Cleavage of π -Deficient Heterocyclic Sulfones. Synthesis of α -Fluoro Esters

Stanislaw F. Wnuk,* Jeannette M. Rios, Jahanzeb Khan, and Ya-Li Hsu

Department of Chemistry, Florida International University, Miami, Florida 33199

Received March 9, 2000

Treatment of ethyl 2-(pyridin-2-ylsulfonyl)hexanoate with tributylstannane and azobis(2-methyl-2-propanitrile) (AIBN) in benzene at reflux for 36 h resulted in hydrogenolysis to give ethyl hexanoate (60%), whereas no reaction was observed after 48 h at reflux with ethyl 2-(phenylsulfonyl)hexanoate. Ethyl 2-(pyrimidin-2-ylsulfonyl)hexanoate underwent quantitative hydrogenolysis within 1 h under these conditions. This represents a mild new methodology for removal of the synthetically useful sulfone moiety. Substitution of Bu_3SnD for Bu_3SnH gave access to α -deuterium-labeled esters. Treatment of the α -(pyrimidin-2-ylsulfonyl) enolates derived from several esters with Selectfluor gave high yields of the 2-fluoro-2-(pyrimidin-2-ylsulfonyl)alkanoates, which were smoothly desulfonylated [Bu_3SnH (2 equiv)/AIBN/benzene/ Δ] to give 2-fluoroalkanoates. "Catalytic" tin hydride, generated from tributyltin chloride (0.15 equiv) and excess polymethylhydrosiloxane in the presence of potassium fluoride, also effected removal of the π -deficient α -(pyrimidin-2-ylsulfonyl) moiety from acid derivatives in high yields. Desulfonylation is suggested to proceed via alkoxy ketyl-type radicals and tin enolates.

Introduction

The sulfone group is a well-established activating moiety for construction of carbon–carbon skeletons and other transformations.¹ During recent work on synthesis of a 6'-deoxy-6'-fluorohomonucleoside phosphonate from uridine, we found that standard procedures for desulfonylation^{1d} [e.g., $\text{Al}(\text{Hg})$ or $\text{Na}(\text{Hg})$; or base-promoted elimination] were ineffective for removal of the pyridin-2-ylsulfonyl group from the α -carbon of phosphonic esters.² We then explored the feasibility of a radical-mediated cleavage of sulfonyl groups from the α -carbon of carboxylic and phosphonic esters with tributyltin hydride. Our initial success prompted us to investigate the broader potential of radical cleavage of π -deficient heterocyclic sulfones.

Tributylstannane is used routinely for hydrogenolysis of carbon–halogen, carbon–sulfur, carbon–selenium, and carbon–nitro bonds,³ but is generally recognized as ineffective for cleavage of saturated sulfones.^{1d} Recently, desulfonylation of β -ketosulfones,⁴ *N*-sulfonylated amides,⁵ and 2-(alkyl and aryl)sulfonylpyrroles with Bu_3SnH ⁶ as well stannodesulfonylations of vinyl sulfones⁷ have been

reported. Desulfonylations of allylic sulfones^{8a} with tributylstannane are known, and sulfonyl radicals are versatile intermediates in organic synthesis.^{8b}

Selective introduction of fluorine into organic molecules often causes significant changes in biological activity.⁹ In particular, α -fluoro carbonyl compounds are important since they have been utilized as diagnostic tools in metabolic processes and serve as building blocks in the synthesis of more complex molecules.^{9a,b,e} α -Fluoro esters have been prepared from toxic fluoroacetate ions,¹⁰ by reaction of α -hydroxy esters with DAST,¹¹ electrophilic fluorination of stabilized carbanions, enolates or silyl enol ethers,¹² metal-catalyzed addition of fluoroiodoacetates to alkenes,¹³ and Reformatsky reactions with bromofluoroacetates.¹⁴ Other methods also exist.⁹

* To whom correspondence should be addressed. E-mail: wnuk@fiu.edu. Fax: (305) 348–3772.

(1) (a) *The Chemistry of Sulphones and Sulfoxides*; Patai, S., Rappoport, Z., Stirling, C., Eds.; John Wiley & Sons: Chichester, 1988. (b) Simpkins, N. S. *Sulphones in Organic Synthesis*; Pergamon Press: Oxford, 1993. (c) Jin, Z.; Vandort, P. C.; Fuchs, P. L. *Phosphorus Sulfur Silicon* **1994**, 95–96, 1–19. (d) Najera, C.; Yus, M. *Tetrahedron* **1999**, 55, 10547–10658.

(2) Wnuk, S. F.; Robins, M. J. *J. Am. Chem. Soc.* **1996**, 118, 2519–2520.

(3) (a) Neumann, W. P. *Synthesis* **1987**, 665–683. (b) Pereyre, M.; Quintard, J.-P.; Rahm, A. *Tin in Organic Synthesis*; Butterworth: London, 1987.

(4) (a) Smith, A. B., III; Hale, K. J.; McCauley, J. P., Jr. *Tetrahedron Lett.* **1989**, 30, 5579–5582. (b) Giovannini, R.; Petri, M. *Synlett* **1995**, 973–974.

(5) Parsons, A. F.; Pettifer, R. M. *Tetrahedron Lett.* **1996**, 37, 1667–1670.

(6) Antonio, Y.; De La Cruz, M. E.; Galeazzi, E.; Guzman, A.; Bray, B. L.; Greenhouse, R.; Kurz, L. J.; Lustig, D. A.; Maddox, M. L.; Muchowski, J. M. *Can. J. Chem.* **1994**, 72, 15–22.

(7) (a) Watanabe, Y.; Ueno, Y.; Araki, T.; Endo, T.; Okawara, M. *Tetrahedron Lett.* **1986**, 27, 215–218. (b) McCarthy, J. R.; Matthews, D. P.; Stermerick, D. M.; Huber, E. W.; Bey, P.; Lippert, B. J.; Snyder, R. D.; Sunkara, P. S. *J. Am. Chem. Soc.* **1991**, 113, 7439–7440. (c) Wnuk, S. F.; Yuan, C.-S.; Borchardt, R. T.; Balzarini, J.; De Clercq, E.; Robins, M. J. *J. Med. Chem.* **1994**, 37, 3579–3587.

(8) (a) Ueno, Y.; Aoki, S.; Okawara, M. *J. Am. Chem. Soc.* **1979**, 101, 5414–5415. (b) Bertrand, M. P. *Org. Prep. Proc. Int.* **1994**, 26, 257–290.

(9) (a) Rozen, S.; Filler, R. *Tetrahedron* **1985**, 41, 1111–1153. (b) Welch, J. T. *Tetrahedron* **1987**, 43, 3123–3197. (c) Resnati, G. *Tetrahedron* **1993**, 49, 9385–9445. (d) *Chemistry of Organic Fluorine Compounds II. A Critical Review*; Hudlicky, M., Pavlath, A. E., Eds.; ACS Monograph 187; American Chemical Society: Washington, DC, 1995. (e) Lal, G. S.; Pez, G. P.; Syvert, R. G. *Chem. Rev.* **1996**, 96, 1737–1755. (f) Taylor, S. D.; Kotoris, C. C.; Hum, G. *Tetrahedron* **1999**, 55, 12431–12477.


(10) (a) Bergmann, E. D.; Szinai, S. *J. Chem. Soc.* **1956**, 1521–1526. (b) Welch, J. T.; Plummer, J. S. *Synth. Commun.* **1989**, 19, 1081–1090.

(11) Middleton, W. J. *J. Org. Chem.* **1975**, 40, 574–578.

(12) (a) Differding, E.; Ofner, H. *Synlett* **1991**, 187–189. (b) Resnati, G.; DesMarteau, D. D. *J. Org. Chem.* **1991**, 56, 4925–4929. (c) Lal, G. S. *J. Org. Chem.* **1993**, 58, 2791–2796. (d) Davis, F. A.; Han, W.; Murphy, C. K. *J. Org. Chem.* **1995**, 60, 4730–4737. (e) Chambers, R. D.; Hutchinson, J. *J. Fluorine Chem.* **1998**, 92, 45–52.

(13) (a) Wang, Y.; Yang, Z.-Y.; Burton, D. J. *Tetrahedron Lett.* **1992**, 33, 2137–2140. (b) Zhi, C.; Chen Q.-Y. *J. Chem. Soc., Perkin Trans. 1* **1996**, 1741–1747.

(14) (a) Linderman, R. J.; Graves, D. M. *J. Org. Chem.* **1989**, 54, 661–668. (b) Katritzky, A. R.; Nichols, D. A.; Qi, M. *Tetrahedron Lett.* **1998**, 39, 7063–7066.

Scheme 1^a


^a Key: (a) Bu₃SnH(D)/AIBN/benzene or toluene/Δ; (b) Bu₃SnCl/PMHS/KF/H₂O/toluene/Δ; (c) KH/Selectfluor/THF/DMF; (d) 1c/RBr/NaH/DMF.

We now report convenient and efficient methodologies for synthesis of carboxylate α-pyrimidin-2-yl sulfones, their α-fluorination with Selectfluor, and their desulfonylation with tributylstannane or a "catalytic" tin equivalent. This provides a facile new route for the preparation of α-[²H], α-[²H₂], and α-fluoro-α-[²H] carbonyl compounds. Mechanistic considerations for this novel radical desulfonylation procedure are suggested.

Results and Discussion

The 2-(pyrimidin-2-ylsulfonyl) **1a–c** and 2-(pyridin-2-ylsulfonyl) **2a** esters were prepared from the corresponding ethyl 2-bromoalkanoates and pyrimidin- or pyridin-2-thiolates, followed by oxidation (*m*-CPBA) of the ethyl 2-(arylsulthio)alkanoates. Alkylation of ethyl 2-(pyrimidin-2-ylsulfonyl)acetate (**1c**) with the corresponding alkyl bromides gave sulfones **1d–h** (Scheme 1). Treatment of ethyl 2-(pyridin-2-ylsulfonyl)hexanoate (**2a**) with Bu₃SnH/AIBN/benzene at reflux for 36 h gave ethyl hexanoate (**3a**, 60%) plus unchanged **2a** and minor decomposition products. Analogous treatment of ethyl 2-(pyrimidin-2-ylsulfonyl)hexanoate (**1a**) gave complete conversion to **3a** within 1 h and in toluene desulfonylation was completed in 30 min. Parallel treatment of standard ethyl 2-(phenylsulfonyl)hexanoate (48 h) caused no observed change in the starting material. Other 2-(pyrimidin-2-ylsulfonyl)alkanoates (e.g., **1b,e–g**) also underwent clean desulfonylation with Bu₃SnH (~1.5–2.0 equiv) to give esters **3b,e–g** (81–91%).

π-Deficient heterocyclic sulfones also were found to be advantageous in reactions that involve generation of sulfonyl-stabilized carbanions (acidifying effects of pyridin- and pyrimidin-2-ylsulfonyl groups on α-carbon are greater than that of the phenylsulfonyl group). Thus, 2-(pyrimidin-2-ylsulfonyl) **1a,b,d–h** and 2-(pyridin-2-ylsulfonyl) **2a** esters were treated with potassium hydride, and the enolates were quenched with Selectfluor^{12c} to give ethyl 2-fluoro-2-(pyrimidin-2-ylsulfonyl)alkanoates (**5a,b,d–h**) and ethyl 2-fluoro-2-(pyridin-2-ylsulfonyl)hexanoate (**6a**) in good yields (72–92%). Tributylstannane-mediated desulfonylation of **5a** (1 h) and **6a** (28 h) gave ethyl 2-fluorohexanoate (**7a**; 95% and 60%, respectively). Treatment of α-fluoro-α-(pyrimidin-2-ylsulfonyl)

Scheme 2^a

^a Key: (a) KH/THF/D₂O; (b) Bu₃SnD/AIBN/benzene/Δ.

esters **5b,d–h** by this procedure gave α-fluoro esters **7b,d–h** (77–91%). Isolated double bonds, a carboxylate ester, and a silyl protected hydroxyl were tolerated under the fluorination and desulfonylation conditions.

It is noteworthy that an α-fluoro substituent has no effect on the time required and yield of the radical desulfonylation reactions in contrast to the impact of the second nitrogen atom in the heterocyclic ring [**1a/5a** (1 h, 95%) versus **2a/6a** (36/28 h, 60%)]. Although removal of the pyridin-2-ylsulfonyl group is less efficient, easy access to the pyridin-2-yl thioethers with the radical-stabilizing group at α-carbon (via Barton's thiohydroamic ester chemistry)^{15,16} enhances the versatility of our mild radical-mediated removal of the pyrimidin-2-ylsulfonyl group. In an attempt to facilitate removal of the pyrimidin-2-ylsulfonyl group, compound **2a** was oxidized to the *N*-oxide. However, treatment of the latter with Bu₃SnH (2 equiv, 1 h) gave **2a** (~80%) plus **3a** (~12%). Deoxygenation of *N*-oxides with tin reagents (including Bu₃SnH)^{17a} is known.¹⁷

Tributylstannane-mediated desulfonylation also gives access to deuterium-labeled¹⁸ esters. Thus, treatment of **1a** and **5a** with Bu₃SnD gave ethyl 2-deuteriohexanoate (**4a**, ~95% [²H]) and 2-fluoro-2-deuteriohexanoate (**8a**, ~90% [²H]), respectively. Quenching enolates derived from **1a** and **1b** with D₂O yielded α-deuterated sulfone **9a** and **9b** (~90% [²H]) which upon treatment with Bu₃SnD gave ethyl 2,2-dideuterioalkanoates **10a** (~80% [²H₂]) and **10b** (~85% [²H₂]), respectively (Scheme 2).

Disadvantages associated with the use of Bu₃SnH are toxicity¹⁹ and purification²⁰ of organotin species. To alleviate these problems, processes that are "catalyzed" by Bu₃SnH have been developed. One approach utilizes the ability of borohydrides to reduce tributyltin halides,^{4b,21} and Fu's procedures exploit silicon hydride reduction of species with a Sn–O (or Sn–N) bond to regenerate Bu₃SnH.²² (TMS)₃SiH also can serve as a substitute for Bu₃SnH in a number of radical-mediated processes.²³ Recently, in situ generation of tin hydride by treatment

(15) (a) Barton, D. H. R.; Crich, D.; Motherwell, W. B. *J. Chem. Soc., Chem. Commun.* **1983**, 939–941. (b) Barton, D. H. R.; Géro, S. D.; Quiclet-Sire, B.; Samadi, M. *Tetrahedron* **1992**, *48*, 1627–1636. (c) Barton, D. H. R. *Pure Appl. Chem.* **1994**, *66*, 1943–1954.

(16) Boivin, J.; Lallemand, J.-Y.; Schmitt, A.; Zard, S. Z. *Tetrahedron Lett.* **1995**, *36*, 7243–7246.


(17) (a) Kozuka, S.; Akasaka, T.; Furumai, S.; Oae, S. *Chem. Ind. (London)* **1974**, 452–453. (b) Jousseume, B.; Chamson, E. *Synthesis* **1987**, 55–56.

(18) Thomas, A. F. *Deuterium Labeling in Organic Chemistry*; Appleton-Century-Crofts: New York, 1971.

(19) Boyer, I. *Toxicology* **1989**, *55*, 253–298.

(20) (a) Crich, D.; Sun, S. *J. Org. Chem.* **1996**, *61*, 7200–7201. (b) Curran, D. P.; Hadida, S.; Kim, S.-Y.; Luo, Z. *J. Am. Chem. Soc.* **1999**, *121*, 6607–6615.

(21) (a) Corey, E. J.; Suggs, J. W. *J. Org. Chem.* **1975**, *40*, 2554–2555. (b) Stork, G.; Sher, P. M. *J. Am. Chem. Soc.* **1986**, *108*, 303–304.

Scheme 3^a

^a Key: (a) $\text{Bu}_3\text{SnH(D)}/\text{AIBN}/\text{benzene}/\Delta$; (b) $\text{Bu}_3\text{SnCl}/\text{PMHS}/\text{KF}/\text{H}_2\text{O}/\text{toluene}/\Delta$.

of catalytic Bu_3SnCl with excess polymethylhydrosiloxane (PMHS) in the presence of potassium fluoride has been reported.²⁴

To reduce the amount of tributyltin hydride, a modification of our stannyl radical-mediated desulfonation has been developed. Treatment of the **1a** or **1b** with catalytic Bu_3SnH or Bu_3SnCl (0.2 equiv) and excess $\text{Bu}_4\text{NBH}_3\text{CN}$ (benzene/AIBN/ Δ) did not effect desulfonation in good yields. The amounts of products formed (**3a** or **3b**) were proportional to the quantities of Bu_3SnH used. Treatment of **1a** and **5a** with Ph_3SiH or $(\text{TMS})_3\text{SiH}$ [benzene/AIBN or $(\text{BzO})_2/\Delta$] also resulted in recovery of unchanged sulfones (80–90%). However, treatment of **1b** with another “catalytic” tin hydride system [Bu_3SnCl (0.15 equiv.)/PMHS/KF/ $\text{H}_2\text{O}/\text{toluene}/\Delta$]²⁴ effected hydrogenolysis to give **3b** (85%) which was readily purified. Analogous treatment of 2-(pyrimidin-2-ylsulfonyl) **1d,f** and 2-fluoro-2-(pyrimidin-2-ylsulfonyl) **5b,f** resulted in smooth desulfonation to give esters **3d,f** and α -fluoro esters **7b,f** (84–89%), respectively.

Possible reaction mechanisms might involve formation of either stabilized α -carboxylate or alkoxy ketyl-type intermediate radicals. The absence of 5-*exo*-trig ring-closure during radical desulfonation of ethyl 2-(pyrimidin-2-sulfonyl)hept-6-enoate (**1f**) to **3f** [Bu_3SnH (84% isolated yield), $\text{Bu}_3\text{SnCl}/\text{PMHS}$ (89%)] argues against an α -carboxylate radical intermediate. Similarly, α -fluoro analogue **5f** produced **7f** (84–91%; ^{19}F NMR of the crude reaction mixture showed peaks only for **7f**). Attack by the tin radical on the carbonyl oxygen of **1** (or **5**) would generate a ketyl-type radical **11**, and β -elimination of a sulfonyl radical would produce the tin enolate **12** (Scheme 3). In a propagation step, hydrogen (deuterium)

transfer^{8b,25} from tributyltin hydride or deuteride to the sulfonyl radical would give the pyrimidin-2-ylsulfinic acid **13** (isolated from the reaction mixture). Protonation of **12** by **13** would yield products **3/4** (or **7/8**). Desulfonylation of β -ketosulfones⁴ and *N*-sulfonylated amides⁵ with Bu_3SnH have been proposed to proceed via analogous ketyl-type radicals and tin-enolates. The ability of PMHS to reduce oxygen–tin bonds is well documented.^{22b,c,24} The fact that ethyl 6-(pyrimidin-2-ylsulfonyl)hexanoate (**14**) did not undergo the radical-mediated desulfonation is consistent with the proposed mechanism. A further possibility might involve single electron transfer from the tin radical to the ester group, again leading to enolate formation.

Reductive cyclization of ketyl radicals arising from δ,ϵ -unsaturated ketones and aldehydes (as well esters having an auxiliary α -hydroxymethyl group)^{26a} are known to give the corresponding substituted cyclopentanols.^{26b} Intramolecular 1,5-cyclization of 5-hexenoyl chloride to 2-methylcyclopentanone^{27a} and reductive cyclization of unsaturated aldehydes and ketones with Bu_3SnH have been reported.^{27b,c} However, desulfonylation of hex-5-enoates **1e** and **5e** occurred without observed formation of cyclization products (**3e** and **7e** were isolated in 81% yields). Apparently, β -elimination of a sulfonyl radical is much faster than intramolecular cyclization involving *O*-stannyl ketyl radicals of type **11**.

In summary, we have developed convenient and efficient methodologies for synthesis of heterocyclic α -sulfones of carboxylate esters, their α -fluorination with Selectfluor, and their desulfonation with tributylstannane or catalytic tin reagents in the presence of polymethylhydrosiloxane. This provides a facile new route for the preparation of α -[^2H], α -[$^2\text{H}_2$], and α -fluoro- α -[^2H] esters. Desulfonylation is suggested to proceed via alkoxy ketyl-type radicals and tin enolates.

Experimental Section

Uncorrected melting points were determined with a capillary tube apparatus. ^1H NMR spectra were determined with solution in CDCl_3 at 200 or 400 MHz, ^{13}C at 100.6 MHz, and ^{19}F (CCl_3F) at 376.4 MHz unless otherwise noted. Mass spectra (MS) were obtained by electron impact (EI), atmospheric pressure chemical ionization (APCI) or CI (CH_4), or fast atom bombardment (FAB, 5% trifluoroacetic acid/thioglycerol matrix) techniques. Reagent-grade chemicals were used, and solvents were dried by reflux over and distillation from CaH_2 under an argon atmosphere. Selectfluor fluorinating reagent (>95% active [F^+]) was purchased from Aldrich. TLC was performed on Merck kieselgel 60-F₂₅₄ with $\text{MeOH}/\text{CHCl}_3$ (1:19) and $\text{EtOAc}/\text{hexane}$ (1:2) as developing systems, and products were detected with 254 nm light or by development of color with I_2 . Merck kieselgel 60 (230–400 mesh) was used for column chromatography. Elemental analyses were determined by Galbraith Laboratories, Knoxville, TN. Purity and identity of the products (crude and/or purified) were established using GC/MS (EI) system [capillary column (30 m \times 0.25 mm \times 25 μm), program: 40 $^\circ\text{C}$ for 1 min with increase

(22) (a) Hays, D. V.; Scholl, M.; Fu, G. C. *J. Org. Chem.* **1996**, *61*, 6751–6752. (b) Lopez, R. M.; Hays, D. V.; Fu, G. C. *J. Am. Chem. Soc.* **1997**, *119*, 6949–6950. (c) Hays, D. V.; Fu, G. C. *J. Org. Chem.* **1998**, *63*, 2796–2797. (d) Tormo, J.; Hays, D. V.; Fu, G. C. *J. Org. Chem.* **1998**, *63*, 5296–5297.

(23) Chatgililoglu, C. *Acc. Chem. Res.* **1992**, *25*, 188–194.

(24) Terstiege, I.; Maleczka, R. E., Jr. *J. Org. Chem.* **1999**, *64*, 342–343.

(25) (a) Culshaw, P. N.; Walton, J. C. *J. Chem. Soc., Perkin Trans. 2* **1991**, 1201–1208. (b) Crich, D. In *Organosulfur Chemistry: Synthetic Aspects*; Page, P., Ed.; Academic Press: London, 1995; pp 49–88.

(26) (a) Cossy, J.; Gille, B.; Bellosta, V. *J. Org. Chem.* **1998**, *63*, 3141–3146. (a) Jasperse, C. P.; Curran, D. P.; Fevig, T. L. *Chem. Rev.* **1991**, *91*, 1237–1286.

(27) (a) Walsh, E. J., Jr.; Messinger, J. M., II; Grudowski, D. A.; Allchin, C. A. *Tetrahedron Lett.* **1980**, *21*, 4409–4412. (b) Ardisson, J.; Ferezou, J. P.; Julia, M.; Pancrazi, A. *Tetrahedron Lett.* **1987**, *28*, 2001–2004. (c) Sugawara, T.; Otter, B. A.; Ueda, T. *Tetrahedron Lett.* **1988**, *29*, 75–78.

15 °C/min to 300 °C]. Typical t_R for the series of octanoate compounds are as follow: **1b** (16.8 min), **3b** (7.6 min), **5b** (16.7 min), and **7b** (7.9 min). CAUTION! All procedures involving benzene and tributyltin hydride should be carried out in a well-ventilated hood.

Ethyl 2-(Pyrimidin-2-ylsulfonyl)hexanoate (1a). **Procedure A.** NaH (1.01 g, 50%/mineral oil, 21 mmol) was washed (dried Et₂O, 1 × 25 mL) and suspended in dried DMF (35 mL) under N₂. Two equal portions of 2-mercaptopyrimidine (2.24 g, 20 mmol) were added slowly at ~0 °C (ice bath). The resulting solution was stirred at ambient temperature for 1 h and cooled to ~0 °C, and ethyl 2-bromohexanoate (3.65 mL, 4.46 g, 20 mmol) was added. After 1 h, the mixture was allowed to warm to ambient temperature, stirred overnight, and evaporated, and the residue was partitioned (EtOAc/H₂O). The organic layer was washed (NaHCO₃/H₂O; brine), dried (MgSO₄), and evaporated to give the viscous thioether (4.83 g, 95%) that was dissolved (CHCl₃, 50 mL), cooled (~-20 °C), and treated dropwise with *m*-CPBA (9.66 g/75% reagent, 42 mmol) in CHCl₃/CH₂Cl₂ (1:1, 100 mL). After 2 h, the mixture was allowed to warm to ambient temperature and stirred for 18 h. Saturated NaHCO₃/H₂O (100 mL) was added, stirring was continued for 30 min, the organic layer was separated, and the aqueous layer was extracted (CHCl₃, 50 mL). The combined organic phase was washed (NaHCO₃/H₂O; brine), dried (MgSO₄), evaporated, and chromatographed (50% hexanes/EtOAc → EtOAc) to give **1a** (5.15 g, 90% overall) as a solidified oil: mp 50–51 °C; ¹H NMR δ 0.90 (t, *J* = 6.6 Hz, 3H), 1.09 (t, *J* = 7.1 Hz, 3H), 1.28–1.51 (m, 4H), 2.16–2.28 (m, 2H), 4.10 (q, *J* = 7.1 Hz, 2H), 4.61 (dd, *J* = 6.2, 8.7 Hz, 1H), 7.60 (t, *J* = 4.9 Hz, 1H), 8.97 (d, *J* = 4.9 Hz, 2H); ¹³C NMR δ 13.8, 13.9, 22.4, 24.8, 29.3, 62.4, 66.1, 124.1, 158.8, 165.3; IR (Nujol) 2924, 1731, 1556, 1456, 1316, 1113 cm⁻¹; MS (CI) *m/z* 287 (100, MH⁺). Anal. Calcd for C₁₂H₁₈N₂O₄S (286.35): C, 50.33; H, 6.34; N, 9.78. Found: C, 50.33; H, 6.15; N, 9.60.

Ethyl 2-(Pyrimidin-2-ylsulfonyl)acetate (1c). The 2-mercaptopyrimidine (1.12 g, 10 mmol) was added to a solution of EtONa in EtOH [Na (330 mg, 14 mmol) in EtOH (30 mL)], and the resulting solution was stirred at ambient temperature for 1 h. Ethyl 2-bromoacetate (1.11 mL, 1.67 g, 10 mmol) was added, stirring was continued overnight, and volatiles were evaporated. The residue was subjected to the remaining part of procedure A to give **1c** (1.95 g, 84%) as a white solid: mp 40–41 °C; ¹H NMR δ 1.14 (t, *J* = 7.2 Hz, 3H), 4.12 (q, *J* = 7.2 Hz, 2H), 4.58 (s, 2H), 7.60 (t, *J* = 5.2 Hz, 1H), 8.96 (d, *J* = 4.9 Hz, 2H); ¹³C NMR δ 14.2, 56.1, 62.7, 124.9, 159.2, 162.7, 165.0; MS (FAB) *m/z* 231 (100, MH⁺). Anal. Calcd for C₈H₁₀N₂O₄S (230.25): C, 41.73; H, 4.38; N, 12.17. Found: C, 41.69; H, 4.43; N, 12.10.

Diethyl 2-(Pyrimidin-2-ylsulfonyl)octanedioate (1d). **Procedure B.** NaH (264 mg, 50%/mineral oil, 5.5 mmol) was washed (dried Et₂O, 1 × 25 mL) and suspended in dried DMF (20 mL) under N₂. Compound **1c** (1.15 g, 5 mmol) was added slowly at ~0 °C (ice bath), and the resulting solution was stirred at ambient temperature for 30 min. Ethyl 6-bromohexanoate (0.98 mL, 1.23 g, 5.5 mmol) was added (syringe), and after being stirred for 16 h the mixture was heated for 2 h at 50 °C. Volatiles were evaporated, and the residue was partitioned (EtOAc/NH₄Cl/H₂O). The organic layer was washed (NaHCO₃/H₂O; brine), dried (MgSO₄), evaporated, and chromatographed (40% hexanes/EtOAc → EtOAc) to give **1d** (1.32 g, 71%) as an oil: ¹H NMR δ 1.04 (t, *J* = 7.1 Hz, 3H), 1.19 (t, *J* = 7.3 Hz, 3H), 1.30–1.62 (m, 6H), 2.11–2.20 (m, 2H), 2.24 (t, *J* = 7.4 Hz, 2H), 4.02–4.09 (m, 4H), 4.59 (dd, *J* = 3.9, 9.6 Hz, 1H), 7.60 (t, *J* = 4.8 Hz, 1H), 8.93 (d, *J* = 4.9 Hz, 2H); ¹³C NMR δ 14.1, 14.6, 24.8, 25.1, 27.1, 28.9, 34.4, 60.6, 62.7, 66.1, 124.5, 159.1, 165.4, 165.5, 173.9; IR (neat) 2938, 1734, 1566 cm⁻¹; MS (EI) *m/z* 327 (30, M⁺ - OEt), 185 (100). Anal. Calcd for C₁₆H₂₄N₂O₆S (372.45): C, 51.60; H, 6.50; N, 7.52. Found: C, 51.87; H, 6.46; N, 7.72.

Ethyl 2-(Pyrimidin-2-ylsulfonyl)hept-6-enoate (1f). Treatment of **1c** (1.15 g, 5 mmol) with 5-bromo-1-pentene (0.77 mL, 968 mg, 6.5 mmol) by procedure B gave **1f** (1.22 g, 82%) as a solidified oil: ¹H NMR δ 1.06 (t, *J* = 7.2 Hz, 3H), 1.54 (quint, *J* = 7.7 Hz, 2H), 2.04–2.26 (m, 4H), 4.08 (q, *J* = 7.2

Hz, 2H), 4.62 (dd, *J* = 6.2, 8.7 Hz, 1H), 4.93 (dm, *J* = 9.9 Hz, 1H), 5.00 (dm, *J* = 16.5 Hz, 1H), 5.73 (ddt, *J* = 16.9, 10.1, 6.5 Hz, 1H), 7.62 (t, *J* = 5.1 Hz, 1H), 8.96 (d, *J* = 5.1 Hz, 2H); ¹³C NMR δ 14.2, 24.8, 26.7, 33.6, 62.8, 66.3, 116.0, 124.3, 137.8, 159.0, 165.5, 165.6; MS (FAB) *m/z* 299 (100, MH⁺). Anal. Calcd for C₁₃H₁₈N₂O₄S (298.37): C, 52.33; H, 6.08; N, 9.39. Found: C, 52.71; H, 6.41; N, 9.08.

Ethyl 2-(Pyridin-2-ylsulfonyl)hexanoate (2a). Treatment of ethyl 2-bromohexanoate (3.65 mL, 4.46 g, 20 mmol) with 2-mercaptopyridine (2.22 g, 20 mmol) and oxidation (20 h) with *m*-CPBA (9.66 g/75% reagent, 42 mmol) by procedure A gave **2a** (5.07 g, 89%; oil): ¹H NMR δ 0.85–0.93 (m, 3H), 1.11 (t, *J* = 7.1 Hz, 3H), 1.31–1.45 (m, 4H), 2.10–2.21 (m, 2H), 4.21 (q, *J* = 7.1 Hz, 2H), 4.46 (dd, *J* = 5.5, 9.4 Hz, 1H), 7.58 (ddd, *J* = 1.3, 4.7, 7.6 Hz, 1H), 7.97 (td, *J* = 7.7, 1.7 Hz, 1H), 8.08 (dt, *J* = 7.8, 1.3 Hz, 1H), 8.78 (ddd, *J* = 1.3, 1.7, 4.8 Hz, 1H); ¹³C NMR δ 14.11, 14.12, 22.6, 25.6, 29.5, 62.6, 67.2, 123.8, 128.2, 138.6, 150.7, 156.8, 165.9; IR (neat) 2961, 1740, 1580, 1325, 1164 cm⁻¹; MS (CI) *m/z* 286 (100, MH⁺). Anal. Calcd for C₁₃H₁₉NO₄S (285.36): C, 54.72; H, 6.71; N, 4.91. Found: C, 54.63; H, 6.52; N 5.09.

Treatment of **2a** (285 mg, 1 mmol) with *m*-CPBA (460 mg/75% reagent, 2 mmol) in CHCl₃/CH₂Cl₂ (1:1, 15 mL) at ambient temperature for 72 h followed by gently heating at reflux for 3 h and workup (procedure A) gave **2a** *N*-oxide (184 mg, 61%) and unchanged **2a** (88 mg, 31%). ¹H NMR spectrum of **2a**-*N*-oxide show significant downfield shift for the proton at C2: δ 5.31 (dd, *J* = 4.3, 9.6 Hz, 1H).

Ethyl hexanoate (3a). **Procedure C.** Argon was bubbled through a solution of **1a** (286 mg, 1.0 mmol) in benzene (5.0 mL) for 30 min, and Bu₃SnH (0.467 mL, 509 mg, 1.75 mmol) was added. Deoxygenation was continued for 5 min, AIBN (25 mg, 0.15 mmol) was added, and the solution was refluxed for 1 h (TLC showed complete conversion of **1a**). Volatiles were evaporated (<25 °C, ~20 mmHg), and the residue was dissolved (EtOAc, 5 mL). The solution was stirred overnight with KF/H₂O (30 mg/0.3 mL), evaporated, and chromatographed (pentane → 3% EtOAc/pentane) to give **3a** (137 mg, 95%) with spectral data identical to those of an authentic sample.²⁸ Evaporation of the reaction mixture and direct column chromatography [hexane (100 mL) → 5% EtOAc/hexane] also gave **3a** with similar yield and purity. A ratio of 1.35 equiv. of Bu₃SnH also gave essentially quantitative conversion of **1a** to **3a** within 1 h.

Treatment of **2a** (285 mg, 1 mmol) with Bu₃SnH (1.65 mmol) by procedure C (30 min) using toluene instead of benzene gave **3a** (131 mg, 91%).

Analogous treatment (36 h, benzene) of **2a** (285 mg, 1 mmol) with Bu₃SnH (2.0 mmol) and AIBN (0.2 mmol) [additional Bu₃SnH (1.0 mmol) and AIBN (0.2 mmol) after 15h] by procedure C gave **3a** [87 mg, 60%; further elution (EtOAc/pentane, 1:1) gave recovered **2a** (85 mg, 30%)]. Treatment of **2a** *N*-oxide (90 mg, 0.3 mmol) by procedure C [1h, Bu₃SnH (2 equiv)] gave **2a** (68 mg, 80%) and **3a** (5 mg, 12%).

Treatment of ethyl 2-(phenylsulfonyl)hexanoate [prepared by procedure A (thiophenol, 10 mmol; 2.41 g, 85%) with data as described]²⁹ by procedure C (48 h, benzene) gave recovered starting material (~95%).

Ethyl Octanoate (3b). **Procedure D.** Nitrogen was bubbled through a solution of **1b** (314 mg, 1 mmol), Bu₃SnCl (49 mg, 0.041 mL, 0.15 mmol), and AIBN (5 mg, 0.03 mmol) in toluene (2 mL) for 15 min. The solution was heated at reflux for 3 h, and PMHS (0.2 mL) and KF [116 mg (2 mmol) in H₂O (0.5 mL)] were added in three equal portions immediately after reaching the boiling point and after 1 and 2 h. Volatiles were evaporated, and the residue was partitioned (EtOAc/NaHCO₃/H₂O). The organic layer was washed (brine), dried (MgSO₄), evaporated, and chromatographed [hexane (100 mL) → 10% EtOAc/hexane] to give **3b** [(147 mg, 85%); GC/MS purity (>99%); MS *m/z* 172 (10, M⁺), 127 (42, M⁺ - OEt), 88 (100)] with spectral data identical to those of an authentic sample.²⁸

(28) *The Aldrich Library of ¹³C and ¹H FT-NMR Spectra*; Pouchert, C. J., Behnke, J., Eds.; Aldrich Chemical: Milwaukee, WI, 1993.

(29) Wang, Y.; Jiang, Y. *Synth. Commun.* **1992**, *22*, 2287–2291.

Further elution (EtOAc/hexane, 1:1) gave recovered **1b** (31 mg, 10%). Evaporation of the reaction mixture and direct column chromatography also gave **3b** with similar yield and purity.

Treatment of **1b** (157 mg, 0.5 mmol) by procedure C gave **3b** (78 mg, 91%).

Diethyl Octanedioate (Diethyl Suberate) (3d). Treatment of **1d** (186 mg, 0.5 mmol) by procedure D gave **3d** (101 mg, 88%) with spectra identical to those of an authentic sample:²⁸ GC/MS purity (>99%); MS m/z 185 (80, $M^+ - OEt$), 143 (100).

Ethyl Hept-6-enoate (3f). Treatment of **1f** (170 mg, 0.57 mmol) with Bu_3SnH (1.14 mmol) by procedure C (chromatography: pentane \rightarrow 10% EtOAc/pentane) gave **3f**³⁰ (75 mg, 84%) contaminated (~5–10%, 1H NMR) by tin compound(s). Hydrolysis (NaOH/ H_2O /MeOH) of this material gave 6-heptenoic acid with spectra identical to those of authentic sample.²⁸

Analogous treatment of **1f** by procedure C in C_6D_6 and direct analysis of the aliquots (0.25 h, 0.5 h, 1 h) by 1H NMR (C_6D_6) showed that integration of the characteristic signals for the vinylic protons at δ 5.54–5.75 (m, 1, H₆) and 4.90–5.02 (m, 3, H_{2,7,7'}) remained constant ($\pm 5\%$) compared with the signals for the methylene protons at δ 3.88 (**1f**) and 4.01 (**3f**) from the ester groups. Washing of the C_6D_6 solution with $D_2O/NaHCO_3$ and 1H NMR of D_2O layer showed only signals for **13**.

Treatment of **1f** (149 mg, 0.5 mmol) by procedure D also gave **3f** (69 mg, 89%); GC/MS purity (99%); MS m/z 156 (5), 88 (100).

Ethyl 2-Deuteriohexanoate (4a). Treatment of **1a** (143 mg, 0.5 mmol) with Bu_3SnD (0.268 mL, 291 mg, 1 mmol) by procedure C (2 h) gave **4a** (65 mg, 90%); 1H NMR spectra corresponded with those of **3a** with 50% reduction in the intensity of signals²⁸ at δ 2.26 (t, $J = 7.7$ Hz, 1H, 2-CHD); MS (CI) m/z 146 (100, MH^+ [$C_8H_{16}DO_2$]).

Ethyl 2-Fluoro-2-(pyrimidin-2-ylsulfonyl)hexanoate (5a). Procedure E. KH [(571 mg, 35%/mineral oil, 5 mmol) or (220 mg, 5.5 mmol, dried/pressed between filter paper)] in a flame-dried flask under Ar was washed (dried hexane, dried Et_2O), and dried THF (25 mL) was added. The suspension was cooled (~0 °C, ice bath), and compound **1a** (1.14 g, 4 mmol) in dried THF (15 mL) was added (syringe). The solution was stirred (0 °C for 15 min, ambient temperature for 60 min, cooled to 0 °C), and Selectfluor (2.13 g, 6 mmol) was added in one portion. After 15 min, dried DMF (15 mL) was added (syringe), the ice bath was removed after 5 min, and stirring was continued at ambient temperature for 2 h. The reaction mixture was cooled to ~0 °C (ice bath), and $CHCl_3$ (30 mL) and saturated NH_4Cl/H_2O (15 mL) were slowly added. The organic layer was separated after 5 min, and the aqueous layer was extracted ($CHCl_3$, 2 \times 25 mL). The combined organic phase was washed (saturated $NaHCO_3/H_2O$, brine), dried ($MgSO_4$), evaporated, and chromatographed (50 \rightarrow 90% EtOAc/hexanes) to give **5a** (1.12 g, 92%) as a slightly yellow solidified oil: 1H NMR δ 0.91 (t, $J = 6.6$ Hz, 3H), 1.30 (t, $J = 7.1$ Hz, 3H), 1.25–1.60 (m, 4H), 2.43–2.72 (m, 2H), 4.35 (q, $J = 7.1$ Hz, 2H), 7.62 (t, $J = 4.9$ Hz, 1H), 8.98 (d, $J = 4.9$ Hz, 2H); ^{13}C NMR δ 14.1, 14.4, 22.8, 25.1, 30.9 (d, $^2J = 19.7$ Hz), 63.9, 107.6 (d, $^1J = 232.8$ Hz), 125.0, 159.3, 163.7, 163.9 (d, $^2J = 25.2$ Hz); ^{19}F NMR δ -158.8 (dd, $^3J_{F-3a} = 10.9$ Hz, $^3J_{F-3b} = 37.8$ Hz); IR (neat) 2963, 1751, 1567, 1347 cm^{-1} ; MS (EI) m/z 304 (42, M^+), 197 (50), 79 (100). Anal. Calcd for $C_{12}H_{17}FN_2O_4S$ (304.3): C, 47.36; H, 5.63; N, 9.20. Found: C, 47.57; H, 5.72; N, 9.19.

Diethyl 2-Fluoro-2-(pyrimidin-2-ylsulfonyl)octanedioate (5d). Treatment of **1d** (186 mg, 0.5 mmol) by procedure E gave **5d** (166 mg, 85%); 1H NMR δ 1.22 (t, $J = 7.1$ Hz, 3H), 1.30 (t, $J = 7.2$ Hz, 3H), 1.40 (quint, $J = 7.3$ Hz, 2H), 1.52–1.66 (m, 4H), 2.26 (t, $J = 7.4$ Hz, 2H), 2.48–2.66 (m, 2H), 4.08 (q, $J = 7.1$ Hz, 2H), 4.34 (q, $J = 7.1$ Hz, 2H), 7.65 (t, $J = 4.8$ Hz, 1H), 8.95 (d, $J = 4.9$ Hz, 2H); ^{13}C NMR δ 14.3, 14.6, 22.6 (d, $^3J = 2.0$ Hz), 24.8, 28.9, 30.9 (d, $^2J = 19.8$ Hz), 34.3, 60.7, 63.9, 107.3 (d, $^1J = 234.4$ Hz), 125.1, 159.3, 163.4, 163.7 (d, $^2J = 25.1$ Hz), 173.8; ^{19}F NMR δ -158.9 (dd, $^3J_{F-3a} = 11.3$ Hz,

$^3J_{F-3b} = 37.6$ Hz); MS (APCI) m/z 391 (100, MH^+). Anal. Calcd for $C_{16}H_{23}FN_2O_6S$ (390.44): C, 49.22; H, 5.94; N, 7.18. Found: C, 48.88; H, 6.30; N, 6.84.

Ethyl 2-Fluoro-2-(pyrimidin-2-ylsulfonyl)hept-6-enoate (5f). Treatment of **1f** (596 mg, 2.0 mmol) by procedure E gave **5f** (556 mg, 88%) as an oil: 1H NMR δ 1.28 (t, $J = 7.1$ Hz, 3H), 1.35–1.68 (m, 2H), 2.08–2.18 (m, 2H), 2.45–2.68 (m, 2H), 4.32 (q, $J = 7.1$ Hz, 2H), 4.95 (dm, $J = 9.9$ Hz, 1H), 5.01 (dm, $J = 16.6$ Hz, 1H), 5.65–5.75 (m, 1H), 7.65 (t, $J = 4.8$ Hz, 1H), 8.98 (d, $J = 4.9$ Hz, 2H); ^{13}C NMR δ 14.4, 22.3 ($^3J = 2.0$ Hz), 30.5 (d, $^2J = 19.8$ Hz), 33.4, 63.9, 107.4 (d, $^1J = 234.2$ Hz), 116.4, 124.9, 137.3, 159.2, 163.5, 163.8 (d, $^2J = 25.3$ Hz); ^{19}F NMR δ -158.6 (dd, $^3J_{F-3a} = 11.9$ Hz, $^3J_{F-3b} = 36.9$ Hz); MS (APCI) m/z 317 (100, MH^+). Anal. Calcd for $C_{13}H_{17}FN_2O_4S$ (316.36): C, 49.36; H, 5.42; N, 8.86. Found: C, 49.39; H, 5.51; N, 8.80.

Ethyl 2-Fluoro-2-(pyridin-2-ylsulfonyl)hexanoate (6a). Treatment of **2a** (428 mg, 1.5 mmol) with KH (2.0 mmol) and Selectfluor (2.5 mmol) (2 h) by procedure E gave **6a** (414 mg, 91%; viscous oil): 1H NMR δ 0.90 (t, $J = 6.8$ Hz, 3H), 1.16–1.52 (m, 7H), 2.30–2.73 (m, 2H), 4.31 (q, $J = 7.2$ Hz, 2H), 7.60 (ddd, $J = 1.4$, 4.7, 7.6 Hz, 1H), 7.98 (dt, $J = 1.7$, 7.6 Hz, 1), 8.09 (dt, $J = 1.1$ Hz, 7.8 Hz, 1), 8.73 (ddd, $J = 1.0$, 1.7, 4.8 Hz, 1); ^{13}C NMR δ 14.2, 14.4, 22.8, 25.1 (d, $^3J = 2.0$ Hz), 30.7 (d, $^2J = 19.7$ Hz), 63.9, 107.5 (d, $^1J = 231.4$ Hz), 126.32, 128.8, 138.68, 150.78, 154.2, 164.0 (d, $^2J = 24.1$ Hz); ^{19}F NMR δ -159.3 (dd, $^3J_{F-3a} = 10.0$ Hz, $^3J_{F-3b} = 38.7$ Hz); IR (neat) 2964, 1758, 1353, 1266, 1174, 734 cm^{-1} ; MS (CI) m/z 304 (100, MH^+). Anal. Calcd for $C_{13}H_{18}FNO_4S$ (303.3): C, 51.47; H 5.98; N, 4.62. Found: C, 51.39; H, 6.12; N, 4.51.

Ethyl 2-Fluorohexanoate (7a). Treatment of **5a** (304 mg, 1 mmol) with Bu_3SnH (1.75 mmol) by procedure C gave **7a** (154 mg, 95%; oil) with data as reported.^{31a}

Analogous treatment of **6a** (303 mg, 1 mmol) with Bu_3SnH (2.0 mmol) and AIBN (0.2 mmol) (28 h) [additional Bu_3SnH (1.0 mmol) and AIBN (0.2 mmol) were added after 14 h] gave **7a** (97 mg, 60%). Further elution (EtOAc/pentane, 1:1) gave recovered **5b** (61 mg, 20%).

Treatment of **5a** (1 mmol) with Bu_3SnH (1.3 mmol) gave **7a** (85%) [TLC: **5a** (~5–10%)].

Ethyl 2-Fluorooctanoate (7b). Treatment of **5b** (664 mg, 2 mmol) with Bu_3SnH (3.5 mmol) by procedure C (chromatography: hexane \rightarrow 10% EtOAc/hexane) gave **7b** (334 mg, 88%; oil) with data as reported:¹³ GC/MS purity (99%); MS m/z 190 (5, M^+), 106 (100).

Treatment of **5b** (332 mg, 1 mmol) by procedure D gave **7b** [163 mg, 86%; GC (>99%)].

Diethyl 2-Fluorooctanedioate (7d). Treatment of **5d** (195 mg, 0.5 mmol) by procedure C [chromatography (hexane \rightarrow 10% EtOAc/hexane)] gave **7d** (109 mg, 88%); 1H NMR δ 1.25 (t, $J = 7.1$ Hz, 3H), 1.31 (t, $J = 7.2$ Hz, 3H), 1.35–1.53 (m, 4H), 1.65 (quint, $J = 7.5$ Hz, 2H), 1.82–1.95 (m, 2H), 2.25 (t, $J = 7.4$ Hz, 2H), 4.12 (q, $J = 7.2$ Hz, 2H), 4.26 (q, $J = 7.1$ Hz, 2H), 4.89 (dt, $J = 49.4$, 5.9 Hz, 1H); ^{13}C NMR δ 14.5, 14.6, 24.4 (d, $^3J = 2.9$ Hz), 25.0, 28.9, 32.6 (d, $^2J = 21.1$ Hz), 34.5, 60.7, 61.8, 89.3 (d, $^1J = 183.9$ Hz), 170.4 (d, $^2J = 23.6$ Hz), 174.0; ^{19}F NMR δ -192.6 (dt, $^2J_{F-2} = 49.0$ Hz, $^3J_{F-3a,b} = 25.0$ Hz); GC/MS purity (>99%); MS m/z 203 (45, $M^+ - OEt$), 88 (100). Anal. Calcd for $C_{12}H_{21}FO_4$ (248.30): C, 58.05; H, 8.53. Found: C, 58.25; H, 8.75.

Ethyl 2-Fluorohept-6-enoate (7f). Treatment of **5f** (158 mg, 0.5 mmol) with Bu_3SnH (1 mmol) by procedure C [chromatography (hexane \rightarrow 10% EtOAc/hexane)] gave **7f** (79 mg, 91%); 1H NMR δ 1.32 (t, $J = 7.1$ Hz, 3H), 1.58 (“quint” $J = 8.2$ Hz, 2H), 1.85–1.96 (m, 2H), 2.11 (“q”, $J = 6.7$ Hz, 2H), 4.29 (q, $J = 7.1$ Hz, 2H), 4.90 (dt, $J = 49.5$, 5.9 Hz, 1H), 4.98 (dm, $J = 9.9$ Hz, 1H), 5.03 (dm, $J = 16.7$ Hz, 1H), 5.79 (ddt, $J = 16.9$, 10.1, 6.6 Hz, 1H); ^{13}C NMR δ 14.6, 23.9 (d, $^3J = 2.8$ Hz), 32.1 (d, $^2J = 21.1$ Hz), 33.5, 61.9, 89.3 (d, $^1J = 183.9$ Hz), 115.7, 138.2, 170.4 (d, $^2J = 23.7$ Hz); ^{19}F NMR δ -192.6 (dt, $^2J_{F-2} = 49.5$ Hz, $^3J_{F-3a,b} = 25.4$ Hz); IR (neat) 2931, 1758, 1738,

(30) Vettel, S.; Vaupel, A.; Knochel, P. *J. Org. Chem.* **1996**, *61*, 7473–7481.

(31) (a) Thenappan, A.; Burton, D. J. *J. Org. Chem.* **1990**, *55*, 2311–2317. (b) Brown, D. J.; Hoskin, J. A. *J. Chem. Soc. (B)* **1971**, 2214–2217.

1640 cm^{-1} ; MS (APCI) m/z 175 (100, MH^+); GC/MS purity (99%); MS m/z 174 (10, M^+). Anal. Calcd for $\text{C}_9\text{H}_{15}\text{FO}_2$ (174.22): C, 62.05; H, 8.68. Found: C, 61.71; H, 8.81.

^{19}F NMR of the crude reaction mixture showed only peaks for **7f**.

Analogous treatment of **5f** (158 mg, 0.5 mmol) by procedure D gave **7f** (73 mg, 84%): GC/MS purity (>99%); MS m/z 174 (10, M^+). ^{19}F NMR of the crude reaction mixture showed peaks for **7f** (~92%) and unchanged **5f** (~8%).

Ethyl 2-Deuterio-2-fluorohexanoate (8a). Treatment of **5a** (152 mg, 0.5 mmol) with Bu_3SnD (0.268 mL, 291 mg, 1.0 mmol) by procedure C gave **8a** (74 mg, 91%; contained ~10% of **7a**). ^1H NMR data for **8a** corresponded to those reported for **7a**^{31a} except for small signals (~10%) at δ 4.86 (dt, $^2J_{\text{F-H}} = 49.2$ Hz, $^3J_{\text{F-H}} = 5.9$ Hz, 2-CHF) and simplification of the multiplet at δ 1.87 (3- CH_2); ^{19}F NMR δ -193.2 (tt, $^2J_{\text{F-D}} = 7.9$ Hz, $^3J_{\text{F-H}} = 24.9$ Hz, 0.9, 2-CDF), -192.5 (dt, $^2J_{\text{F-H}} = 49.2$ Hz, $^3J_{\text{F-H}} = 24.9$ Hz, 0.1, 2-CHF); MS (CI) m/z 164 (100, MH^+ [$\text{C}_8\text{H}_{15}\text{DFO}_2$]).

Ethyl 2-Deuterio-2-(pyrimidin-2-ylsulfonyl)hexanoate (9a). **Procedure F.** Compound **1a** (356 mg, 1.25 mmol) was treated with KH (1.75 mmol) as described in procedure E. The solution was stirred at 0 °C for 15 min followed by 90 min at ambient temperature and then was cooled to -60 °C, and D_2O (1 mL) was added dropwise. After 5 min, the reaction mixture was allowed to warm to ~0 °C and solid NH_4Cl (1 g) followed by saturated $\text{NH}_4\text{Cl}/\text{H}_2\text{O}$ (15 mL) and EtOAc (25 mL) were added. The organic layer was separated, and the aqueous layer was extracted (EtOAc). The combined organic phase was washed (saturated $\text{NaHCO}_3/\text{H}_2\text{O}$, brine), dried (MgSO_4), evaporated, and chromatographed (50 \rightarrow 85% hexanes/EtOAc) to give **9a** (271 mg, 76%): ^1H NMR spectra corresponded with those of **1a** with ~90% reduction in the intensity of signals at δ 4.61 (0.1H) and simplification of multiplet at δ 2.16–2.28 (2H); MS (CI) m/z 288 (100, MH^+ [$\text{C}_{12}\text{H}_{17}\text{DN}_2\text{O}_4\text{S}$]).

Ethyl 2,2-Dideuteriohexanoate (10a). Treatment of **9a** (29 mg, 0.1 mmol) with Bu_3SnD (2 equiv) by procedure C gave **10a** (11 mg, 76%). ^1H NMR spectra corresponded with those of **3a** with ~80% reduction in the intensity of signals at δ 2.28 (t, $J = 7.7$ Hz, 0.2H, 2- CD_2); MS (CI) m/z 147 (100, MH^+ [$\text{C}_8\text{H}_{15}\text{D}_2\text{O}_2$]).

Pyrimidin-2-ylsulfonic Acid (13). A sample of **1a** (286 mg, 1 mmol) was treated with Bu_3SnH as described in procedure C, and the crude benzene solution was partitioned (EtOAc/ $\text{NaHCO}_3/\text{H}_2\text{O}$). The aqueous phase was washed (EtOAc) and was evaporated to give a white solid. Extraction of this material with MeOH and evaporation gave **13** as a sodium salt (148 mg, 89%): ^1H NMR (D_2O) δ 7.60 (t, $J = 4.9$ Hz, 1H), 8.88 (d, $J = 4.9$ Hz, 2H); ^{13}C NMR ($\text{D}_2\text{O}/\text{Me}_2\text{SO}-d_6$) δ 123.2, 159.1, 177.0. ^1H NMR spectrum of **13** parallels with that reported for potassium pyrimidin-2-ylsulfonate.^{31b}

Analogous treatment of **1f** (298 mg, 1 mmol) by procedure C also gave **13** (144 mg, 87%).

Ethyl 6-(Pyrimidin-2-ylsulfonyl)hexanoate (14). Treatment of ethyl 6-bromohexanoate (1.82 mL, 2.23 g, 10 mmol) with 2-mercaptopyrimidine (1.12 g, 10 mmol) and oxidation (20 h) with *m*-CPBA (4.8 g/75% reagent, 21 mmol) by procedure A gave **14** (2.43 g, 85%): ^1H NMR δ 1.23 (t, $J = 7.1$ Hz, 3H), 1.43–1.95 (m, 6H), 2.28 (t, $J = 7.1$ Hz, 3H), 3.52 (dd, $J = 7.4$, 8.2 Hz, 2H), 4.09 (q, $J = 7.1$ Hz, 2H), 7.58 (t, $J = 4.9$ Hz, 1H), 8.95 (d, $J = 4.9$ Hz, 2H); ^{13}C NMR δ 14.7, 22.3, 24.7, 28.3, 34.3, 51.5, 60.8, 124.4, 159.2, 166.2, 173.7; MS (CI) m/z 287 (100, MH^+). Anal. Calcd for $\text{C}_{12}\text{H}_{18}\text{N}_2\text{O}_4\text{S}$ (286.35): C, 50.33; H, 6.34; N, 9.78. Found: C, 50.40; H, 6.27; N, 9.56.

Treatment of **14** with Bu_3SnH (2 equiv) by procedure C gave unchanged **14** (89%) contaminated (~10%, ^1H NMR) by tin compound(s).

Acknowledgment. We thank Alberto J. Sabucedo at Advanced Mass Spectrometry Facility at FIU for his contribution and the Florida International University Foundation for support. S.F.W. is grateful to Professor Morris J. Robins (Brigham Young University) for his encouragement and many stimulating discussions.

Supporting Information Available: Experimental procedures and characterization data for compounds **1b**, **e**, **g**, **h**, **3e**, **g**, **5b**, **e**, **g**, **h**, **7e**, **g**, **h**, **9b**, and **10b** as well as copies of ^1H and ^{13}C NMR spectra of compound **13**. This material is available free of charge via the Internet at <http://pubs.acs.org>

JO000342N